

Endothelin-converting enzyme-1 degrades internalized somatostatin-14

Article

Published Version

Roosterman, D., Kempkes, C., Cottrell, G. S. ORCID: <https://orcid.org/0000-0001-9098-7627>, Padilla, B. E., Bunnett, N. W., Turck, C. W. and Steinhoff, M. (2008) Endothelin-converting enzyme-1 degrades internalized somatostatin-14. *Endocrinology*, 149 (5). pp. 2200-2207. ISSN 1945-7170 doi: <https://doi.org/10.1210/en.2007-1628> Available at <https://centaur.reading.ac.uk/30261/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

To link to this article DOI: <http://dx.doi.org/10.1210/en.2007-1628>

Publisher: Endocrine Society

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online

Endothelin-Converting Enzyme-1 Degrades Internalized Somatostatin-14

Dirk Roosterman, Cordula Kempkes, Graeme S. Cottrell, Benjamin E. Padilla, Nigel W. Bunnett, Christoph W. Turck, and Martin Steinhoff

Department of Dermatology (D.R., C.K., M.S.), Interdisziplinäres Zentrum für Klinische Forschung Münster, and Ludwig Boltzmann Institute for Cell and Immunobiology of the Skin, University Münster, D-48149 Münster, Germany; Departments of Surgery and Physiology (G.S.C., B.E.P., N.W.B.), University of California, San Francisco, San Francisco, California 94143-0660; and Max Planck Institute of Psychiatry, Proteomics and Biomarkers (C.W.T.), D-80804 Munich, Germany

Agonist-induced internalization of somatostatin receptors (sst) determines subsequent cellular responsiveness to peptide agonists and influences sst receptor scintigraphy. To investigate sst2A trafficking, rat sst2A tagged with epitope was expressed in human embryonic kidney cells and tracked by antibody labeling. Confocal microscopical analysis revealed that stimulation with sst and octreotide induced internalization of sst2A. Internalized sst2A remained sequestered within early endosomes, and 60 min after stimulation, internalized sst2A still colocalized with β -arrestin1-enhanced green fluorescence protein (EGFP), endothelin-converting enzyme-1 (ECE-1), and rab5a. Internalized ^{125}I -Tyr¹¹-SST-14 was rapidly hydrolyzed by endosomal endopeptidases, with radioactive metabolites being released from the cell. Internalized ^{125}I -Tyr¹-octreotide accumulated as an intact peptide and was released from the cell as an intact peptide ligand. We have identified ECE-1 as one of the endopeptidases responsible for inactivation of internalized SST-14. ECE-1-mediated

cleavage of SST-14 was inhibited by the specific ECE-1 inhibitor, SM-19712, and by preventing acidification of endosomes using bafilomycin A₁. ECE-1 cleaved SST-14 but not octreotide in an acidic environment. The metallopeptidases angiotensin-1 converting enzyme and ECE-2 did not hydrolyze SST-14 or octreotide. Our results show for the first time that stimulation with SST-14 and octreotide induced sequestration of sst2A into early endosomes and that endocytosed SST-14 is degraded by endopeptidases located in early endosomes. Furthermore, octreotide was not degraded by endosomal peptidases and was released as an intact peptide. This mechanism may explain functional differences between octreotide and SST-14 after sst2A stimulation. Moreover, further investigation of endopeptidase-regulated trafficking of neuropeptides may result in novel concepts of neuropeptide receptor inactivation in cancer diagnosis. (*Endocrinology* 149: 2200–2207, 2008)

THE TETRADECAPEPTIDE somatostatin (SST)-14 and the N-terminally extended form, SST-28, are major inhibitory peptides with broad endocrine, exocrine, and neuronal functions (1). sst acts via six sst receptor subtypes (sst1, sst2A, sst2B, sst3, sst4, and sst5) that are expressed in the central nervous system, and endocrine and immune systems.

The high density of sst receptors on neuroendocrine tumors allows the use of stable sst analogs (octreotide) to inhibit hormonal hypersecretion (2). Moreover, this dense distribution of receptors has made the development of sst receptor scintigraphy for tumor imaging possible (3).

Recently, we investigated agonist-induced endocytosis of rat sst1 (4). We demonstrated that sst1 internalized after agonist exposure. Internalized SST-14 was not transported to lysosomes for degradation but was recycled to the medium as an intact and biologically active peptide. The ability to

internalize and recycle intact SST-14 has also been reported for sst2A (5). In contrast to sst1, sst2A binds β -arrestins with high affinity, suggesting that sst2A belongs to the receptor family of “class B” receptors that form prolonged interactions with β -arrestins (6).

Analysis of agonist-mediated β -arrestin translocation to multiple G protein-coupled receptors (GPCRs) has identified two major classes of receptors. “Class A” receptors (e.g. β_2 adrenergic receptor, μ -opioid receptor, endothelin type A receptor, sstr3, sstr5, dopamine D_{1A} receptor, and $\alpha_1\text{b}$ adrenergic receptor) bind β -arrestin2 with higher affinity than β -arrestin1 (6, 7). In contrast, “class B” receptors [e.g. angiotensin II type 1A receptor, neurotensin receptor 1, vasopressin V2 receptor, calcitonin receptor-like receptor (CLR), TSH-releasing hormone receptor, and neurokinin receptor 1 (NK₁R)] bind both isoforms of β -arrestin with similar high affinities. “Class B” receptors are structurally characterized by clusters of serine/threonine residues within their carboxy-terminal tails. They are functionally characterized by a long-lasting sequestration after stimulation with high concentrations of the agonist (8, 9). Internalized “class B” receptors are sequestered with β -arrestins in hollow core vesicles, which are positive for the early endosome marker proteins, early endosomal antigen-1 and rab5a (9). Recently, we demonstrated that endothelin-converting enzyme (ECE)-1 also colocalizes with the internalized “class B” receptors for substance P (SP) (NK₁R) (10) and the receptor for

First Published Online February 14, 2008

Abbreviations: ACE, Angiotensin converting enzyme; CGRP, calcitonin gene-related peptide; CLR, calcitonin receptor-like receptor; ECE, endothelin-converting enzyme; EGFP, enhanced green fluorescence protein; GPCR, G protein-coupled receptor; GTP, guanosine 5'-triphosphate; HBSS, Hanks' buffered saline solution; HEK, human embryonic kidney; NK₁R, neurokinin receptor 1; rh, recombinant human; SP, substance P; sst, somatostatin receptor; TFA, trifluoroacetic acid.

Endocrinology is published monthly by The Endocrine Society (<http://www.endo-society.org>), the foremost professional society serving the endocrine community.

the calcitonin gene-related peptide (CGRP) (CRL) (11). ECE-1 is a metalloendopeptidase related to neprilysin (12). Its four isoforms (ECE-1a–d) share a catalytic domain, but differences in the cytosolic tail determine their subcellular location (13–15). All four isoforms are present within vesicles inside the cell, and ECE-1a as well as ECE-1c are also present at the plasma membrane. We observed that ECE-1 degrades and inactivates SP and CGRP. Endosomal degradation of ligands by ECE-1 allowed recycling and resensitization of the receptors. These experiments identified a new role for ECE-1 in regulating receptor trafficking.

Recent studies of sst2A have shown that the β -arrestin-dependent trafficking of the sst2A sst receptor resembled that of a “class B” receptor (6). Furthermore, investigation of sst2A-mediated uptake of ^{125}I -SST-14 did not clearly show if the internalized ligand is released as an intact peptide or as a metabolite of ^{125}I -SST-14 (5). Therefore, the effects of the natural agonist SST-14 and the peptidase-resistant and clinically usefully agonist octreotide on the subcellular localization of sst2A have not been fully evaluated, and the fate of the endocytosed peptides are unknown.

The aim of the present study was to characterize the fate of endocytosed sst2A and its associated ligands SST-14 and octreotide. We found that stimulation of sst2A with SST-14 or octreotide induced internalization of sst2A and sequestration of the receptor within early endosomes. Internalized ^{125}I -Tyr¹¹-SST-14 is degraded by endosomally located peptidases. Interestingly, inhibition of the activity of the endosomally located peptidases ECE-1 with SM-19712 or preventing acidification of endosomes using bafilomycin A₁ only partially inhibits degradation of SST-14, indicating that other peptidases participate in the degradation of internalized SST-14. Two iodinated metabolites of SST-14 accumulated within the cell during stimulation, whereas mainly one peptide metabolite was detected in the supernatant. Internalized ^{125}I -Tyr¹¹-SST-14 was not routed to lysosomes for degradation. Interestingly, octreotide accumulated as an intact peptide in the cells and was slowly released as an intact peptide from the cells. Further investigation of the metallopeptidases mediating degradation of sst showed that SST-14 was not degraded at acidic pH or neutral pH by recombinant human (rh) ECE-2 and rh angiotensin-1 converting enzyme (ACE-1), whereas ECE-1 just degraded SST-14 at an acidic pH. In contrast, octreotide was not degraded by rhECE-1, rhECE-2, or rhACE.

Materials and Methods

Materials

SST-14 and octreotide were from Bachem (Torrance, CA). ^{125}I -Tyr¹¹-SST-14 was from Amersham Biosciences (Braunschweig, Germany), and ^{125}I -Tyr¹-octreotide was from Anawa (Cologne, Germany). Mouse antibody to the T7 epitope was from Novagen (San Diego, CA). Donkey antimouse coupled to fluorescein isothiocyanate or Texas Red was from Jackson ImmunoResearch Laboratories, Inc. (West Grove, PA). Biotinylated goat anti-ECE-1 antibody, rhACE, rhECE-1, and rhECE-2 were from R&D Systems, Inc. (Minneapolis, MN). The ECE-1 substrate McaBK2 [(7-methoxycoumarin-4-yl)acetyl-Arg-Pro-Gly-Phe-Ser-Ala-Phe-Lys(2,4-dinitrophenyl)], ECE-1 inhibitor SM-19712 (4-chloro-N-[[[4-cyano-3-methyl-1-phenyl-1H-pyrazol-5-yl]amino]carbonyl] benzenesulfonamide, monosodium salt), and bafilomycin A were from Sigma Chemical Co. (St. Louis, MO). Lipofectamine 2000 was from Invitrogen (Karlsruhe, Germany). Media, sera, and plastics were from

Life Technologies, Inc./BRL (Eggenstein, Germany). Other reagents were from Sigma Chemical. The plasmid construct of rat sst2A containing the amino-terminal T7 epitope tag sequence MASMTGGQQMG in pcDNA3.1 has been described previously (6). Constructs for β -arrestin1-EGFP and rab5a-EGFP were generated as described (16, 17).

Cell culture and transfection

Human embryonic kidney (HEK) 293 cells were grown in DMEM supplemented with 10% fetal calf serum in a humidified atmosphere containing 10% CO₂. Cells were transfected using Lipofectamine 2000 according to the manufacturer's instructions. HEK293 cells stably expressing T7-sst2A were selected in the presence of 400 $\mu\text{g}/\text{ml}$ G418 (Invitrogen).

Microscopy and immunofluorescence

Cells were incubated in DMEM plus 0.1% BSA with 100 nM SST-14 or octreotide for 10 min at 37 C, washed, and incubated for 0–60 min at 37 C. Cells were fixed with 4% paraformaldehyde in 100 mM PBS (pH 7.4), for 20 min. Cells were incubated for 30 min in Hanks' buffered saline solution (HBSS), 5% normal goat serum, and 0.1% saponin. The sst2A was detected using mouse T7 antibody (1:10,000, overnight at 4 C) and Texas Red-conjugated goat antimouse IgG (1:200, 2 h, room temperature). β -Arrestin1 and rab5a were detected using EGFP. ECE-1 was detected using biotinylated goat anti-ECE-1 antibody (1:400) and fluorescein isothiocyanate-coupled streptavidin (1:500). Cells were observed using a 510 Meta confocal microscope (Zeiss, Germany) with a Plan Apo $\times 63$ (numerical aperture 1.4) objective.

Binding assays

Cells grown in 24-well dishes were stimulated with SST-14 (100 nM) or octreotide (100 nM) in DMEM (0.1% BSA) for 0–10 min at 37 C. Cells were washed two times with acidic buffer [HBSS-acetic acid (pH 4.8)] and once with HBSS (0.1% BSA). Cells were incubated for 0–120 min at 37 C, placed on ice, and incubated with 100,000 cpm ^{125}I -Tyr¹¹-SST-14 for 120 min at 4 C. Unbound ^{125}I -Tyr¹¹-SST-14 was washed off with HBSS (0.1% BSA). Cells were lysed in 1 ml NaOH, and bound ^{125}I -Tyr¹¹-SST-14 was collected and determined in a γ -counter (Canberra Packard, Dreieich, Germany) (18).

HPLC analysis of ^{125}I -Tyr¹¹-SST-14 and ^{125}I -Tyr¹-octreotide

Cells were incubated with 100,000 cpm/0.35 ml ^{25}I -Tyr¹¹-SST-14 or ^{125}I -Tyr¹-octreotide mixed with 1 nM unlabeled peptide in HBS (0.1% BSA) for 0–15 min at 37 C. After 15-min stimulation, cells were washed twice with warmed acidic buffer (pH 4.8) (18). In some experiments the cells were preincubated with SM-19712 (10 μM) or bafilomycin A₁ (1 μM) for 30 min. Cells were washed once with HBSS (0.1% BSA), incubated for 15 min at 37 C in 0.5 ml HBSS (0.1% BSA), washed, and incubated for 60 min. The supernatants were collected, acidified by adding 50 μl trifluoroacetic acid (TFA) (10%), and centrifuged (5 min, 13,000 $\times g$). Cell lysates were acidified with 0.5 ml 0.08% TFA and centrifuged (5 min, 13,000 $\times g$). Samples were fractionated by reversed-phase HPLC using a C-18 column (2 \times 25 mm). A separating gradient of 0–40% acetonitrile, 0.08% TFA, 25 min, 1 ml/min was used (HPLC-Model Akta purifier; General Healthcare, Munich, Germany). HPLC fractions were collected every minute and counted (4, 19, 20).

rhACE, rhECE-2, and rhECE-1 enzymatic activity

Peptidase activity was measured using McaBK2 (21). McaBK2 (6 μM) was incubated with 0.5 μg rhACE, rhECE-2, or rhECE-1 in 50 mM 2-(N-morpholino)ethanesulfonic acid (pH 5.5) or 50 mM Tris/HCl (pH 7.4) at 37 C. Fluorescence was measured at λ_{ex} 320 nm and λ_{em} 405 nm.

Peptide degradation by rhACE, rhECE-2, and rhECE-1

SST-14 or octreotide (250 μM) was incubated with rhACE, rhECE-2, or rhECE-1 (0.5 μg) in 150 μl of 50 mM (2-N-morpholino)ethanesulfonic acid (pH 5.8) or 50 mM Tris/HCl (pH 7.4) for 0–90 min at 37 C. Reactions

were stopped by boiling for 3 min. Reaction products were separated by reversed-phase HPLC using a C-18 column equilibrated in 0.1% TFA in water, and eluted using a linear gradient of 0.1% TFA in acetonitrile (0–40% acetonitrile, 25 min) at 1 ml/min. Peptides were detected by spectrophotometry (A256 nm). Products were collected and analyzed by time-of-flight mass spectrometry.


Results

Agonists of sst2A induce receptor internalization and association with β -arrestin1

“Class B” receptors form stable and high-affinity interactions with β -arrestin1 and β -arrestin2 at the cell surface and in endosomes (7, 22). To further characterize the internalization of rat somatostatin receptor 2A (rsst2A), we examined the colocalization of β -arrestin1 with internalized rsst2A. A fusion protein of β -arrestin1 and EGFP (β -arrestin1-EGFP) and T7-rsst2A were transiently coexpressed in HEK293 cells (23). In unstimulated cells, β -arrestin1-EGFP was evenly distributed in the cytoplasm and nucleus (Fig. 1). Stimulation with SST-14 or octreotide for 10 min induced translocation of β -arrestin1-EGFP from the cytosol to the cell membrane, where it colocalized with

rsst2A. At later times SST-14 and octreotide induced endocytosis of rsst2A and β -arrestin1-EGFP to small vesicles close to the cell membrane. The distribution and size of the vesicles are similar to those found in studies analyzing rsst2A-mediated internalization of fluorescent SST-14 (24). Internalized rsst2A then translocated to hollow core vesicles that also contained β -arrestin1-EGFP (Fig. 1, A and B, arrowheads). Interestingly, in cells stimulated with SST-14, β -arrestin1 is associated with internalized rsst2A (Fig. 1, A and B, arrowheads) and at the cell membrane (Fig. 1, A and B, arrows). In cells stimulated with octreotide, β -arrestin1 is mainly associated with internalized rsst2A. The fluorescence intensity scan (Fig. 1, A and B, yellow arrows) of the cells is shown in Fig. 1, E and F. The fluorescence signal of β -arrestin1-EGFP is more intensified at the plasma membrane in cells stimulated with SST-14 (Fig. 1, E and F, green arrows) than in cells stimulated with octreotide. Thus, stimulation of rsst2A with agonist results in translocation of β -arrestin1 to the plasma membrane and subsequent association with rsst2A for prolonged periods indicative of a “class B” GPCR.

FIG. 1. Subcellular distribution of T7-rsst2A. HEK293 cells were transiently transfected with T7-rsst2A, β -arrestin1-EGFP (A and B) or rab5a-EGFP (C). Cells were stimulated with 10 nM SST-14 or 10 nM octreotide, washed, and incubated for 15 or 60 min at 37°C. T7-rsst2A was detected using a mouse anti-T7 antibody and Texas Red-conjugated antimouse antibody. β -Arrestin1-EGFP and rab5a-EGFP were detected with EGFP. ECE-1 was detected using goat anti-ECE-1 antibody. In unstimulated cells, T7-rsst2A was detected at the cell surface. β -Arrestin1-EGFP was detected as a diffuse protein located within the cytosol. Stimulation with the peptide ligand SST-14 for 10 min induced receptor internalization. A and B, Internalized T7-rsst2A was strongly associated with β -arrestin1. Incubation for 15 or 60 min induced the formation of hollow core vesicles (arrowheads). β -Arrestin remained strongly associated with internalized T7-rsst2A even 60 min after cell stimulation. In cells stimulated with SST-14, β -arrestin1 is also located at the cell membrane (arrows). The yellow arrows show the area of the intensity scans. C, Colocalization of internalized T7-rsst2A with rab5a-EGFP 60 min after cell stimulation with SST-14 (10 nM), indicating the localization of internalized T7-rsst2A in early endosomes. D, Colocalization of internalized T7-rsst2A with endogenously expressed ECE-1 60 min after stimulation of the cells with SST-14 (10 nM). Fluorescence intensity scan of cells stimulated with SST-14 (E) and octreotide (F) after 60 min recovery. The fluorescence signal of β -arrestin1-EGFP is more intensified at the plasma membrane (green arrows) in cells stimulated with SST-14 than in cells stimulated with octreotide.



Internalized rsst2A is sequestered in early endosomes

After stimulation with agonist, “class B” GPCRs such as NK₁R, CLR, and angiotensin II type 1A receptor are sequestered within early endosomes for prolonged periods (9, 25, 26). To determine the subcellular location of internalized rsst2A, HEK293 cells were transiently transfected with T7-rsst2A and a marker protein for early endosomes, rab5a-EGFP, and their localization was examined using immunofluorescence and confocal microscopy. In unstimulated cells, rsst2A was present at the plasma membrane, and rab5a was present in vesicles within the cell (data not shown). After stimulation with SST-14 (10 nM, 10 min) and incubation in SST-14-free medium for 60 min, internalized rsst2A colocalized with rab5a-EGFP in small and hollow core vesicles (Fig. 1C, arrowheads). Thus, after stimulation with SST-14, rsst2A traffics to early endosomes and remained there for at least 60 min after stimulation.

Internalized rsst2A colocalizes with endogenous ECE-1

HEK293 cells endogenously express all four isoforms of ECE-1 (10). Within the acidic environment of early endosomes, ECE-1 has inactivated the peptide ligands of some “class B” GPCRs (10, 11). We have previously demonstrated that SP and CGRP are inactivated by ECE-1 endogenously expressed by HEK293 cells. HEK293 cells were transiently transfected with rsst2A and the receptor and ECE-1 localized by immunofluorescence and confocal microscopy. In unstimulated cells, rsst2A was localized at the plasma membrane, and ECE-1 was distributed in cytoplasmic vesicles. After stimulation rsst2A was localized at the plasma membrane and colocalized with ECE-1 in vesicles. Thus, rsst2A is transported from the plasma membrane to early endosomes where it remains colocalized with ECE-1 (Fig. 1D).

Recovery of agonist-binding sites after sst2A stimulation

Another characteristic of “class B” GPCRs is that they slowly recycle back to the cell surface because they were sequestered within early endosomes. Confocal microscopical analysis of sst2A expressed in rat insulinoma cell line 1046–38 or in HEK293 cells suggests that sst2A recycled to the plasma membrane within 1–2 h (6, 18). Therefore, we analyzed sst2A recycling by determining the re-appearance of cell surface binding sites after stimulation with SST-14 or octreotide. Rat sst2A-HEK293 cells were stimulated with SST-14 or octreotide, and the recovery of surface binding sites was determined by incubation with ¹²⁵I-SST-14. We found that stimulation with SST-14 or octreotide rapidly induced a marked reduction of surface binding sites (Fig. 2). After stimulation (10 min, 37 C), surface binding sites of SST-14 (51 ± 3%) and octreotide (45 ± 4%) were markedly reduced. Interestingly, the binding sites did not recover during 2-h incubation at 37 C. These findings indicate that sst2A remained internalized for at least 2 h after stimulation with both peptide ligands. This slow recovery of binding sites is indicative of the classical trafficking of a class B GPCR.





FIG. 2. Agonist-induced loss of cell surface binding sites. HEK293 cells stably expressing T7-sst2A were stimulated for 10 min with SST-14 (100 nM) (squares) or octreotide (100 nM) (diamonds). Cells were washed with acidic buffer to remove extracellular ligands, and were incubated for 0, 60, and 120 min at 37 C. Cells were set on ice, and recovery of surface binding sites was determined using ¹²⁵I-SST-14. Surface-binding sites did not recover 120 min after washing the cells. The binding sites remain at approximately 50% of unstimulated cells.

Internalized ¹²⁵I-Tyr¹¹-SST-14 but not ¹²⁵I-Tyr¹-octreotide is degraded by endosomal endopeptidases

Because the receptor-ligand complex of “class B” receptors is sequestered within acidic early endosomes, endosomal located peptidases are capable of mediating the degradation of the endocytosed peptide ligands (10, 11). We analyzed the degradation of ¹²⁵I-SST-14 and ¹²⁵I-octreotide in HEK293 cells stably expressing sst2A. Cells were incubated for 0–15 min with 100,000 cpm radiolabeled peptide, mixed with 1 nM unlabeled peptide in HBSS (0.1% BSA) at 37 C, washed twice with acidic HBSS (pH 4.8), and once with HBSS (0.1% BSA), and incubated for 0–90 min at 37 C. Cell lysates and supernatants were collected and analyzed by HPLC. Analysis of cell lysates from SST-14-treated cells indicates that SST-14 was rapidly endocytosed by sst2A-HEK293 cells (Fig. 3A, closed diamonds). SST-14 was rapidly degraded into two radioactive labeled peptide metabolites that accumulated within the cell. After 15-min incubation with ¹²⁵I-SST-14, only 42.9% of total radioactivity was found to be cell associated as intact ¹²⁵I-SST-14. Of the remaining radioactivity, 19.3% was peptide A (retention time, 9 min), and 37.7% was peptide B (retention time, 11 min) (Fig. 3A). Incubation for longer periods at 37 C induced further degradation of the internalized SST-14. At 90 min after stimulation, only 7.3% of the total radioactivity remained as intact SST-14, whereas 13.7% of the total radioactivity was peptide B, and 2.6% of the total radioactivity was peptide A. Interestingly, most of the radioactivity was released from the cells (Fig. 3C). HPLC analysis of supernatants demonstrated that two peptide metabolites were released from the cells. The major product was peptide B (69.2% of the total radioactivity), with the minor product, peptide A, only being 4.9% of the total radioactivity. The small amount of peptide A detected into the supernatant (4.9%, even after 90 min) is in marked contrast to the maximal amount associated with the cell after 15 min (19.3% of total radioactivity). These findings suggest that peptide A is an intermediate degradation product before being further processed to peptide B.

We found that incubation of sst2A cells with ¹²⁵I-octreotide

FIG. 3. Fate of internalized radiolabeled SST-14 and octreotide (octre). HEK293 cells stably expressing T7-sst2A were incubated for 5, 10, and 15 min with ^{125}I -SST-14 (100,000 cpm) mixed with unlabeled SST-14 (1 nM) (A and C) or with ^{125}I -octreotide (100,000 cpm) mixed with unlabeled octreotide (1 nM) (B and D). Cells were washed with an acidic buffer and incubated for 30, 60, and 90 min at 37 C. Radioactivity in the supernatant and cells was analyzed by HPLC. Internalized ^{125}I -SST-14 (closed diamonds) was rapidly degraded by peptidases during 15-min incubation with the ligand. The amount of cell-associated intact SST-14 decreased to 42.9%. Further incubation at 37 C allowed degradation of internalized SST-14 after 90-min incubation. Of the total radioactivity, 7.3% was intact ^{125}I -SST-14. A, Two radioactive degradation products accumulated in the cells (closed triangles and closed circles) during the incubation with ^{125}I -SST-14. C, The degradation products were rapidly released from the cells. Intact ^{125}I -SST-14 was not detected in the supernatant. The degradation product with the retention time of 11 min was the major degradation product detected in the supernatant. B, During incubation of cells, octreotide accumulated as an intact peptide within the cells. No degradation products were detected as cell associated. After washing, the amount of cell-associated octreotide decreased to 83%. D, Only intact octreotide was detected in the supernatant. E, A representative HPLC profile of cell-associated ^{125}I -SST-14 after 5-min incubation with ^{125}I -SST-14. F, The influence of the ECE-1 inhibitor SM-19712 and bafilomycin A on the degradation of ^{125}I -SST-14. Cells were pretreated with SM-19712 (10 μM) or bafilomycin A (1 μM) for 30 min, stimulated with ^{125}I -SST-14 for 15 min, washed, and incubated for 60 min in the presence of the inhibitors. In control cells, 7.3% of the total radioactivity was intact SST-14. In cells pretreated with SM-19712, 24% of the total radioactivity was intact SST-14. In cells pretreated with bafilomycin A, 26% of the total radioactivity was intact SST-14. **, $P < 0.01$ to control. G, The sequences of SST-14 and octreotide. The peptides were digested with rhECE-1, and the products were purified by HPLC and identified by mass spectrometry.



induced accumulation of intact octreotide in the cells (Fig. 3B). Ninety minutes after cells were washed, 83% of the internalized octreotide was still cell associated, and 17% of the radioactivity was found in the supernatant as intact octreotide (Fig. 3D).

As shown in a representative HPLC chromatogram (Fig. 3E), intact ^{125}I -Tyr¹¹-SST-14 eluted by 16 min, and two radioactive degradation products (peptide A and peptide B) were recognized by HPLC as having eluted by 9 and 11 min, respectively. The ^{125}I -Tyr eluted by 3–4 min and could not be detected in the cell lysate or the supernatant of rsst2A-HEK cells, indicating that SST-14 was not routed to lysosomal degradation (11).

HEK293 cells endogenously express all four isoforms of ECE-1 (10). Therefore, in our next set of experiments, we tried to inhibit intracellular degradation of internalized ^{125}I -SST-14 by preincubating the cells with the ECE-1-specific inhibitor, SM-19712 (10 μM), and by neutralizing acidic ves-

icles by preincubating with bafilomycin A1 (1 μM). Both inhibitors had been previously tested for the ECE-1-mediated degradation of internalized SP and CGRP (10, 11). We preincubated cells for 15 min with 100,000 cpm ^{125}I -Tyr¹¹-SST-14 mixed with 1 nM SST-14 in the presence of the inhibitors, washed, and incubated the cells for 60 min at 37 C. Without inhibitor, 7.3% (4) of the total radioactivity was intact ^{125}I -Tyr¹¹-SST-14. In cells preincubated with SM-19712, 24 \pm 3% of the total radioactivity was intact ^{125}I -Tyr¹¹-SST-14, and in cells pretreated with bafilomycin A₁, 26 \pm 4% was intact ^{125}I -Tyr¹¹-SST-14 (Fig. 3F). These findings indicate that SM-19712 significantly blocked the intracellular degradation of ^{125}I -Tyr¹¹-SST-14 and that the degradation is sensitive to neutralization of acidic vesicles. Interestingly, whereas the intracellular degradation of SP or CGRP could be almost completely blocked with SM-19712 and bafilomycin A₁ (10, 11), the degradation of ^{125}I -Tyr¹¹-SST-14 was only partially blocked with the inhibitors, indicating that besides ECE-1,

other endopeptidases participate in the endosomal degradation of SST-14. Figure 3G shows the sequences of SST-14 and octreotide. The arrows mark the sites of cleavage by ECE-1 inferred from the product masses.

We next sought to determine whether other endopeptidase were able to degrade SST-14. Therefore, we tested rhECE-1, rhECE-2, and rhACE-1 to see if they degraded SST-14 and octreotide. The activity of the metallopeptidases was measured with the synthetic fluorescence substrate McaBK2. We tested the degradation of McaBK2 at neutral pH and at pH 5.8, which is equal to the pH of acidic early endosomes (10). The metallopeptidases are active at pH 5.8 (data not shown). At neutral pH, rhACE degrades McaBK2 10 times faster than at pH 5.5, whereas rhECE-2 did not degrade McaBK2 at pH 7.4 (data not shown). While testing the degradation of SST-14 and octreotide, we found that ECE-1 did not degrade octreotide in an acidic environment but did degrade SST-14 (Fig. 4, A and B). Time-of-flight mass spectroscopy showed that SST-14 was degraded at position Asn⁵-Phe⁶ and at position Thr¹⁰-Phe¹¹, leading to two degradation products: the peptide SST-14 (1–5)–SST-14 (11–14) and to SST-14 (6–10).

Figure 4C shows the time course of rhECE-1 mediated degradation of SST-14 in an acidic and neutral environment. At pH 5.8, rhECE-1 rapidly degraded SST-14. After 30-min incubation, 50% of the SST-14 was degraded. Interestingly, prolonged incubation with rhECE-1 did not result in a complete degradation of the peptide. A similar incomplete degradation profile was observed for the ECE-1 mediated degradation of CGRP (11).

Figure 4D shows the amount of intact peptide for octreotide and SST-14 after 90-min incubation with rhECE-1, rhECE-2, and rhACE. Of note, octreotide was not degraded by any of these endopeptidases. SST-14 was only degraded by ECE-1 in an acidic environment by approximately 50%


(Fig. 4D). None of the other tested endopeptidases hydrolyzed SST-14.

Discussion

The first aim of this investigation was to determine the subcellular localization of internalized sst2A. Our results show that stimulation with SST-14 or octreotide induced internalization of sst2A into early endosomes. Even 60 min after stimulation, sst2A still colocalized with the early endosomal marker protein rab5a. These results, including the strong association of internalized rsst2A with β -arrestin1-EGFP, indicate that sst2A trafficking resembles the trafficking mechanism of a “class B” receptor (9, 26). Interestingly, β -arrestin1 is located at the cell membrane and at early endosomes when the cells are stimulated with SST-14, whereas β -arrestin1 is mainly associated with early endosomes after stimulation with octreotide. The data let us suggest that degradation of SST-14 allowed dissociation of β -arrestin1 from internalized rsst2A and translocation of β -arrestin1 to the cell membrane. Our results are in line with previous results analyzing β -arrestin trafficking and endosomal sorting of other sst receptor subtypes (6, 18). Interestingly, whereas confocal analysis of the recycling behavior of sst2A in rat insulinoma 1046–38 cells or HEK293 suggests that sst2A recycled within 1–2 h, we were unable to observe recycling of rsst2A 60 min after stimulation. That we were able to determine the reappearance of surface binding sites indicates that sst2A did not recycle 2 h after stimulation with the cognitive ligand.

Another characteristic of “class B” GPCRs is the formation of hollow core vesicles. This has been clearly demonstrated for the AT(1A)R and the NK₁R (9, 26). Hollow core vesicles are enlarged vesicles, which are formed by the constitutively active rab5a mutant rab5aQ79L. Other studies found that rab5a is a binding protein of the AT(1A)R and suggested that

FIG. 4. Degradation of SST-14 and octreotide with rhACE, rhECE-2, and rhECE-1. A, The HPLC profile of octreotide (250 μ M) incubated with ECE-1 (0.5 μ g) for 90 min at 37 C (pH 5.8). Octreotide was not degraded by ECE-1. B, The HPLC profile of SST-14 incubated with ECE-1 for 90 min at 37 C (pH 5.8). C, The time dependence of ECE-1-mediated degradation of SST-14 at pH 5.8 and pH 7.4. At neutral pH, ECE-1 did not degrade SST-14. SST-14 was rapidly degraded by ECE-1 at pH 5.8. After 30-min incubation, 47% of SST-14 was degraded. Longer incubation of the peptide with ECE-1 did not result in further degradation of the peptide. D, The degradation of octreotide and SST-14 by rhACE, rhECE-2, and rhECE-1 at a neutral pH (gray bars) and at pH 5.8 (black bars). Only SST-14 was degraded by ECE-1 in an acidic environment.



an intrinsic guanosine 5'-diphosphate-guanosine 5'-triphosphate (GTP) exchange element is encoded in the carboxy-terminal tail of the AT(1A)R (25, 26). The direct interaction of phosphorylated receptor and activated (GTP-bound) rab5a may lead to the homotypical fusion of early endosomes and the subsequent formation of hollow core vesicles. A detailed analysis of deletion mutants of AT(1A)R demonstrated that truncation of the C-terminal tail of the receptor inhibits the formation of hollow core vesicles but does not inhibit internalization of the receptor (26). Previous studies showing that the guanosine 5'-diphosphate-GTP-exchange element is coded at the far end of the C terminus of the AT(1A)R let us suggest that the naturally occurring splice variant of *sst2*, *sst2B*, demonstrates a different pattern of internalization, resensitization, and trafficking compared with *sst2A* (27, 28).

Our confocal microscopic observation of T7-rsst2A and β -arrestin1-EGFP also shows that stimulation with SST-14 and octreotide induced the formation of β -arrestin1 and coated hollow core vesicles. These results indicate that rsst2A trafficking is blocked in early endosomes. The localization within early endosomes could be also demonstrated by the colocalization of internalized rsst2A with rab5a-EGFP. Because the receptor is not immediately transported to late endosomes, and the ligand is not routed into lysosomal degradation, endopeptidases with an acidic pH optimum for degradation of small peptides, like ECE-1 or ECE-2, should be involved in the hydrolysis and inactivation process of the endocytosed peptide ligands (29, 30). Recently, the endopeptidase ACE was also cotransported with internalized receptors (31), suggesting that this degrading enzyme may not necessarily be constitutively located within early endosomes. The degradation of an internalized peptide agonist by early endosomally located endopeptidases was recently demonstrated for the NK₁R/SP and for the CLR/CGRP system (10, 11). The same studies showed that for both class B receptors, intracellular ECE-1 inactivates the internalized ligand within early endosomes and allowed dissociation of β -arrestin1 from the internalized receptor. They also showed that the degradation of ligand by ECE-1 is an essential step in the process of neuropeptide receptor trafficking and resensitization. In particular, ECE-1 cleaved SP at positions Gln⁶-Phe⁷ and Gly⁹-Leu¹⁰ to form the metabolites SP[1–6] and SP[1–9]. Similar to the degradation of SP and CGRP, that of SST-14 by ECE-1 is also sensitive to a specific pH.

The metabolites of ECE-1-mediated degradation of SP, SP[1–6] and SP[1–9], accumulated within the cells and were slowly released into the supernatant. In contrast, the metabolites of CGRP transiently occurred within the cells and were most probably transported toward lysosomal degradation. Here, we demonstrate that the preferred cleavage sites of SST-14 are at positions Asn⁵-Phe⁶ and Thr¹⁰-Phe¹¹. ECE-1 hydrolyzed peptides N terminal from hydrophobic amino acids. Thus, SST-14 has four potential degradation sites for ECE-1. The degradation analysis of ¹²⁵I-Tyr¹¹-SST-14 revealed that internalized ¹²⁵I-SST-14 was processed to two radioactively labeled products, suggesting that ECE-1 cleaved ¹²⁵I-SST-14 at two positions. One of these degradation products accumulates within the cell, and, subsequently,

only parts of this product could be detected in the supernatant after prolonged incubation.

It is unknown whether the intermediate peptide is further processed by hydrolysis or by a reduction/oxidation process of the peptide. Intermediate degradation products, which are further degraded by hydrolysis, were reported for the ECE-1-mediated degradation of internalized SP and CGRP (10, 11). However, the reduction of the peptide will release the hydrophilic N-terminal part of SST-14 (AGSKN) from the hydrophobic radioactively labeled C-terminal end. This fact can explain the prolonged retention time of the major degradation product. Oxidation of the intermediate peptide also has an impact on the retention time. Following this hypothesis, it remains uncertain if ¹²⁵I-SST-14 is specifically hydrolyzed at two positions or, alternatively, is merely hydrolyzed once and further converted by a reduction or oxidation process that may occur in the supernatant during the incubation time at 37 C.

Because iodination of the tyrosine changes the elution time of the peptide, we were unable to determine the sequence of the iodinated degradation product. Our inability to detect ¹²⁵I-Tyr as a degradation product suggests that SST-14 is not transported to lysosomal degradation. In support of these findings, others have shown that *sst2A* stimulation in Chinese hamster ovary cells leads to the release of the radiolabeled agonist as an iodinated peptide (5). However, in these studies the degradation products were not analyzed by HPLC. Therefore, it was still unknown whether or not the peptide was released as an intact peptide or as a metabolite. The release of intact SST-14 was recently reported for *sst1*. Therefore, it was important to determine the fate of internalized SST-14 by HPLC (4). Our experiments clearly demonstrate that SST-14 was not released as an intact peptide, whereas the synthetic peptide, octreotide, accumulated as an intact peptide within the cells and was subsequently released as an intact peptide from the cells. We further showed that rsst2A internalizes and colocalizes with ECE-1 within early endosomes. Furthermore, ECE-1 was able to degrade SST-14 in an acidic environment of early endosomes, but not at a neutral pH.

In our study, preventing acidification of endosomes with bafilomycin A₁ or inhibition of ECE-1 activity by SM-19712 prevented the intracellular degradation of SP and CGRP, whereas the intracellular degradation of SST-14 was only partially blocked by inhibitors. This observation strongly suggests also that other endopeptidases besides ECE-1 participate in the endosomal degradation process of SST-14. Therefore, we tested whether rhECE-2 and rhACE were able to degrade SST-14 and octreotide. Interestingly, both peptidases were unable to degrade SST-14. Alternatively, an endosomal location for cathepsin B has been recently reported (32). Therefore, we tested whether the cathepsin B inhibitor, CA074-ME, was able to prevent endosomal degradation of SST-14. In fact, pretreatment of cells with CA074-ME before SST-14 stimulation did not influence the endosomal degradation of SST-14 (data not shown).

In the present study, we compared the fate of a natural ligand, SST-14, and the long-lasting synthetic *sst* analog octreotide in cells expressing rsst2A. Both peptides induced internalization and sequestration of *sst2A* within early en-

osomes. In early endosomes, SST-14 was hydrolyzed and inactivated by different endopeptidases. The endopeptidase-generated degradation products were released as peptide fragments from these cells. At the same time, the internalized receptor remained intracellularly located and was not recycled. Furthermore, the synthetic analog (octreotide) remained as an intact ligand for a prolonged time within the cells, where it colocalized with the internalized receptor sst2A. The finding that an internalized ligand of sst2A can be degraded within early endosomes while not routed into lysosomal degradation will help to develop novel radiolabeled sst or other neuropeptide ligand analogs useful for medical diagnostics such as tumor imaging as well as molecular imaging for *in vivo* models of inflammation (33, 34).

Acknowledgments

We thank Dr. H.-J. Kreienkamp for the kind gift of T7-somatostatin receptor 2A cDNA.

Received November 28, 2007. Accepted January 24, 2008.

Address all correspondence and requests for reprints to: Dr. Dirk Roosterman, Department of Dermatology, Interdisziplinäres Zentrum für Klinische Forschung Münster, and Ludwig Boltzmann Institute for Cell and Immunobiology of the Skin, University Münster, D-48149 Münster, Germany. E-mail: roosterman@gmx.net.

This work was supported by grants from Interdisziplinäres Zentrum für Klinische Forschung Münster (STE12/076/06), SFB 293 (A14), and SFB 492 (B13), CERIES (Paris) (to M.S.), Deutsche Forschungsgemeinschaft STE 1014/2-2, Rosacea Foundation (to M.S. and D.R.), International Monetary Fund (RO 120611) (to D.R.), and National Institutes of Health Grants DK39957 and DK43207 (to N.W.B.).

Disclosure Statement: The authors have nothing to declare.

References

- Olias G, Viollet C, Kusserow H, Epelbaum J, Meyerhof W 2004 Regulation and function of somatostatin receptors. *J Neurochem* 89:1057–1091
- Lamberts SW, Oosterom R, Neufeld M, del Pozo E 1985 The somatostatin analog SMS 201–995 induces long-acting inhibition of growth hormone secretion without rebound hypersecretion in acromegalic patients. *J Clin Endocrinol Metab* 60:1161–1165
- Lamberts SW, Bakker WH, Reubi JC, Krenning EP 1990 Somatostatin-receptor imaging in the localization of endocrine tumors. *N Engl J Med* 323:1246–1249
- Roosterman D, Kreuzer OJ, Brune N, Cottrell GS, Bunnett NW, Meyerhof W, Steinhoff M 2007 Agonist-induced endocytosis of rat somatostatin receptor 1. *Endocrinology* 148:1050–1058
- Koenig JA, Kaur R, Dodgeon I, Edwardson JM, Humphrey PP 1998 Fates of endocytosed somatostatin sst2 receptors and associated agonists. *Biochem J* 336(Pt 2):291–298
- Tulipano G, Stumm R, Pfeiffer M, Kreienkamp HJ, Holt V, Schulz S 2004 Differential beta-arrestin trafficking and endosomal sorting of somatostatin receptor subtypes. *J Biol Chem* 279:21374–21382
- Oakley RH, Laporte SA, Holt JA, Barak LS, Caron MG 1999 Association of beta-arrestin with G protein-coupled receptors during clathrin-mediated endocytosis dictates the profile of receptor resensitization. *J Biol Chem* 274:32248–32257
- Oakley RH, Laporte SA, Holt JA, Barak LS, Caron MG 2001 Molecular determinants underlying the formation of stable intracellular G protein-coupled receptor-beta-arrestin complexes after receptor endocytosis. *J Biol Chem* 276:19452–19460
- Roosterman D, Cottrell G, Schmidlin F, Steinhoff M, Bunnett NW 2004 Recycling and resensitization of the neurokinin 1 receptor: influence of agonist concentration and RAB GTPases. *J Biol Chem* 279:30670–30679
- Roosterman D, Cottrell GS, Padilla BE, Muller L, Eckman CB, Bunnett NW, Steinhoff M 2007 Endothelin-converting enzyme 1 degrades neuropeptides in endosomes to control receptor recycling. *Proc Natl Acad Sci USA* 104:11838–11843
- Padilla B, Cottrell GS, Roosterman D, Muller L, Steinhoff M, Bunnett NW 2007 Endothelin-converting enzyme-1 regulates endosomal sorting of calcitonin receptor-like receptor and beta-arrestins. *J Cell Biol* 179:981–997
- Turner AJ, Isaac RE, Coates D 2001 The neprilysin (NEP) family of zinc metalloendopeptidases: genomics and function. *Bioessays* 23:261–269
- Schweizer A, Valdenaire O, Nelbock P, Deuschle U, Dumas Milne Edwards JB, Stumpf JG, Löffler BM 1997 Human endothelin-converting enzyme (ECE-1): three isoforms with distinct subcellular localizations. *Biochem J* 328(Pt 3):871–877
- Shimada K, Takahashi M, Ikeda M, Tanzawa K 1995 Identification and characterization of two isoforms of an endothelin-converting enzyme-1. *FEBS Lett* 371:140–144
- Shimada K, Takahashi M, Tanzawa K 1994 Cloning and functional expression of endothelin-converting enzyme from rat endothelial cells. *J Biol Chem* 269:18275–18278
- Schmidlin F, Dery O, DeFea KO, Slice L, Patierno S, Sternini C, Grady EF, Bunnett NW 2001 Dynamin and Rab5a-dependent trafficking and signaling of the neurokinin 1 receptor. *J Biol Chem* 276:25427–25437
- Roosterman D, Schmidlin F, Bunnett NW 2003 Rab5a and rab11a mediate agonist-induced trafficking of protease-activated receptor 2. *Am J Physiol Cell Physiol* 284:C1319–C1329
- Roosterman D, Roth A, Kreienkamp HJ, Richter D, Meyerhof W 1997 Distinct agonist-mediated endocytosis of cloned rat somatostatin receptor subtypes expressed in insulinoma cells. *J Neuroendocrinol* 9:741–751
- Lucius R, Mentlein R 1991 Degradation of the neuropeptide somatostatin by cultivated neuronal and glial cells. *J Biol Chem* 266:18907–18913
- Mentlein R, Dahms P 1994 Endopeptidases 24.16 and 24.15 are responsible for the degradation of somatostatin, neurotensin, and other neuropeptides by cultivated rat cortical astrocytes. *J Neurochem* 62:27–36
- Johnson GD, Ahn K 2000 Development of an internally quenched fluorescent substrate selective for endothelin-converting enzyme-1. *Anal Biochem* 286:112–118
- Oakley RH, Laporte SA, Holt JA, Caron MG, Barak LS 2000 Differential affinities of visual arrestin, beta-arrestin1, and beta-arrestin2 for G protein-coupled receptors delineate two major classes of receptors. *J Biol Chem* 275:17201–17210
- Dery O, Thoma MS, Wong H, Grady EF, Bunnett NW 1999 Trafficking of proteinase-activated receptor-2 and beta-arrestin-1 tagged with green fluorescent protein. beta-Arrestin-dependent endocytosis of a proteinase receptor. *J Biol Chem* 274:18524–18535
- Nouel D, Gaudriault G, Houle M, Reisine T, Vincent JP, Mazella J, Beaudet A 1997 Differential internalization of somatostatin in COS-7 cells transfected with SST1 and SST2 receptor subtypes: a confocal microscopic study using novel fluorescent somatostatin derivatives. *Endocrinology* 138:296–306
- Dale LB, Seachrist JL, Babwah AV, Ferguson SS 2004 Regulation of angiotensin II type 1A receptor intracellular retention, degradation, and recycling by Rab5, Rab7, and Rab11 GTPases. *J Biol Chem* 279:13110–13118
- Seachrist JL, Laporte SA, Dale LB, Babwah AV, Caron MG, Anborgh PH, Ferguson SS 2002 Rab5 association with the angiotensin II type 1A receptor promotes Rab5 GTP binding and vesicular fusion. *J Biol Chem* 277:679–685
- Vanetti M, Kouba M, Wang X, Vogt G, Holt V 1992 Cloning and expression of a novel mouse somatostatin receptor (SSTR2B). *FEBS Lett* 311:290–294
- Holliday ND, Tough IR, Cox HM 2007 A functional comparison of recombinant and native somatostatin sst2 receptor variants in epithelia. *Br J Pharmacol* 152:132–140
- Pan H, Mzhavia N, Devi LA 2004 Endothelin converting enzyme-2: a processing enzyme involved in the generation of novel neuropeptides. *Protein Pept Lett* 11:461–469
- Fahnoe DC, Knapp J, Johnson GD, Ahn K 2000 Inhibitor potencies and substrate preference for endothelin-converting enzyme-1 are dramatically affected by pH. *J Cardiovasc Pharmacol* 36(5 Suppl 1):S22–S25
- Chen Z, Deddish PA, Minshall RD, Becker RP, Erdos EG, Tan F 2006 Human ACE and bradykinin B2 receptors form a complex at the plasma membrane. *FASEB J* 20:2261–2270
- Authier F, Kouach M, Briand G 2005 Endosomal proteolysis of insulin-like growth factor-I at its C-terminal D-domain by cathepsin B. *FEBS Lett* 579:4309–4316
- Roosterman D, Goerge T, Schneider SW, Bunnett NW, Steinhoff M 2006 Neuronal control of skin function: the skin as a neuroimmunoenocrine organ. *Physiol Rev* 86:1309–1379
- Cevikbas F, Steinhoff A, Homey B, Steinhoff M 2007 Neuroimmune interactions in allergic skin diseases. *Curr Opin Allergy Clin Immunol* 7:365–373