The reversibility of sea level rise

Article

Published Version

Bouttes, N., Gregory, J. M. and Lowe, J. A. (2013) The reversibility of sea level rise. *Journal of Climate*, 26 (8). pp. 2502-2513. ISSN 1520-0442 doi: <https://doi.org/10.1175/JCLI-D-12-00285.1> Available at <https://centaur.reading.ac.uk/34191/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

To link to this article DOI: <http://dx.doi.org/10.1175/JCLI-D-12-00285.1>

Publisher: American Meteorological Society

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online

The Reversibility of Sea Level Rise

N. BOUTTES

NCAS-Climate, University of Reading, Reading, United Kingdom

J. M. GREGORY

NCAS-Climate, University of Reading, Reading, and Met Office Hadley Center, Exeter, United Kingdom

J. A. LOWE

Met Office Hadley Center, Exeter, United Kingdom

(Manuscript received 17 May 2012, in final form 20 September 2012)

ABSTRACT

During the last century, global climate has been warming, and projections indicate that such a warming is likely to continue over coming decades. Most of the extra heat is stored in the ocean, resulting in thermal expansion of seawater and global mean sea level rise. Previous studies have shown that after CO₂ emissions cease or CO₂ concentration is stabilized, global mean surface air temperature stabilizes or decreases slowly, but sea level continues to rise. Using idealized CO₂ scenario simulations with a hierarchy of models including an AOGCM and a step-response model, the authors show how the evolution of thermal expansion can be interpreted in terms of the climate energy balance and the vertical profile of ocean warming. Whereas surface temperature depends on cumulative CO₂ emissions, sea level rise due to thermal expansion depends on the time profile of emissions. Sea level rise is smaller for later emissions, implying that targets to limit sea level rise would need to refer to the rate of emissions, not only to the time integral. Thermal expansion is in principle reversible, but to halt or reverse it quickly requires the radiative forcing to be reduced substantially, which is possible on centennial time scales only by geoengineering. If it could be done, the results indicate that heat would leave the ocean more readily than it entered, but even if thermal expansion were returned to zero, the geographical pattern of sea level would be altered. Therefore, despite any aggressive CO₂ mitigation, regional sea level change is inevitable.

1. Introduction

Model-based studies have explored the evolution of climate in scenarios where atmospheric CO₂ concentrations rise at various rates, following which CO₂ is stabilized or the emissions are stopped (Meehl et al. 2005; Plattner et al. 2008; Solomon et al. 2009; Frölicher and Joos 2010; Gillett et al. 2011). In the latter case, global mean surface air temperature subsequently stays approximately constant or declines slowly as CO₂ is removed from the atmosphere by natural processes. In some scenarios, including the later part of the policy-relevant Representative Concentration Pathway 2.6

(RCP2.6; Moss et al. 2010), which will be assessed in the next report of the Intergovernmental Panel on Climate Change (IPCC), CO₂ is assumed to be artificially removed from the atmosphere at greater rates than natural uptake, producing a faster reduction in forcing and a faster cooling. However, sea level from thermal expansion continues to rise for many centuries following a stabilization of CO₂ concentration in the atmosphere or a cessation of CO₂ emissions (Meehl et al. 2005; Lowe et al. 2006; Plattner et al. 2008; Solomon et al. 2009; Frölicher and Joos 2010; Gillett et al. 2011). This raises the question of what it would take to reverse sea level rise.

In this work, we use a 3D atmosphere–ocean general circulation model (AOGCM) to explore how the emissions pathway affects the peak sea level and the subsequent decline. For simplicity, we consider CO₂-only scenarios, since CO₂ is the dominant forcing. We consider only the global mean thermosteric sea level, and

---

Corresponding author address: Nathaëlle Bouttes, NCAS-Climate, Meteorology Department, University of Reading, Reading, RG66BB, United Kingdom.
E-mail: n.bouttes@reading.ac.uk

not the additional component from glaciers and the ice sheets of Greenland and Antarctica (Charbit et al. 2008; Ridley et al. 2009; Vizcaíno et al. 2010; Huybrechts et al. 2011). The thermosteric term is influenced by ocean interior transport processes, including large-scale oceanic circulations such as the Atlantic meridional overturning circulation (AMOC). In the deep ocean, temperature changes caused by these processes occur on a hundred- to thousand-year time scale. Although this is slow, the thermosteric sea level change may be reversed more quickly than ice sheet mass loss, which would require a longer time scale of thousands of years (Charbit et al. 2008; Ridley et al. 2009). Likewise, reversing the sea level change induced by glaciers and ice caps would be a slower process. Thermosteric sea level change is subject to less (though still substantial) scientific uncertainty than ice sheets and glaciers, and is sufficient alone to cause significant sea level rise impacts for many centuries into the future.

2. Method

We run simulations with the Fast Met Office/UK Universities Simulator (FAMOUS) AOGCM (Smith et al. 2008). FAMOUS is a low-resolution version of the third climate configuration of the Met Office Unified Model (HadCM3) AOGCM (Gordon et al. 2000): its atmosphere component runs on a 5° latitude by 7.5° longitude grid with 11 levels and its ocean component on a 2.5° latitude by 3.75° longitude grid with 20 levels. It is structurally almost identical to HadCM3 and produces climate and climate-change simulations that are reasonably similar to HadCM3, but runs about 20 times faster and is hence particularly useful for investigations involving many long integrations. Despite the coarse resolution, the present-day climatology of FAMOUS is adequate without flux correction.

The simulations employ idealized scenarios in which atmospheric CO_2 is first increased and then either held constant or abruptly decreased. The increase follows three pathways (Fig. 1a): an instantaneous increase of CO_2 to 4 times the preindustrial value ($4\times\text{CO}_2$; the preindustrial value is 280 ppm), a 1% increase of $\text{CO}_2 \text{ yr}^{-1}$ ($1\%\text{CO}_2$) for 140 yr, and a 2% increase of $\text{CO}_2 \text{ yr}^{-1}$ ($2\%\text{CO}_2$) for 70 yr. Both the $1\%\text{CO}_2$ and $2\%\text{CO}_2$ ramps end at 4 times the preindustrial value. Thus, at $t = 140$ yr the CO_2 concentration is the same in all scenarios, but the three pathways that have led to that level differ. Beyond $t = 140$ yr the CO_2 concentration is either kept constant at this value or instantaneously reduced, to either the preindustrial value ($1\times\text{CO}_2$) or half of it ($0.5\times\text{CO}_2$), and then held constant at the reduced value; in section 4c we also consider a scenario in which CO_2 is ramped down. We consider these idealized

FIG. 1. Time series of radiative forcing due to the change of atmospheric CO_2 and climate response. (a) Evolution of radiative forcing due to the atmospheric CO_2 change in the simulations (W m^{-2}), (b) evolution of the global mean surface air temperature anomaly ($^\circ\text{C}$), (c) evolution of the global mean thermosteric sea level anomaly (m), and (d) evolution of the time integral of the radiative forcing ($\text{W m}^{-2} \text{ yr}^{-1}$). The anomalies are between the simulation and the control.

scenarios because they give a clear demonstration of the qualitative behavior of sea level. In particular, this is the reason for choosing the $0.5\times\text{CO}_2$ concentration (140 ppm), which is of course unrealistic for natural carbon sinks, being lower than the concentration at glacial maxima such as during the Last Glacial Maximum (atmospheric CO_2 concentration of around 190 ppm approximately 21 000 yr ago).

As the CO_2 concentrations scenarios are idealized, they are not evaluated in terms of feasibility or corresponding emissions. For the latter, a coupled carbon–climate model would be needed, which would include the fact that if

emissions are cut to zero, the biosphere and ocean would progressively take up less carbon as they saturate. This would suggest that to achieve the assumed rates of CO_2 reduction, actively removing CO_2 from the atmosphere by artificial means may be needed, since the CO_2 would otherwise stay in the atmosphere for a long time (several thousand years; Eby et al. 2009). The reduction of forcing does not necessarily require CO_2 reduction and could possibly be obtained by other suggested geo-engineering mechanisms (Shepherd et al. 2009), but the feasibility of this is not further discussed here.

We calculate sea level rise due to thermal expansion from ocean temperature change diagnosed from the model. As found previously (Russell et al. 2000), global mean thermosteric sea level rise η and global mean ocean temperature rise T_o (equivalent to the increase in ocean heat content) are in a nearly linear relationship (Fig. 2) and so we speak of them interchangeably. In the following sections, we interpret the qualitative features of results of the AOGCM for surface air temperature and sea level change using a range of simpler models (summarized in Table 1).

3. Surface air temperature change

During the first part of the simulations (the “high- CO_2 phase”), while the atmospheric CO_2 concentration is

FIG. 2. Sea level change (m) as a function of the global mean ocean temperature change ($^{\circ}\text{C}$). Slope = $0.66 \text{ m } ^{\circ}\text{C}^{-1}$, for the $4\times\text{CO}_2$, $2\%\text{CO}_2$, and $1\%\text{CO}_2$ simulations. The dashed lines indicate the results for the corresponding $0.5\times\text{CO}_2$ simulations described in Fig. 1.

increasing, the change of global mean surface air temperature T_s tracks the CO_2 concentration (Figs. 1a,b). In the case where the CO_2 concentration is instantaneously increased, T_s takes a couple of decades to respond. After 140 yr, T_s is the same in the three simulations. It has similarly been shown that, under scenarios in which emissions peak and decline, the peak warming attained depends only on the cumulative CO_2 emission and is largely insensitive to the emissions pathway up to the time of maximum temperature (Allen et al. 2009a; Matthews et al. 2009; Zickfeld et al. 2009; Bowerman et al. 2011).

TABLE 1. Summary of the main results from the models.

Model	Main features of the results	
FAMOUS AOGCM (Smith et al. 2008)	Sea level evolution depends on the emission pathway Sea level continues to rise after stabilization Sea level rise is reversed if forcing becomes zero or negative Sea level rise and fall are not symmetric	
	Successfully explains	Fails to explain
Zero-layer model (Gregory and Mitchell 1997; Raper et al. 2002; Gregory and Forster 2008; Held et al. 2010) Surface climate energy balance	Sea level rise depends on the emission pathway Sea level continues to rise after stabilization	Sea level rise can be reversed without negative forcing (the model assumes heat loss into an infinite heat sink that does not warm up and therefore does not give back heat spontaneously)
Two-layer model (Gregory 2000; Held et al. 2010) Has an upper layer with small heat capacity and a lower layer with large heat capacity; the layers are thermally coupled	Features explained successfully by the zero-layer model Sea level rise is reversible: it falls during low- CO_2 phase (the model is able to distinguish surface temperature from heat content and can lose heat spontaneously)	Asymmetry of the sea level fall and rise (the single lower layer has too much heat capacity and does not warm up very much, so heat flux depends almost entirely on the upper temperature)
Step model (Good et al. 2011) Uses the diagnosed 3D response of FAMOUS to $4\times\text{CO}_2$	Features explained successfully by the two-layer model. Part of the asymmetry between the sea level fall and rise (because of multiple layers in the ocean)	Remaining part of the asymmetry (due to the different penetration of heat during warming and cooling, because the step model does not have any functional dependence on vertical stability)

The second part of the simulations (the “low-CO₂ phase”) begins at year 141, when the CO₂ concentration is either stabilized or lowered. After CO₂ stabilizes, the rate of surface air temperature increase is greatly reduced (Fig. 1b). If the radiative forcing abruptly returns to zero, T_s falls most of the way to zero within around two decades. If CO₂ is set to a value smaller than the initial one, the radiative forcing becomes negative, and T_s falls below its initial value. It is apparent that T_s depends roughly linearly on the forcing (Figs. 1a and 1b are similar in form), as can be understood from simple “zero-layer” heat balance equations for time-dependent climate change (Gregory and Mitchell 1997; Raper et al. 2002; Gregory and Forster 2008; Held et al. 2010; see Table 1):

$$F = (\alpha + \kappa)T_s \Rightarrow T_s = \frac{F}{\alpha + \kappa}, \quad (1)$$

where α is the climate feedback parameter, κ is the ocean heat uptake efficiency, and the radiative forcing F depends roughly linearly (actually logarithmically) on the atmospheric CO₂ concentration C . In this zero-layer model, there is no feedback of ocean temperature on the rate of heat uptake and the ocean is effectively infinitely deep (i.e., it can absorb heat indefinitely).

C is equivalent to the increase in the CO₂ burden of the atmosphere (1 GtC $\equiv$ 0.47 ppm). Hence C is determined uniquely at each time by the cumulative CO₂ emissions E through the relationship $C = AE$, provided the airborne fraction A is constant. Observations indicate that A has been approximately constant during recent decades (Knorr 2009), but it could change in the future (Friedlingstein et al. 2006). How much it might change appears to depend on both the emissions scenario (Gloor et al. 2010) and the behavior of the carbon cycle (Friedlingstein et al. 2006), which is uncertain. For the shapes of scenario considered here, and assuming that a linear model provides a reasonable first-order representation of the carbon cycle (Joos et al. 1996), then we would expect a constant value of A to also be an acceptable approximation. If so, T_s will track E (Matthews et al. 2009; Gregory et al. 2009).

4. Sea level change

a. Dependence of sea level rise on the pathway of emissions

Sea level behaves qualitatively differently from T_s . The evolution of sea level rise due to thermal expansion depends on the pathway of emissions (Stouffer and Manabe 1999; Zickfeld et al. 2012) and reaches different values at 140 years despite the CO₂ concentration and T_s

being the same at that time (Fig. 1c). Comparing Figs. 1c and 1d, we see that sea level rise depends almost linearly on the integral of the radiative forcing while the forcing is positive (high-CO₂ phase). This is consistent with the zero-layer model, in which the rate of ocean heat uptake N is proportional to T_s ,

$$N = \kappa T_s = F \times \frac{\kappa}{\alpha + \kappa}, \quad (2)$$

and hence the *rate* of sea level depends on F . However, the *actual* sea level rise is given by

$$\eta \propto \int N dt = \int F \times \frac{\kappa}{\alpha + \kappa} dt = \frac{\kappa}{\alpha + \kappa} \int F dt, \quad (3)$$

provided κ and α are constant (Gregory and Forster 2008).

Thus, because the thermosteric sea level is proportional to the time integral of the radiative forcing, the longer the forcing lasts, the bigger the change in thermosteric sea level rise (and oceanic temperature). The sea level change is the largest in the simulation where the largest fractional increase in the atmospheric burden of CO₂ occurs the earliest, which implies where CO₂ has been emitted the earliest. Because of a focus on impacts related to peak warming, it has been proposed that policy targets for avoiding dangerous climate change might be set in terms of cumulative CO₂ emission (Allen et al. 2009b). However, if we are concerned with mitigating sea level impacts, targets must be set on the rate of emission as well.

b. Long-term commitment to sea level rise

If CO₂ is stabilized, ocean temperature and sea level continue to rise (Fig. 1c), in contrast to T_s (Fig. 1b). After the CO₂ concentration is stabilized, because T_s is approximately the same in the three stabilization simulations (Fig. 1b; $\sim 5.5^\circ\text{C}$), the rate of thermosteric sea level rise is nearly the same in all three. That is, the three solid lines in Fig. 1c are nearly parallel; the rate is $\sim 4 \text{ mm yr}^{-1}$, much higher than the recent rate of 0.9 mm yr^{-1} for 1993–2008 (Church et al. 2011). Thus the zero-layer model still explains the evolution of both temperature and sea level in the AOGCM for at least a century after stabilization of the CO₂ concentration.

c. Reversibility of sea level rise

The experiments with stabilized CO₂ may give an impression that sea level rise is irreversible, but if CO₂ is returned instantaneously to its initial value, sea level immediately falls, albeit at a slow and decreasing rate so that after 160 years it has fallen only half the way to its initial level. Any practical measures to remove CO₂

from the atmosphere would produce a gradual rather than an instantaneous decrease; if CO_2 is ramped down, the oceanic temperature initially continues to rise, then gradually stabilizes, and then begins to fall. This is illustrated by a simulation in which CO_2 increases at $1\% \text{ yr}^{-1}$ for 140 yr and then decreases at $1\% \text{ yr}^{-1}$ for the next 140 yr (cyan lines in Fig. 1). In this experiment the time-integral forcing is the same as in the scenario of $4\times\text{CO}_2$ followed by $1\times\text{CO}_2$ (dotted red lines); the later reduction in CO_2 means that sea level is higher following the ramp-down. If the radiative forcing becomes negative (dashed lines), sea level falls back more rapidly to its initial value. Thus, sea level rise is in principle reversible, provided that the radiative forcing is reduced sufficiently. A long-term multi-century commitment to rising thermal expansion is not physically inevitable, although it may be practically so.

However, the zero-layer model fails to explain the sea level reversibility in the AOGCM. Since the sea level rise is proportional to the integral of the forcing F , it predicts that sea level will remain constant if the forcing is returned to zero, and will fall only if a negative forcing is applied. The zero-layer model assumes heat loss into an infinite heat sink that does not warm up and therefore does not give back heat spontaneously. It cannot account for sea level falling under zero forcing because it neglects changes happening below the ocean surface layer. In reality, the ocean is finite and more than one time scale is relevant. The rapid time scale of one or two decades relates to the ocean surface layer and accounts for most of the response of surface air temperature to the radiative forcing following abrupt change in CO_2 . The time scale of the “recalcitrant” response (Held et al. 2010) is another time scale that applies to the evolution of sea level.

This can be illustrated with a two-layer model (Gregory 2000; Held et al. 2010; Table 1). The two-layer model is composed of an ocean with an upper layer of thickness d_u (100 m) and temperature T_u , and a deep layer of thickness d_l (2000 m) and temperature T_l , where both temperature are perturbations from an equilibrium with zero surface heat flux. The upper layer has little heat capacity and plays the role of a regulator; it absorbs the forcing, determines the radiation of heat to space, and is thermally coupled to the lower layer, which has large heat capacity and provides inertia. The heat flux between the two is proportional to their temperature difference. The temperatures thus follow the two equations

$$cd_u \frac{dT_u}{dt} = F - ck \frac{(T_u - T_l)}{0.5 \times (d_l + d_u)} - \alpha T_u \quad \text{and}$$

$$cd_l \frac{dT_l}{dt} = ck \frac{(T_u - T_l)}{0.5 \times (d_l + d_u)},$$

FIG. 3. Time series of the climate response to the same radiative forcing change as in Fig. 1 with the two-layer model. (a) Evolution of the upper ocean layer temperature anomaly ($^{\circ}\text{C}$), (b) evolution of the lower ocean layer temperature anomaly ($^{\circ}\text{C}$), and (c) lower ocean layer temperature change ($^{\circ}\text{C}$) as a function of the integral of the radiative forcing during the phase with high CO_2 (solid lines) and continuing under $0.5\times\text{CO}_2$ (dashed lines).

with $c = 4.218 \times 10^6 \text{ J m}^{-3} \text{ K}^{-1}$ being the volumetric heat capacity, $F (\text{W m}^{-2})$ the surface heat flux, $k = 1 \times 10^{-4} \text{ m}^2 \text{ s}^{-1}$ the thermal diffusivity, and $\alpha = 0.81 \text{ W m}^{-2} \text{ K}^{-1}$ the climate feedback parameter.

When the radiative forcing F changes abruptly, T_u rapidly reaches a new near-equilibrium (Fig. 3a), in which F is nearly balanced by transfer of heat to or from the lower layer. If F reverts to zero, the upper layer rapidly becomes cooler than the lower layer, which therefore loses heat (Fig. 3b, dotted lines). If a negative forcing is applied, the upper layer gets still colder, the temperature difference between the two layers is greater, the transfer of heat from the lower to the upper ocean is faster, and the lower layer cools down more rapidly (Fig. 3b, dashed lines). Unlike the zero-layer model,

FIG. 4. (a),(b) Atmospheric CO_2 (ppm), (c),(d) global mean surface air temperature ($^{\circ}\text{C}$) and (e),(f) global mean oceanic temperature ($^{\circ}\text{C}$) evolution in additional simulations in which CO_2 is either (left) raised to $4\times\text{CO}_2$ and held constant for 140 yr or (right) ramped up at $1\% \text{ yr}^{-1}$ for 140 yr, and in either case lowered to $0.5\times\text{CO}_2$ at different times between 10 and 140 yr and then held constant.

the two-layer model reproduces remarkably well these features of the evolution of surface temperature and thermosteric sea level in the much more complex AOGCM (Figs. 3a and 3b are qualitatively very similar to Figs. 1b and 1c). The two-layer model thus accounts for the reversibility of sea level rise when the radiative forcing diminishes because it has a finite ocean and the model is able to distinguish surface temperature from heat content so that it can lose heat spontaneously.

d. Asymmetry of sea level evolution

In the zero-layer model, the ocean temperature and sea level change should be back to zero when the integral of F returns to zero (i.e., after a negative forcing of the same time integral as the one during the high- CO_2 phase). For example, for the $1\times\text{CO}_2$ scenario followed

by $0.5\times\text{CO}_2$, this is at 280 yr, but actually the sea level in the AOGCM reaches zero sooner, at 224 yr.

To quantify better the link between the ocean temperature change and the radiative forcing, we have run additional AOGCM simulations (Fig. 4), beginning with a high- CO_2 phase of either constant $4\times\text{CO}_2$ or a $1\% \text{ CO}_2$ ramp-up. In these simulations, CO_2 is reduced abruptly to $0.5\times\text{CO}_2$ after various times between 10 and 100 yr (instead of 140 yr as in the scenario of Fig. 1).

During the low- CO_2 phase, the time for the oceanic temperature to cool down to its initial value depends linearly on the oceanic warming achieved during the high- CO_2 phase (Fig. 5a), and hence on the integral of the forcing during that phase (Fig. 5b). However, it takes less negative integral radiative forcing F (or less time with a symmetrical forcing evolution) for the oceanic

FIG. 5. Relations between (a) the time for the global mean ocean temperature change T_o to return to its initial value under $0.5 \times \text{CO}_2$ and T_o at the time when CO_2 is lowered to $0.5 \times \text{CO}_2$, (b) the time for T_o to return to its initial value under $0.5 \times \text{CO}_2$ and the time integral of the radiative forcing during the phases with high CO_2 (F-up), and (c) the integral of the radiative forcing during the phases with high CO_2 (F-up) and during the phase with the low CO_2 (F-down) up to the time at which T_o returns to 0°C . Each symbol is the result of one of the simulations in Fig. 3. The dotted line indicates where the symbols would align if the warming and cooling were symmetrical. The solid line is the linear fit with a slope of -0.58 . (d) Results from the simulations of Fig. 1 for global mean ocean temperature change ($^\circ\text{C}$) as a function of the integral of the radiative forcing during the phase with high CO_2 and stabilization (solid lines) and continuing under $0.5 \times \text{CO}_2$ (dashed lines).

temperature to go back to its initial value (Fig. 5c) ($|\int F_{\text{down}} dt| < |\int F_{\text{up}} dt|$). Thus, the relationship between the oceanic temperature change and the integral of the forcing is different during the high- CO_2 and low- CO_2 phases (Fig. 5d). The amount of negative radiative forcing needed for the sea level to go back to its initial value is linearly related to the amount of positive radiative forcing:

$$\left| \int F_{\text{down}} dt \right| \approx 0.58 \left| \int F_{\text{up}} dt \right|.$$

The numerical factor depends on the scenario and model.

Although it exhibits reversibility, the two-layer model cannot explain why the ocean cools down more effectively than it warms up. In the two-layer model, the relationship between ocean temperature and $\int F dt$ is the same in the high- CO_2 and low- CO_2 phases (Fig. 3c), unlike in the AOGCM (Fig. 5d). The lower layer is a very large well-mixed heat capacity that sets the long time scale for approach to a steady state. Consequently, the lower layer warms much less than the upper layer in the high phase (at 140 yr in the 1% scenario, $T_u = 4.4^\circ\text{C}$ in Fig. 3a and $T_l = 0.4^\circ\text{C}$ in Fig. 3b). Because T_u closely tracks F , and $T_l \ll T_u$, the heat flux $\propto T_u - T_l$ between the layers has roughly the same time profile as F , and $T_l \propto \int F dt$. Because the single lower layer has too much heat capacity and does not warm up very much, the heat flux depends almost entirely on the upper temperature and hence on the integral of the radiative forcing. More layers would be needed to explain the faster loss of heat during the low- CO_2 phase.

To account for the asymmetry, we apply the step-response model of Good et al. (Good et al. 2011; Table 1). In this model, the evolution of any climate variable $X(t)$ is estimated by regarding it as the linear superposition of the responses to a succession of small instantaneous step-forcing changes $\delta F(t)$ in forcing, such that $F(t) = \int_0^t \delta F(t') dt'$. We carry out a single AOGCM experiment in which CO_2 is quadrupled at $t = 0$, giving forcing $F(4 \times \text{CO}_2)$, and then held constant. If $X = X_4(t)$ in this experiment, we estimate that $X(t) = \int_0^t X_4(t - t') \delta F(t') / F(4 \times \text{CO}_2) dt'$ —that is, a convolution of the response to the $4 \times \text{CO}_2$ step with the forcing time series. Good et al. (2011) show that this method is accurate for surface temperature and has skill at reproducing ocean heat uptake. We use the step model to study the experiment with 1% CO_2 followed by $0.5 \times \text{CO}_2$ as an example.

The step model successfully reproduces the evolution of the surface air temperature T_s and ocean temperature T_o during the high- CO_2 phase (Figs. 6a,b). (Note that T_s from the step model lacks the variability of the AOGCM because the convolution tends to smooth it

FIG. 6. Results for the climate response of the AOGCM and the step model in response to 1%CO₂ for 140 yr, then 0.5×CO₂. (top) Time series of (a) global mean surface air temperature and (b) global mean oceanic temperature. (c)–(f) zonal mean ocean temperature at (left) $t = 140$ yr and (right) $t = 224$ yr (when the mean oceanic temperature change is back to 0°C), for (c),(d) the AOGCM and (e),(f) the step model.

FIG. 7. Global mean ocean temperature change ($^{\circ}\text{C}$) as a function of the integral of the radiative forcing during the phase with high CO_2 ($1\%\text{CO}_2$, solid lines) and continuing under $0.5\times\text{CO}_2$ (dashed lines).

out.) The step model estimate of the ocean temperature change latitude–depth distribution is also very good (cf. Figs. 6c,e), showing the warming spreading downward from the surface, with deeper penetration at high latitudes.

The step model gives a better approximation of the evolution during the low- CO_2 phase than the zero- and two-layer models (Table 1). In particular, it correctly predicts that T_o returns to its initial value before year 280; it is qualitatively superior to the two-layer model in partly reproducing the asymmetry between the warming and cooling phases (Fig. 7). This is because the step model, like the AOGCM, has many layers and time scales. Unlike in the two-layer model, the ocean below the surface is not well mixed. The shallower layers have relatively small heat capacity and warm substantially during the high- CO_2 phase, and as time passes the warming spreads to deeper layers. During the low- CO_2 phase, the shallow layers likewise cool rapidly. In the AOGCM, this causes a large temperature contrast between the newly cooled surface and previously warmed subsurface layers, which forces heat out of the ocean more rapidly than it entered. In the step model, the large

temperature contrast and rapid heat loss are predicted as a consequence of the large negative step in forcing from $4\times\text{CO}_2$ to $0.5\times\text{CO}_2$. In both models, as a consequence of the high- CO_2 phase, additional heat is still being transported to deeper layers during the low- CO_2 phase at the same time as the cooling penetrates into the surface. In the global mean, similar qualitative behavior would be obtained from a vertical diffusion model of ocean temperature (e.g., Marčelja 2010). The advantages of the step model are that it can be applied in 3D, and that it emulates the AOGCM without any tuning of parameters being required.

However, the step model shows some inaccuracies. In particular, T_s falls too quickly at the start (Fig. 6a), and after about 30 years the step model underestimates the oceanic cooling progressively more seriously (Fig. 6b). These phenomena can be linked: both are related to an insufficient heat flux from lower layers toward the surface. Correspondingly, the cooling anomaly does not penetrate as deeply at high latitudes in the step model as in the AOGCM simulation, while in middle and low latitudes the thermocline does not cool as quickly (cf. Figs. 6d and 6f). The asymmetry between high- CO_2 and low- CO_2 phases is thus not as pronounced as in the AOGCM (Fig. 7).

We infer that these remaining discrepancies arise from nonlinear behavior of the AOGCM that the step model does not capture. In particular, the responses of the AOGCM to positive and negative step changes in forcing of equal magnitude are not equal and opposite (Fig. 8). This asymmetry cannot be captured by the step model, which is constructed by using the response to a positive step only, and presumably arises from the dependence of vertical heat transport on stability in the AOGCM (Stouffer and Manabe 1999). In particular, for the same absolute value of radiative forcing, the warming penetrates relatively deeper in the Arctic whereas

FIG. 8. Zonal-mean ocean temperature anomaly ($^{\circ}\text{C}$) as a function of depth after 70 yr of simulation, (a) under $4\times\text{CO}_2$ (warming) and (b) under $0.25\times\text{CO}_2$ (cooling). Both simulations start from the same initial state. The anomaly is with respect to the control simulation.

the cooling goes deeper in the Southern Ocean (Fig. 8). This different penetration of the heat anomaly during warming and cooling simulations has previously been described by Manabe et al. (1991), with similar results in the Southern Ocean but slightly different penetration of heat in the North Atlantic. After a sequence of warming ($1\%CO_2$ experiment) followed by a cooling ($0.5\times CO_2$ experiment), this different penetration of the heat anomaly results in a warmer Arctic and cooler Southern Ocean (except in the surface) in the AOGCM than in the step model at the time (224 years) when T_o in the AOGCM is back to its initial value.

e. Regional sea level changes

Impacts of sea level change arise from its regional distribution. As shown in Fig. 6, even when the global mean ocean temperature is back to its initial value, heat in the ocean is distributed differently, with some areas warmer and others colder. This means that there is still a regional pattern of thermosteric sea level change when T_o returns to $0^\circ C$ (Fig. 9). Although the global mean η is 0 m, sea level is higher than in the initial state in the Arctic and Atlantic south of $45^\circ N$, and lower in the North Atlantic and southern Indian Ocean. This distribution is likely to be model dependent as the regional sea level change changes from model to model (Pardaens et al. 2011), but the same qualitative point would hold. Thus, even if it were practical to eliminate the commitment to global thermosteric sea level change by negative radiative forcing, the commitment to regional sea level change would be yet more recalcitrant.

5. Conclusions

We have studied the future evolution of global mean sea level rise due to thermal expansion (i.e., not including contributions from ice sheets and glaciers) with an AOGCM under idealized CO_2 scenarios. Unlike surface temperature change, sea level change depends not only on the cumulative emission of CO_2 but also on the emission pathway. A greater rise in sea level results from earlier emissions than from later, for the same cumulative emission. Hence, targets to limit sea level rise would need to refer to rates of emissions as well as the total.

Thermal expansion will continue for many centuries if CO_2 emissions cease or if CO_2 concentration is stabilized. However, it is in principle reversible. Reducing the radiative forcing sufficiently would halt or reverse it, and a negative forcing would reverse it more quickly. Of course, reducing the forcing would require removal of CO_2 from the atmosphere or other geoengineering, which is not yet technologically feasible and might have

FIG. 9. Thermosteric sea level change (m) when the global mean ocean temperature is back to its initial value at $t = 224$ yr. The scenario followed is $1\%CO_2$ for 140 yr, then $0.5\times CO_2$.

side effects, while a large negative forcing would produce a climate colder than the preindustrial. Even if sea level returned to its preindustrial global mean, the geographical pattern would be different from the initial one because the penetration of heat is different during warming and cooling.

We have interpreted the behavior of thermal expansion in the AOGCM by comparison with a range of simpler models. The important qualitative difference between surface temperature and sea level is that the former depends on the prevailing radiative forcing and hence the cumulative CO_2 emission, while the latter depends on the time integral of radiative forcing and hence the time profile of CO_2 emission. The reversibility of thermal expansion cannot be explained without this distinction, which arises because surface temperature relates to the temperature of the upper ocean only, and thermal expansion to the full depth of the ocean. The AOGCM shows that sea level rise and fall are not symmetrical with respect to forcing. When the radiative forcing is reduced, heat leaves the ocean more readily than it entered. This is partly due to the vertical profile of the ocean temperature change, which retains a memory of the time profile of radiative forcing, and partly due to the dependence of vertical heat transport processes on temperature gradients. The success of the step-response model in largely reproducing the AOGCM results indicates that this method could be a useful way to continue investigations into AOGCM ocean heat uptake in a scenario-independent way although consideration of the nonlinear behavior may be necessary.

Acknowledgments. We thank A. Pardaens and P. Good for helpful discussion, and three anonymous reviewers for their constructive comments that helped improve the manuscript. J. A. Lowe acknowledges support from the

AVOID programme (DECC and Defra) under contract GA0215. The research leading to these results has received funding from the European Research Council under the European Community's Seventh Framework Programme (FP7/2007-2013), ERC Grant Agreement 247220, project "Seachange."

REFERENCES

- Allen, M. R., D. J. Frame, C. Huntingford, C. D. Jones, J. A. Lowe, M. Meinshausen, and N. Meinshausen, 2009a: Warming caused by cumulative carbon emissions towards the trillionth tonne. *Nature*, **458**, 1163–1166, doi:10.1038/nature08019.
- , and Coauthors, 2009b: The exit strategy. *Nat. Rep. Climate Change*, **3**, 56–58, doi:10.1038/climate.2009.38.
- Bowerman, N. H. A., D. J. Frame, C. Huntingford, J. A. Lowe, and M. R. Allen, 2011: Cumulative carbon emissions, emissions floors and short-term rates of warming: Implications for policy. *Philos. Trans. Roy. Soc.*, **369A**, 45–66, doi:10.1098/rsta.2010.0288.
- Charbit, S., D. Paillard, and G. Ramstein, 2008: Amount of CO₂ emissions irreversibly leading to the total melting of Greenland. *Geophys. Res. Lett.*, **35**, L12503, doi:10.1029/2008GL033472.
- Church, J. A., and Coauthors, 2011: Revisiting the Earth's sea-level and energy budgets from 1961 to 2008. *Geophys. Res. Lett.*, **38**, L18601, doi:10.1029/2011GL048794.
- Eby, M., K. Zickfeld, A. Montenegro, D. Archer, K. J. Meissner, and A. J. Weaver, 2009: Lifetime of anthropogenic climate change: Millennial time scales of potential CO₂ and surface temperature perturbations. *J. Climate*, **22**, 2501–2511.
- Friedlingstein, P., and Coauthors, 2006: Climate–carbon cycle feedback analysis: Results from the C⁴MIP model intercomparison. *J. Climate*, **19**, 3337–3353.
- Frölicher, T. L., and F. Joos, 2010: Reversible and irreversible impacts of greenhouse gas emissions in multi-century projections with the NCAR global coupled carbon cycle–climate model. *Climate Dyn.*, **35**, 1439–1459, doi:10.1007/s00382-009-0727-0.
- Gillett, N. P., V. K. Arora, K. Zickfeld, S. J. Marshall, and W. J. Merryfield, 2011: Ongoing climate change following a complete cessation of carbon dioxide emissions. *Nat. Geosci.*, **4**, 83–87, doi:10.1038/ngeo1047.
- Gloor, M., J. L. Sarmiento, and N. Gruber, 2010: What can be learned about carbon cycle climate feedbacks from the CO₂ airborne fraction? *Atmos. Chem. Phys.*, **10**, 7739–7751, doi:10.5194/acp-10-7739-2010.
- Good, P., J. M. Gregory, and J. A. Lowe, 2011: A step-response simple climate model to reconstruct and interpret AOGCM projections. *Geophys. Res. Lett.*, **38**, L01703, doi:10.1029/2010GL045208.
- Gordon, C., C. Cooper, C. A. Senior, H. Banks, J. M. Gregory, T. C. Johns, J. F. B. Mitchell, and R. A. Wood, 2000: The simulation of SST, sea ice extents and ocean heat transports in a version of the Hadley Centre coupled model without flux adjustments. *Climate Dyn.*, **16**, 147–168.
- Gregory, J. M., 2000: Vertical heat transports in the ocean and their effect on time-dependent climate change. *Climate Dyn.*, **16**, 501–515, doi:10.1007/s003820000059.
- , and J. F. B. Mitchell, 1997: The climate response to CO₂ of the Hadley Centre coupled AOGCM with and without flux adjustment. *Geophys. Res. Lett.*, **24**, 1943–1946, doi:10.1029/97GL01930.
- , and P. M. Forster, 2008: Transient climate response estimated from radiative forcing and observed temperature change. *J. Geophys. Res.*, **113**, D23105, doi:10.1029/2008JD010405.
- , C. D. Jones, P. Cadule, and P. Friedlingstein, 2009: Quantifying carbon-cycle feedbacks. *J. Climate*, **22**, 5232–5250.
- Held, I. M., M. Winton, K. Takahashi, T. Delworth, F. Zeng, and G. K. Vallis, 2010: Probing the fast and slow components of global warming by returning abruptly to preindustrial forcing. *J. Climate*, **23**, 2418–2427.
- Huybrechts, P., H. Goelzer, I. Janssens, E. Driesschaert, T. Fichefet, H. Goosse, and M.-F. Loutre, 2011: Response of the Greenland and Antarctic ice sheets to multi-millennial greenhouse warming in the earth system model of intermediate complexity LOVECLIM. *Surv. Geophys.*, **32**, 397–416, doi:10.1007/s10712-011-9131-5.
- Joos, F., M. Bruno, R. Fink, U. Siegenthaler, T. F. Stocker, C. Le Quéré, and J. L. Sarmiento, 1996: An efficient and accurate representation of complex oceanic and biospheric models of anthropogenic carbon uptake. *Tellus*, **48B**, 397–417, doi:10.1034/j.1600-0889.1996.t01-2-00006.x.
- Knorr, W., 2009: Is the airborne fraction of anthropogenic CO₂ emissions increasing? *Geophys. Res. Lett.*, **36**, L21710, doi:10.1029/2009GL040613.
- Lowe, J. A., J. M. Gregory, J. K. Ridley, P. Huybrechts, R. J. Nicholls, and M. Collins, 2006: The role of sea-level rise and the Greenland ice sheet in dangerous climate change: Implications for the stabilisation of climate. *Avoiding Dangerous Climate Change*, H. J. Schellnhuber et al., Eds., Cambridge University Press, 29–36.
- Manabe, S., R. J. Stouffer, M. J. Spelman, and K. Bryan, 1991: Transient responses of a coupled ocean–atmosphere model to gradual changes of atmospheric CO₂. Part I: Annual mean response. *J. Climate*, **4**, 785–818.
- Marčelja, S., 2010: The timescale and extent of thermal expansion of the global ocean due to climate change. *Ocean Sci.*, **6**, 179–184.
- Matthews, H. D., N. P. Gillett, P. A. Stott, and K. Zickfeld, 2009: The proportionality of global warming to cumulative carbon emissions. *Nature*, **459**, 829–832, doi:10.1038/nature08047.
- Meehl, G. A., W. M. Washington, W. D. Collins, J. M. Arblaster, A. Hu, L. E. Buja, W. G. Strand, and H. Teng, 2005: How much more global warming and sea level rise? *Science*, **307**, 1769–1772, doi:10.1126/science.1106663.
- Moss, R. H., and Coauthors, 2010: The next generation of scenarios for climate change research and assessment. *Nature*, **463**, 747–756, doi:10.1038/nature08823.
- Pardaens, A. K., J. M. Gregory, and J. A. Lowe, 2011: A model study of factors influencing projected changes in regional sea level over the twenty-first century. *Climate Dyn.*, **36**, 2015–2033, doi:10.1007/s00382-009-0738-x.
- Plattner, G.-K., and Coauthors, 2008: Long-term climate commitments projected with climate–carbon cycle models. *J. Climate*, **21**, 2721–2751.
- Raper, S. C. B., J. M. Gregory, and R. J. Stouffer, 2002: The role of climate sensitivity and ocean heat uptake on AOGCM transient temperature response. *J. Climate*, **15**, 124–130.
- Ridley, J., J. M. Gregory, P. Huybrechts, and J. Lowe, 2009: Thresholds for irreversible decline of the Greenland ice sheet. *Climate Dyn.*, **35**, 1049–1057, doi:10.1007/s00382-009-0646-0.
- Russell, G. L., V. Gornitz, and J. R. Miller, 2000: Regional sea-level changes projected by the NASA/GISS atmosphere–ocean model. *Climate Dyn.*, **16**, 789–797, doi:10.1007/s003820000090.

- Shepherd, J., and Coauthors, 2009: Geoengineering the climate: Science, governance and uncertainty. Royal Society Rep., 81 pp. [Available online at <http://royalsociety.org/policy/publications/2009/geoengineering-climate/>.]
- Smith, R. S., J. M. Gregory, and A. Osprey, 2008: A description of the FAMOUS (version XDBUA) climate model and control run. *Geosci. Model Dev.*, **1**, 53–68, doi:10.5194/gmd-1-53-2008.
- Solomon, S., G.-K. Plattner, R. Knutti, and P. Friedlingstein, 2009: Irreversible climate change due to carbon dioxide emissions. *Proc. Natl. Acad. Sci. USA*, **106**, 1704–1709, doi:10.1073/pnas.0812721106.
- Stouffer, R. J., and S. Manabe, 1999: Response of a coupled ocean–atmosphere model to increasing atmospheric carbon dioxide: Sensitivity to the rate of increase. *J. Climate*, **12**, 2224–2237.
- Vizcaíno, M., U. Mikolajewicz, J. Jungclaus, and G. Schurgers, 2010: Climate modification by future ice sheet changes and consequences for ice sheet mass balance. *Climate Dyn.*, **34**, 301–324, doi:10.1007/s00382-009-0591-y.
- Zickfeld, K., M. Eby, H. D. Matthews, and A. J. Weaver, 2009: Setting cumulative emissions targets to reduce the risk of dangerous climate change. *Proc. Natl. Acad. Sci. USA*, **106**, 16 129–16 134, doi:10.1073/pnas.0805800106.
- , V. K. Arora, and N. P. Gillett, 2012: Is the climate response to CO₂ emissions path dependent? *Geophys. Res. Lett.*, **39**, L05703, doi:10.1029/2011GL050205.