

The Brownian traveller on manifolds

Article

Accepted Version

Kolb, M. and Krejcirik, D. (2013) The Brownian traveller on manifolds. *Journal of Spectral Theory*, 4 (2). pp. 235-281. ISSN 1664-0403 doi: 10.4171/JST/69 Available at <https://centaur.reading.ac.uk/34321/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

To link to this article DOI: <http://dx.doi.org/10.4171/JST/69>

Publisher: European Mathematical Society

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online

The Brownian traveller on manifolds

Martin Kolb^a and David Krejčířík^{b,c}

- a) *Department of Mathematics and Statistics, University of Reading, Whiteknights, PO Box 220, Reading RG6 6AX, United Kingdom; m.kolb@reading.ac.uk*
- b) *Department of Theoretical Physics, Nuclear Physics Institute ASCR, 25068 Řež, Czech Republic; krejcirik@ujf.cas.cz*
- c) *IKERBASQUE, Basque Foundation for Science, 48011 Bilbao, Kingdom of Spain*

9 January 2013

Abstract

We study the influence of the intrinsic curvature on the large time behaviour of the heat equation in a tubular neighbourhood of an unbounded geodesic in a two-dimensional Riemannian manifold. Since we consider killing boundary conditions, there is always an exponential-type decay for the heat semigroup. We show that this exponential-type decay is slower for positively curved manifolds comparing to the flat case. As the main result, we establish a sharp extra polynomial-type decay for the heat semigroup on negatively curved manifolds comparing to the flat case. The proof employs the existence of Hardy-type inequalities for the Dirichlet Laplacian in the tubular neighbourhoods on negatively curved manifolds and the method of self-similar variables and weighted Sobolev spaces for the heat equation.

1 Introduction

The intimate intertwining between properties of Brownian motion (or alternatively the heat flow) on a Riemannian manifold and the curvature properties of the manifold are a classical research question that has been investigated extensively (see, *e.g.*, [20, 21, 12, 13, 31, 18]) and has led to deep results and new methods, which turned out to be also of importance in other fields of mathematics. For a more complete overview and further references we refer to recent monograph [14]. One of the main themes here is to characterize probabilistic properties via geometric ones and vice versa. Thinking of the Brownian particle as a ‘traveller’ in a curved space we continue this line of research and investigate the influence of the curvature on its large time behaviour.

However, in contrast to previous works, we restrict the motion of the Brownian particle to a tubular neighbourhood of a curve in the Riemannian manifold and kill it when it leaves this quasi-one-dimensional subset. This line of research seems to have its origin in the mathematical physics literature, where

one aims to describe the dynamics of quantum particles in very thin almost one-dimensional waveguides. The constraint on the Brownian motion to the quasi-one-dimensional subsets leads to additional effects not present in the case of an unrestricted stochastic conservative motion. In particular it will turn out that the behaviour of the Brownian particle in the tube-like set is sensitive to local perturbations of the geometry.

A more precise description of our setting is the following. Let the ambient space of the Brownian traveller be a complete non-compact *two-dimensional* Riemannian manifold \mathcal{A} (not necessarily embedded in the Euclidean space \mathbb{R}^3) with Gauss curvature K . We restrict to the case of *locally perturbed traveller* by assuming that K is compactly supported.

We further assume that the motion is *quasi-one-dimensional* in the sense that the Brownian traveller is forced to move along an infinite curve Γ on the surface \mathcal{A} . To focus on the effects induced by the intrinsic curvature K itself, we suppress side effects induced by the curvature of the curve by assuming that Γ is a geodesic.

The constraint to move along the geodesic curve is introduced by imposing *killing boundary conditions* on the boundary of the tubular neighbourhood

$$\Omega := \{q \in \mathcal{A} \mid \text{dist}(q, \Gamma) < a\}, \quad (1.1)$$

where a is a positive (not necessarily small) number. That is, the Brownian traveller ‘dies’ whenever it hits the boundary $\partial\Omega$ of the strip Ω .

The problem is mathematically described by the diffusion equation

$$\begin{cases} \partial_t u - \Delta_q u = 0 & \text{in } \Omega \times (0, \infty), \\ u = 0 & \text{on } \partial\Omega \times (0, \infty), \\ u = u_0 & \text{on } \Omega \times \{0\}, \end{cases} \quad (1.2)$$

in the space time variables $(q, t) \in \Omega \times (0, \infty)$, where u_0 is an initial datum. More specifically, for the Dirac distribution $u_0(q) = \delta(q - q_0)$, the solution $u(q, t)$ is related to the density of the transition probability of the Brownian motion starting at $q_0 \in \Omega$ as follows. Let us denote by \mathbb{E}_q (respectively, \mathbb{P}_q) the expectation (respectively, probability) of a stochastic process $(X_t)_{t \geq 0}$ on the manifold \mathcal{A} started at $q \in \mathcal{A}$ which is given as the solution of the martingale problem associated to the operator Δ . By a slight abuse of notation we call this process Brownian motion despite the fact that the generator of the ‘true’ Brownian motion is $\frac{1}{2}\Delta$. (This should not lead to any confusion as the relation between both processes is given by a simple time change.) Moreover, let $\tau_\Omega := \inf\{t > 0 \mid X_t \in \partial\Omega\}$ denote the first exit time. Then

$$u(q, t) = \mathbb{E}_q[u_0(X_t), \tau_\Omega > t] \quad (1.3)$$

solves equation (1.2). If $u_0 = \chi_B$ for some measurable set $B \subset \Omega$, we get

$$u(q, t) = \mathbb{P}_q(X_t \in B, \tau_\Omega > t), \quad (1.4)$$

which is the probability that the Brownian particle survived up to time t and is in B at time t .

Now imagine a Brownian traveller in Ω and we imagine that he/she reached his/her goal when hitting the boundary. The ultimate question we would like

to address in this paper is to decide *which geometry is better to travel*. By the ‘good geometry’ we understand that which enables the Brownian traveller to reach his/her goal as soon as possible or ‘to escape from his/her starting point as far as possible’. More precisely, we are interested in quantifying the large time of (1.4) for bounded sets $B \subset \Omega_0$.

In any case, the question is related to the large time decay of the solutions of (1.2) as regards the curvature K . We mainly study a Hilbert-space version of the problem by analysing the asymptotic behaviour of the heat semigroup on $L^2(\Omega)$ associated with (1.2). Nevertheless, we establish some pointwise results about the large time behaviour of $u(q, t)$ as well.

Our results are informally summarized in Table 1.

curvature	positive	zero	negative
transport	<i>bad</i>	<i>critical</i>	<i>good</i>
probability decay	$e^{\gamma t} e^{-E_1 t}$	$t^{-1/2} e^{-E_1 t}$	$t^{-3/2} e^{-E_1 t}$

Table 1: An informal summary of our results.

There $E_1 := \pi^2/(2a)^2$ denotes the lowest Dirichlet eigenvalue of the strip cross-section $(-a, a)$ and γ is a positive number. As explained above, the vague statements about transport in Table 1 should be understood in the spirit of the large time decay of the solutions to (1.2) stated there. It turns out that the solutions of (1.2) has worse (respectively, better) decay properties if K is non-negative (respectively, non-positive) as a consequence of the existence of stationary solutions (respectively, Hardy-type inequalities). More general results, involving surfaces with sign changing curvatures, are established in this paper.

The effect of curvature on the transience/recurrence of a Brownian particle have been extensively studied (see [12] for a nice review). It turned out that on manifolds with ‘large’ negative curvature Brownian motion leaves compact subsets faster than on manifolds with non-negative curvature. But local changes of the Riemannian metric cannot change transience to recurrence or vice versa. Observe that for the results presented in Table 1 this is not longer true. In probabilistic literature this corresponds to the R -recurrence/ R -transience dichotomy (see [46, 45]) or in analytic literature to the critical/subcritical dichotomy (see, *e.g.*, [36], or [34, 35] for a brief overview). Indeed, in our setting the Brownian motion in the negatively curved tube with *compactly supported* curvature is E_1 -transient in contrast to the case of no curvature.

The organization of this paper is as follows. In the forthcoming Sections 2 and 3 we properly define the configuration space of the Brownian traveller and the associated heat equation (1.2), respectively. The case of zero curvature is briefly mentioned in Section 4. In Section 5 we consider direct consequences in a more general situation when the curvature vanishes at infinity. The influence of positive curvature on the Brownian traveller is studied in Section 6. The main part of the paper consists of Section 7, where we establish the existence of Hardy-type inequalities in negatively curved manifolds and develop the method of self-similar variables for the heat equation to reveal the subtle effect of negative curvature. The paper is concluded by Section 8 where we summarize our results and refer to some open problems.

2 Geometric preliminaries

We start by imposing some natural hypotheses to give an instructive geometrical interpretation of the configuration space Ω of the Brownian traveller. The conditions will be considerably weakened later when we reconsider the problem in an abstract setting.

2.1 The configuration space

Let us assume that the Riemannian manifold \mathcal{A} is of class C^2 and that its Gauss curvature K is continuous. The latter holds under the additional assumption that \mathcal{A} is either of class C^3 (by Gauss's Theorema Egregium) or that it is embedded in \mathbb{R}^3 (by computing principal curvatures).

Any geodesic curve Γ on \mathcal{A} is C^2 -smooth and, without loss of generality, we may assume that it is parameterized by arc-length. To enable the traveller to propagate to infinity, we consider unbounded geodesics Γ only. For a moment, we make the strong hypothesis that $\Gamma : \mathbb{R} \rightarrow \mathcal{A}$ is an embedding.

Since Γ is parameterized by arc-length, the derivative $T := \dot{\Gamma}$ defines the unit tangent vector field along Γ . Let N be the unit normal vector field along Γ which is uniquely determined as the C^1 -smooth mapping from \mathbb{R} to the tangent bundle of \mathcal{A} by requiring that $N(s)$ is orthogonal to $T(s)$ and that $\{T(s), N(s)\}$ is positively oriented for all $s \in \mathbb{R}$ (cf [40, Sec. 7.B]).

The feature of our model is that the Brownian traveller is assumed to be confined to the strip-like a -tubular neighbourhood (1.1). By definition, Ω is the set of points q in \mathcal{A} for which there exists a geodesic of length less than a from q meeting Γ orthogonally. More precisely, we introduce a mapping \mathcal{L} from the flat strip

$$\Omega_0 := \mathbb{R} \times (-a, a) \quad (2.1)$$

(considered as a subset of the tangent bundle of \mathcal{A}) to the manifold \mathcal{A} by setting

$$\mathcal{L}(x_1, x_2) := \exp_{\Gamma(x_1)}(N(x_1) x_2), \quad (2.2)$$

where \exp_q is the exponential map of \mathcal{A} at $q \in \mathcal{A}$. Then we have

$$\Omega = \mathcal{L}(\Omega_0). \quad (2.3)$$

Note that $x_1 \mapsto \mathcal{L}(x_1, x_2)$ traces the curves parallel to Γ at a fixed distance $|x_2|$, while the curve $x_2 \mapsto \mathcal{L}(x_1, x_2)$ is a geodesic orthogonal to Γ for any fixed x_1 . See Figure 1.

2.2 The Fermi coordinates

Making the hypothesis that

$$\mathcal{L} : \Omega_0 \rightarrow \Omega \text{ is a diffeomorphism,} \quad (2.4)$$

we get a convenient parametrization of Ω via the (Fermi or geodesic parallel) 'coordinates' $x = (x_1, x_2)$ determined by (2.2), cf Figure 1. We refer to [10, Sec. 2] and [16] for the notion and properties of Fermi coordinates. In particular,




Figure 1: The parametrization of the strip Ω via the Fermi coordinates $x = (x_1, x_2)$ defined by (2.2).

it follows by the generalized Gauss lemma that the metric G induced by (2.2) acquires the diagonal form:

$$G = \begin{pmatrix} f^2 & 0 \\ 0 & 1 \end{pmatrix}, \quad (2.5)$$

where f is continuous, has continuous partial derivatives $\partial_2 f$, $\partial_2^2 f$ and satisfies the Jacobi equation

$$\partial_2^2 f + Kf = 0 \quad \text{with} \quad \begin{cases} f(\cdot, 0) = 1, \\ \partial_2 f(\cdot, 0) = 0. \end{cases} \quad (2.6)$$

Here K is considered as a function of the Fermi coordinates (x_1, x_2) .

By the inverse function theorem, a sufficient condition to ensure (2.4) is that \mathcal{L} is injective and f positive. The latter can always be achieved for sufficiently small a as the following lemma shows.

Lemma 2.1. *Let $K \in L^\infty(\Omega_0)$ and $\|K\|_\infty a^2 < 1$. For every $x \in \Omega_0$, we have*

$$1 - \frac{\bar{K}(x_1)a^2}{1 - \bar{K}(x_1)a^2} \leq f(x) \leq 1 + \frac{\bar{K}(x_1)a^2}{1 - \bar{K}(x_1)a^2}, \quad (2.7)$$

where $\|\cdot\|_\infty := \|\cdot\|_{L^\infty(\Omega_0)}$ and

$$\bar{K}(x_1) := \operatorname{esssup}_{x_2 \in (-a, a)} |K(x_1, x_2)|.$$

Proof. Integrating (2.6), we arrive at the identity

$$\forall x \in \Omega_0, \quad \partial_2 f(x) = - \int_0^{x_2} (Kf)(x_1, \xi) d\xi.$$

Consequently,

$$|\partial_2 f(x)| \leq a \bar{K}(x_1) \bar{f}(x_1), \quad \text{with} \quad \bar{f}(x_1) := \sup_{\xi \in (-a, a)} |f(x_1, \xi)|, \quad (2.8)$$

for all $x \in \Omega_0$. By the mean value theorem, we deduce the bounds

$$\forall x \in \Omega_0, \quad 1 - a^2 \bar{K}(x_1) \bar{f}(x_1) \leq f(x) \leq 1 + a^2 \bar{K}(x_1) \bar{f}(x_1). \quad (2.9)$$

Taking the supremum over $x_2 \in (-a, a)$, the upper bound leads to the upper bound of (2.7). Finally, using the upper bound of (2.7) to estimate \bar{f} in the lower bound of (2.9), we conclude with the lower bound of (2.7). \square

2.3 The abstract setting

It follows from the preceding subsection that, under the hypothesis (2.4), we can identify $\Omega \subset \mathcal{A}$ with the Riemannian manifold (Ω_0, G) . However, the assumption (2.4) is not really essential provided that one is ready to abandon the geometrical interpretation of Ω as a tubular neighbourhood embedded in \mathcal{A} .

Indeed, (Ω_0, G) , with the metric G determined by (2.5) and (2.6), can be considered as an abstract Riemannian manifold for which the boundedness of K and a restriction of a are the only important hypotheses. More specifically, we assume

$$K \in L^\infty(\Omega_0) \quad \text{and} \quad \|K\|_\infty a^2 < \frac{1}{2}. \quad (2.10)$$

Then the Jacobi equation (2.6) admits a solution $f(x_1, \cdot) \in H^2((-a, a))$ for every $x_1 \in \mathbb{R}$ and it follows from Lemma 2.1 that f is bounded and uniformly positive on Ω_0 .

In the sequel, we therefore allow for self-intersections and low regularity of Ω by considering (Ω_0, G) as an abstract configuration space of the Brownian traveller. The mere boundedness of the metric G is sufficient to establish the desired results.

3 Analytic and probabilistic preliminaries

In this section, we give a precise meaning to the evolution problem (1.2).

3.1 The generator of motion

The meaning of $-\Delta_q u$ in (1.2) should be understood as an action of the Laplace-Beltrami operator $-\Delta$ in the Riemannian manifold Ω . In the Fermi coordinates, considering $-\Delta$ as a differential expression in Ω_0 , we have

$$-\Delta = -|G|^{-1/2} \partial_i |G|^{1/2} G^{ij} \partial_j = -f^{-1} \partial_1 f^{-1} \partial_1 - f^{-1} \partial_2 f \partial_2. \quad (3.1)$$

Here the first identity is a general formula for the Laplace-Beltrami operator in a manifold equipped with the metric G , with the usual notation for the determinant $|G| := \det(G)$ and the coefficients G^{ij} of the inverse metric G^{-1} , and using the Einstein summation convention. The second identity employs the special form of the metric (2.5) in the Fermi coordinates.

The objective of this subsection is to associate to the differential expression (3.1) a self-adjoint operator H_K in the Hilbert space

$$L_f^2(\Omega_0) := L^2(\Omega_0, f(x) dx), \quad (3.2)$$

a space isomorphic to $L^2(\Omega)$ via the Fermi coordinates. In order to implement the Dirichlet boundary conditions of (1.2), we introduce H_K as the Friedrichs

extension of (3.1) initially defined on smooth functions of compact support in Ω_0 (cf [2, Sec. 6]). That is, H_K is the unique self-adjoint operator associated on (3.2) with the quadratic form

$$h_K[\psi] := (\partial_i \psi, G^{ij} \partial_j \psi)_f, \quad \psi \in \mathfrak{D}(h_K) := H_0^1(\Omega_0, G). \quad (3.3)$$

Here $(\cdot, \cdot)_f$ denotes the inner product in (3.2) and $H_0^1(\Omega_0, G)$ denotes the completion of $C_0^\infty(\Omega_0)$ with respect to the norm $\|\cdot\|_{\mathfrak{D}(h_K)} := (h_K[\cdot] + \|\cdot\|_f^2)^{1/2}$, with $\|\cdot\|_f$ denoting the norm in (3.2). The dependence of H_K on the curvature K is understood through the dependence of f on K , cf (2.6).

Under our hypothesis (2.10), it follows from Lemma 2.1 that $\|\cdot\|_f$ is equivalent to the usual norm $\|\cdot\| := \|\cdot\|_1$ in $L^2(\Omega_0) = L_1^2(\Omega_0)$ (i.e. $f = 1$) and, moreover, the $\mathfrak{D}(h_K)$ -norm is equivalent to the usual norm in the Sobolev space $H^1(\Omega_0)$. Consequently,

$$\mathfrak{D}(h_K) = H_0^1(\Omega_0).$$

However, it is important to keep in mind that, although $H^1(\Omega_0, G)$ and $H_0^1(\Omega_0)$ coincide as vector spaces, their topologies are different.

Remark 3.1. Under extra regularity assumptions involving derivatives of f , it is possible to show that H_K acts as (3.1) on the domain $H_0^1(\Omega_0) \cap H^2(\Omega_0)$. However, we shall not need these facts, always considering H_K in the form sense described above.

3.2 The dynamics

As usual, we consider the weak formulation of the parabolic problem (1.2). We say a Hilbert space-valued function $u \in L_{\text{loc}}^2((0, \infty); H_0^1(\Omega_0, G))$, with the weak derivative $u' \in L_{\text{loc}}^2((0, \infty); [H_0^1(\Omega_0, G)]^*)$, is a (global) solution of (1.2) provided that

$$\langle v, u'(t) \rangle_f + h_K(v, u(t)) = 0 \quad (3.4)$$

for each $v \in H_0^1(\Omega_0, G)$ and a.e. $t \in [0, \infty)$, and $u(0) = u_0$. Here $h_K(\cdot, \cdot)$ denotes the sesquilinear form associated with (3.3) and $\langle \cdot, \cdot \rangle_f$ stands for the pairing of $H_0^1(\Omega_0, G)$ and its dual $[H_0^1(\Omega_0, G)]^*$. With an abuse of notation, we denote by the same symbol u both the function on $\Omega_0 \times (0, \infty)$ and the mapping $(0, \infty) \rightarrow H_0^1(\Omega_0, G)$.

Standard semigroup theory implies that there indeed exists a unique solution of (3.4) that belongs to $C^0([0, \infty); L_f^2(\Omega_0))$. More precisely, the solution is given by $u(t) = e^{-tH_K} u_0$, where e^{-tH_K} is the semigroup associated with H_K .

It is easy to see that the real and imaginary parts of the solution u of (1.2) evolve separately. By writing $u = \Re(u) + i \Im(u)$ and solving (1.2) with initial data $\Re(u_0)$ and $\Im(u_0)$, we may therefore reduce the problem to the case of a real function u_0 , without restriction. This reflects the fact that e^{-tH_K} is positivity preserving. Consequently, the functional spaces can be considered to be real when investigating the heat equation (1.2).

Indeed, the quadratic form h_K is a Dirichlet form, to which we can associate a strong Markov process with continuous paths (which as we mentioned call Brownian motion on (Ω_0, G) even though our normalization of the heat equation differs by a factor 1/2 from the probabilistic one). In order to do so let us first

extend f to \mathbb{R}^2 by setting it equal to 1 outside Ω_0 . Moreover, let us define the Dirichlet form \tilde{h}_K in $L^2(\mathbb{R}^2, f(x) dx)$ by

$$\tilde{h}_K[\psi] := \int_{\mathbb{R}^2} \overline{\partial_i \psi(x)} G^{ij}(x) \partial_j \psi(x) f(x) dx, \quad \psi \in \mathfrak{D}(\tilde{h}_K) := H^1(\mathbb{R}^2).$$

Then there exists a strong Markov process $(X_t)_{t \geq 0}$ with continuous paths, which is associated to \tilde{h}_K . According to Theorem 4 in [42] the process is conservative. We use \mathbb{E}_x (respectively, \mathbb{P}_x) to denote the expectation (respectively, probability) conditional on $X_0 = x$. Since Dirichlet boundary conditions correspond to killing in the probabilistic picture, we have the following probabilistic representation

$$e^{-tH_K} u_0(x) = \mathbb{E}_x[u_0(X_t), \tau_{\Omega_0} > t] \quad (3.5)$$

for almost every $x \in \Omega_0$.

3.3 Basic properties

In our first proposition we collect some fundamental properties of the stochastic process $(X_t)_{t \geq 0}$.

Proposition 3.1. *Assume (2.10).*

- *The stochastic process $(X_t)_{t \geq 0}$ has the strong Feller property and is therefore well-defined for every $x \in \Omega_0$. In particular, the right hand side of (3.5) is continuous for every $u_0 \in L^\infty(\Omega_0)$.*
- *The stochastic process $(X_t)_{t \geq 0}$ has a continuous transition function $k_t(\cdot, \cdot)$ with respect to $f(x) dx$, which satisfies a Gaussian bound, i.e. for some constants $C_1 > 0$, $C_2 > 0$, one has*

$$\forall x, y \in \Omega_0, \quad k_t(x, y) \leq \frac{C_1}{t} e^{-\frac{|x-y|^2}{C_2 t}}.$$

Proof. The first assertion follows immediately from the second one by a standard use of Lebesgue's dominated convergence theorem.

In order to prove the second assertion, let us denote by \tilde{H}_K the unique self-adjoint operator associated to \tilde{h}_K . Observe that according to [32, Thm. 1.1] the semigroup $e^{-t\tilde{H}_K}$ has an integral kernel, satisfying a Gaussian upper bound. As e^{tH_K} is dominated by $e^{-t\tilde{H}_K}$ (using either [33] or the probabilistic representation), this bound for \tilde{H}_K carries over to H_K . In order to prove the regularity assertion concerning the transition kernel, observe that the Dirichlet form \tilde{h}_K corresponds to a uniformly elliptic operator (in the sense of [41, Sec. 4]) on the subset Ω_0 of the Riemannian manifold \mathbb{R}^2 with Euclidean metric. Thus, according to the remark below Theorem 6.3 in [41] (compare also [43]), it therefore follows that the transition kernel is locally Hölder continuous. \square

In this work we are mainly interested in the large time behaviour of the stochastic process $(X_t)_{\tau_{\Omega_0} > t \geq 0}$, which is well-known to be connected to spectral properties of its generator H_K . The spectral mapping theorem yields

$$\|e^{-tH_K}\|_{L_f^2(\Omega_0) \rightarrow L_f^2(\Omega_0)} = e^{-\lambda_K t} \quad (3.6)$$

for each time $t \geq 0$, where λ_K denotes the lowest point in the spectrum of H_K , *i.e.*, $\lambda_K := \inf \sigma(H_K)$. Hence, it is important to understand the low-energy properties of H_K in order to study the large time behaviour of the solutions of (1.2).

From equation (3.6) and Proposition 3.1 we deduce the following result showing that the exponential rate of decay of $\mathbb{P}_x(X_t \in B, \tau_{\Omega_0} > t)$ is given by the lowest point in the spectrum.

Proposition 3.2. *Assume (2.10). For any open subset $B \subset \Omega_0$ and every $x \in \Omega_0$,*

$$-\lim_{t \rightarrow \infty} \frac{1}{t} \log \mathbb{P}_x(X_t \in B, \tau_{\Omega_0} > t) = \lambda_K.$$

We omit the proof which can be modelled by following the arguments from [39] and [38] used there in the context of Schrödinger operators. A strongly related assertion with a related proof can be found in Theorem 10.14 in [14].

One of the main themes is that a better understanding of low-energy properties of H_K leads to much more precise estimates.

4 Flat manifolds

We say that (a submanifold of) \mathcal{A} is *flat* if its Gauss curvature K is identically equal to zero (on the submanifold). The Brownian motion in a flat ambient space is easy to understand because Ω coincides with the straight Euclidean strip Ω_0 , *i.e.* G is identity, for which the heat equation (1.2) can be solved by separation of variables.

4.1 Separation of variables

By the ‘separation of variables’ mentioned above we mean precisely that the Dirichlet Laplacian $H_0 = -\Delta_D^{\Omega_0}$ on $L^2(\Omega_0)$ can be identified with the decomposed operator

$$(-\Delta^{\mathbb{R}}) \otimes 1 + 1 \otimes (-\Delta_D^{(-a,a)}) \quad \text{in} \quad L^2(\mathbb{R}) \otimes L^2((-a,a)). \quad (4.1)$$

Here we denote by $-\Delta_D^U$ the Dirichlet Laplacian on $L^2(U)$ for any open Euclidean set U , suppress the subscript D if the boundary of U is empty, and 1 stands for the identity operators in the appropriate spaces. In a probabilistic language, (4.1) is essentially a reformulation of the fact that the horizontal and the vertical component of $(X_t)_{t \geq 0}$ are independent.

The eigenvalues and (normalized) eigenfunctions of $-\Delta_D^{(-a,a)}$ are respectively given by $(n = 1, 2, \dots)$

$$E_n := \left(\frac{n\pi}{2a}\right)^2, \quad \mathcal{J}_n(x_2) := \sqrt{\frac{1}{a}} \sin \left[\sqrt{E_n} (x_2 + a) \right], \quad (4.2)$$

while the spectral resolution of $-\Delta^{\mathbb{R}}$ is obtained by the Fourier transform. Then it is easy to see that the heat semigroup e^{-tH_0} is an integral operator with kernel

$$s_0(x, x', t) := \sum_{n=1}^{\infty} e^{-E_n t} \mathcal{J}_n(x_2) p(x_1, x'_1, t) \mathcal{J}_n(x'_2), \quad (4.3)$$

where

$$p(x_1, x'_1, t) := \frac{e^{-(x_1 - x'_1)^2 / (4t)}}{\sqrt{4\pi t}}$$

is the well known heat kernel of $-\Delta^{\mathbb{R}}$.

4.2 The decay rate

Concerning the large time behaviour of e^{-tH_0} , it follows from the decomposition (4.1) that

$$\sigma(H_0) = \sigma_{\text{ess}}(H_0) = [E_1, \infty), \quad (4.4)$$

and therefore, as a consequence of (3.6),

$$\|e^{-tH_0}\|_{L^2(\Omega_0) \rightarrow L^2(\Omega_0)} = e^{-E_1 t} \quad (4.5)$$

for each time $t \geq 0$. Consequently, any solution of (1.2) satisfies the global decay estimate $\|u(t)\| \leq e^{-E_1 t} \|u_0\|$ for every $t \geq 0$.

However, it is possible to obtain an extra polynomial decay for solutions with initial data decaying sufficiently fast at the infinity of the strip Ω_0 . To see it, let us consider the weight function

$$w(x) := e^{x_1^2/4} \quad (4.6)$$

and restrict the class of initial data to those u_0 which belong to the weighted space $L_w^2(\Omega_0)$ defined in the same way as (3.2). Then we have the improved decay estimate $\|u(t)\| \leq C t^{-1/4} e^{-E_1 t} \|u_0\|_w$ for every $t \geq 1$. This is a consequence of the following result.

Proposition 4.1. *There exists a positive constant C such that for every $t \geq 1$,*

$$C^{-1} t^{-1/4} e^{-E_1 t} \leq \|e^{-tH_0}\|_{L_w^2(\Omega_0) \rightarrow L^2(\Omega_0)} \leq C t^{-1/4} e^{-E_1 t}.$$

Moreover, for every bounded open set $B \subset \Omega_0$ and $x \in \Omega_0$ there is a constant $C_{B,x}$ such that for $t \geq 1$,

$$C_{B,x}^{-1} t^{-1/2} e^{-E_1 t} \leq \mathbb{P}_x(X_t \in B, \tau_{\Omega_0} > t) \leq C_{B,x} t^{-1/2} e^{-E_1 t}.$$

Proof. The second assertion is a rather immediate consequence of (4.3). In order to see this, observe that

$$\begin{aligned} \mathbb{P}_x(X_t \in B, \tau_{\Omega_0} > t) &= \sum_{n=1}^{\infty} e^{-E_n t} \mathcal{J}_n(x_2) \int_B p(x_1, x'_1, t) \mathcal{J}_n(x'_2) dx \\ &= e^{-E_1 t} \mathcal{J}_1(x_2) \int_B p(x_1, x'_1, t) \mathcal{J}_1(x'_2) d(x'_1, x'_2) + R_B(t, x_1, x_2), \end{aligned} \quad (4.7)$$

where $R_B(t, x_1, x_2)$ satisfies $|R_B(t, x_1, x_2)| \leq e(x_1, x_2) e^{-E_2 t}$ ($t \geq 1$) for some locally bounded function $e : \Omega_0 \rightarrow \mathbb{R}_+$. Thus there exists $t_0 = t_0(x_1, x_2, B) \geq 1$ such that for every $t \geq t_0$ one has

$$|R_B(t, x_1, x_2)| \leq \frac{1}{2} e^{-E_1 t} \mathcal{J}_1(x_2) \int_B p(x_1, x'_1, t) \mathcal{J}_1(x'_2) dx'.$$

Therefore from (4.7) we conclude that for $t \geq t_0$

$$\begin{aligned} \frac{1}{2} e^{-E_1 t} \mathcal{J}_1(x_2) \int_B p(x_1, x'_1, t) \mathcal{J}_1(x'_2) dx' &\leq \mathbb{P}_x(X_t \in B, \tau_{\Omega_0} > t) \\ &\leq \frac{3}{2} e^{-E_1 t} \mathcal{J}_1(x_2) \int_B p(x_1, x'_1, t) \mathcal{J}_1(x'_2) dx', \end{aligned}$$

which, using the explicit form of p , gives the assertion for $t \geq t_0$. Adjusting the constants $C_{B,x}$ allows to extend this to $t \geq 1$.

Let us now consider the first assertion. Using the Schwarz inequality, we get

$$\begin{aligned} \|e^{-tH_0} u_0\|^2 &\leq \|u_0\|_w^2 \int_{\Omega_0 \times \Omega_0} s_0(x, x', t)^2 w(x')^{-1} dx dx' \\ &= \|u_0\|_w^2 e^{-2E_1 t} \sum_{n=1}^{\infty} e^{-2(E_n - E_1)t} \int_{\mathbb{R} \times \mathbb{R}} p(x_1, x'_1, t)^2 e^{-x_1'^2/4} dx_1 dx'_1 \end{aligned}$$

for every $u_0 \in L_w^2(\Omega_0)$ and $t \geq 0$. Here the sum can be estimated by a constant independent of $t \geq 1$ and the integral (computable explicitly) is proportional to $t^{-1/2}$. This establishes the upper bound of the proposition.

To get the lower bound, we may restrict to the class of initial data of the form $u_0(x) = \varphi(x_1) \mathcal{J}_1(x_2)$ with $\varphi \in L_w^2(\mathbb{R})$ (here w is considered as a function on \mathbb{R}). Then it is easy to see from (4.3) that

$$\|e^{-tH_0}\|_{L_w^2(\Omega_0) \rightarrow L^2(\Omega_0)} \geq e^{-E_1 t} \|e^{t\Delta^{\mathbb{R}}}\|_{L_w^2(\mathbb{R}) \rightarrow L^2(\mathbb{R})}$$

for every $t \geq 0$. The lower bound with $t^{-1/4}$ is well known for the heat semigroup of $-\Delta^{\mathbb{R}}$ (or can be easily established by taking $\varphi = w^{-\alpha}$ with any $\alpha > 1/2$ and evaluating the integrals with the kernel p explicitly). \square

Remark 4.1. It is clear from the proof that the bounds hold in less restrictive weighted spaces. Indeed, it is enough to have a corresponding result for the one-dimensional heat semigroup $e^{t\Delta^{\mathbb{R}}}$.

For the following Corollary we recall the definition of the elementary conditional probability. If the measurable subset B satisfies $\mathbb{P}_x(B) > 0$, then $\mathbb{P}_x(A | B) := \mathbb{P}_x(A \cap B) / \mathbb{P}_x(B)$. The concept of conditional probabilities allows to focus on the polynomial decay factors, as the exponential terms cancel each other.

Corollary 4.1. *Let $K = 0$. For every bounded open subset $B \in \Omega_0$ and every $x \in \Omega_0$ there exists a constant $c_{B,x} > 0$ such that*

$$c_{B,x}^{-1} t^{-\frac{1}{2}} \leq \mathbb{P}_x(X_t \in B | \tau_{\Omega_0} > t) \leq c_{B,x} t^{-\frac{1}{2}}$$

for every $t \geq 1$.

Proof. The inequalities follow from Proposition 4.1 and the fact that for every $x = (x_1, x_2) \in \Omega_0$ by independence of the horizontal and vertical components of (X_t) (in the flat case)

$$\lim_{t \rightarrow \infty} e^{E_1 t} \mathbb{P}_x(\tau_{\Omega} > t) = \mathcal{J}_1(x_2) \int_{(-a,a)} \mathcal{J}_1(x_2) dx_2.$$

From the definition of the conditional probability, we see that the exponential cancel and we remain with the polynomial decay as stated in the assertion. \square

As a consequence of this result, we get that conditioned on not hitting the boundary Ω_0 the Brownian particle will escape to infinity.

Proposition 4.1 establishes the decay rate for zero curvature as announced in Table 1.

4.3 The criticality of the transport

Let us now explain what we mean by the vague statement in Table 1 that the transport is ‘critical’ on flat surfaces.

We say that the transport is *critical* if the spectral threshold of H_K is not ‘stable against local attractive perturbations’, *i.e.*,

$$\forall V \in C_0^\infty(\Omega_0), V \neq 0, V \leq 0, \quad \inf \sigma(H_K + V) < \lambda_K. \quad (4.8)$$

Then we also say that H_K is critical. As a consequence of the spectral mapping theorem, we get

$$\|e^{-t(H_K+V)}\|_{L_f^2(\Omega_0) \rightarrow L_f^2(\Omega_0)} = e^{\gamma t} e^{-\lambda_K t}$$

for each time $t \geq 0$, where $\gamma := \lambda_K - \inf \sigma(H_K + V)$ is positive. That is, the criticality leads to an exponential slow-down in the decay of the perturbed semigroup.

Property (4.8) is well known for H_0 and is equivalent to the fact that the first component of $(X_t)_{t \geq 0}$ – a one-dimensional Brownian motion – is recurrent. For some results concerning this connection in a more abstract context we refer to [30].

Proposition 4.2. *H_0 is critical.*

Proof. By the variational characterization of the spectral threshold, it is enough to construct a test function ψ from $H_0^1(\Omega_0)$ such that

$$Q[\psi] := \|\nabla \psi\|^2 - E_1 \|\psi\|^2 - \| |V|^{1/2} \psi \|^2 < 0.$$

For every $n \geq 1$, we define $\psi_n(x) := \varphi_n(x_1) \mathcal{J}_1(x_2)$, with $\varphi_n := w^{-\frac{1}{n}}$, where w is the weight (4.6) (considered as a function on \mathbb{R}). Due to the normalization of \mathcal{J}_1 , we have

$$Q[\psi_n] = \|\dot{\varphi}_n\|_{L^2(\mathbb{R})}^2 - \|v \varphi_n\|_{L^2(\mathbb{R})}^2,$$

where $v(x_1) := \| |V(x_1, \cdot)|^{1/2} \mathcal{J}_1 \|^2_{L^2((-a,a))}$. By hypothesis, $v \in L^1(\mathbb{R})$ and the integral $\|v\|_{L^1(\mathbb{R})}$ is positive. Finally, an explicit calculation yields $\|\dot{\varphi}_n\|_{L^2(\mathbb{R})} \sim n^{-1/4}$. By the dominated convergence theorem, we therefore have

$$Q[\psi_n] \xrightarrow{n \rightarrow \infty} -\|v\|_{L^1(\mathbb{R})}.$$

Consequently, taking n sufficiently large, we can make $Q[\psi_n]$ negative. □

In Section 6 we shall show that the spectrum of H_0 is unstable against purely geometric deformations characterized by positive curvature, too.

5 Asymptotically flat manifolds

We say that the strip Ω is *asymptotically flat* if its Gauss curvature K vanishes at infinity, *i.e.*,

$$\lim_{|x_1| \rightarrow \infty} \operatorname{esssup}_{x_2 \in (-a, a)} |K(x)| = 0. \quad (5.1)$$

In this paper, we are interested in a ‘locally perturbed traveller’ by usually assuming a stronger hypothesis that K is compactly supported, *i.e.*,

$$\operatorname{supp}(K) \cap \Omega_0 \text{ is bounded.} \quad (5.2)$$

It follows from (5.2) that there exists a positive R such that $K(x) = 0$ for all $|x_1| > R$. Then, as a consequence of (2.6),

$$|x_1| > R \implies f(x) = 1. \quad (5.3)$$

Of course, (5.1) trivially holds for the strips satisfying (5.2). Nevertheless, let us state the following result under the more general hypothesis (5.1).

Theorem 5.1. *Assume (2.10) and (5.1). Then*

$$\sigma_{\operatorname{ess}}(H_K) = [E_1, \infty).$$

Proof. The fact that the threshold of the essential spectrum does not descend below the energy E_1 has been proved in [24, Thm. 1] by means of a Neumann bracketing argument. Let us therefore only show that $[E_1, \infty)$ belongs to the essential spectrum of H_K .

Our proof is based on the Weyl criterion adapted to quadratic forms in [4] and applied to quantum waveguides in [27]. By this general characterization of essential spectrum and since the set $[E_1, \infty)$ has no isolated points, it is enough to find for every $\lambda \in [E_1, \infty)$ a sequence $\{\psi_n\}_{n=1}^\infty \subseteq \mathfrak{D}(h_K)$ such that

- (i) $\forall n \in \mathbb{N} \setminus \{0\}, \quad \|\psi_n\|_f = 1,$
- (ii) $\|(H_K - \lambda)\psi_n\|_{[\mathfrak{D}(h_K)]^*} \xrightarrow{n \rightarrow \infty} 0.$

Here $\|\cdot\|_{[\mathfrak{D}(h_K)]^*}$ denotes the norm in the dual space $[\mathfrak{D}(h_K)]^*$ of $\mathfrak{D}(h_K)$. Let $n \in \mathbb{N} \setminus \{0\}$. Given $k \in \mathbb{R}$, we set $\lambda = E_1 + k^2$.

Since Ω is asymptotically flat, a good candidate for the sequence are plane waves in the x_1 -direction modulated by the ground-state eigenfunction \mathcal{J}_1 in the x_2 -direction and ‘localized at infinity’:

$$\psi_n(x) := \varphi_n(x_1) \mathcal{J}_1(x_2) e^{ikx_1}.$$

Here $\varphi_n(x_1) := \varphi(x_1/n - n)^{-1/2}$ with φ being a non-zero C^∞ -smooth function with compact support in the interval $(-1, 1)$. Note that $\operatorname{supp} \varphi_n \subset (n^2 - n, n^2 + n)$. We further assume that φ is normalized to 1 in $L^2(\mathbb{R})$, so that the norm of φ_n is 1 as well.

Clearly, $\psi_n \in H_0^1(\Omega_0) = \mathfrak{D}(h_K)$. To satisfy (i), one can redefine ψ_n by dividing it by its norm $\|\psi_n\|_f$. However, since

$$\|\psi_n\|_f^2 \geq 1 - \frac{\|K\|_\infty a^2}{1 - \|K\|_\infty a^2} > 0$$

due to Lemma 2.1 and the normalizations of φ and \mathcal{J}_1 , it is enough to verify the condition (ii) directly for our unnormalized functions ψ_n .

By the definition of the dual norm, we have

$$\|(H_K - \lambda)\psi_n\|_{[\mathfrak{D}(h_K)]^*} = \sup_{\phi \in H_0^1(\Omega_0) \setminus \{0\}} \frac{|h_K(\phi, \psi_n) - \lambda(\phi, \psi_n)_f|}{\|\phi\|_{\mathfrak{D}(h_K)}}. \quad (5.4)$$

An explicit computation using integrations parts yields

$$\begin{aligned} h_K(\phi, \psi_n) - \lambda(\phi, \psi_n)_f &= (\phi, [-\ddot{\varphi}_n - 2ik\dot{\varphi}_n]\mathcal{J}_1 e^{ikx_1}) \\ &\quad + (\partial_1 \phi, [f^{-1} - 1]\partial_1 \psi_n) - k^2 (\phi, [f - 1]\psi_n) - (\phi, [\partial_2 f]\partial_2 \psi_n). \end{aligned}$$

Using the Schwarz inequality, we estimate the individual terms on the right hand side of the identity as follows

$$\begin{aligned} |(\phi, [-\ddot{\varphi}_n - 2ik\dot{\varphi}_n]\mathcal{J}_1 e^{ikx_1})| &\leq \|\phi\|_{\mathfrak{D}(h_K)} \sqrt{\|\ddot{\varphi}_n\|_{L^2(\mathbb{R})}^2 + 4k^2 \|\dot{\varphi}_n\|_{L^2(\mathbb{R})}^2} \|f^{1/2}\|_\infty, \\ |(\partial_1 \phi, [f^{-1} - 1]\partial_1 \psi_n)| &\leq \|\phi\|_{\mathfrak{D}(h_K)} \|\dot{\varphi}_n\|_{L^2(\mathbb{R})} \operatorname{esssup}_{\operatorname{supp} \varphi_n} (f^{1/2} |f^{-1} - 1|), \\ |(\phi, [f - 1]\psi_n)| &\leq \|\phi\|_{\mathfrak{D}(h_K)} \|\varphi_n\|_{L^2(\mathbb{R})} \operatorname{esssup}_{\operatorname{supp} \varphi_n} (f^{-1/2} |f - 1|), \\ |(\phi, [\partial_2 f]\partial_2 \psi_n)| &\leq \|\phi\|_{\mathfrak{D}(h_K)} \|\varphi_n\|_{L^2(\mathbb{R})} E_1 \operatorname{esssup}_{\operatorname{supp} \varphi_n} (f^{-1/2} |\partial_2 f|). \end{aligned}$$

Hence, the dual norm (5.4) can be bounded from above by a constant multiplied by a sum of terms containing either $\|\dot{\varphi}_n\|_{L^2(\mathbb{R})}$, $\|\ddot{\varphi}_n\|_{L^2(\mathbb{R})}$ or the suprema involving f over the support of φ_n . By hypothesis (5.1), the suprema tend to zero as $n \rightarrow \infty$ due to Lemma 2.1 and (2.8). The remaining terms tend to zero as $n \rightarrow \infty$ because

$$\|\dot{\varphi}_n\|_{L^2(\mathbb{R})} = n^{-1} \|\dot{\varphi}\|_{L^2(\mathbb{R})}, \quad \|\ddot{\varphi}_n\|_{L^2(\mathbb{R})} = n^{-2} \|\ddot{\varphi}\|_{L^2(\mathbb{R})}.$$

□

Theorem (5.1) implies that we always have $\lambda_K \leq E_1$ for asymptotically flat strips. Therefore, as a consequence of (3.6),

$$\|e^{-tH_K}\|_{L_f^2(\Omega_0) \rightarrow L_f^2(\Omega_0)} \geq e^{-E_1 t}$$

for each time $t \geq 0$.

6 Positively curved manifolds

We say that a manifold is *positively curved* if K is non-zero and non-negative (in the sense of a measurable function on the manifold). In this section we give a meaning to the vague statement of Table 1 that the ‘positive curvature is bad for transport’. It is based on the following result, which we adopt from [24].

Theorem 6.1. *Assume (2.10) and $K \in L^1(\Omega_0)$. We have*

$$(\mathcal{J}_1, K\mathcal{J}_1)_f > 0 \quad \implies \quad \inf \sigma(H_K) < E_1.$$

Remark 6.1. Recall that \mathcal{J}_1 is the first transverse eigenfunction introduced in (4.2). Here, not to burden the notation, we denote by the same symbol \mathcal{J}_1 the function $x \mapsto \mathcal{J}_1(x_2)$ on Ω_0 .

Proof. The proof of the theorem is very similar to that of Proposition 4.2. By the variational characterization of the spectral threshold of H_K , it is enough to construct a test function ψ from $H_0^1(\Omega_0)$ such that

$$Q_K[\psi] := h_K[\psi] - E_1 \|\psi\|_f^2 < 0.$$

Using the same sequence of functions $\psi_n(x) = \varphi_n(x_1)\mathcal{J}_1(x_2)$ as in the proof of Proposition 4.2, we arrive at

$$Q_K[\psi_n] = (\partial_1 \psi_n, f^{-1} \partial_1 \psi_n) - \frac{1}{2} (\psi_n, K \psi_n)_f. \quad (6.1)$$

Here the first (positive) integral on the right hand side vanishes as $n \rightarrow \infty$ because

$$(\partial_1 \psi_n, f^{-1} \partial_1 \psi_n) \leq \frac{1 - \|K\|_\infty a^2}{1 - 2\|K\|_\infty a^2} \|\dot{\varphi}_n\|_{L^2(\mathbb{R})}^2,$$

due to Lemma 2.1 and the normalization of \mathcal{J}_1 , and $\|\dot{\varphi}_n\|_{L^2(\mathbb{R})} \sim n^{-1/4}$. Using, at the same time, the dominated convergence theorem in the second integral on the right hand side of (6.1), we finally get

$$Q_K[\psi_n] \xrightarrow{n \rightarrow \infty} -\frac{1}{2} (\mathcal{J}_1, K \mathcal{J}_1)_f.$$

Since the limit is negative by hypothesis, we can make $Q[\psi_n]$ negative by taking n sufficiently large. \square

Remark 6.2. The integrability of K is just a technical assumption in Theorem 6.1. It is only important to give a meaning to the integral $(\mathcal{J}_1, K \mathcal{J}_1)_f$, the value $+\infty$ being admissible in principle. For instance, it is enough to assume that K is non-trivial and non-negative on Ω_0 for the present proof to work.

Combining Theorem 6.1 with Theorem 5.1, we get that H_K possesses at least one discrete eigenvalue below the essential spectrum under the hypotheses. In view of the criticality notion introduced in Section 4.3, the result of Theorem 6.1 can be also interpreted in the sense that H_0 is not stable against geometric perturbations characterized by the presence of positive curvature.

In any case, regardless of whether the spectral threshold of H_K represents an eigenvalue or the bottom of the essential spectrum, Theorem 6.1 implies that the gap $\gamma := E_1 - \lambda_K$ is always positive for positively curved strips. If K vanishes at infinity, then the bottom of the spectrum has to be an isolated eigenvalue. Therefore, as a consequence of (3.6) and [37], we conclude with

Corollary 6.1. *Assume (2.10), $K \in L^1(\Omega_0)$ and $(\mathcal{J}_1, K \mathcal{J}_1)_f > 0$. Then*

$$\|e^{-tH_K}\|_{L_f^2(\Omega_0) \rightarrow L_f^2(\Omega_0)} = e^{\gamma t} e^{-E_1 t}$$

for each time $t \geq 0$, where γ is positive. Moreover, if additionally (5.1) is satisfied then there exists a unique non-negative normalized $\phi_0 \in L_f^2(\Omega_0)$ such that for every bounded measurable set $B \subset \Omega_0$ and every $x \in \Omega_0$

$$\lim_{t \rightarrow \infty} e^{-(\gamma - E_1)t} \mathbb{P}_x(X_t \in B, \tau_{\Omega_0} > t) = \phi_0(x) \int_B \phi_0(y) f(y) dy.$$

That is, the presence of positive curvature clearly slows down the decay of the heat semigroup, even without the need to work with the weighted space $L_{wf}^2(\Omega_0)$. A Brownian traveller should avoid ‘mountains’ satisfying $(\mathcal{J}_1, K\mathcal{J}_1)_f > 0$, if he/she wants to make sure that he/she is able to reach his goal early and wants to avoid spending too much time in a given bounded region.

The following Corollary (again a rather direct consequence of Theorems 6.1 and 5.1) shows that in contrast to the flat case the Brownian traveller – conditioned on not-hitting the boundary $\partial\Omega_0$ – might not have been able to have left a bounded region forever.

Corollary 6.2. *Assume that (5.1) and the conditions of Theorem 6.1 are satisfied. Then for almost every $x \in \Omega_0$ we have*

$$\lim_{t \rightarrow \infty} \mathbb{P}_x(X_t \in \cdot \mid \tau_{\Omega_0} > t) = \frac{\int_{\Omega_0} \phi_0(y) f(y) dy}{\int_{\Omega_0} \phi_0(y) f(y) dy},$$

where the convergence is with respect to the total variation distance.

Proof. By definition of the total variation distance we have to prove that

$$\lim_{t \rightarrow \infty} \sup_{B \subset \Omega_0} \left| \mathbb{P}_x(X_t \in B \mid \tau_{\Omega_0} > t) - \frac{\int_B \phi_0(y) f(y) dy}{\int_{\Omega_0} \phi_0(y) f(y) dy} \right| = 0.$$

Observe that we do not assume that the sets B are bounded and that the assertions of Corollary 6.1 do not suffice to prove the desired assertion.

According to general spectral theory we know, that the eigenfunction $\phi_0 \in L_f^2(\Omega_0)$ does not change sign and that the eigenspace is one-dimensional. In the first step we show that ϕ_0 actually also belongs to $L_f^1(\Omega_0)$, with the notation $L_f^p(\Omega_0) := L^p(\Omega_0, f(x) dx)$. This will allow us to interpret the ground state as a probability distribution. Of course, many results concerning the decay properties are known, but we have not been able to find a reference covering our setting. Observe first that due to the probabilistic interpretation the semigroup $(e^{-tH_K})_{t \geq 0}$ in $L_f^2(\Omega_0)$ gives rise to a consistent strongly continuous semigroups $(T_t^p)_{t \geq 0}$ in $L_f^p(\Omega_0)$ for $1 \leq p < \infty$. Moreover, due to the Gaussian bound from Proposition 3.1, these semigroups are analytic with angle $\pi/2$. Let the generators be denoted by H_K^p . Due to the consistence of the semigroups, by taking Laplace transforms of the semigroups, we conclude that the resolvents $F^p(z) := (H_K^p - z)^{-1}$ ($z \in \rho(H_K^p) := \mathbb{C} \setminus \sigma(H_K^p)$) are as well consistent in the sense that for every $z \in \rho(H_K^p) \cap \rho(H_K^q)$

$$F^p(z) \upharpoonright L_f^p(\Omega_0) \cap L_f^q(\Omega_0) = F^q(z) \upharpoonright L_f^p(\Omega_0) \cap L_f^q(\Omega_0).$$

Since according to Theorem 5 in [3] we have $\sigma(H_K^p) = \sigma(H_K^2) (= \sigma(H_K))$ for every $1 \leq p < \infty$ and since λ_K is an isolated eigenvalue for H_K^2 , we conclude by Corollary 1.4 in [17] that λ_K is an isolated point of $\sigma(H_K^1)$ and that the eigenvector ϕ_0 of H_K^2 is also an eigenvector of H_K^1 , i.e., in particular, $\phi_0 \in L_f^1(\Omega_0)$.

Observe now that $(X_t)_{\tau_{\Omega_0} \geq t \geq 0}$ is λ_K -recurrent in the sense of [44] and we also conclude that the measure $\pi(dx) = \phi_0(y) f(y) dy$ is finite and due to reversibility with respect to the measure $f(x) dx$ satisfies

$$\mathbb{P}_\pi(X_t \in A, \tau_{\Omega_0} > t) = e^{-\lambda_K t} \int_A \phi_0(x) f(x) dx$$

for every measurable set $A \subset \Omega_0$. As $\phi \in L^2_f(\Omega_0)$ we conclude that (X_t) is λ_K -positive recurrent in the sense of [44] (product-critical in the sense of [36]). Applying Theorem 7 in [44], we are thus able to conclude the assertion of the Corollary. More precisely, formula (5.9) in [44] shows that for almost all $x \in \Omega_0$

$$\lim_{t \rightarrow \infty} \sup_{B \subset \Omega_0} \left| e^{\lambda_K t} \mathbb{P}_x(X_t \in B, \tau_{\Omega_0} > t) - \phi_0(x) \int_B \phi_0(y) f(y) dy \right| = 0. \quad (6.2)$$

Therefore we have for almost all $x \in \Omega_0$

$$\begin{aligned} & \sup_{B \subset \Omega_0} \left| \mathbb{P}_x(X_t \in B \mid \tau_{\Omega_0} > t) - \frac{\int_B \phi_0(z) f(z) dz}{\int_{\Omega_0} \phi_0(y) f(y) dy} \right| \\ & \leq (e^{\lambda_K t} \mathbb{P}_x(\tau_{\Omega_0} > t))^{-1} \sup_{B \subset \Omega_0} \left| e^{\lambda_K t} \mathbb{P}_x(X_t \in B) - \phi_0(x) \int_B \phi_0(y) f(y) dy \right| \\ & \quad + \sup_{B \subset \Omega_0} \left| \frac{\phi_0(x) \int_B \phi_0(y) f(y) dy}{e^{\lambda_K t} \mathbb{P}_x(\tau_{\Omega_0} > t)} - \frac{\int_B \phi_0(y) f(y) dy}{\int_{\Omega_0} \phi_0(y) f(y) dy} \right| \\ & = (e^{\lambda_K t} \mathbb{P}_x(\tau_{\Omega_0} > t))^{-1} \sup_{B \subset \Omega_0} \left| e^{\lambda_K t} \mathbb{P}_x(X_t \in B) - \phi_0(x) \int_B \phi_0(y) f(y) dy \right| \\ & \quad + \left(\sup_{B \subset \Omega_0} \int_B \phi_0(z) dz \right) \left| \frac{\phi_0(x)}{e^{\lambda_K t} \mathbb{P}_x(\tau_{\Omega_0} > t)} - \frac{1}{\int_{\Omega_0} \phi_0(y) f(y) dy} \right|. \end{aligned}$$

Two applications of (6.2) complete the proof. \square

7 Negatively curved manifolds

In analogy with positively curved manifolds, we say that a manifold is *negatively curved* if K is non-zero and non-positive. In this section, on the contrary, we show that the presence of negative curvature improves the decay of the heat semigroup, supporting in this way the vague statement of Table 1 that the ‘negative curvature is good for transport’. First, however, we have to explain why the negative sign of curvature is much more delicate for the study of large time properties of (1.2).

Recall that the positivity of the curvature K pushes the spectrum below E_1 (cf Theorem 6.1). The objective of this subsection is to show that the effect of negative curvature is rather opposite: it ‘has the tendency’ to push the spectrum above E_1 . This effect is more subtle because $[E_1, \infty)$ belongs to the spectrum of H_K , irrespectively of the sign of the curvature, as long as the curvature vanishes at infinity (cf Theorem 5.1).

The way how to understand this ‘repulsive tendency’ is to replace the Poincaré-type inequality requirement $H_K - E_1 \geq \text{const} > 0$ (which is false for the asymptotically flat manifolds) by a weaker, *Hardy-type inequality*:

$$H_K - E_1 \geq \rho > 0. \quad (7.1)$$

Here $\rho : \Omega_0 \rightarrow (0, \infty)$ is assumed to be merely a *positive function* (necessarily vanishing at the infinity of Ω_0 for the asymptotically flat manifolds).

By Theorem 6.1, (7.1) is false for positively curved manifolds. It is also violated for flat manifolds because of the criticality result of Proposition 4.2. In this subsection, we show that (7.1) typically holds for negatively curved manifolds.

7.1 Hardy-type inequality and the large time behaviour

For completeness we first sketch an abstract elementary argument from [22], relating the Hardy inequality to low energy properties of the Hamiltonian and the large time behaviour of the semigroup. If the semigroup is associated to a stochastic process, then the validity of a Hardy-type inequality is related to the concept of R -transience of the stochastic process.

Assume that there exists a positive function ρ , with a locally bounded inverse ρ^{-1} , such that the inequality (7.1) holds true for the selfadjoint non-negative operator $L_K := H_K - E_1$. Then according to Theorem 8.31 in [47] we conclude that for all $\lambda < 0$ and every $h \in L^2_f(\Omega_0)$ we have

$$(h, (L_K - \lambda)^{-1}h)_f \leq (h, (M_\rho - \lambda)^{-1}h)_f, \quad (7.2)$$

where M_ρ denotes the maximal multiplication operator acting via multiplication with the function ρ . If h satisfies $(h, \rho^{-1}h)_f < \infty$, then (7.2) implies

$$\forall \lambda < 0, \quad \int_{(E_1, \infty)} (\nu - \lambda)^{-1} d\|E_\nu^{L_K} h\|_f^2 \leq (h, \rho^{-1}h)_f < \infty, \quad (7.3)$$

where $(E_\nu^{L_K})_\nu$ denotes the spectral resolution of L_K . Using monotone convergence, we get for all h with $(h, \rho^{-1}h)_f < \infty$ (in particular for all continuous h with compact support in Ω_0)

$$\int_0^\infty (h, e^{-t(H_K - E_1)}h)_f dt < \infty. \quad (7.4)$$

Observe that (7.4) – which in the probabilistic literature such as [44], [46] and [45] might be called E_1 -transience – does not hold in the case of positively curved and flat manifolds.

Property (7.3) is related to the low energy behaviour of the spectral measure $E^{L_K}(\cdot)$ in the sense that it implies that for all $r \in (0, 1]$ and $-1 \leq \lambda < 0$

$$\|E^{L_K}((0, r))h\|_f^2 = \int_0^r d\|E_\nu^{L_K} h\|_f^2 \leq \int_0^r \frac{r - \lambda}{\nu - \lambda} d\|E_\nu^{L_K} h\|_f^2. \quad (7.5)$$

where we used that $\frac{r - \lambda}{\nu - \lambda} \geq 1$ for $\nu \in (0, r)$ and negative λ . Sending λ to 0 and using (7.3), we conclude that there is $C > 0$ such that for h with $(h, \rho^{-1}h)_f \leq 1$ and $r \in (0, 1)$

$$\|E^{L_K}((0, r))h\|_f^2 \leq C r.$$

This insight can easily be translated into an assertion concerning the large time behaviour.

Proposition 7.1. *Assume that $H_K - E_1$ satisfies the Hardy-type inequality (7.1) with a positive function ρ satisfying $\rho^{-1} \in L^\infty_{\text{loc}}(\Omega_0)$. Then*

$$\sup_{(h, \rho^{-1}h)_f < 1} \|e^{-t(H_K - E_1)}h\|_f^2 \leq \frac{1}{t} (1/2 + 2e^{-2}).$$

Proof. For the proof we again set $L_K := H_K - E_1$ and denote by μ_h the spectral measure corresponding to L_K and h . Via the spectral theorem, integration by

parts and (7.5), we obtain

$$\begin{aligned}
\|e^{-tL_K} h\|_f^2 &= \int_0^\infty e^{-2\nu t} d\mu_h(\nu) = 2t \int_0^1 e^{-2\nu t} \mu_h(\nu) d\nu + 2t \int_1^\infty e^{-2\nu t} \mu_h(\nu) d\nu \\
&\leq 2t \int_0^1 e^{-2\nu t} \nu d\mu_h + 2t \int_1^\infty e^{-2\nu t} \mu_h(\nu) d\nu \\
&\leq 2t \int_0^\infty e^{-2\nu t} \nu d\nu + t \int_1^\infty e^{-2\nu t} d\nu \leq \frac{\Gamma(1)}{2t} + 2te^{-2t} \\
&= \frac{1}{t} (1/2 + 2t^2 e^{-2t}).
\end{aligned}$$

Observing that $\max_{t>0} (2t^2 e^{-2t}) = 2e^{-2}$ yields the desired assertion. \square

Observe again that, under weak conditions on the Hardy weight ρ , Proposition 7.1 already gives an accelerated decay rate when compared with the one in the flat case given in Proposition 4.1.

7.2 The Hardy inequality for negatively curved manifolds

In this subsection, we show that (7.1) typically holds for negatively curved manifolds.

One way how to establish (7.1) is to generalize the method of [25]. It works as follows:

1. *Transverse ground-state estimate.* Recalling the structure of our operator (3.1), we clearly have

$$H_K - E_1 \geq -f^{-1} \partial_1 f^{-1} \partial_1 + \mu_K \quad (7.6)$$

in the form sense on $L_f^2(\Omega_0)$, where $x \mapsto \mu_K(x_1)$ denotes the lowest eigenvalue of the one-dimensional shifted ‘transverse’ operator $-f^{-1} \partial_2 f \partial_2 - E_1$ on the Hilbert space $L^2((-a, a), f(x_1, x_2) dx_2)$, subject to Dirichlet boundary conditions, with x_1 being considered as a parameter in the one-dimensional eigenvalue problem. More specifically, we have

$$\mu_K(x_1) = \inf_{\varphi \in H_0^1((-a, a)) \setminus \{0\}} \frac{\int_{-a}^a |\dot{\varphi}(x_2)|^2 f(x_1, x_2) dx_2}{\int_{-a}^a |\varphi(x_2)|^2 f(x_1, x_2) dx_2} - E_1. \quad (7.7)$$

With an abuse of notation, we denote by the same symbol μ_K both the function on \mathbb{R} and its natural extension $x \mapsto \mu_K(x_1)$ to Ω_0 .

2. *Longitudinal Hardy-type estimate.* Now we regard the right hand side of (7.6) as a one-dimensional Schrödinger-type operator on the Hilbert space $L^2((-a, a), f(x_1, x_2) dx_1)$, with x_2 being considered as a parameter and μ_K playing the role of potential. We assume that each of the x_2 -dependent family of operators satisfies a Hardy-type inequality, so that

$$-f^{-1} \partial_1 f^{-1} \partial_1 + \mu_K \geq \rho_K > 0 \quad (7.8)$$

in the form sense on $L_f^2(\Omega_0)$, with some positive function $\rho_K : \Omega_0 \rightarrow (0, \infty)$. Then (7.1) holds as a consequence of (7.8) and (7.6).

In this way, we have reduced the problem to ensuring the existence of one-dimensional Hardy-type inequalities (7.8). However, the criticality of one-dimensional Schrödinger operators is well studied, at least if $f = 1$. We present two sufficient conditions which guarantee the validity of (7.8) and confirm thus that (7.1) typically holds for negatively curved manifolds.

7.2.1 Positivity of the ground-state estimates

Since the kinetic part of the Schrödinger-type operator on the left hand side of (7.8) is a non-negative operator, we get a trivial estimate

$$-f^{-1}\partial_1 f^{-1}\partial_1 + \mu_K \geq \mu_K \quad (7.9)$$

in the form sense on $L_f^2(\Omega_0)$. As a consequence of (7.6), $H_K - E_1 \geq \mu_K$.

This represents a *local* Hardy-type inequality provided that μ_K is non-zero and non-negative. By ‘local’ we mean that the function μ_K is compactly supported for manifolds with compactly supported curvature K , which is a typical hypothesis of the present paper. Hence it does not fit to the initial definition (7.1), which can be called *global* Hardy-type inequality. However, it is known that local Hardy-type inequalities imply global ones.

Theorem 7.1 (Hardy inequality for non-negative μ_K). *Assume (2.10). If μ_K is non-zero and non-negative in some bounded open subinterval $J \subset \mathbb{R}$, then there exists a positive constant c_K , depending on a and properties of K , such that*

$$-f^{-1}\partial_1 f^{-1}\partial_1 + \mu_K \geq \frac{c_K}{1 + \delta^2} \quad (7.10)$$

in the form sense on $L_f^2(\Omega_0)$. Here $\delta(x) := |x_1 - x_1^0|$, with x_1^0 being the mid-point of J . As a consequence of (7.6), the Hardy-type inequality (7.1) holds.

Proof. The proof follows by a modification of the proof of [25, Thm. 3.1]. For the clarity of the exposition, we divide it into several steps.

1. *A consequence of the classical Hardy inequality.* The main ingredient in the proof is the following Hardy-type inequality for a Schrödinger operator in the strip Ω_0 with a characteristic-function potential:

$$\|(1 + \delta^2)^{-1/2}\psi\|^2 \leq 16 \|\partial_1 \psi\|^2 + (2 + 64/|J|^2) \|\chi_J \psi\|^2 \quad (7.11)$$

for every $\psi \in H^1(\Omega_0)$. Here J is any bounded open subinterval of \mathbb{R} and χ_J denotes the characteristic function of the set $J \times (-a, a) \subset \Omega_0$. This inequality can be established quite easily (cf [6, Sec. 3.3]) by means of Fubini’s theorem and the classical one-dimensional Hardy inequality $\int_0^b s^{-2} |\varphi(s)|^2 ds \leq 4 \int_0^b |\dot{\varphi}(s)|^2 ds$ valid for any $\varphi \in H^1((0, b))$, $b > 0$, satisfying $\varphi(0) = 0$.

Using Lemma 2.1, (7.11) can be cast into the form

$$\|f^{-1}\partial_1 \psi\|_f^2 + \|\mu_K^{1/2} \psi\|_f^2 \geq c \|(1 + \delta^2)^{-1/2}\psi\|_f^2 - C \|\chi_J \psi\|_f^2, \quad (7.12)$$

where the constants are given by

$$c := \frac{1 - \|K\|_\infty a^2}{16}, \quad C := \left(\frac{1}{8} + \frac{4}{|J|^2} \right) \left[1 - \left(\frac{\|K\|_\infty a^2}{1 - \|K\|_\infty a^2} \right)^2 \right]^{-1}.$$

2. *A Poincaré-type inequality in a bounded strip.* For every $\psi \in H^1(\Omega_0)$, we have

$$\begin{aligned} \|f^{-1}\partial_1\psi\|_f^2 + \|\mu_K^{1/2}\psi\|_f^2 &\geq \|\chi_J f^{-1}\partial_1\psi\|_f^2 + \|\chi_J \mu_K^{1/2}\psi\|_f^2 \\ &\geq \lambda_J \|\chi_J \psi\|_f^2, \end{aligned} \quad (7.13)$$

where λ_J denoted the lowest eigenvalue of the operator $-f^{-1}\partial_1 f^{-1}\partial_1 + \mu_K$ on $L_f^2(J \times (-a, a))$, subject to Neumann-type (*i.e.* no in the form setting) boundary conditions at $(\partial J) \times (-a, a)$. We claim that λ_J can be bounded from below by a positive constant which depends exclusively on properties of μ_K . Indeed, assume $\lambda_J = 0$. By the variational characterization of λ_J , it follows that

$$\|\chi_J f^{-1}\partial_1\psi_J\|_f^2 = 0 \quad \text{and} \quad \|\chi_J \mu_K^{1/2}\psi_J\|_f^2 = 0,$$

where $\psi_J \in H^1(J \times (-a, a))$ is an eigenfunction corresponding to λ_J . Recalling Lemma 2.1, we conclude that $\|\mu_K\|_{L^1(J)} = 0$, which contradicts the hypothesis that μ_K is non-trivial on J .

3. *Some interpolation.* Combining (7.12) with (7.13), we eventually arrive at

$$\|f^{-1}\partial_1\psi\|_f^2 + \|\mu_K^{1/2}\psi\|_f^2 \geq c\epsilon \|(1 + \delta^2)^{-1/2}\psi\|_f^2 + [(1 - \epsilon)\lambda_J - C\epsilon] \|\chi_J \psi\|_f^2$$

for every $\psi \in H^1(\Omega_0)$ and any $\epsilon \in (0, 1)$. Choosing ϵ in such a way that the term with the square brackets vanishes, we get the Hardy-type inequality of the theorem with $c_K := c\lambda_J/(\lambda_J + C)$. \square

7.2.2 On the positivity of the ground-state eigenvalue

Since the fundamental hypothesis of Theorem 7.1 is the non-negativity of μ_K , let us comment on its relation to the non-positivity of K .

We claim that the function μ_K is typically positive for negatively curved manifolds. Indeed, for any fix $x_1 \in \mathbb{R}$, let us make the change of test function $\phi := \sqrt{f(x_1, \cdot)} \varphi$ in (7.7). Integrating by parts and using (2.6), one easily arrives at

$$\mu_K(x_1) = \inf_{\phi \in H_0^1((-a, a)) \setminus \{0\}} \frac{\int_{-a}^a \left(|\dot{\phi}(x_2)|^2 - E_1 |\phi(x_2)|^2 + V(x) |\phi(x_2)|^2 \right) dx_2}{\int_{-a}^a |\phi(x_2)|^2 dx_2} \quad (7.14)$$

with

$$V := -\frac{1}{2}K + \frac{1}{4} \left(\frac{\partial_2 f}{f} \right)^2. \quad (7.15)$$

By the Poincaré inequality for the Dirichlet Laplacian in $L^2((-a, a))$, we therefore get

$$\mu_K(x_1) \geq \operatorname{ess\,inf}_{x_2 \in (-a, a)} V(x_1, x_2). \quad (7.16)$$

Let us assume for a moment that K is continuous. Then, for every $x_1 \in \mathbb{R}$ fixed, it follows from (2.6) that

$$\lim_{a \rightarrow 0} V(x) = -\frac{1}{2}K(x_1, 0).$$

Hence, if $K(x_1, x_2)$ is negative for every $x_2 \in (-a, a)$ and x_1 from a compact interval J , there exists a positive half-width a such that $\mu_K(x_1)$ is positive for every $x_1 \in J$. For merely bounded curvature K , we replace the pointwise non-positivity requirement on the curve Γ by the hypothesis that the function

$$k(x_1) := \lim_{a \rightarrow 0} \operatorname{ess\,inf}_{x_2 \in (-a, a)} K(x_1, x_2) \quad (7.17)$$

is non-zero and non-positive.

It is less obvious how to get uniform lower bounds, *i.e.* to ensure that, for a given a , $\mu_K(x_1)$ is non-negative for almost every $x_1 \in \mathbb{R}$. An example of manifolds for which the uniform non-negativity is possible to check is given by strips on ruled surfaces studied in [25].

Example 7.1 (Ruled strips). Let Γ be a straight line in \mathbb{R}^3 ; without loss of generality, we may assume that $\Gamma(x_1) = (x_1, 0, 0)$. Given a C^1 -smooth function $\theta : \mathbb{R} \rightarrow \mathbb{R}$, let us define $\mathcal{L}(x) := (x_1, x_2 \cos \theta(x_1), x_2 \sin \theta(x_1))$. The image (2.3) is a ruled surface, composed of segments of length $2a$ translated and rotated along Γ . It is straightforward to check that the corresponding metric G admits the block form (2.5) with the explicit formulae

$$f(x) = \sqrt{1 + \dot{\theta}(x_1)^2 x_2^2}, \quad K(x) = -\frac{\dot{\theta}(x_1)^2}{f(x)^4}.$$

The *ad hoc* defined mapping \mathcal{L} represents an explicit parametrization of Ω via the exponential map (2.2). The hypothesis (2.10) is satisfied for every a provided that we assume that $\dot{\theta}$ is bounded. The strip Ω is asymptotically flat if $\dot{\theta}(x_1)$ tends to zero as $|x_1| \rightarrow \infty$. Finally, an explicit calculation yields

$$V(x) = \frac{\dot{\theta}(x_1)^2 [2 - \dot{\theta}(x_1)^2 x_2^2]}{4 f(x)^4}. \quad (7.18)$$

It follows that V is non-zero and non-negative provided that $\dot{\theta}$ is non-zero and the half-width a is so small that $\|\dot{\theta}\|_\infty a < \sqrt{2}$. Consequently, under the same assumptions about a and $\dot{\theta}$, the quantity μ_K is non-zero and non-negative, too. We refer to [25] for more geometric and spectral properties of the ruled strips.

7.2.3 Thin strips

The second sufficient condition which guarantees the validity of (7.8) is based on the ideas of the previous subsection.

Theorem 7.2 (Hardy inequality for thin strips). *Assume (2.10) and (5.2). Let the function k defined in (7.17) be non-zero and non-positive. Then there exists a positive number a_0 , depending on properties of K , such that*

$$-f^{-1} \partial_1 f^{-1} \partial_1 + \mu_K \geq \frac{\tilde{c}_K}{1 + x_1^2} \quad (7.19)$$

holds in the form sense on $L_f^2(\Omega_0)$ for all $a \leq a_0$ with some constant \tilde{c}_K depending on properties of K . As a consequence of (7.6), the Hardy-type inequality (7.1) holds for all $a \leq a_0$.

Proof. In view of (7.16), Lemma 2.1, (2.8) and (5.3), it is easy to show that

$$\mu_K(x_1) \geq -\frac{1}{2}k - C(\|K\|_\infty a^2) \chi_{[-R,R]}(x_1),$$

for almost every $x_1 \in \mathbb{R}$, where

$$C(\xi) := \frac{1}{4}\xi^2 \left(1 + \frac{\xi^2}{1 - \xi^2}\right)^2 \left(1 - \frac{\xi^2}{1 - \xi^2}\right)^{-2}.$$

Hence, $\mu_K \rightarrow \mu_K^0 := -\frac{1}{2}k$ as $a \rightarrow 0$. For every $\psi \in H^1(\Omega_0)$, we write

$$\|f^{-1}\partial_1\psi\|_f^2 + (\psi, \mu_K\psi)_f = \|f^{-1}\partial_1\psi\|_f^2 + (\psi, \mu_K^0\psi)_f + (\psi, [\mu_K - \mu_K^0]\psi)_f.$$

Applying Theorem 7.1 to the first two terms on the right hand side of this identity, we get

$$\begin{aligned} & \|f^{-1}\partial_1\psi\|_f^2 + (\psi, \mu_K\psi)_f \\ & \geq \int_{\Omega_0} \left[\frac{c_K}{1+x_1^2} - C(\|K\|_\infty a^2) \chi_{[-R,R]}(x_1) \right] |\psi(x)|^2 f(x) dx. \end{aligned}$$

It is important to notice that c_K can be bounded from below by a positive constant independent of a (cf proof of Theorem 7.1). On the other hand, $C(\|K\|_\infty a^2)$ tends to zero as $a \rightarrow 0$. Then the result follows by estimating the characteristic function by $(1+x_1^2)^{-1}$ multiplied by a constant smaller than c_K for all sufficiently small a . \square

Remark 7.1. The positive function ρ on the right hand side of (7.1) can in principle vanish on the boundary of $\partial\Omega_0$. The objective of this remark is to show that, if (7.1) holds, with an arbitrary positive function ρ , there is also an inequality of the type (7.8) with the right hand side which is independent of the ‘transverse’ variable x_2 . This can be seen as follows. Assume (7.8) and (5.2). For any $\psi \in H_0^1(\Omega_0)$ and $\epsilon \in (0, 1)$, we write

$$\begin{aligned} h_K[\psi] - E_1 \|\psi\|_f^2 &= \epsilon (h_K[\psi] - E_1 \|\psi\|_f^2) + (1 - \epsilon)(h_K[\psi] - E_1 \|\psi\|_f^2) \\ &\geq \epsilon (\|\partial_2\psi\|_f^2 - E_1 \|\psi\|_f^2) + (1 - \epsilon) \|\rho^{1/2}\psi\|_f^2 \\ &= \epsilon (\|\partial_2\phi\|^2 - E_1 \|\phi\|^2 + (\phi, V\phi)) + (1 - \epsilon) \|\rho^{1/2}\phi\|^2 \\ &\geq \epsilon (\psi, [V + \lambda_\epsilon]\psi). \end{aligned}$$

Here the last equality follows by the change of test function $\phi := \sqrt{f}\psi$, as in Section 7.2.2, and $x \mapsto \lambda_\epsilon(x_1)$ denotes the lowest eigenvalue of the one-dimensional operator $-\partial_2^2 - E_1 + (1 - \epsilon)\epsilon^{-1}\rho(x_1, \cdot)$ on $L^2((-a, a))$, subject to Dirichlet boundary conditions, with x_1 considered as a parameter. More specifically, we have

$$\lambda_\epsilon(x_1) := \inf_{\varphi \in H_0^1((-a, a)) \setminus \{0\}} \frac{\int_{-a}^a (|\dot{\varphi}(x_2)|^2 - E_1 |\varphi(x_2)|^2 + \frac{1-\epsilon}{\epsilon} \rho(x_1, x_2) |\varphi(x_2)|^2) dx_2}{\int_{-a}^a |\phi(x_2)|^2 dx_2}.$$

Since K has bounded support, it is also true for V , cf (5.3). On the other hand, since $\rho(x)$ is positive for almost every $x \in \Omega_0$, $\lambda_\epsilon(x_1)$ is positive for almost every $x_1 \in \mathbb{R}$. Furthermore, $\lambda_\epsilon(x_1)$ tends to infinity as $\epsilon \rightarrow 0$ for almost every $x_1 \in \mathbb{R}$. Consequently, for sufficiently small ϵ , $V + \lambda_\epsilon$ can be bounded from below by a positive function which depends on x_1 only.

Finally, let us emphasize that Theorem 7.2 covers a very general class of manifolds, not necessarily negatively curved. It is only important that the manifold is ‘negatively curved in the vicinity of the reference curve’ Γ , cf (7.17).

7.3 The fine decay rate

As in the flat case in Proposition 4.1, we again restrict the class of initial data to the weighted spaces of $L_{wf}^2(\Omega_0) \subset L_f^2(\Omega_0)$ and consider the following (polynomial) *decay rate* quantity:

$$\Gamma_K := \sup \left\{ \Gamma \in \mathbb{R} \mid \exists C_\Gamma > 0, \forall t \geq 0, \right. \\ \left. \|e^{-(H_K - E_1)t}\|_{L_{wf}^2(\Omega_0) \rightarrow L_f^2(\Omega_0)} \leq C_\Gamma (1+t)^{-\Gamma} \right\}. \quad (7.20)$$

Sections 7.1 and 7.2 already imply that the heat semigroup decays faster than in the flat case provided that the Hardy-type inequality (7.1) holds. It follows from Proposition 4.1 that we have $\Gamma_0 = 1/4$ (i.e. for $K = 0$), whereas Proposition 7.1 gives $\Gamma_K \geq 1/2$ if (7.1) is satisfied.

The abstract arguments leading to Proposition 7.1 do not give the precise additional polynomial decay rate. The objective of the following subsections is to show that Γ_K is in fact three times bigger whenever the curvature K is non-zero and non-positive.

In probabilistic terms we are interested in the precise decay exponent

$$\gamma_K(x, B) := \sup \left\{ \gamma \in \mathbb{R} \mid \exists \tilde{C}_\gamma > 0, \forall t \geq 0, \right. \\ \left. \mathbb{P}_x(X_t \in B, \tau_{\Omega_0} > t) \leq C_\gamma (1+t)^{-\gamma} \right\}. \quad (7.21)$$

where $x \in \Omega_0, B \subset\subset \Omega_0$. Again we find that the non-zero and non-positive situation differs from the straight manifold by a factor 3. This is the meaning of the last item in Table 1.

In view of (7.20), it is more convenient to study the shifted heat equation

$$\begin{cases} \partial_t u + H_K u - E_1 u = 0 & \text{in } \Omega_0 \times (0, \infty), \\ u = u_0 & \text{on } \Omega_0 \times \{0\}, \end{cases} \quad (7.22)$$

in the functional setting on $L_f^2(\Omega_0)$ as explained in Section 3.2. Indeed, (7.22) is obtained from (1.2) by the replacement $u(t) \mapsto e^{-E_1 t} u(t)$, with help of the Fermi coordinates.

7.4 The self-similarity transformation

Our method to study the asymptotic behaviour of the heat equation (1.2) in the presence of curvature is to adapt the technique of self-similar solutions used in the case of the heat equation in the whole Euclidean space by Escobedo and Kavian [7] to the present problem. We closely follow the approach of the recent papers [28, 29], where the technique is applied to twisted waveguides in three and two dimensions, respectively.

We perform the self-similarity transformation in the first (longitudinal) space variable only, while keeping the other (transverse) space variable unchanged.

More precisely, given $s \in (0, \infty)$, let us consider the change of function defined by

$$(U_s \psi)(y) := e^{s/4} \psi(e^{s/2} y_1, y_2).$$

It defines a unitary transformation from $L_f^2(\Omega_0)$ to $L_{f_s}^2(\Omega_0)$, where

$$f_s(y) := f(e^{s/2} y_1, y_2). \quad (7.23)$$

Now we associate to every solution $u \in L_{\text{loc}}^2((0, \infty), dt; L_f^2(\Omega_0))$ of (1.2) a ‘self-similar’ solution $\tilde{u}(s) := U_s[u(e^s - 1)]$ in a new s -time weighted space $L_{\text{loc}}^2((0, \infty), e^s ds; L_{f_s}^2(\Omega_0))$. We have

$$\tilde{u}(y_1, y_2, s) = e^{s/4} u(e^{s/2} y_1, y_2, e^s - 1) \quad (7.24)$$

and the inverse change of variables is given by

$$u(x_1, x_2, t) = (t+1)^{-1/4} \tilde{u}((t+1)^{-1/2} x_1, x_2, \log(t+1)).$$

Note that the original space-time variables (x, t) are related to the ‘self-similar’ space-time variables (y, s) via the relations

$$\begin{aligned} (x_1, x_2, t) &= (e^{s/2} y_1, y_2, e^s - 1), \\ (y_1, y_2, s) &= ((t+1)^{-1/2} x_1, x_2, \log(t+1)). \end{aligned} \quad (7.25)$$

Hereafter we consistently use the notation for respective variables to distinguish the two space-times.

It is easy to check that this change of variables transfers the weak formulation of (7.22) to the evolution problem

$$\langle \tilde{v}, \tilde{u}'(s) - \frac{1}{2} y_1 \partial_1 \tilde{u}(s) \rangle_{f_s} + \tilde{a}_s(\tilde{v}, \tilde{u}(s)) = 0, \quad (7.26)$$

for each $\tilde{v} \in H_0^1(\Omega_0)$ and a.e. $s \in [0, \infty)$, with $\tilde{u}(0) = \tilde{u}_0 := U_0 u_0 = u_0$. Here $\langle \cdot, \cdot \rangle_{f_s}$ stands for the pairing of $H_0^1(\Omega_0, G_s)$ and its dual $[H_0^1(\Omega_0, G_s)]^*$, where G_s is the metric of the form (2.5) with f being replaced by f_s , and $\tilde{a}_s(\cdot, \cdot)$ denotes the sesquilinear form associated with

$$\tilde{a}_s[\tilde{u}] := \|f_s^{-1} \partial_1 \tilde{u}\|_{f_s}^2 + e^s \|\partial_2 \tilde{u}\|_{f_s}^2 - e^s E_1 \|\tilde{u}\|_{f_s}^2 - \frac{1}{4} \|\tilde{u}\|_{f_s}^2, \quad (7.27)$$

$$\tilde{u} \in \mathfrak{D}(\tilde{a}_s) := H_0^1(\Omega_0).$$

More specifically, $H_0^1(\Omega_0, G_s)$ denotes the completion of $C_0^\infty(\Omega_0)$ with respect to the norm $\|\cdot\|_{\mathfrak{D}(h_{K_s})} := (h_{K_s}[\cdot] + \|\cdot\|_{f_s}^2)^{1/2}$, where h_{K_s} is defined as (3.3) with f being replaced by f_s .

Remark 7.2. Note that (7.26) is a parabolic equation with s -time-dependent coefficients. The same occurs and has been previously analysed for the heat equation in the twisted waveguides [28, 29], for the heat equation in the plane with magnetic field [26] and also for a convection-diffusion equation in the whole space but with a variable diffusion coefficient [8, 5]. A careful analysis of the behaviour of the underlying elliptic operators as s tends to infinity leads to a sharp decay rate for its solutions. An important difference of the present problem with respect to the previous works is that also the Hilbert space becomes time-dependent after the self-similarity transformation, which makes the analysis substantially more difficult.

7.5 The setting in weighted Sobolev spaces

Since U_s acts as a unitary transformation, it preserves the space norm of solutions of (7.22) and (7.26), *i.e.*,

$$\|u(t)\|_f = \|\tilde{u}(s)\|_{f_s}. \quad (7.28)$$

This means that we can analyse the asymptotic time behaviour of the former by studying the latter.

However, the natural space to study the evolution (7.26) is not $L^2_{f_s}(\Omega_0)$ but rather the weighted space $L^2_{wf_s}(\Omega_0)$ with the Gaussian weight (4.6). Following the approach of [28] based on a theorem of J. L. Lions [1, Thm. X.9] about weak solutions of parabolic equations with time-dependent coefficients, it can be shown that (7.26) is well posed in the scale of Hilbert spaces

$$H^1_0(\Omega_0, wG_s) \subset L^2_{wf_s}(\Omega_0) \subset [H^1_0(\Omega_0, wG_s)]^*. \quad (7.29)$$

Here $H^1_0(\Omega_0, wG_s)$ denotes the completion of $C^\infty_0(\Omega_0)$ with respect to the norm $(h_{K_s}[w^{1/2}\cdot] + \|\cdot\|_{wf_s}^2)^{1/2}$.

More precisely, choosing $\tilde{v} := wv$ for the test function in (7.26), where $v \in C^\infty_0(\Omega_0)$ is arbitrary, we can formally cast (7.26) into the form

$$\langle v, \tilde{u}'(s) \rangle_w + a_s(v, \tilde{u}(s)) = 0. \quad (7.30)$$

Here $\langle \cdot, \cdot \rangle_w$ denotes the pairing of $H^1_0(\Omega_0, wG_s)$ and $[H^1_0(\Omega_0, wG_s)]^*$ and $a_s(\cdot, \cdot)$ denotes the sesquilinear form associated with

$$\begin{aligned} a_s[\tilde{u}] &:= \|f_s^{-1} \partial_1 \tilde{u}\|_{wf_s}^2 + e^s \|\partial_2 \tilde{u}\|_{wf_s}^2 - e^s E_1 \|\tilde{u}\|_{wf_s}^2 - \frac{1}{4} \|\tilde{u}\|_{wf_s}^2 \\ &\quad + \frac{1}{2} (y_1 \tilde{u}, [f_s^{-2} - 1] \partial_1 \tilde{u})_{wf_s}, \end{aligned} \quad (7.31)$$

$$\mathfrak{D}(a_s) := H^1_0(\Omega_0, w),$$

with $H^1_0(\Omega_0, w)$ denoting the closure of $C^\infty_0(\Omega_0)$ with respect to the weighted Sobolev norm $(\|\nabla \cdot\|_w^2 + \|\cdot\|_w^2)^{1/2}$. Note the appearance of the extra term with respect to (7.27) (it makes the form a_s non-symmetric if the Hilbert space $L^2_{wf}(\Omega_0)$ is considered to be complex).

By ‘formally’ we mean that the formulae are meaningless in general, because the solution $\tilde{u}(s)$ and its derivative $\tilde{u}'(s)$ may not belong to $H^1_0(\Omega_0, wG_s)$ and $[H^1_0(\Omega_0, wG_s)]^*$, respectively. The justification of (7.26) being well posed in the scale (7.29) consists basically in checking the boundedness and a coercivity of the form a_s defined on $\mathfrak{D}(a_s)$ and in noticing that the time-dependent spaces $L^2_{wf_s}(\Omega_0)$ and $H^1_0(\Omega_0, wG_s)$ coincide with $L^2_w(\Omega_0)$ and $H^1_0(\Omega_0, w)$, respectively, as vector spaces. It is straightforward by using (2.10) and Lemma 2.1.

7.6 Reduction to a spectral problem

Choosing $v := \tilde{u}(s)$ in (7.30), we arrive at the identity

$$\frac{1}{2} \frac{d}{ds} \|\tilde{u}(s)\|_{wf_s}^2 = -\hat{l}_s[\tilde{u}(s)], \quad (7.32)$$

where $\hat{l}_s[\tilde{u}] := \Re\{a_s[\tilde{u}]\}$, $\tilde{u} \in \mathfrak{D}(\hat{l}_s) := \mathfrak{D}(a_s) = H^1_0(\Omega_0, w)$ (independent of s as a vector space). It remains to analyse the coercivity of \hat{l}_s .

More precisely, as usual for energy estimates, we replace the right hand side of (7.32) by the spectral bound, valid for each fixed $s \in [0, \infty)$,

$$\forall \tilde{u} \in \mathfrak{D}(\hat{l}_s), \quad \hat{l}_s[\tilde{u}] \geq \nu_K(s) \|\tilde{u}\|_{wf_s}^2, \quad (7.33)$$

where $\nu_K(s)$ denotes the lowest point in the spectrum of the self-adjoint operator \hat{L}_s associated on $L_{wf_s}^2(\Omega_0)$ with \hat{l}_s ; it depends on the curvature K through the dependence on f . Then (7.32) together with (7.33) implies the exponential bound

$$\forall s \in [0, \infty), \quad \|\tilde{u}(s)\|_{wf_s} \leq \|\tilde{u}_0\|_{wf_0} e^{-\int_0^s \nu_K(r) dr}. \quad (7.34)$$

Finally, recall that the exponential bound in s transfers to a polynomial bound in the original time t , cf (7.25). In this way, the problem is reduced to a spectral analysis of the family of operators $\{\hat{L}_s\}_{s \geq 0}$.

7.7 Removing the weight

In order to investigate the operator \hat{L}_s on $L_{wf_s}^2(\Omega_0)$, we first map it into a unitarily equivalent operator $L_s := \mathcal{U} \hat{L}_s \mathcal{U}^{-1}$ on $L_{f_s}^2(\Omega_0)$ via the unitary transform

$$\mathcal{U} \tilde{u} := w^{1/2} \tilde{u}.$$

By definition, L_s is the self-adjoint operator associated on $L_{f_s}^2(\Omega_0)$ with the quadratic form $l_s[v] := \hat{l}_s[\mathcal{U}^{-1}v]$, $v \in \mathfrak{D}(l_s) := \mathcal{U} \mathfrak{D}(\hat{l}_s)$. A straightforward calculation yields

$$\begin{aligned} l_s[v] &= \|f_s^{-1} \partial_1 v\|_{f_s}^2 + e^s \|\partial_2 v\|_{f_s}^2 - e^s E_1 \|v\|_{f_s}^2 - \frac{1}{4} \|v\|_{f_s}^2 \\ &\quad - \frac{1}{2} (y_1 v, \partial_1 v)_{f_s} + \frac{1}{16} (y_1 v, [2 - f_s^{-2}] y_1 v)_{f_s}. \end{aligned} \quad (7.35)$$

Here and in the sequel, we assume that v is real, which is justified by the positivity preserving property of the heat equation as explained in Section 3.2.

For everywhere vanishing curvature, *i.e.* $K = 0$, we have that f is identically equal to one. Consequently, $f_s = 1$ for all $s \geq 0$. Then, integrating by parts in the first term on the second line of (7.35), we get that l_s coincides with the form l_s^0 on $L^2(\Omega_0)$ defined by

$$\begin{aligned} l_s^0[v] &:= \|\partial_1 v\|^2 + e^s \|\partial_2 v\|^2 - e^s E_1 \|v\|^2 + \frac{1}{16} \|y_1 v\|^2, \\ \mathfrak{D}(l_s^0) &:= H_0^1(\Omega_0) \cap L^2(\Omega_0, y_1^2 dy). \end{aligned} \quad (7.36)$$

In order to specify the domain of l_s for any curvature, we assume (5.2) and consider l_s as a perturbation of l_s^0 . It follows from (5.3) that

$$|y_1| > e^{-s/2} R \implies f_s(y) = 1. \quad (7.37)$$

In particular, $f_s(y) = 1$ for all $|y_1| > R$.

Lemma 7.1. *Assume (2.10) and (5.2). Then*

$$\mathfrak{D}(l_s) = \mathfrak{D}(l_s^0) = H_0^1(\Omega_0) \cap L^2(\Omega_0, y_1^2 dy).$$

Proof. Using some rearrangement and integration by parts, it is convenient to rewrite (7.35) as follows

$$l_s[v] = \|f_s^{-1}\partial_1 v\|_{f_s}^2 + e^s \|\partial_2 v\|_{f_s}^2 - e^s E_1 \|v\|_{f_s}^2 + \frac{1}{16} \|y_1 v\|_{f_s}^2 + r_s[v], \quad (7.38)$$

where

$$r_s[v] := -\frac{1}{4} (v, [f_s - 1]v) - \frac{1}{2} (y_1 v, [f_s - 1]\partial_1 v) - \frac{1}{16} (y_1 v, [f_s^{-1} - f_s]y_1 v).$$

Using Lemma 2.1, it is easy to see that for each $s \geq 0$ there exists a positive constant $C = C(s, \|K\|_\infty, a)$ such that

$$C^{-1} l_s^0[v] \leq l_s[v] - r_s[v] \leq C l_s^0[v]$$

for every $v \in \mathfrak{D}(l_s^0)$. Consequently (see, e.g., [2, Corol. 4.4.3]), the quadratic form $l_s - r_s$ is closed on the domain $\mathfrak{D}(l_s^0)$ given by (7.36). It remains to show that r_s is a relatively bounded perturbation of l_s^0 with relative bound smaller than one. It is clear for the first term of r_s which is in fact a bounded perturbation in view of Lemma 2.1. We employ (7.37) to deal with the remaining terms. For the second term we have,

$$\begin{aligned} |(y_1 v, [f_s - 1]\partial_1 v)| &\leq \|f - 1\|_\infty \int_{\{|y_1| < e^{-s/2} R\}} |y_1| |v(y)| |\partial_1 v(y)| dy \\ &\leq \|f - 1\|_\infty e^{-s/2} R \|v\| \|\partial_1 v\| \\ &\leq \|f - 1\|_\infty R (\epsilon^{-1} \|v\|^2 + \epsilon \|\partial_1 v\|^2) \end{aligned}$$

for every $v \in \mathfrak{D}(l_s^0)$. and any $\epsilon \in (0, 1)$. Similarly,

$$|(y_1 v, [f_s^{-1} - f_s]y_1 v)| \leq \|f^{-1} - f\|_\infty R^2 \|v\|^2.$$

for every $v \in \mathfrak{D}(l_s^0)$. □

Remark 7.3. The proof of the lemma represents a direct way how to show that the form (7.35) is closed on the domain $\mathfrak{D}(l_s^0)$. In view of the unitary equivalence \mathcal{U} , it also *a posteriori* establishes the closedness of the form (7.31).

As a consequence of Lemma 7.1, we get that L_s (and therefore \hat{L}_s) has compact resolvent and thus purely discrete spectrum for all $s \geq 0$. In particular, $\nu_K(s)$ represents the lowest eigenvalue of L_s .

7.8 The strong-resolvent convergence

In order to study the decay rate via (7.34), we need information about the limit of the eigenvalue $\nu_K(s)$ as the time s tends to infinity. This can be deduced from the asymptotic properties of the resolvent of L_s for large s .

In view of (5.2), the function $y \mapsto f_s(y)$ converges to one locally uniformly in $|y_1| > 0$, $y_2 \in (-a, a)$, as $s \rightarrow \infty$. Moreover, the scaling of the transverse variable in (7.35) corresponds to considering the operator L_0 in the shrinking strip $\mathbb{R} \times (-e^{-s/2}a, e^{-s/2}a)$. This suggests that L_s will converge, in a suitable sense, to the one-dimensional harmonic-oscillator operator

$$h := -\frac{d^2}{dy_1^2} + \frac{1}{16} y_1^2 \quad \text{on} \quad L^2(\mathbb{R}) \quad (7.39)$$

(*i.e.* the Friedrichs extension of this operator initially defined on $C_0^\infty(\mathbb{R})$), potentially subjected to an extra condition at the origin. For further purposes, let us note that the spectrum of h is known explicitly (see any book on quantum mechanics, *e.g.*, [11, Sec. 2.3])

$$\sigma(h) = \left\{ \frac{1}{2} \left(n + \frac{1}{2} \right) \right\}_{n=0}^\infty. \quad (7.40)$$

We shall see that the difference between the negatively curved and flat case consists in that the limit operator for the former is subjected to an extra Dirichlet boundary condition at $y_1 = 0$. Thus, simultaneously to h introduced in (7.39), let us consider the self-adjoint operator h_D in $L^2(\mathbb{R})$ whose quadratic form acts in the same way as that of h but has a smaller domain

$$\mathfrak{D}(h_D^{1/2}) := \{ \varphi \in \mathfrak{D}(h^{1/2}) \mid \varphi(0) = 0 \}.$$

To make this singular operator limits mentioned above rigorous (L_s and h act on different spaces), we decompose the Hilbert space $L^2(\Omega_0)$ into an orthogonal sum

$$L^2(\Omega_0) = \mathfrak{H}_1 \oplus \mathfrak{H}_1^\perp, \quad (7.41)$$

where the subspace \mathfrak{H}_1 consists of functions of the form $\psi_1(y) = \varphi(y_1)\mathfrak{J}_1(y')$. Recall that \mathfrak{J}_1 denotes the positive eigenfunction of $-\Delta_D^{(-a,a)}$ corresponding to E_1 , normalized to 1 in $L^2((-a,a))$, *cf* (4.2). Given any $\psi \in L^2(\Omega_0)$, we have the decomposition $\psi = \psi_1 + \psi^\perp$ with $\psi_1 \in \mathfrak{H}_1$ as above and $\psi^\perp \in \mathfrak{H}_1^\perp$. The mapping $\pi : \varphi \mapsto \psi_1$ is an isomorphism of $L^2(\mathbb{R})$ onto \mathfrak{H}_1 . Hence, with an abuse of notations, we may identify any operator h on $L^2(\mathbb{R})$ with the operator $\pi h \pi^{-1}$ acting on $\mathfrak{H}_1 \subset L^2(\Omega_0)$.

Finally, we mention that the Hilbert spaces $L^2(\Omega_0)$ and $L_{f_s}^2(\Omega_0)$ can be identified as vector sets because their norms are equivalent. More specifically, in view of Lemma 2.1 and the definition (7.23), we have

$$1 - \frac{\|K\|_\infty a^2}{1 - \|K\|_\infty a^2} \leq \frac{\|\psi\|_{f_s}^2}{\|\psi\|^2} \leq 1 + \frac{\|K\|_\infty a^2}{1 - \|K\|_\infty a^2}, \quad (7.42)$$

for every non-zero $\psi \in L^2(\Omega_0)$ and all $s \geq 0$.

In the flat case, *i.e.* $K = 0$, it is readily seen that the operator L_s^0 associated with the form (7.36) can be identified with the decomposed operator

$$h \otimes 1 + 1 \otimes (-e^s \Delta_D^{(-a,a)} - e^s E_1) \quad \text{in} \quad L^2(\mathbb{R}) \otimes L^2((-a,a)), \quad (7.43)$$

where 1 denotes the identity operators in the appropriate spaces. Using (7.40), it follows that $\nu_0(s) = 1/4$ for all $s \in [0, \infty)$. Consequently,

$$\nu_0(\infty) := \lim_{s \rightarrow \infty} \nu_0(s) = 1/4. \quad (7.44)$$

Moreover, (7.43) can be used to show that L_s^0 converges to $h \oplus 0^\perp$ in the norm-resolvent sense as $s \rightarrow \infty$, where 0^\perp denotes the zero operator on \mathfrak{H}_1^\perp .

It is more difficult (and more interesting) to establish the asymptotic behaviour of $\nu_K(s)$ for $K \neq 0$. A fine analysis of its limit leads to the key observation of the paper, ensuring a gain of 1/2 in the decay rate in the negatively curved case. This can be understood from the following proposition, which represents the main auxiliary result of the present paper.

Proposition 7.2. *Assume (2.10) and (5.2). Let the Hardy-type inequality (7.1) holds. Then the operator L_s converges to $h_D \oplus 0^\perp$ in the strong-resolvent sense as $s \rightarrow \infty$, i.e.,*

$$\forall F \in L^2(\Omega_0), \quad \lim_{s \rightarrow \infty} \left\| (L_s + i)^{-1} F - \left[(h_D + i)^{-1} \oplus 0^\perp \right] F \right\| = 0.$$

Proof. For the clarity of the exposition, we divide the proof into several steps. The equivalence of norms (7.42) and other consequences of Lemma 2.1 are widely used in the present proof.

1. *The resolvent equation for L_s .* Let $F \in L^2(\Omega_0)$. Then also $F \in L^2_{f_s}(\Omega_0)$ for every $s \geq 0$ due to (7.42). Let z be a sufficiently large positive number to be specified later. We set $\psi_s := (L_s + z)^{-1} F$. In other words, ψ_s satisfies the resolvent equation

$$\forall v \in \mathfrak{D}(l_s), \quad l_s(v, \psi_s) + z(v, \psi_s)_{f_s} = (v, F)_{f_s}. \quad (7.45)$$

In particular, choosing ψ_s for the test function v in (7.45), we have

$$\forall v \in \mathfrak{D}(l_s), \quad l_s[\psi_s] + z \|\psi_s\|_{f_s}^2 = (\psi_s, F)_{f_s} \leq \|\psi_s\|_{f_s} \|F\|_{f_s}. \quad (7.46)$$

2. *Boundedness of ψ_s .* Our primary objective is to deduce from (7.46) that $\{\psi_s\}_{s \geq 0}$ is a bounded family in the space $\mathfrak{D}_0 := H_0^1(\Omega_0) \cap L^2(\Omega_0, y_1^2 dy)$ equipped with the intersection topology.

We search a lower bound to the operator $L_s + z$. Using the convenient form (7.38) for $l_s[\psi_s]$ and proceeding as in the proof of Lemma 7.1, we easily check that

$$|r_s[\psi_s]| \leq C (\epsilon \|f_s^{-1} \partial_1 \psi_s\|_{f_s}^2 + \epsilon^{-1} \|\psi_s\|_{f_s}^2). \quad (7.47)$$

with any $\epsilon \in (0, 1)$, where C is a positive constant depending on $\|K\|_\infty a^2$ and R . Hence

$$\begin{aligned} l_s[\psi_s] + z \|\psi_s\|_{f_s}^2 &\geq (1 - 2\epsilon) \|f_s^{-1} \partial_1 \psi_s\|_{f_s}^2 + e^s \|\partial_2 \psi_s\|_{f_s}^2 - e^s E_1 \|\psi_s\|_{f_s}^2 \\ &\quad + \epsilon \|f_s^{-1} \partial_1 \psi_s\|_{f_s}^2 + \frac{1}{16} \|y_1 \psi_s\|_{f_s}^2 + (z - C \epsilon^{-1}) \|\psi_s\|_{f_s}^2. \end{aligned} \quad (7.48)$$

If we choose z larger than $C \epsilon^{-1}$, all the terms on the second line are non-negative.

To get a non-negative lower bound to the first line on the right hand side of (7.48), we introduce a new function u_s by $\psi_s(y) = e^{s/4} u_s(e^{s/2} y_1, y_2)$ (cf the self-similarity transformation (7.24)). Making the change of variables $(x_1, x_2) = (e^{s/2} y_1, y_2)$, recalling the definition (7.7) and using the Hardy-type inequality (7.1), we obtain

$$\begin{aligned} (1 - 2\epsilon) \|f_s^{-1} \partial_1 \psi_s\|_{f_s}^2 + e^s \|\partial_2 \psi_s\|_{f_s}^2 - e^s E_1 \|\psi_s\|_{f_s}^2 \\ = e^s (1 - 2\epsilon) \|f^{-1} \partial_1 u_s\|_f^2 + e^s \|\partial_2 u_s\|_f^2 - e^s E_1 \|u_s\|_f^2 \\ \geq e^s (1 - 2\epsilon) \|\rho^{1/2} u_s\|_f^2 + e^s 2\epsilon (u_s, \mu_K u_s)_f. \end{aligned} \quad (7.49)$$

Here ρ is a positive function and, as pointed out in Remark 7.1, we may assume that it depends on x_1 only. On the other hand, μ_K has compact support due to (5.3). Hence, we can choose ϵ sufficiently small so that the new Hardy weight

$\tilde{\rho}(x_1) := (1 - 2\epsilon)\rho(x_1) + 2\epsilon\mu_K(x_1)$ is positive for almost every $x_1 \in \mathbb{R}$. Coming back to our coordinates y , we thus conclude from (7.49)

$$(1 - 2\epsilon) \|f_s^{-1} \partial_1 \psi_s\|_{f_s}^2 + e^s \|\partial_2 \psi_s\|_{f_s}^2 - e^s E_1 \|\psi_s\|_{f_s}^2 \geq e^s \|\tilde{\rho}_s^{1/2} \psi_s\|_f^2,$$

where $\tilde{\rho}_s(y_1) := \rho(e^{s/2} y_1)$.

Using the last inequality in (7.48) and employing Lemma 2.1, we eventually arrive at

$$l_s[\psi_s] + z \|\psi_s\|_{f_s}^2 \geq c \left(e^s \|\tilde{\rho}_s^{1/2} \psi_s\|^2 + \|\partial_1 \psi_s\|^2 + \|y_1 \psi_s\|^2 + (z - C\epsilon^{-1}) \|\psi_s\|^2 \right), \quad (7.50)$$

where c is a positive constant depending on $\|K\|_\infty a^2$. Comparing this inequality with (7.46), we see that there exists a constant z_0 , depending on a and properties of K , such that for all $z \geq z_0$

$$\|\psi_s\| \leq C \|F\|, \quad \|y_1 \psi_s\| \leq C \|F\|, \quad \|\partial_1 \psi_s\| \leq C \|F\|, \quad (7.51)$$

and

$$e^s \|\tilde{\rho}_s^{1/2} \psi_s\|^2 \leq C \|F\|^2, \quad (7.52)$$

with some constant C depending on a and properties of K . Furthermore, directly from (7.48) and (7.46) with help of (7.51), we also get

$$\|\partial_2 \psi_s\| \leq C \|F\|, \quad (7.53)$$

The estimate (7.50) also shows that $L_s + z$ is invertible for all $z \geq z_0$. This, *a posteriori*, justifies the definition of ψ_s as the unique solution of (7.45).

From (7.51) and (7.53), we conclude that $\{\psi_s\}_{s \geq 0}$ is a bounded family in \mathfrak{D}_0 . Therefore it is precompact in the weak topology of \mathfrak{D}_0 . Let ψ_∞ be a weak limit point, *i.e.*, for an increasing sequence of positive numbers $\{s_n\}_{n \in \mathbb{N}}$ such that $s_n \rightarrow \infty$ as $n \rightarrow \infty$, $\{\psi_{s_n}\}_{n \in \mathbb{N}}$ converges weakly to ψ_∞ in \mathfrak{D}_0 . Actually, we may assume that it converges strongly in $L^2(\Omega_0)$ because \mathfrak{D}_0 is compactly embedded in $L^2(\Omega_0)$.

3. Transverse mode decomposition of ψ_s . Now we employ the Hilbert space decomposition (7.41) and write $\psi_s(y) = \varphi_s(y_1) \mathcal{J}_1(y_2) + \psi_s^\perp(y)$, where $\psi_s^\perp \in \mathfrak{H}_1^\perp$, *i.e.*,

$$(\mathcal{J}_1, \psi_s^\perp(y_1, \cdot))_{L^2((-a, a))} = 0 \quad (7.54)$$

for a.e. $y_1 \in \mathbb{R}$. It follows from (7.51), (7.53) and (7.54) that also $\{\psi_s^\perp\}_{s \geq 0}$ is a bounded family in \mathfrak{D}_0 and that $\{\varphi_s\}_{s \geq 0}$ is a bounded family in $H^1(\mathbb{R}) \cap L^2(\mathbb{R}, y_1^2 dy_1)$ equipped with the intersection topology. We denote by ψ_∞^\perp and φ_∞ the respective limit points.

We come back to (7.46) with (7.48) and focus on the inequality

$$e^s \|\partial_2 \psi_s\|_{f_s}^2 - e^s E_1 \|\psi_s\|_{f_s}^2 \leq C \|F\|^2 \quad (7.55)$$

we have already used to get (7.53). In the same way as we proceeded to get (7.14), we write $\phi_s := \sqrt{f_s} \psi_s$ and obtain

$$\|\partial_2 \psi_s\|_{f_s}^2 - E_1 \|\psi_s\|_{f_s}^2 = \|\partial_2 \phi_s\|^2 - E_1 \|\phi_s\|^2 + (\phi_s, V_s \phi_s), \quad (7.56)$$

where V_s is defined in the same way as (7.15) but with K and f being replaced by K_s and f_s , respectively. Using (7.37), it is possible to check that $\{\phi_{s_n}\}_{n \in \mathbb{N}}$ is strongly converging in $L^2(\Omega_0)$; in fact,

$$\lim_{n \rightarrow \infty} \|\phi_{s_n} - \psi_\infty\| = 0. \quad (7.57)$$

Using the fact that the scaled potential V_s in (7.56) vanishes for all $|y_1| > e^{-s/2}R$ together with the strong convergence of $\{\phi_{s_n}\}_{n \in \mathbb{N}}$, it is easy to see that the integral containing the potential tends to zero as $n \rightarrow \infty$, after passing to the subsequence $\{s_n\}_{n \in \mathbb{N}}$. Multiplying (7.55) by e^{-s_n} and putting the asymptotically vanishing integral on the right hand side of the inequality, we thus get

$$\lim_{n \rightarrow \infty} (\|\partial_2 \phi_{s_n}\|^2 - E_1 \|\phi_{s_n}\|^2) = 0. \quad (7.58)$$

Using in addition the Hilbert space decomposition (7.41) of ϕ_s , *i.e.* $\phi_s(y) = \eta_s(y_1)\mathcal{J}_1(y_2) + \phi_s^\perp(y)$, we see that the same limit (7.58) holds for $\phi_{s_n}^\perp \in \mathfrak{H}_1^\perp$ as well. In that limit, we use the estimate $\|\partial_2 \phi_{s_n}^\perp\|^2 \geq E_2 \|\phi_{s_n}^\perp\|^2$, where $E_2 = 4E_1$ denotes the second eigenvalue of $-\Delta_D^{(-a,a)}$, and conclude that $\|\phi_{s_n}^\perp\|$ tends to zero as $n \rightarrow \infty$. The latter together with (7.57) finally yields

$$\lim_{n \rightarrow \infty} \|\psi_{s_n}^\perp\| = 0 \quad \text{and} \quad \lim_{n \rightarrow \infty} \|\eta_{s_n} - \varphi_\infty\|_{L^2(\mathbb{R})} = 0. \quad (7.59)$$

That is, $\psi_\infty \in \mathfrak{H}_1$.

4. The Dirichlet condition at zero. Now we come back to the inequality (7.52). Recall that $\tilde{\rho}_s(y_1) = \tilde{\rho}_s(e^{s/2}y_1)$ and that $\tilde{\rho}$ is positive (although necessarily vanishing at infinity). Without loss of generality, we may assume that $\tilde{\rho}$ in (7.52) belongs to $L^1(\mathbb{R})$ (since we can always replace the estimate using a smaller function). Then $e^{s/2}\tilde{\rho}_s$ converges in the sense of distributions on \mathbb{R} to a Dirac delta at $y_1 = 0$. We want to use this heuristic consideration to show that $\varphi_\infty(0) = 0$.

To do so, first, we use the Hilbert space decomposition (7.41) of ψ_s and notice that the left hand side of (7.52) splits into a sum of two non-negative parts, the mixed term being zero due to (7.54). Second, multiplying the obtained inequality for the term involving the \mathfrak{H}_1 -part of ψ_s by $e^{-s/2}$, passing to the subsequence $\{s_n\}_{n \in \mathbb{N}}$ and taking the limit $n \rightarrow \infty$, we arrive at

$$|\varphi_\infty(0)|^2 \int_{\mathbb{R}} \tilde{\rho}(x_1) dx_1 = 0.$$

The limiting procedure is justified by recalling that $\{\varphi_{s_n}\}_{n \in \mathbb{N}}$ converges weakly in $H^1(\mathbb{R})$ and therefore strongly in $H^1(J)$, which is compactly embedded in $C^{0,\lambda}(J)$ for every $\lambda \in (0, 1/2)$, where J is any bounded interval of \mathbb{R} .

Since the integral of $\tilde{\rho}$ is positive by our hypotheses, we thus verify that φ_∞ satisfies the extra Dirichlet condition

$$\varphi_\infty(0) = 0.$$

5. The resolvent equation for L_s as $s \rightarrow \infty$. Let us summarize our results. We have obtained that the solutions ψ_{s_n} of (7.45) converge in the weak topology of \mathfrak{D}_0 and in the strong topology of $L^2(\Omega_0)$ to some ψ_∞ . Moreover, the limiting solution ψ_∞ belongs to \mathfrak{H}_1 , so that $\psi_\infty(y) = \varphi_\infty(y_1)\mathcal{J}_1(y_2)$ with some $\varphi_\infty \in$

$H^1(\mathbb{R}) \cap L^2(\mathbb{R}, y_1^2 dy_1) = \mathfrak{D}(h)$. Finally, $\varphi_\infty(0) = 0$, so that actually $\varphi_\infty \in \mathfrak{D}(h_D)$.

Recall that the set $C_0^\infty(\mathbb{R} \setminus \{0\})$ is dense in $\mathfrak{D}(h_D)$. Let $\varphi \in C_0^\infty(\mathbb{R} \setminus \{0\})$ be arbitrary. We take $v(y) := \varphi(y_1)\mathcal{J}_1(y_2)$ as the test function in (7.45) and note that φ and $f_s - 1$ have disjoint support for sufficiently large s due to (7.37). Sending n to infinity in (7.45) with s being replaced by s_n , we thus easily check that

$$(\dot{\varphi}, \dot{\varphi}_\infty)_{L^2(\mathbb{R})} + \frac{1}{16} (y_1 \varphi, y_1 \varphi_\infty)_{L^2(\mathbb{R})} + z (\varphi, \varphi_\infty)_{L^2(\mathbb{R})} = (\varphi, f)_{L^2(\mathbb{R})},$$

where $f(y_1) := (\mathcal{J}_1, F(y_1, \cdot))_{L^2((-a, a))}$. That is, $\varphi_\infty = (h_D + z)^{-1} f$, for *any* weak limit point of $\{\varphi_s\}_{s \geq 0}$.

In conclusion, we have shown that ψ_s converges strongly to ψ_∞ in $L^2(\Omega_0)$ as $s \rightarrow \infty$, where $\psi_\infty(y) := \varphi_\infty(y_1)\mathcal{J}_1(y_2) = [(h_D + z)^{-1} \oplus 0^\perp] F(y)$.

6. The strong convergence for other values of the spectral parameter. Finally, let us argue that we can replace the real number z by any non-real number. This is actually a consequence of [19, Thm. VIII.1.3], the fact that L_s is self-adjoint on $L_{f_s}^2(\Omega_0)$ and the equivalence of this Hilbert space with $L^2(\Omega_0)$, to which we consider the limit of the strong convergence, due to (7.42). \square

Remark 7.4. The crucial step in the proof is certainly the usage of the Hardy inequality (7.1). Indeed, it enables us, first, to ensure the non-negativity of the right hand side of (7.48) and, second, to establish the extra Dirichlet condition at zero.

7.9 Spectral consequences

Assume for a moment that Proposition 7.2 stated that the operator L_s converges to $h_D \oplus 0^\perp$ in the *norm-resolvent* sense as $s \rightarrow \infty$. Then we would immediately know that $\nu_K(s)$ converges to the first eigenvalue of h_D as $s \rightarrow \infty$. In view of the symmetry, the first eigenvalue of h_D coincides with the second eigenvalue of h , which is $3/4$ due to (7.40). Hence, under the hypotheses of proposition 7.2, we would have that the limit of $\nu_K(s)$ as $s \rightarrow \infty$ is three-times larger than the same limit in the flat case (7.44).

Unfortunately, the strong-resolvent convergence of Proposition 7.2 is not sufficient to guarantee the convergence of spectra. In general, this is true for eigenvalues of the limiting operator which are *stable* under the perturbation (*cf* [19, Sec. VIII.1]). In our case, however, the spectral convergence can be established directly using the fact that both L_s and h_D are operators with compact resolvents. Using the compactness, the convergence of eigenvalues follow by a straightforward modification of the proof of Proposition 7.2. In particular, we have the following result for the lowest eigenvalue, exactly as we would have under the norm-resolvent convergence described above.

Corollary 7.1. *Under the hypotheses of Proposition 7.2, one has*

$$\nu_K(\infty) := \lim_{s \rightarrow \infty} \nu_K(s) = 3/4.$$

Proof. First of all, let us notice that $\nu_K(s)$ remains bounded as $s \rightarrow \infty$. This is easily seen by the Rayleigh-Ritz variational formula for the lowest eigenvalue

of L_s , in which we use the trial function of the form $\psi(y) := \varphi(y_1)\mathcal{J}_1(y_2)$, where $\varphi \in C_0^\infty(\mathbb{R})$ is supported outside $\text{supp}(f-1) \supseteq \text{supp}(f_s-1)$ (cf (7.37)). Indeed,

$$\nu_K(s) \leq \frac{l_s[\psi]}{\|\psi\|_{f_s}^2} = \frac{\|\dot{\varphi}\|_{L^2(\mathbb{R})}^2 + \frac{1}{16} \|y_1 \varphi\|_{L^2(\mathbb{R})}^2}{\|\varphi\|_{L^2(\mathbb{R})}^2}, \quad (7.60)$$

irrespectively of the properties of K .

Now, let ψ_s be the positive eigenfunction of L_s corresponding to $\nu_K(s)$, normalized to 1 in $L_{f_s}^2(\Omega_0)$. ψ_s is a solution of the problem (7.45) with $F := (\nu_K(s) + z)\psi_s$. It is important that F is uniformly bounded in s as an element of $L_{f_s}^2(\Omega_0)$, due to (7.60) and the normalization of ψ_s . Then we can proceed exactly as in the proof of Proposition 7.2.

We show that $\{\psi_s\}_{s \geq 0}$ contains a subsequence $\{\psi_{s_n}\}_{n \in \mathbb{N}}$ which is weakly converging to some ψ_∞ in \mathfrak{D}_\circ . Since \mathfrak{D}_\circ is compactly embedded in $L^2(\Omega_0)$, we know that $\{\psi_{s_n}\}_{n \in \mathbb{N}}$ converges to ψ_∞ strongly in $L^2(\Omega_0)$. In particular, $\|\psi_\infty\| = 1$, so that we know that ψ_∞ is non-trivial. At the same time, we show that $\psi_\infty \in \mathfrak{H}_1$ and that $\varphi_\infty(y_1) := (\mathcal{J}_1, \psi_\infty(y_1, \cdot))_{L^2((-a, a))}$ vanishes at $y_1 = 0$.

Taking $v(y) := \varphi(y_1)\mathcal{J}_1(y_2)$ with $\varphi \in C_0^\infty(\mathbb{R} \setminus \{0\})$ as the test function in the weak formulation of the eigenvalue problem (7.45), with s being replaced by s_n , and sending n to infinity, we eventually find that φ_∞ is an eigenfunction of $h_D + z$ with the eigenvalue $\nu_K(\infty) + z$. Since ψ_∞ is obtained as a limit of positive functions, we know that φ_∞ is positive as well. Hence, $\nu_K(\infty)$ represents the lowest eigenvalue of h_D .

It remains to recall that the first eigenvalue of h_D coincides with the second eigenvalue of h , which is $3/4$ due to (7.40). \square

7.10 A spectral bound to the decay rate

We come back to (7.34). Assume $K = 0$ or that there exists a Hardy-type inequality (7.1). Recalling (7.44) and Corollary 7.1, we know that for arbitrarily small positive number ε there exists a (large) positive time s_ε such that for all $s \geq s_\varepsilon$, we have $\nu_K(s) \geq \nu_K(\infty) - \varepsilon$. Hence, fixing $\varepsilon > 0$, we have

$$\begin{aligned} - \int_0^s \nu_K(r) dr &\leq - \int_0^{s_\varepsilon} \nu_K(r) dr - [\nu_K(\infty) - \varepsilon](s - s_\varepsilon) \\ &\leq \int_0^{s_\varepsilon} |\nu_K(r)| dr + [\nu_K(\infty) - \varepsilon]s_\varepsilon - [\nu_K(\infty) - \varepsilon]s \end{aligned}$$

for all $s \geq s_\varepsilon$. At the same time, assuming $\varepsilon \leq 1/4$, we trivially have

$$- \int_0^s \nu_K(r) dr \leq \int_0^{s_\varepsilon} |\nu_K(r)| dr + [\nu_K(\infty) - \varepsilon]s_\varepsilon - [\nu_K(\infty) - \varepsilon]s$$

also for all $s \leq s_\varepsilon$. Summing up, for every $s \in [0, \infty)$, we have

$$\|\tilde{u}(s)\|_{wf_s} \leq C_\varepsilon e^{-[\nu_K(\infty) - \varepsilon]s} \|\tilde{u}_0\|_{wf_0}, \quad (7.61)$$

where $C_\varepsilon := e^{\int_0^{s_\varepsilon} |\nu_K(r)| dr + [\nu_K(\infty) - \varepsilon]s_\varepsilon}$.

Now we return to the original variables (x, t) via (7.25). Using (7.28) together with the point-wise estimate $1 \leq w$, and recalling that $f_0 = f$ and $\tilde{u}_0 = u_0$, it follows from (7.61) that

$$\|u(t)\|_f = \|\tilde{u}(s)\|_{f_s} \leq \|\tilde{u}(s)\|_{wf_s} \leq C_\varepsilon (1+t)^{-[\nu_K(\infty) - \varepsilon]} \|u_0\|_{wf}$$

for every $t \in [0, \infty)$. Consequently, we conclude with

$$\|e^{-(H_K - E_1)t}\|_{L^2_{wf}(\Omega_0) \rightarrow L^2_f(\Omega_0)} = \sup_{u_0 \in L^2_{wf}(\Omega_0) \setminus \{0\}} \frac{\|u(t)\|_f}{\|u_0\|_{wf}} \leq C_\varepsilon (1+t)^{-[\mu_\theta(\infty) - \varepsilon]}$$

for every $t \in [0, \infty)$. Since ε can be made arbitrarily small, this bound implies

$$\Gamma_K \geq \nu_K(\infty). \quad (7.62)$$

7.11 The improved decay rate

Now we arrive to the main result of this paper. It follows from Proposition 4.1 that $\Gamma_0 = 1/4$ (i.e. $K = 0$). The lower bound $\Gamma_0 \geq 1/4$ alternatively follows from (7.62) using (7.44). The following theorem states that the decay rate is three times better in the presence of a Hardy-type inequality (7.1).

Theorem 7.3. *Assume (2.10) and (5.2). If (7.1) holds, then*

$$\Gamma_K = 3/4.$$

Proof. The assertion $\Gamma_K \geq 3/4$ follows from (7.62) using Corollary 7.1. In order to prove the $\Gamma_K \leq 3/4$ it is sufficient to show, that for some suitable function $\varphi \in C_0^\infty(\Omega_0)$, some constant $c_\varphi > 0$ and some constant $t_0 \geq 0$

$$\forall t \geq t_0, \quad \|e^{-tH_K}\varphi\|_{L^2_f(\Omega_0)} \geq c_\varphi t^{-3/4} e^{E_1 t}. \quad (7.63)$$

Due to (5.2) the support of f is contained in a rectangle $\Omega_R := (-R, R) \times (-a, a)$ for $R > 0$. We choose $\varphi \in C_0^\infty(\Omega_0)$ such that $\text{supp}(\varphi) \subset \Omega_0 \setminus \Omega_R$. Recall that \mathbb{E}_x (respectively, \mathbb{P}_x) denote the expectation (respectively, probability measure) corresponding to the Markov process $(X_t)_{t \geq 0}$ associated to the Dirichlet form h_k . Define the stopping times τ_{Ω_0} and τ_{Ω_R} by

$$\tau_{\Omega_0} = \inf\{t \geq 0 \mid X_t \in \Omega_0\} \quad \text{and} \quad \tau_{\Omega_R} = \inf\{t \geq 0 \mid X_t \in \partial\Omega_R\}.$$

The process $(X_t)_{0 \leq t < \tau_{\Omega_0}}$ is called Brownian motion on Ω_0 killed at the boundary. For every $x = (x_1, x_2) \in Q_0$ we then conclude

$$e^{-tH_K}\varphi(x) = \mathbb{E}_x[\varphi(X_t), ; \tau_{\Omega_0}] \geq \mathbb{E}_x[\varphi(X_t), \tau_{\Omega_R} \wedge \tau_{\Omega_0} > t], \quad (7.64)$$

where $\tau_{\Omega_R} \wedge \tau_{\Omega_0}$ denotes the minimum of the stopping times τ_{Ω_0} and τ_{Ω_R} . Integration of (7.64) and using $K \upharpoonright \Omega_0 \setminus \Omega_R = 0$ and hence – by Lemma 2.1 – $f = 1$ in $\Omega_0 \setminus \Omega_R$ yields

$$\begin{aligned} \|e^{-tH_K}\varphi\|_{L^2_f(\Omega_0)}^2 &\geq \int_{(R, \infty) \times (-a, a)} |\mathbb{E}_x[\varphi(X_t), \tau_{\Omega_R} \wedge \tau_{\Omega_0} > t]|^2 dx \\ &= \int_{(R, \infty) \times (-a, a)} |\mathbb{E}_x[\varphi(X_t), \tau_{\Omega_0} > t]|^2 dx \end{aligned} \quad (7.65)$$

where $\Omega_{0R} := (R, \infty) \times (-a, a)$ and

$$\tau_{\Omega_{0R}} = \inf\{t \geq 0 \mid X_t \in \partial\Omega_{0R}\}.$$

Due to $f = 1$ in $\Omega_0 \setminus \Omega_R$ the stochastic process $(X_t)_{\tau_{\Omega_0 R} > t \geq 0}$ is a (deterministically time changed by the factor 2) Brownian motion killed, when exiting the set Ω_{0R} . Due to independence of the coordinates we have

$$\mathbb{E}_x[\varphi(X_t), \tau_{\Omega_R} \wedge \tau_{\Omega_0} > t] = \sum_{n=1}^{\infty} e^{-E_n t} \mathcal{J}_n(x_1) \int_{\Omega_{0,R}} p_0(t, x_1, y_1) \mathcal{J}_n(2) \varphi(y_1, y_2) dy, \quad (7.66)$$

where

$$p_0(t, x, y) := \frac{1}{\sqrt{4\pi t}} \left(e^{-\frac{(x-y)^2}{4t}} - e^{-\frac{(x+y)^2}{4t}} \right)$$

is the transition function of a one-dimensional Brownian motion killed when hitting 0. Using (7.65) and (7.66) an elementary calculation gives assertion (7.63). \square

Observe that the proof of Theorem 7.3 demonstrates that the ‘transient’ effect of negative curvature on the survival properties of a Brownian particle is as strong as if we kill a particle when entering the curved region.

7.12 From normwise to pointwise bounds

Theorem 7.3 may be reformulated in terms of certain pointwise assertions.

Corollary 7.2. *Assume (2.10), (5.2) as well as the (7.1). Let $x \in \Omega_0$, $\delta > 0$ and a measurable bounded subset $B \subset \Omega_0$ be given. Then there exists a constant $C_{B,\delta,x} > 0$ such that*

$$\mathbb{P}_x(X_t \in B, \tau_{\Omega_0} > t) \leq C_{B,\delta,x} e^{-E_1 t} t^{-\frac{3}{2}+\delta}.$$

Proof. We use that according to Proposition 3.1 the integral kernel $e^{-tH_K}(x, y)$ of e^{-tH_K} satisfies the following Gaussian upper bound

$$e^{-tH_K}(x, y) \leq \frac{c_1}{\sqrt{4\pi t}} e^{-\frac{(x-y)^2}{4c_2 t}}$$

for some constants $c_1, c_2 > 0$. For fixed x set

$$\psi_{x,\varepsilon}(y) = \frac{c_1}{\sqrt{4\pi\varepsilon}} e^{-\frac{(x-y)^2}{4c_2\varepsilon}},$$

where ε is chosen small enough such that $\psi_{x,\varepsilon} \in L^2_{wf}(\Omega_0)$. Then for $t > \varepsilon$ we have for some constant $C_\delta > 0$

$$\begin{aligned} \mathbb{P}_x(X_t \in B, \tau_{\Omega_0} > t) &= e^{-tH_K} \chi_B(x) = e^{-\varepsilon H_K} e^{-(t-\varepsilon)H_K} \chi_B(x) \\ &= (\psi_{x,\varepsilon}, e^{-(t-\varepsilon)H_K} \chi_B)_f \\ &= (e^{-\frac{t-\varepsilon}{2}H_K} \psi_{x,\varepsilon}, e^{-\frac{t-\varepsilon}{2}H_K} \chi_B)_f \\ &\leq \|\psi_{x,\varepsilon}\|_{wf} \|\chi_B\|_{wf} \left[C_\delta \left(\frac{t-\varepsilon}{2} \right)^{-(\frac{3}{4}-\delta/2)} e^{-E_1 \frac{t-\varepsilon}{2}} \right]^2, \end{aligned}$$

where the last inequality follows using the Cauchy-Schwarz inequality and Theorem 7.3 have been used. This implies the assertion of the Corollary. \square

Remark 7.5. In the case of positively curved manifolds satisfying hypotheses (2.10) and (5.2), the decay rate of $\mathbb{P}_x(X_t \in B, \tau_{\Omega_0} > t)$ is exactly exponential, whereas in the situation of a flat manifold one has $t^{-1/2}e^{-E_1 t}$.

In terms of Tweedie's R -theory (see [45] and [46]) one can therefore conclude that a Brownian particle in a positively curved tube satisfying condition (2.10) and (5.1) is E_1 -positive recurrent, in a flat manifold the Brownian particle is E_1 -null recurrent and in the negatively curved tube satisfying (2.10) and (5.2) the Brownian motion is E_1 -transient.

Let us finally reformulate our findings in the negatively curved case in another way using conditional probabilities, again.

Corollary 7.3. *Assume (2.10) and (5.2). Let $x \in \Omega_0$, $\delta > 0$ and a bounded measurable subset $B \subset \Omega_0$ be given. Then there exists a constant $\tilde{C}_{B,\delta,x} > 0$ such that*

$$\mathbb{P}_x(X_t \in B \mid \tau_{\Omega_0} > t) \leq \tilde{C}_{B,\delta,x} t^{-1+\delta}$$

Proof. This follows directly from Corollary 7.2 together with the fact that for a suitable constant c_x

$$\mathbb{P}_x(\tau_{\Omega_0} > t) \geq c_x e^{-\lambda_{\kappa} t} t^{-\frac{1}{2}}.$$

The latter assertion can be proved by adding a Dirichlet boundary as was done in the proof of Theorem 7.3. \square

8 Conclusions

The objective of this paper was to investigate the interplay between the curvature and the properties of Brownian motion in the simplest non-trivial case, when the ambient space is two-dimensional and the motion in fact quasi-one-dimensional. More precisely, we were interested in the large time behaviour of the solution to the heat equation in tubular neighbourhoods of unbounded geodesics in a two-dimensional Riemannian manifold, subject to Dirichlet boundary conditions.

Our results are schematically summarized in Table 1. The corresponding precise statements can be found in: Propositions 4.1 and 4.2 for flat manifolds; Corollary 6.1 for positively curved manifolds; and Theorem 7.3 for negatively curved manifolds. The moral of the story is that the negative curvature is 'better for travelling', in the sense that the heat semigroup gains an extra polynomial, geometrically induced decay rate. The latter is in fact a consequence of the existence of Hardy-type inequalities in negatively curved manifolds, which play a central role in our proof. Though the proofs are mainly analytic some effort has been made in order to connect our findings with notions and results available in the probabilistic literature, *e.g.* on Markov chains.

The present paper can be considered as a contribution to recent works on the consequences of the existence of Hardy inequality on large-time behaviour of the heat semigroup for quantum waveguides [28, 29, 15, 22] and magnetic Schrödinger operators [23, 26]. More generally, recall that we expect that there is always an improvement of the decay rate for the heat semigroup of an operator satisfying a Hardy-type inequality (*cf* [28, Conjecture in Sec. 6] and [9, Conjecture 1]). The present paper confirms the general conjecture in the particular case of the Dirichlet Laplace-Beltrami operator in the strip-like surfaces.

As pointed out in the body of the paper, the Hardy inequality is essentially equivalent to transience properties. Thus it is reasonable to expect that a combination of available probabilistic and analytic methods might be necessary in order to make progress towards a solution of the above mentioned conjectures.

Open Problem: One of the characteristic hypotheses of the present paper was that the curvature K has compact support. We expect the same decay rates if this assumption is replaced by its fast decay at infinity. However, it is quite possible that a slow decay of curvature at infinity will improve the decay of the heat semigroup even further. In particular, can Γ_K be strictly greater than $3/4$ if K decays to zero very slowly at infinity?

Acknowledgement

M.K. would like to thank Laurent Saloff-Coste for clarifying remarks concerning his work [41] and the continuity of transition kernels. This work has been partially supported by the Czech Ministry of Education, Youth, and Sports within the project LC06002 and by the GACR grant No. P203/11/0701.

References

- [1] H. Brézis, *Analyse fonctionnelle: Théorie et applications*, Dunod, 2002.
- [2] E. B. Davies, *Spectral theory and differential operators*, Camb. Univ Press, Cambridge, 1995.
- [3] E. B. Davies, *L^p spectral independence and L^1 analyticity*, J. London Math. Soc. **52** (1995), 177–184.
- [4] Y. Dermenjian, M. Durand, and V. Iftimie, *Spectral analysis of an acoustic multistratified perturbed cylinder*, Commun. in Partial Differential Equations **23** (1998), no. 1&2, 141–169.
- [5] G. Duro and E. Zuazua, *Large time behavior for convection-diffusion equations in \mathbb{R}^N with asymptotically constant diffusion*, Commun. in Partial Differential Equations **24** (1999), 1283–1340.
- [6] T. Ekholm, H. Kovařík, and D. Krejčířík, *A Hardy inequality in twisted waveguides*, Arch. Ration. Mech. Anal. **188** (2008), 245–264.
- [7] M. Escobedo and O. Kavian, *Variational problems related to self-similar solutions of the heat equation*, Nonlinear Anal.-Theor. **11** (1987), 1103–1133.
- [8] M. Escobedo and E. Zuazua, *Large time behavior for convection-diffusion equations in R^N* , J. Funct. Anal. **100** (1991), 119–161.
- [9] M. Frass, D. Krejčířík, and Y. Pinchover, *On some strong ratio limit theorems for heat kernels*, Discrete Contin. Dynam. Systems A **28** (2010), 495–509.
- [10] A. Gray, *Tubes*, Addison-Wesley Publishing Company, New York, 1990.
- [11] D. J. Griffiths, *Introduction to quantum mechanics*, Prentice Hall, Upper Saddle River, NJ, 1995.

- [12] A. Grigoryan, *Analytic and geometric background of recurrence and non-explosion of the Brownian motion on Riemannian manifolds*, Bull. Amer. Math. Soc. (N.S.) **36** (1999), 135–249.
- [13] A. Grigor'yan and L. Saloff-Coste, *Hitting probabilities for Brownian motion on Riemannian manifolds*, J. Math. Pures Appl. **81** (2002), no. 2, 115142.
- [14] A. Grigor'yan, *Heat kernel and analysis on manifolds*, AMS/IP Studies in Advanced Mathematics, 47. American Mathematical Society, Providence, RI; International Press, Boston, MA, 2009.
- [15] G. Grillo, H. Kovařík and Y. Pinchover, *Sharp two-sided heat kernel estimates of twisted tubes and applications*, submitted; preprint on arXiv:1105.0842v1 [math.AP] (2011).
- [16] P. Hartman, *Geodesic parallel coordinates in the large*, American Journal of Mathematics **86** (1964), 705–727.
- [17] R. Hempel and J. Voigt, *On the L_p -spectrum of Schrödinger operators*, J. Math. Anal. Appl. **121** (1987), 138–159.
- [18] E. P. Hsu and G. Qin, *Volume growth and escape rate of Brownian motion on a complete manifold*, Ann. Probab. **38** (2010), 1570–1582.
- [19] T. Kato, *Perturbation theory for linear operators*, Springer-Verlag, Berlin, 1966.
- [20] W. S. Kendall, *Brownian motion, negative curvature, and harmonic maps*, Stochastic integrals (Proc. Sympos., Univ. Durham, Durham, 1980) Lecture Notes in Math., 851, 479–491, Springer, Berlin, 1981,
- [21] W. S. Kendall, *Stochastic differential geometry: an introduction*. Acta Appl. Math. **9** (1987), 29–60.
- [22] M. Kolb and A. Wübker, *Brownian motion in twisted Dirichlet-Neumann tubes*, preprint (2011), submitted
- [23] H. Kovařík, *Heat kernels of two dimensional magnetic Schrödinger and Pauli operators*, Calc. Var. Partial Differential Equations, to appear; preprint on arXiv:1007.1851v1 [math-ph] (2010).
- [24] D. Krejčířík, *Quantum strips on surfaces*, J. Geom. Phys. **45** (2003), no. 1–2, 203–217.
- [25] ———, *Hardy inequalities in strips on ruled surfaces*, J. Inequal. Appl. **2006** (2006), Article ID 46409, 10 pages.
- [26] ———, *The improved decay rate for the heat semigroup with local magnetic field in the plane*, Calc. Var. Partial Differential Equations, to appear; preprint on arXiv:1101.1806 [math.AP] (2011).
- [27] D. Krejčířík and J. Kříž, *On the spectrum of curved quantum waveguides*, Publ. RIMS, Kyoto University **41** (2005), no. 3, 757–791.
- [28] D. Krejčířík and E. Zuazua, *The Hardy inequality and the heat equation in twisted tubes*, J. Math. Pures Appl. **94** (2010), 277–303.
- [29] ———, *The asymptotic behaviour of the heat equation in a twisted Dirichlet-Neumann waveguide*, J. Differential Equations **250** (2011), 2334–2346.

- [30] I. McGillivray and E. M. Ouhabaz, *Some spectral properties of recurrent semigroups*, Arch. Math. **66** (1996), 233–242
- [31] R. Neel, *The small time asymptotics of the heat kernel at the cut locus*, Comm. Anal. Geom. **15** (2007), 845–890
- [32] J. R. Norris, *Long-Time Behavior of Heat Flow: Global Estimates and exact Asymptotics*, Arch. Rational Mech. Anal. **140** (1997), 161–195
- [33] E.-M. Ouhabaz, *Invariance of closed convex sets and domination criteria for semigroups*, Potential Anal. **5** (1996), 611–625.
- [34] Y. Pinchover, *Large time behavior of the heat kernel*, J. Funct. Anal. **206** (2004), 191–209.
- [35] Y. Pinchover, *Topics in the theory of positive solutions of second-order elliptic and parabolic partial differential equations*, Spectral Theory and Mathematical Physics: A Festschrift in Honor of Barry Simon’s 60th Birthday (F. Gesztesy, et al., ed.), Proc. Sympos. Pure Math., vol. 76, Amer. Math. Soc., Providence, RI, 2007, pp. 329–356.
- [36] R. G. Pinsky, *Positive harmonic functions and diffusion*, Cambridge Studies in Advanced Mathematics, 45. Cambridge University Press, Cambridge, 1995. xvi+474 pp.
- [37] B. Simon, *Large time behavior of the heat kernel: on a theorem of Chavel and Karp*, Proc. Amer. Math. Soc. **118** (1993), 513–514.
- [38] B. Simon, *Brownian motion, L^p properties of Schrödinger operators and the localization of binding*, J. Funct. Anal. **35** (1980), 215–229.
- [39] B. Simon, *Large Time Behavior of the L^p Norm of Schrödinger Semigroups*, J. Funct. Anal. **40** (1981), 66–83
- [40] M. Spivak, *A comprehensive introduction to differential geometry*, vol. IV, Publish or Perish, Boston, Mass., 1975.
- [41] L. Saloff-Coste, *Uniformly elliptic operators on Riemannian manifolds*, J. Differential Geometry, **36** (1992), 417–450
- [42] K. T. Sturm, *Analysis on local Dirichlet Spaces – I. Recurrence, conservativeness and L^p -Liouville*, J. reine angew. Math. **456** (1994), 173–196
- [43] K. T. Sturm, *Analysis on local Dirichlet Spaces – II. Upper Gaussian Estimates for the Fundamental Solutions of Parabolic Equations*, Osaka J. Math **32** (1995), 275–312
- [44] P. Tuominen and R. L. Tweedie, *Exponential Decay and Ergodicity of General Markov Processes and Their Discrete Skeletons*, Adv. in Appl. Probab. **11** (1979), 784–803
- [45] R. L. Tweedie, *R-theory for Markov chains on a general state space. II. R-subinvariant measures for R-transient chains*. Ann. Probability **2** (1974), 865–878.
- [46] R. L. Tweedie, *R-theory for Markov chains on a general state space. I. Solidarity properties and R-recurrent chains*. Ann. Probability **2** (1974), 840–864.
- [47] J. Weidmann, *Lineare Operatoren im Hilbertraum: Teil 1 Grundlagen*, Teubner, Stuttgart, 2000