

STX1A and Asperger syndrome: a replication study

Article

Published Version

Creative Commons: Attribution 3.0 (CC-BY)

Durdiaková, J., Warrier, V., Banerjee-Basu, S., Baron-Cohen, S. and Chakrabarti, B. ORCID: <https://orcid.org/0000-0002-6649-7895> (2014) STX1A and Asperger syndrome: a replication study. *Molecular Autism*, 5 (1). 14. ISSN 2040-2392 doi: <https://doi.org/10.1186/2040-2392-5-14> Available at <https://centaur.reading.ac.uk/36091/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

Published version at: <http://dx.doi.org/10.1186/2040-2392-5-14>

To link to this article DOI: <http://dx.doi.org/10.1186/2040-2392-5-14>

Publisher: Springer

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online



RESEARCH

Open Access

STX1A and Asperger syndrome: a replication study

Jaroslava Durdiaková¹, Varun Warriar¹, Sharmila Banerjee-Basu⁴, Simon Baron-Cohen^{1,2*} and Bhismadev Chakrabarti^{1,3*}

Abstract

Background: Autism spectrum conditions (ASC) are a group of conditions characterized by difficulties in communication and social interaction, alongside unusually narrow interests and repetitive, stereotyped behaviour. Genetic association and expression studies have suggested an important role for the GABAergic circuits in ASC. Syntaxin 1A (*STX1A*) encodes a protein involved in regulation of serotonergic and GABAergic systems and its expression is altered in autism.

Methods: In this study, the association between three single nucleotide polymorphisms (SNPs) (rs4717806, rs941298 and rs6951030) in *STX1A* gene and Asperger syndrome (AS) were tested in 650 controls and 479 individuals with AS, all of Caucasian ancestry.

Results: rs4717806 ($P = 0.00334$) and rs941298 ($P = 0.01741$) showed a significant association with AS, replicating previous results. Both SNPs putatively alter transcription factor binding sites both directly and through other variants in high linkage disequilibrium.

Conclusions: The current study confirms the role of *STX1A* as an important candidate gene in ASC. The exact molecular mechanisms through which *STX1A* contributes to the etiology remain to be elucidated.

Keywords: Asperger syndrome, Serotonergic system, Single nucleotide polymorphism, *STX1A*

Background

Autism spectrum conditions (ASC) are characterized by difficulties in social interaction and impaired communication, alongside unusually narrow interests and repetitive, stereotyped behaviour (DSM-IV). Asperger syndrome (AS), a subcategory of ASC, differs from other autism conditions by lack of any delay in language and cognitive development [1]. ASC has a prevalence of 1 in 100 [2] and heritability of around 80% [3]. Several genes involved in neural transmission, and in particular GABAergic neurotransmission, have been implicated in ASC [4-7].

Syntaxin 1A (*STX1A*) is an important component of the GABAergic system, due to its involvement in the transport of glutamate and its conversion to gamma-aminobutyric

acid (GABA) [8]. The protein *STX1A* is expressed pre-synaptically and is involved in neurotransmitter release and synaptic vesicle docking and thus is an essential factor for neurotransmission. It also interacts with serotonin and alters its subcellular localization and expression [9]. Atypical function in both serotonergic and GABAergic neurotransmission systems has been reported in autism [10]. GABAergic signaling during embryonic development contributes to the excitatory/inhibitory signaling balance in neural networks. Dysregulation of GABAergic system during development is thought to account for some of the sensory sensitivity in people with ASC [6].

The gene *STX1A* consists of 10 exons (Figure 1) spanning a 20.42 kb region (<http://www.genome.ucsc.edu>). It lies on 7q11.23, a locus that has been reported to be duplicated in some individuals with autism, and associated with deletions in Williams syndrome [11]. A possible role for *STX1A* in the development of autism has previously been suggested. Three single nucleotide polymorphisms (SNPs) (rs2293485 and rs4717806 in the Caucasian population,

* Correspondence: sb205@cam.ac.uk; b.chakrabarti@reading.ac.uk

¹Autism Research Centre, Department of Psychiatry, University of Cambridge, 18b Trumpington Road, Cambridge CB2 8AH, UK

³Centre for Integrative Neuroscience and Neurodynamics, School of Psychology and Clinical Language Sciences, University of Reading, Reading RG6 6AL, UK
Full list of author information is available at the end of the article




Figure 1 Genomic structure of *STX1A* and the location of tested SNPs. Exons are indicated by boxes; introns are indicated by lines; SNP positions are denoted by arrows (<http://www.genome.ucsc.edu>).

and rs6951030 in the Japanese population) are nominally associated with high functioning autism (HFA) [12,13]. rs4717806 has also been related to early developmental differences in people with autism [12]. Another SNP, rs1569061, is nominally associated with ASC in a Caucasian population [11]. Additionally, *STX1A* mRNA expression is higher in lymphocytes of people with HFA [12], and lower in the anterior cingulate gyrus in post-mortem brain tissue [13], compared to matched controls. This makes *STX1A* an interesting candidate gene for ASC. In this study, we examined the association of three SNPs in *STX1A* with AS, in a Caucasian sample, in an effort to replicate previous findings.

Methods

All individuals enrolled in the current study were adults of Caucasian origin from the same geographic region (the United Kingdom). Participants were asked to complete an online version of the Autism Spectrum Quotient (AQ) test [14], which is a measure of autistic traits. A score of 32 and above is an excellent predictor of ASC [15], and the mean AQ score in the general population is 16.4 (SD = 6.3) [14]. Individuals with a clinical diagnosis of AS ($n = 479$; 97 females, 382 males) took part and achieved a mean AQ score of 35.8 ± 8.3 . The control group comprised participants ($n = 650$; 367 females, 284 males) chosen to have an AQ score below 24, to ensure a balanced representation of individuals from both ends of the autistic trait continuum. The mean AQ score within the control population was 14.8 ± 5.2 . Consent was obtained from all the participants. This study was approved by the University of Cambridge Psychology Research Ethics Committee and the NHS National Research Ethics Service.

Based on previously published data, three SNPs (rs4717806, rs941298, rs6951030) were selected and analyzed in this study. rs4717806 and rs6951030 have been reported to be nominally associated with HFA in the Caucasian and Japanese population, respectively [12,13]. The third SNP, rs941298, is in high linkage disequilibrium (LD) with rs4717806 ($R^2 = 0.96$) and in moderate LD with an exonic SNP, rs2293485 ($R^2 = 0.67$), which is nominally associated with HFA in the Caucasian population [12]. LD values between SNPs of interest in the HapMap CEPH European samples of the Utah Residents with Northern and Western European Ancestry (CEU) population data were calculated using SNAP ([\[broadinstitute.org/mpg/snap/\]\(http://www.broadinstitute.org/mpg/snap/\)\). The minor allele frequency \(MAF\) for the tested SNPs was above 0.05 in the CEPH CEU population as calculated from the dbSNP database \(<http://www.ncbi.nlm.nih.gov/projects/SNP/>\). LD plots for the sample studied were created using Haploview \[16\].](http://www.</p></div><div data-bbox=)

DNA was extracted from buccal swabs and anonymized. SNP genotyping was performed using TaqMan SNP Genotyping Assays (Applied Biosystems Inc., CA, USA) according to the previously described protocol [17]. Allelic association testing was performed using Plink v1.07 (<http://pngu.mgh.harvard.edu/~purcell/plink/>) [18]. Bonferroni correction was performed to correct for multiple SNPs tested after evaluating the number of completely independent SNPs and determination of a new alpha value using SNPSpD (<http://gump.qimr.edu.au/general/daleN/SNPSpD/>) [19]. This was done as Bonferroni correction is over-corrective when the SNPs are in LD. P values were reported as significant if they were below the alpha threshold.

Functional annotation was performed using Haploreg (<http://www.broadinstitute.org/mammals/haploreg/haploreg.php>) [20], SNPnexus (<http://snp-nexus.org/>) [21], and F-SNP (<http://compbio.cs.queensu.ca/F-SNP/>) [22].

Results

In the association analyses, rs4717806 ($P = 0.003$) and rs941298 ($P = 0.017$) showed significant association with AS. These results remained significant after correcting for multiple testing. The estimated average number of independent loci was 2. The SNPSpD determined threshold P value was 0.023. None of the SNPs deviated from the Hardy-Weinberg equilibrium at $P < 0.05$. Genotyping rate was above 98%. The test statistics are summarized in Table 1. In our sample, all three SNPs are part of a common LD block as calculated by Haploview (Figure 2).

Discussion

In this study, three SNPs in *STX1A* were tested for association with AS to replicate findings previously reported in the literature. We found that two of these SNPs (rs4717806 and rs941298) were significantly associated with a clinical diagnosis of AS. The odds ratio for the two significant SNPs was below 1, indicating the protective role of the minor allele. rs4717806 and rs6951030 have been reported to be nominally associated with HFA in previous reports [12,13]. The direction of effect is in agreement

Table 1 Single SNP association analyses

dbSNp ID	Alleles ^a	MAF ^b	F_A ^c	F_U ^d	CHISQ ^e	Odds ratio	P value ^f	Alpha ^g
rs6951030	T/G	0.18	0.19	0.17	2.40	1.24	0.06	0.023
rs4717806	T/A	0.33	0.29	0.36	9.49	0.76	0.003	0.023
rs941298	G/A	0.33	0.29	0.35	6.09	0.80	0.017	0.023

^aCommon allele is listed first.

^bCalculated by Plink v1.07 in analysed sample.

^cThe frequency of this variant in cases.

^dThe frequency of this variant in controls.

^eThe χ^2 statistic for this test (1 df).

^fComputed on the basis of likelihood ratio test.

^gDetermined after evaluating the number of completely independent SNPs using SNPSpD.

with our findings. Both SNPs have been under-transmitted from non-affected parents to affected probands. rs941298 is in high LD with rs4717806 ($R^2 = 0.96$) and in moderate LD with exonic rs2293485 ($R^2 = 0.67$), which is also nominally associated with HFA [12].

We identified significant association between rs4717806 and AS, supporting previously published data [12]. rs941298 was also significantly associated with AS in the

current study. We were not able to replicate the nominal association between rs69510130 and HFA [13] in our sample. However, rs69510130 was nominally significant in a Japanese sample in the previous study, whereas we tested for association in a Caucasian sample.

Both SNPs (rs4717806, rs941298) that showed significant association with AS in our study are intronic SNPs. rs4717806 lies in intron 9 and rs941298 lies in intron 2; they are 8,786 bp apart, in high LD in our sample ($R^2 = 0.92$), and part of a common LD block as calculated by Haploview (Figure 2). Both the SNPs alter various transcription binding sites both directly and through other SNPs in LD (<http://www.broadinstitute.org/mammals/haploreg/>). There are no copy number polymorphisms or miRNA binding sites associated with either of the SNPs (<http://www.snp-nexus.org/>). rs4717806 is highly conserved across species according to F-SNP. It is likely that the expression of *STXA1* is regulated by these SNPs along with additional cis- and trans-acting factors.

While this study replicates the results from previous studies [12,13], it is different in that we focused on AS, rather than HFA or classic autism. Further, the previous studies used Transmission Disequilibrium Tests to identify over-transmitted alleles in family trios of individuals with ASC and HFA [12]. The current study design is population based and with a moderate sample size; hence, these results need to be replicated in a larger sample. It is worth mentioning that females are overrepresented in the control sample compared to the AS group. Men typically score higher than women in the self-reported AQ test [14], which may account for the lower AQ score describing the control population. In the sex-stratified analyses, none of the SNPs are significant after Bonferroni correction indicating that the over-representation of females in controls is not driving the association (data not shown).

Additionally, only three SNPs were tested, chosen specifically to replicate previous results. The high LD between rs4717806 and rs941298 must be taken into account when interpreting the results. It is likely that both the SNPs are tagging one single causative locus, which is reflected in the significant *P* values of the two SNPs. rs941298 would not remain significant if traditional Bonferroni correction




Figure 2 LD structure of *STXA1* for AS case-control dataset.

LD structure of the region investigated in the AS case-control dataset, as plotted on Haplotype is shown. Three SNPs which were genotyped have been plotted; they fall into one LD block.

was applied. A single gene is unlikely to have a major effect in complex conditions like autism, and many other genes are likely to contribute to the phenotype. This study, in combination with previous studies, provides evidence for *STX1A* as involved in the etiology of ASC. Further cellular, molecular, and neurocognitive studies are required to elucidate the role of this gene in ASC.

Conclusions

rs4717806 ($P = 0.00334$) and rs941298 ($P = 0.01741$) showed significant association with AS, replicating findings in previous studies in the same direction. Both SNPs putatively alter transcription factor binding sites both directly and through other variants in high linkage disequilibrium. The current study confirms the role of *STX1A* as an important candidate gene in ASC. The exact molecular mechanisms through which *STX1A* contributes to the etiology remain to be elucidated.

Abbreviations

AQ: Autism spectrum quotient; AS: Asperger syndrome; ASC: Autism spectrum conditions; CEU: European samples of Utah Residents with Northern and Western European Ancestry, GABA, Gamma-aminobutyric acid; HFA: High-functioning autism; LD: Linkage disequilibrium; MAF: Minor allele frequency; SNP: Single nucleotide polymorphism; *STX1A*: Syntaxin 1A.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

BC and SBC co-designed the study. SBC obtained funding for the study. SBB suggested the mapping of *STX1A* as a relevant candidate gene. VW and JD conducted the analysis. JD wrote the first draft of the paper revised by all authors. All authors read and approved the final manuscript.

Acknowledgements

This study was funded by grants to SBC by Target Autism Genome, the Autism Research Trust, the MRC UK, and the Max Planck Institute for Psycholinguistics, Nijmegen. We are grateful to Jonathan Breidbord, Allen Chan, Agnese Di Napoli, Laura Murphy, and Simon Fisher for help at various stages of the project.

Author details

¹Autism Research Centre, Department of Psychiatry, University of Cambridge, 18b Trumpington Road, Cambridge CB2 8AH, UK. ²CLASS Clinic, Cambridgeshire and Peterborough NHS Foundation Trust (CPFT), Fulbourn, Cambridge CB21 5EF, UK. ³Centre for Integrative Neuroscience and Neurodynamics, School of Psychology and Clinical Language Sciences, University of Reading, Reading RG6 6AL, UK. ⁴Mindspec Inc, 9656 Blake Ln, Fairfax, VA 22031, USA.

Received: 11 December 2013 Accepted: 5 February 2014

Published: 18 February 2014

References

1. McPartland J, Klin A: **Asperger's syndrome**. *Adolesc Med Clin* 2006, **17**(3):771–788.
2. Baron-Cohen S, Scott FJ, Allison C, Williams J, Bolton P, Matthews FE, Brayne C: **Prevalence of autism-spectrum conditions: UK school-based population study**. *Br J Psychiatry* 2009, **194**(6):500–509.
3. Klei L, Sanders SJ, Murtha MT, Hus V, Lowe JK, Willsey AJ, Moreno-De-Luca D, Yu TW, Fombonne E, Geschwind D, Grice DE, Ledbetter DH, Lord C, Mane SM, Martin CL, Martin DM, Morrow EM, Walsh CA, Melhem NM, Chaste P, Sutcliffe JS, State MW, Cook EH Jr, Roeder K, Devlin B: **Common genetic variants, acting additively, are a major source of risk for autism**. *Mol Autism* 2012, **3**(1):9.

4. Buxbaum JD, Silverman JM, Smith CJ, Greenberg DA, Kilifarski M, Reichert J, Cook EH Jr, Fang Y, Song CY, Vitale R: **Association between a GABRB3 polymorphism and autism**. *Mol Psychiatry* 2002, **7**(3):311–316.
5. Fatemi SH, Reutiman TJ, Folsom TD, Thuras PD: **GABA(A) receptor downregulation in brains of subjects with autism**. *J Autism Dev Disord* 2009, **39**(2):223–230.
6. Pizzarelli R, Cherubini E: **Alterations of GABAergic signaling in autism spectrum disorders**. *Neural Plast* 2011, **2011**:297153.
7. Tavassoli T, Auyeung B, Murphy LC, Baron-Cohen S, Chakrabarti B: **Variation in the autism candidate gene GABRB3 modulates tactile sensitivity in typically developing children**. *Mol Autism* 2012, **3**(1):6.
8. Yu YX, Shen L, Xia P, Tang YW, Bao L, Pei G: **Syntaxin 1A promotes the endocytic sorting of EAAC1 leading to inhibition of glutamate transport**. *J Cell Sci* 2006, **119**(Pt 18):3776–3787.
9. Haase J, Killian AM, Magnani F, Williams C: **Regulation of the serotonin transporter by interacting proteins**. *Biochem Soc Trans* 2001, **29**(Pt 6):722–728.
10. Coghill S, Horder J, Inkster B, Mendez MA, Murphy DG, Nutt DJ: **GABA system dysfunction in autism and related disorders: from synapse to symptoms**. *Neurosci Biobehav Rev* 2012, **36**(9):2044–2055.
11. Malenfant P, Liu X, Hudson ML, Qiao Y, Hrynychak M, Riendeau N, Hildebrand MJ, Cohen IL, Chudley AE, Forster-Gibson C, Mickelson EC, Rajcan-Separovic E, Lewis ME, Holden JJ: **Association of GTF2i in the Williams-Beuren syndrome critical region with autism spectrum disorders**. *J Autism Dev Disord* 2012, **42**(7):1459–1469.
12. Nakamura K, Anitha A, Yamada K, Tsujii M, Iwayama Y, Hattori E, Toyota T, Suda S, Takei N, Iwata Y, Suzuki K, Matsuzaki H, Kawai M, Sekine Y, Tsuchiya KJ, Sugihara G, Ouchi Y, Sugiyama T, Yoshikawa T, Mori N: **Genetic and expression analyses reveal elevated expression of syntaxin 1A (STX1A) in high functioning autism**. *Int J Neuropsychopharmacol* 2008, **11**(8):1073–1084.
13. Nakamura K, Iwata Y, Anitha A, Miyachi T, Toyota T, Yamada S, Tsujii M, Tsuchiya KJ, Iwayama Y, Yamada K, Hattori E, Matsuzaki H, Matsumoto K, Suzuki K, Suda S, Takebayashi K, Takei N, Ichikawa H, Sugiyama T, Yoshikawa T, Mori N: **Replication study of Japanese cohorts supports the role of STX1A in autism susceptibility**. *Prog Neuropsychopharmacol Biol Psychiatry* 2011, **35**(2):454–458.
14. Baron-Cohen S, Wheelwright S, Skinner R, Martin J, Clubley E: **The autism-spectrum quotient (AQ): evidence from Asperger syndrome/high-functioning autism, males and females, scientists and mathematicians**. *J Autism Dev Disord* 2001, **31**(1):5–17.
15. Woodbury-Smith MR, Robinson J, Wheelwright S, Baron-Cohen S: **Screening adults for Asperger Syndrome using the AQ: a preliminary study of its diagnostic validity in clinical practice**. *J Autism Dev Disord* 2005, **35**(3):331–335.
16. Barrett JC, Fry B, Maller J, Daly MJ: **Haploview: analysis and visualization of LD and haplotype maps**. *Bioinformatics* 2005, **21**(2):263–265.
17. Chakrabarti B, Dudbridge F, Kent L, Wheelwright S, Hill-Cawthorne G, Allison C, Banerjee-Basu S, Baron-Cohen S: **Genes related to sex steroids, neural growth, and social-emotional behavior are associated with autistic traits, empathy, and Asperger syndrome**. *Autism Res* 2009, **2**(3):157–177.
18. Purcell S, Neale B, Todd-Brown K, Thomas L, Ferreira MA, Bender D, Maller J, Sklar P, de Bakker PI, Daly MJ, Sham PC: **PLINK: a tool set for whole-genome association and population-based linkage analyses**. *Am J Hum Genet* 2007, **81**(3):559–575.
19. Nyholt DR: **A simple correction for multiple testing for single-nucleotide polymorphisms in linkage disequilibrium with each other**. *Am J Hum Genet* 2004, **74**(4):765–769.
20. Ward LD, Kellis M: **HaploReg: a resource for exploring chromatin states, conservation, and regulatory motif alterations within sets of genetically linked variants**. *Nucleic Acids Res* 2012, **40**(Database issue):D930–934.
21. Chelala C, Khan A, Lemoine NR: **SNPNexus: a web database for functional annotation of newly discovered and public domain single nucleotide polymorphisms**. *Bioinformatics* 2009, **25**(5):655–661.
22. Lee PH, Shatkay H: **An integrative scoring system for ranking SNPs by their potential deleterious effects**. *Bioinformatics* 2009, **25**(8):1048–1055.

doi:10.1186/2040-2392-5-14

Cite this article as: Durdiaková et al.: *STX1A* and Asperger syndrome: a replication study. *Molecular Autism* 2014 **5**:14.