Size and molecular flexibility affect the binding of ellagitannins to bovine serum albumin

Article

Accepted Version

Dobreva, M., Green, R.J., Mueller-Harvey, I., Salminen, J.-P., Howlin, B.J. and Frazier, R. (2014) Size and molecular flexibility affect the binding of ellagitannins to bovine serum albumin. *Journal of Agricultural and Food Chemistry*, 62 (37). pp. 9186-9194. ISSN 0021-8561 doi: <https://doi.org/10.1021/jf502174r> Available at <https://centaur.reading.ac.uk/37617/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

To link to this article DOI: <http://dx.doi.org/10.1021/jf502174r>

Publisher: American Chemical Society

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online

Size and Molecular Flexibility affect the Binding of Ellagitannins to Bovine Serum Albumin

Marina A. Dobрева,[†] Rebecca J. Green,^{*,‡} Irene Mueller-Harvey,^{*,†} Juha-Pekka Salminen,[§] Brendan J. Howlin,[#] Richard A. Frazier[‡]

[†]School of Agriculture, Policy and Development, University of Reading, Earley Gate, PO Box 236, Reading, RG6 6AT, UK;

[‡]School of Chemistry, Food and Pharmacy, University of Reading, Whiteknights, PO Box 224, Reading, RG6 6AP, UK;

[§]Department of Chemistry, University of Turku, Vatselankatu 2, Turun Yliopisto, Turku, FI-20014, Finland.

[#]Chemistry Department, FEPS, University of Surrey, Guildford, Surrey, GU2 7XH, UK.

AUTHOR INFORMATION:

*Phone: +44-118-378-8446. Fax: +44-118-378-6562.

E-mail: rebecca.green@reading.ac.uk.

*Phone: +44-118-378-6619. Fax: +44-118-935-2421.

E-mail: i.mueller-harvey@reading.ac.uk.

ABSTRACT: Binding to bovine serum albumin of monomeric (vescalagin and pedunculagin) and dimeric ellagitannins (roburin A, oenothetin B and gemitin A) was investigated by isothermal titration calorimetry and fluorescence spectroscopy, which indicated two types of binding sites. Stronger and more specific sites exhibited affinity constants, K_1 , of 10^4 to 10^6 M⁻¹ and stoichiometries, n_1 , of 2 to 13 and dominated at low tannin concentrations. Weaker and less-specific binding sites had K_2 -constants of 10^3 to 10^5 M⁻¹ and stoichiometries, n_2 , of 16 to 30, and dominated at higher tannin concentrations. Binding to stronger sites appeared dependent on tannin flexibility and presence of free galloyl groups. Positive entropies for all but gemitin A indicated that hydrophobic interactions dominated during complexation. This was supported by an exponential relationship between the affinity, K_1 , and the modeled, hydrophobic accessible surface area and by a linear relationship between K_1 and the Stern-Volmer quenching constant, K_{SV} .

KEYWORDS: *ellagitannins, isothermal titration calorimetry, fluorescence, molecular flexibility, molecular size, hydrophobic accessible surface area*

INTRODUCTION

Many plants, herbal drugs and foods contain not only the widely-studied condensed tannins but also ellagitannins (ETs).^{1,2} In fact, ellagitannins are much more common than previously recognized (Salminen – unpublished data). We consume ellagitannins in nuts, berries, fruit juices and wines³ and interest is growing in the medicinal and health effects of these polyphenols.³⁻⁸

Dietary polyphenols may prevent diseases through their ability to quench free radicals, which are generated under oxidative stress.³ Whilst ETs that contain hexahydroxydiphenoyl (HHDP) groups are rapidly metabolized into ellagic acid and urolithins after consumption, C-glucosidic ETs contain a nonahydroxytriphenoyl (NHTP) group and are difficult to hydrolyse.⁵ Pedunculagin and gemin A are examples of HHDP-containing ETs and vescalagin and roburin A contain NHTP group(s) (Figure 1). Many ETs have high water solubility and surprisingly high bioavailability.⁹ They are also capable of interacting with various molecular targets that affect signaling pathways¹⁰ and can inhibit tumor promotion.¹¹ For example vescalagin enters cells rapidly,⁴ strongly inhibits DNA topoisomerase¹²⁻¹⁴ and interacts selectively with actin filaments.⁵ Oenothien B is one of the most active ETs in terms of host-mediated antitumor activity and generates an immune response by stimulating interleukin 1 production.¹ Oenothien B and ET metabolites such as ellagic acid and urolithins can also influence histone acetylation/deacetylation and thus inflammatory responses,¹⁵ which play an important role in the development of age-related diseases.

These and many other examples illustrate that tannins are not just non-specific protein-precipitating agents but can also be involved in specific¹⁶ and targeted interactions with certain amino acids.¹⁷ Tannin-protein interactions have been investigated by a wide range of physico-chemical techniques and were summarized previously.^{5,18,19} These included competitive binding assays, NMR spectroscopy, circular dichroism, mass spectrometry, infra-red spectroscopy, dynamic light scattering, small angle X-ray scattering, transmission electron microscopy, equilibrium dialysis, nephelometry, calorimetry, fluorescent quenching, ion mobility,²⁰ and surface plasmon resonance.²¹ Isothermal titration calorimetry (ITC) provides information not only on thermodynamic parameters and the strengths of the tannin-protein interaction but also on the stoichiometry of the resulting complex. Other work has shown that both hydrophobic effects⁵ and hydrogen bonding are involved.^{16,22} Which of these is the dominant factor depends on the precise polyphenol/protein interactions and experimental conditions.^{5,23} Both the flexibility and size of the tannins and proteins appear to affect these interactions.^{16,24,25} When gallotannin (GT)- and ET-binding to gelatin and BSA were compared, it was found that equilibrium binding constants for flexible GTs with the globular BSA were between 10^4 to 10^5 M⁻¹ and for ETs interacting with the highly flexible gelatin these were ca. 10^6 M⁻¹.¹⁹

ETs are unique amongst tannins as i) they can be isolated as pure compounds (unlike most condensed tannins, which tend to be obtained as closely related mixtures); ii) they possess a wide range of molecular flexibilities (in contrast to gallotannins or condensed tannins); and iii) their water solubilities are inversely correlated to their protein precipitation capacities.⁹ ETs, therefore, offer an opportunity for investigating

68 the relative importance of molecular flexibility and size within a series of matched
69 tannin compounds.

70

71 We chose two monomeric and three dimeric ETs that differ in molecular flexibility
72 (i.e. their bond rotational flexibility) and water solubility. Vescalagin and its dimer,
73 roburin A, represent ETs with NHTP groups that are linked to an acyclic glucose.
74 They are consumed in wine that has been stored in oak barrels. Walnuts are a rich
75 source of pedunculagin, which represents ETs with HHDP groups and a cyclic
76 glucose. The dimers, oenothien B and gemin A can be found in several European and
77 Japanese herbal medicines^{1,26} and differ in their molecular flexibilities. The water
78 solubilities of these ETs decrease in the following order: vescalagin > pedunculagin $\approx$
79 roburin A > oenothien B > gemin A.⁹

80

81 The present study investigated the effects of flexibility and size on the
82 thermodynamics of the ET-protein interaction using isothermal titration calorimetry
83 and fluorescence spectroscopy to probe their interaction mechanisms and binding
84 sites. We choose bovine serum albumin (BSA) as the protein-binding partner for two
85 reasons. Firstly, BSA is a widely used model for globular proteins such as Rubisco,
86 which is a major dietary protein.^{27,28} Recent studies have shown that the binding
87 affinity in tannin-protein complexes was negatively correlated with digestibility.²⁹⁻³¹
88 Secondly, albumins are involved in transporting dietary polyphenols^{32,33} and drugs to
89 their molecular targets,^{34,35} and exhibit a large sequence homology, *e.g.* 76% in the
90 case of BSA and human serum albumin.³⁶ It is, therefore, important to understand
91 how ellagitannins interact with globular proteins, as this may also impact on protein-
92 drug interactions in the digestive tract and in the blood plasma.³⁵

MATERIALS AND METHODS

Ellagitannins. Two monomeric ellagitannins (pedunculagin and vescalagin) and three dimeric ellagitannins (oenothetin B, roburin A and gemin A) were extracted from plants and purified as described previously.³⁷

Isothermal Titration Calorimetry. Titrations of BSA (purity $\geq 99\%$, essentially globulin free, 66 kDa; Sigma, Poole, Dorset, U.K.) with ellagitannins were performed using a NanoITC instrument (TA Instruments Ltd., Crawley, West Sussex, U.K.) as described previously^{19,28} with few adaptations. All solutions were prepared in 50 mM citrate buffer at pH 6 and were degassed under vacuum prior to use. In a typical experiment, buffered BSA solution (20 μM) was placed in the 950 μL sample cell of the calorimeter and buffered ellagitannin solution (3 mg mL^{-1}) was loaded into the injection syringe. Ellagitannin was titrated into the sample cell at 298 K as a sequence of 24 injections of 10 μL aliquots for monomeric ellagitannins (pedunculagin and vescalagin) and 48 injections of 10 μL aliquots for dimeric ellagitannins (oenothetin B, roburin A and gemin A). The time delay (to allow equilibration) between successive injections was 360 s. The contents of the sample cell were stirred throughout the experiment to ensure thorough mixing.

Raw data were obtained as plots of heat (μJ) against injection number and featured a series of peaks for each injection. These raw data peaks were transformed using the instrument software to obtain a plot of observed enthalpy change per mole of injectant (ΔH_{obs} , kJ mol^{-1}) against molar ratio (see ITC Data Analysis).

Control experiments included the titration of buffered ellagitannin solutions into buffer, buffer into BSA and buffer into buffer; controls were repeated for each BSA concentration. Control experiments of buffer titrated into buffer or protein solutions both resulted in small and equal enthalpy changes for each successive injection of buffer and, therefore, were not further considered in the data analysis.³⁸ Experimental data were analyzed after subtraction of the tannin into buffer control data from the sample data. Ellagitannin molecules tend to self-associate into aggregates due to hydrophobic groups; therefore, when injected from the syringe into buffer the ellagitannin molecules undergo an endothermic process of deaggregation, analogous to surfactant demicellization.³⁹ The extent of deaggregation depends inversely on the concentration of ellagitannin already present in the sample cell; therefore, successive injections of ellagitannin into buffer lead to the observation of progressively lower endothermic enthalpy changes as has been illustrated in earlier work.³⁹ The data are shown after subtraction of the effects of ellagitannin deaggregation, which means that the assumption is made that ellagitannins dissociate prior to binding.

ITC Data Analysis. Estimated binding parameters were obtained from ITC data using the Bindworks™ (Version 3.1.13, Applied Thermodynamics, Hunt Valley, MD, U.S.A.) data analysis program. Data fits were obtained using a model for two independent sets of multiple binding sites. For this model, the analytical solution for the total heat measured (Q) is determined by the formula:

$$Q = V[M] \left\{ \frac{n_1 \Delta H_1 K_1 [L]}{1 + K_1 [L]} + \frac{n_2 \Delta H_2 K_2 [L]}{1 + K_2 [L]} \right\} \quad (\text{Equation 1})$$

where n_1 and n_2 are the molar ratios of interacting species, ΔH_1 and ΔH_2 are the enthalpies, and K_1 and K_2 are the equilibrium binding constants (*syn.* affinity

constants) for each of the two sets of multiple binding sites.⁴⁰ The quality of fit was determined by calculation of χ^2 ; the data fits were acceptable in each case since the χ^2 values were less than the critical values for the appropriate degree of freedom.

Tryptophan Fluorescence Quenching. Fluorescence intensity was recorded using an FP-6200 spectrofluorimeter (JASCO UK Ltd., Great Dunmow, Essex, U.K.) with selective excitation of tryptophan residues in BSA at 295 nm. The excitation and emission slits were 5 nm and the emission spectrum was recorded between 300 and 500 nm. The measured fluorescence intensities are reported without correction for inner filter effects.⁴¹

Titration of BSA (5 μ M, 2 mL) with ellagitannin (3 mg mL⁻¹, 500 μ L) was carried out as a sequence of aliquots (10 μ L up to volume of 100 μ L, 20 μ L up to a volume of 460 μ L). For the calculation of quenching constants, the data were plotted as a Stern-Volmer plot of F_0/F against $[Q]$ and the quenching constant, K_{SV} , calculated by linear regression.⁴² According to the Stern-Volmer equation:

$$\frac{F_0}{F} = 1 + k_q \tau_0 [Q] = 1 + K_{SV} [Q] \quad (\text{Equation 2})$$

where, F_0 and F are the fluorescence intensities before and after the addition of the quencher, respectively, k_q is the bimolecular quenching constant, τ_0 is the lifetime of the fluorophore in the absence of quencher, $[Q]$ is the concentration of the quencher (in this case tannin) and K_{SV} is the Stern-Volmer quenching constant.

However, in the case of combined dynamic and static quenching, where the Stern-Volmer plot exhibits an upward curvature (concave towards the y-axis) an alternative equation can be used for fitting:⁴²

$$\frac{F}{F_o} = 1 + K_{app}[Q] \quad \text{(Equation 3)}$$

where,

$$K_{app} = \left[\frac{F}{F_o} - 1 \right] \frac{1}{[Q]} = (K_D + K_S) + K_D K_S [Q] \quad \text{(Equation 4)}$$

The actual plot of K_{app} or $((F/F_o-1)/Q)$ *versus* $[Q]$ yields a straight line with an intercept of K_D+K_S and a slope of $K_D K_S$.

Molecular Modeling. Molecular modeling was performed with MOE2011.10 (Chemical Computing Ltd., Montreal, Canada). Atomistic models of the respective tannins were built from chemical structures on an atom by atom basis using the Builder Module of MOE. All models were energy minimized to convergence (rms gradient of 0.05 Kcal/mol) using the MMX94x force field by smart minimization (steepest descents, followed by conjugate gradient and truncated Newton-Raphson methods). Physical parameters were calculated from these models using the QuasAR module in MOE. Parameters calculated were accessible surface area (ASA), hydrophobic accessible surface area (ASA_H). Molecular flexibility was calculated by performing a conformational energy search about the torsion angles of each model using Lowmode Molecular Dynamics in the Conformational Analysis module of MOE. Results from conformational analysis were stored in databases of conformations for each tannin indexed by final energy.

RESULTS AND DISCUSSION

This study investigated the interaction between BSA and five ellagitannin compounds using isothermal titration calorimetry and fluorescence spectroscopy. The molecular flexibility and surface areas of five ellagitannins and pentagalloyl glucose were

modelled in order to improve our understanding of these interactions.

Molecular Flexibility of Ellagitannins. Molecular simulation was used to assess molecular flexibility in terms of numbers of low energy conformations located and energy of these conformers (Table 1). Modeling experiments⁴³ demonstrated previously that the nonahydroxytriphenoyl (NHTP) group is much less flexible than the hexahydroxydiphenoyl (HHDP) group (Figure 1) and the low energy conformers of the monomers clearly reflect the extent of these structural constraints: pentagalloyl glucose conformers (n=21) ranged from 110-116 kcal mol⁻¹ (no NHTP or HHDP groups), pedunculagin conformers (n=4) from 158-164 kcal mol⁻¹ (2 HHDP groups) and vescalagin conformers (n=18) from 229 to 236 kcal mol⁻¹ (1 NHTP, 1 HHDP group). The low energy conformers of the dimers followed a similar trend: oenotherin B conformers (n=2) had 278-280 kcal mol⁻¹ (2 HHDP groups), gemin A (n=1) had 323 kcal mol⁻¹ (3 HHDP groups) and roburin A conformers (n=4) had 406-412 kcal mol⁻¹ (2 NHTP, 2 HHDP groups).

Isothermal Titration Calorimetry of Ellagitannin-BSA Interactions and Data

Fitting. ITC experiments for ellagitannin titration into BSA were carried out for five ellagitannins (depicted in Figure 1). Figure 2 shows the ITC binding isotherms for the interaction with BSA of vescalagin and its dimer, roburin A, consisting of the experimental data (as points) and the binding model that best fitted to each data set (as lines). For both ellagitannins, the binding interaction led to an exothermic response, which decreased in magnitude with successive titration injections and signifies a saturation of binding sites on BSA.

As found in previous studies,^{24,39} small but significant exothermic peaks were observed at higher tannin:protein molar ratios at and beyond the point of apparent binding site saturation. This means that titration experiments did not reach zero enthalpy change, even after removing effects from control experiments (*e.g.* tannin titration into buffer). Therefore, we have assumed that the interaction between ellagitannin and BSA has not reached completion at the higher ellagitannin:protein ratios measured. As a consequence, upon fitting the data to a binding model, this long tail of exothermic peaks prevents the use of a model based on a single set of multiple binding sites (one-site model), and, therefore, all data were fitted to a two independent binding site model. A previous study of the interaction between pentagalloyl glucose with BSA or Rubisco described the selection of the two independent binding site model in more detail.²⁸

Figure 3 shows the ITC binding isotherms for pedunculagin and the related dimers oenothetin B and gemin A. As observed for the vescalagin and roburin A isotherms, the binding isotherm for the dimers shows a clear two-phase shape to the isotherm that signifies two or more binding sites are involved in the interaction. For the oenothetin B and gemin A ellagitannins, the titration experiments were extended by increasing the number of titration injections to observe the interaction at higher molar ratios (see Methods for more details). The thermodynamic binding parameters for the ellagitannin-BSA interaction were obtained from a two-site model and are shown in Table 2, which also includes data from our previous study on pentagalloyl glucose binding to BSA for comparison.²⁸

Thermodynamic Parameters of Ellagitannin-BSA Interactions in Relation to

Molecular Flexibility. In these studies, the excellent fits of the ITC data have enabled us to probe the effects of different structural characteristics of the ellagitannins on their interaction with BSA. The results will be discussed in terms of monomer-dimer pairs and molecular flexibilities. The weakness of the second binding site allows, however, for some uncertainties in exact values for these fits, but does confirm our previous observations, which included related ellagitannins and suggested that these were able to bind (adsorb) non-selectively.^{24,28,39}

The ellagitannin monomers, pendunculagin and vescalagin, show equivalent strengths of interaction ($K_1 = 4.2$ and $5.2 \times 10^4 \text{ M}^{-1}$) and so do the roburin A and oenothien B dimers ($K_1 = 2.3$ and $6.5 \times 10^5 \text{ M}^{-1}$) (Table 2). In contrast, the K_1 affinity constants of pentagalloyl glucose and gemin A were one order of magnitude larger than for the other monomers and dimers ($K_1 = 1.8 \times 10^5$ and $1.8 \times 10^6 \text{ M}^{-1}$, respectively). This coincides with the fact, as pointed out above that both have low conformational energies and are flexible molecules. Taken together gemin A has several features, *i.e.* single bonds and ‘free’ galloyl groups, which may allow selective binding to specific amino acids on the protein surface. These findings are in line with two other studies that reported weaker binding by vescalagin⁴⁴ and by its isomer, castalagin,²⁹ to BSA compared to pentagalloyl glucose. It appears that pentagalloyl glucose and gemin A have optimal structures for BSA binding within the group of tannins studied here.

For both binding sites the interaction in terms of the affinity constants, K_1 and K_2 , is approximately one order of magnitude different between monomers and dimers (Table 2). For each of the comparable ellagitannin monomer-dimer pairs, the dimer was shown to bind more strongly than the monomer. It can also be seen that K_2 is one

order of magnitude weaker than K_I . This weaker second binding site is likely to correspond to non-specific adsorption of tannins that may lead to coating of the protein surface^{28,45} and eventual precipitation of the tannin-protein complex in some systems. The binding isotherms in Figures 2 and 3 for the dimers show a greater exothermic response than for the corresponding monomers, *e.g.* roburin A *vs* vescalagin ($\Delta H_1 = -15.3$ *vs* -7.5 kJ mol⁻¹), oenothien B *vs* pedunculagin ($\Delta H_1 = -21.6$ *vs* -10.9 kJ mol⁻¹) and gemin A *vs* pentagalloyl glucose ($\Delta H_1 = -47.4$ *vs* -29 kJ mol⁻¹) (Table 2).

A closer look at the monomer series and the dimer series (each having tannins with similar molecular weights) suggests that the strength of binding is inversely related to the minimized energies and increases with a greater potential for rotational flexibility as follows (Table 1). The low energy conformers and binding constants, K_1 , of the monomers are: 229-236 kcal/mol and 5.2×10^4 M⁻¹ for vescalagin, 158-164 and 4.2×10^4 for pedunculagin, and 110-116 and 1.8×10^5 for pentagalloyl glucose. For the dimer series the corresponding values are: 406-412 kcal/mol and 2.3×10^5 M⁻¹ for roburin A, 278-280 and 6.5×10^5 for oenothien B, 323 and 1.8×10^6 for gemin A.

The number of strong (type 1) binding sites increases for ellagitannins with more flexibility and ‘free’ galloyl groups from $n_1=1.9$ for roburin A to 4.1 for oenothien B and 13 for gemin A. The binding parameters for two of the monomer-dimer pairs, *i.e.* the vescalagin-roburin A and pedunculagin-oenothien B pairs, show similarities in the stoichiometry of the interaction. For the vescalagin-roburin A pair the stronger binding site n_1 is 2.5 for the monomer and 1.9 for the dimer; and for the second weaker binding site n_2 is 30 for the monomer and 16 for the dimer. For the

pedunculagin-oenothetin B pair n_1 is 2.1 for the monomer and 4.1 for the dimer; and n_2 is 30 for the monomer and 20 for the dimer. However, for the most flexible monomer-dimer pair, which will be able to better support H-bonding, the stoichiometric values were much higher for the stronger binding site: n_1 was 26 for pentagalloyl glucose and 13 for gemin A. The difference reflects the fact that gemin A is twice the molecular size of pentagalloyl glucose. We reported previously that the sum of n_1 and n_2 indicated that 52 pentagalloyl glucose molecules are bound to BSA and this suggested surface coating as the BSA surface can theoretically accommodate 40 to 120 pentagalloyl glucose molecules.²⁸ Similarly, the sum of n_1 and n_2 for gemin A indicated that 38 gemin A dimers were bound (which equates to 76 'monomers') and matches with the calculated surface area. It also suggests that gemin A due to its flexibility is capable of coating the whole BSA protein just like pentagalloyl glucose.²⁸ Comparable stoichiometries were found for commercial preparations of the flexible taratannins when assuming a molecular weight of 1500: *i.e.* 48:1 for the taratannin:BSA ratio.³⁹

It is, however, of note that this molecular size effect was not observed to the same extent for the other pairs as monomers and dimers had similar n_1 -values (2 to 4) and similar sums of $n_1 + n_2$ (*i.e.* 33 for vescalagin/18 for roburin A and 32 for pedunculagin/24 for oenothetin B). This indicates that the less flexible structures and especially roburin A were less able to stretch out on the BSA surface.

Many more binding constants are available for flavonoids and condensed tannins than for ellagitannins. Nevertheless, pure flavonoid compounds tend to have comparable affinities as the ellagitannins investigated here. K_a constants with BSA of 10^2 to 10^5

316 M^{-1} were reported for monomeric and polymeric flavanols^{19,31} and of 10^4 to $10^5 M^{-1}$
317 for flavones and flavonols.⁴⁶⁻⁴⁹ Examples of polyphenol complexes with human
318 serum albumin include epigallocatechin gallate and diosmetin interactions that
319 exhibited comparable Stern-Volmer affinity constants of $\sim 10^5 M^{-1}$ from fluorescence
320 data.⁵⁰⁻⁵¹

321

322 The findings also match results from previous studies, where less pure tannins had
323 been used, and confirm that the tannin components of those fractions did indeed
324 dominate the interaction observed.^{19,24,39} The present study used a two-site binding
325 model and found K_1 of 10^4 to 10^5 and K_2 of 10^3 to $10^4 M^{-1}$ for the monomers in BSA
326 complexes. In comparison, data obtained with a single-site model gave K values of
327 $\sim 10^3 M^{-1}$ for chestnut- and myrabolan-BSA complexes.²⁴

328

329 Quideau *et al*⁵ reported that most polyphenols, including ellagitannins, when bound at
330 protein surfaces, exhibited affinity constants that were rarely above the micromolar
331 range, *i.e.* K_a of $\leq 10^6 M^{-1}$. However, Xiao *et al*⁵² also reported much larger constants
332 of up to $10^8 M^{-1}$ for some flavanoid and flavonoid - human serum albumin complexes.
333 Not surprisingly, some receptor and enzyme targets had particularly high selectivities
334 with affinity constants at the subnanomolar scale, and varied between ellagitannins
335 even if these were of similar molecular weights.^{5,53}

336

337 The data in Table 2 showed positive entropies at both types of binding sites for all but
338 one ellagitannin, *i.e.* gemin A. Positive ΔS_1 values ranged from 33 to 65 and ΔS_2
339 values from 36 to 54 $J mol^{-1} K^{-1}$ and are indicative of a dominance of hydrophobic
340 interactions.^{19,50,54} Gemin A was the only ellagitannin giving a negative entropy, -39 J

mol⁻¹ K⁻¹. For comparison purposes, we have included here also the entropies for pentagalloyl glucose, which shows that the highly flexible pentagalloyl glucose and gemin A had noticeably lower entropies at the stronger binding site than the other ellagitannins and were, therefore, apparently better able to form hydrogen bonds.⁵⁴ The positive entropies for all ellagitannins at the second binding site are consistent with hydrophobic stacking on the BSA surface, although some uncertainty exists with these values as the affinities were very low for some tannins ($K_2 = 10^3 \text{ M}^{-1}$). The literature contains examples of both hydrogen bonding and hydrophobic interactions in tannin-protein complexes.^{16,19,22,50,55} Taken together, the present study also suggests that both types of bonding interactions occur in these ellagitannin-BSA complexes and slight structural differences appear to dictate, which of these will dominate.

Quenching of the BSA Tryptophan Fluorescence by Ellagitannins. To

complement the ITC measurements, ellagitannin binding to BSA was also investigated by fluorescence spectroscopy. These experiments measure the quenching of the intrinsic tryptophan fluorescence of BSA³² after binding with each of the tannins. Two tryptophan residues are found near the BSA surface, one is within its hydrophobic binding pocket (Trp 212) and the other within the surface region of the protein (Trp 134).⁴⁹ Figure 4(a) shows Stern-Volmer plots that describe tryptophan quenching as a function of tannin concentration. In these plots F_0/F increases as fluorescence intensity decreases in the presence of quencher (F) relative to the native fluorescence in the absence of quencher (F_0). The data suggest that the rate of decrease in fluorescence with respect to quencher concentration depends on the molecular size and flexibility of the tannin. The data in Figure 4(a) revealed that four of the tannins fall into either monomer or dimer groups: the monomers, vescalagin

and pedunculagin, exhibit similar quenching behavior, and the dimers, roburin A and oenothien B, also behave similarly to one another. The dimers were more efficient quenchers than the monomers, requiring approximately half the molar tannin concentration to produce equivalent levels of quenching. Indeed, if the quenching data are normalised, as in Figure 4(b) where Stern-Volmer plots are given as a function of 'tannin monomer' concentration, thus removing effects of molecular weight, it can be seen that related monomer-dimer pairs overlay, which demonstrates similar quenching efficiency per monomer unit. In contrast to these four tannins, the most flexible dimer, gemin A, is a much more efficient quencher at much lower concentrations. Pentagalloyl glucose is also a more efficient quencher, $K_{SV} \sim 2 \times 10^6 \text{ M}^{-1}$,²⁸ than vescalagin and pedunculagin, but pentagalloyl glucose and gemin A are not a true monomer-dimer pair and thus do not overlay exactly.

The purpose of the Stern-Volmer plot of fluorescence quenching data is to generate a linearised data set that allows assessment of the ability of tannins to quench the BSA tryptophan residues. The slope of the straight line of best fit is termed the Stern-Volmer quenching constant (K_{SV}). From the data shown in Figure 4(a), K_{SV} -values of 10^4 to 10^5 M^{-1} have been calculated from a straight-line plot through the linear region at low tannin concentrations (Table 3) and are comparable with K_{SV} -values of 10^4 M^{-1} for tea, coffee and cocoa extracts³¹ and with $6.85 \times 10^4 \text{ M}^{-1}$ for quenching of human serum albumin by epigallocatechin gallate (EGCg).⁵⁰ These fluorescence data complement the ITC data by providing additional support to the findings outlined above. The trend in terms of the quenching constants, K_{SV} , matches that seen from ITC measurements of the affinity constants, K_I , with a linear correlation ($R^2 = 0.988$) between K_{SV} and the binding affinity constant, K_I (Tables 2 and 3).

391
392 However, it should be noted that each of the Stern-Volmer plots is non-linear and
393 curves towards the y-axis with higher tannin concentration. This is not unusual in
394 Stern-Volmer plots and provides information on the type of quenching. An upward
395 slope suggests a combination of both static and dynamic quenching in these
396 ellagitannin-BSA systems.⁴² Static quenching is associated with a strong coupling
397 (*e.g.* formation of a ground-state complex) between interacting molecules, whereas
398 dynamic quenching is associated with weak coupling (*e.g.* through-space interactions
399 or collisions).⁴² To further confirm that the curvature was indeed due to combined
400 static and dynamic quenching, a modified form of the Stern-Volmer equation was
401 used (see Methods) that is second order with respect to [Q], and this gave as predicted
402 a linear plot (Figure 4(c)).

403

404 **Polarity of the Tryptophan Environment.** Fluorescence spectra can also yield
405 information on another parameter, *i.e.* the position of the emission peak maximum
406 (λ_{em}), which can reflect the polarity of the tryptophan environment.⁵⁰ The presence of
407 ellagitannins led to a bathochromic or red-shift in λ_{em} (Table 3), suggesting that the
408 tryptophan environment became more polar upon interaction with ellagitannins. The
409 observed red-shift may suggest a change to the protein's tertiary structure and thus
410 account for the associated increase in the polarity of the tryptophan environment.²⁸
411 However, it is difficult to conclude this based solely on the data reported here. A shift
412 to a more polar environment could suggest unfolding of the protein and thus greater
413 solvent exposure or more H-bonding between BSA and the tannins. However, Li *et*
414 *al*³⁰ reported a slight blue-shift upon binding of tannic acid (*i.e.* a mixture of
415 gallotannins containing either a central glucose or quinic acid) to ferritin, whereas

Dobreva *et al*²⁸ measured a red-shift of 5 nm when pure pentagalloyl glucose bound with BSA. The magnitude of the red shift appears to be linked to tannin structure as related monomer-dimer pairs showed some similarities: vescalagin and roburin A generated a 8-9 nm shift, pedunculagin and oenothien B a 4-5 nm shift, but gemin A caused the greatest shift in λ_{em} of 10.3 nm. A similarly large red shift of 12 nm was observed for Trp during EGCg binding to human serum albumin.⁵⁰ Other experiments will be needed to establish the precise reasons for the observed differences in red-shifts.

In conclusion, our ITC data suggest that there are two types of binding sites for ellagitannin binding to BSA. A stronger binding site may well be more selective and dependent on tannin flexibility and the presence of free galloyl groups. Indeed, our previous study suggested a link between the number of aromatic residues on the surface of a globular protein, Rubisco, and pentagalloyl glucose binding.²⁸ This could explain the relationship between K_1 and the hydrophobic accessible surface area, ASA_H (Figure 5), *i.e.* $K_1 = 0.2125 \times e^{0.0114ASA_H}$ ($R^2 = 0.965$), and points to the fact that hydrophobic interactions are an important driver for affinity during BSA-ellagitannin complex formation as supported by positive entropies for 5 of the 6 ellagitannins studied here.

A second weaker binding site was required to fit our data in order to allow for the binding interactions observed at high tannin:protein molar ratios. This second binding site was associated with high values of n_2 for all tannins and suggests non-selective adsorption of tannin to the protein surface. Such findings suggest tannin binding to BSA is concentration dependent, where specific binding might dominate at low

concentrations and non-selective adsorption at higher concentrations of tannin.¹⁶
These studies have shown that ellagitannin binding to BSA increases in strength and
affinity for the larger tannins (dimers) compared to their monomer forms. Bond
rotational flexibility of the tannin also plays a role by increasing the strength of
interaction and number of stronger (possibly hydrogen bonding) binding sites on the
protein surface.

ABBREVIATIONS

BSA, bovine serum albumin; ET, ellagitannins; GT, gallotannins

Funding

This study was supported by the EU Marie Curie Research Training Network
'HealthyHay' (contract: MRTN-CT-2006-035805) and Academy of Finland (grant
258992 to JPS).

ACKNOWLEDGMENTS

Kristiina Lehtonen is acknowledged for help with ellagitannin purification.

REFERENCES

- (1) Okuda, T.; Ito, H. Tannins of constant structure in medicinal and food plants – hydrolysable tannins and polyphenols related to tannins. *Molecules* **2011**, 16, 2191–2217.
- (2) Vrhovsek, U.; Guella, G.; Gasperotti, M.; Pojer, E.; Zancato, M.; Mattivi, F. Clarifying the identity of the main ellagitannins in the fruit of the strawberry, *Fragaria vesca* and *Fragaria ananassa* Duch. *J. Agric. Food Chem.* **2012**, 60, 2507–2516.
- (3) Larrosa, M.; Garcia-Conesa, M. T.; Espin, J. C.; Tomas-Barberan, F. A. Ellagitannins, ellagic acid and vascular health. *Mol. Aspects Med.* **2010**, 31, 513–539.
- (4) Quideau, S.; Genot, E.; Saltel, F.; Douat-Casassus, C.; Delannoy Lopez, D. M. 2012. C-glucosidic ellagitannin compounds for use for altering the supramolecular arrangement of actin and for the treatment of osteoporosis, cancer, bacterial infection and viral infection. International Patent Application WO 2012/113835 A1 (PCT/EP2012/053017),
- (5) Quideau, S.; Deffieux, D.; Douat-Casassus, C.; Pouységu, L. Plant polyphenols: chemical properties, biological activities, and synthesis. *Angew. Chem. Int. Ed.* **2011a**, 50, 5099–5104.
- (6) Scalbert, A.; Andres-Lacueva, C.; Arita, M.; Kroon, P.; Manach, C.; Urpi-Sarda, M.; Wishart, D. Databases on food phytochemicals and their health-promoting effects. *J. Agric. Food Chem.* **2011**, 59, 4331–4348.
- (7) Rothwell, J. A.; Urpi-Sarda, M.; Boto-Ordóñez, M.; Knox, C.; Llorach, R.; Eisner, R.; Cruz, J.; Neveu, V.; Wishart, D.; Manach, C.; Andres-Lacueva, C.; Scalbert, A. Phenol-Explorer 2.0: a major update of the Phenol-Explorer database

484 integrating data on polyphenol metabolism and pharmacokinetics in humans and
 485 experimental animals. *Database* **2012**, doi:10.1093/database/bas031.

486 (8) Fotschki, B.; Milala, J.; Jurgonski, A.; Karlińska, E.; Zdunczyk, Z.;
 487 Juskiewicz, J.. Strawberry ellagitannins [thwarted](#) the positive effects of dietary
 488 fructooligosaccharides in rat cecum. *J. Agric. Food Chem.* **2012**, 62, 5871–5880.

489 (9) Salminen, J. P.; Karonen, M.; Sinkkonen, J. Chemical ecology of tannins:
 490 recent developments in tannin chemistry reveal new structures and structure–activity
 491 patterns. *Chem. Eur. J.* **2011**, 17, 2806–2816.

492 (10) Quideau, S.; Douat-Casassus, C.; Delannoy López, D. M.; Di Primo, C.;
 493 Chassain, S.; Jacquet, R.; Saltel, F.; Genot, E. Binding of filamentous actin and
 494 winding into fibrillar aggregates by the polyphenolic C-glucosidic ellgitannin
 495 vescalagin. *Angew. Chem. Int. Ed.* **2011b**, 50, 5099–5104.

496 (11) Okuda, T. Systematics and health effects of chemically distinct tannins in
 497 medicinal plants. *Phytochemistry* **2005**, 66, 2012–2031.

498 (12) Kashiwada, Y.; Nonaka, G.-I.; Nishioka, I.; Lee, K. J.-H.; Bori, I.;
 499 Fukushima, Y.; Bastow, K. F.; Lee, K.-H. Tannins as potent inhibitors of DNA
 500 topoisomerase-II *in vitro*. *J. Pharm. Sci.* **1993**, 82, 487–492.

501 (13) Quideau, S.; Jourdes, M.; Saucier, C.; Glories, Y.; Pardon, P.; Baudry, C.
 502 DNA topoisomerase inhibitor acutissimin A and other flavano-ellagitannins in red
 503 wine. *Angew. Chem. Int. Ed.* **2003**, 42, 6012–6014.

504 (14) Auzanneau, C.; Montaudon, D.; Jacquet, R.; Puyo, S.; Pouységu, L.;
 505 Deffieux, D.; Elkaoukabi-Chaibi, A.; De Giorgi, F.; Ichas, F.; Quideau, S.; Pourquier,
 506 P. The polyphenolic ellagitannin vescalagin acts as a preferential catalytic inhibitor of
 507 the α isoform of human DNA topoisomerase II. *Mol. Pharmacol.* **2012**, 82, 124–141.

- 508 (15) Kiss, A. K.; Granica, S.; Stolarczyk, M.; Melzig, M. F. Epigenetic modulation
509 of mechanisms involved in inflammation: influence of selected polyphenolic
510 substances on histone acetylation state. *Food Chem.* **2012**, 131, 1015–1020.
- 511 (16) Cala, O.; Dufourc, E.J.; Fouquet, E.; Manigand, C.; Laguerre, M.; Pianet, I.
512 The colloidal state of tannins impacts the nature of their interaction with proteins: the
513 | [v](#)case of salivary proline-rich protein/procyanidins binding. *Langmuir* **2012**, 28,
514 17410–17418.
- 515 (17) Fraga, C. G.; Galleano, M.; Verstraeten S. V.; Oteiza, P. I. Basic biochemical
516 mechanisms behind the health benefits of polyphenols. *Mol. Aspects Med.* **2010**, 31,
517 435–445.
- 518 (18) Bacon, J.R.; Rhodes, M.J.C. Binding affinity of hydrolyzable tannins to
519 parotid saliva and to proline-rich proteins derived from it. *J. Agric. Food Chem.*,
520 **2000**, 48, 838-843.
- 521 (19) Frazier, R. A.; Deaville, E. R.; Green, R. J.; Willoughby, I.; Stringano, E.;
522 Plant, J.; Mueller-Harvey, I. Interactions of tea tannins and condensed tannins with
523 proteins. *J. Pharm. Biomed. Anal.* **2010**, 51, 490–495.
- 524 (20) Canon, F.; Ballivian, R.; Chirot, F.; Antoine, R.; Sarni-Manchado, P.;
525 Lemoine, J.; Dugourd, P. Folding of a salivary intrinsically disordered protein upon
526 binding to tannins. *J. Am. Chem. Soc.* **2011**, 133, 7847–7852.
- 527 (21) Delannoy, D.; Deffieux, D.; Di Primo, C.; Quideau S. Study in real time of
528 polyphenol-protein interactions by surface plasmon resonance. *Polyphenols Commun.*
529 **2012**, 1, 59–60.
- 530 (22) Simon, C.; Barathieu, K.; Laguerre, M.; Schmitter, J.-M.; Fouquet, E.;
531 Pianet, I.; Dufourc, E. J. Three-dimensional structure and dynamics of wine tannin-

532 saliva protein complexes. A multitechnique approach. *Biochemistry* **2003**, 42, 10385–
 533 10395.

534 (23) Hagerman, A.E. Fifty years of polyphenol-protein complexes. Chapter 3 in
 535 *Recent Advances in Polyphenol Research*. Cheynier, V.; Sarni-Manchado, P.;
 536 Quideau, S., Eds; John Wiley & Sons Ltd, Oxford, UK. **2012**, 3, 71-97.

537 (24) Deaville, E. R.; Green, R. J.; Mueller-Harvey, I.; Willoughby, I.; Frazier, R.
 538 A. Hydrolysable tannin structures influence relative binding strengths to globular or
 539 random coil proteins. *J. Agric. Food Chem.* **2007**, 55, 4554–4561.

540 (25) Hagerman, A. E; Butler, L. G. The specificity of proanthocyanidin-protein
 541 interactions. *J. Biol. Chem.* **1981**, 256, 4494–4497.

542 (26) Bazylo, A.; Kiss, K. K.; Kowalski, J. High-performance thin-layer
 543 chromatography method for quantitative determination of oenothien B and quercetin
 544 glucuronide in aqueous extract of *Epilobii angustifolii* herba. *J. Chromatogr. A* **2007**,
 545 1173, 146-150.

546 (27) Giner-Chavez, B-I.; Van Soest, P.J.; Robertson, J.B.; Lascano, C.; Pell, A.N.
 547 Comparison of the Precipitation of Alfalfa Leaf Protein and Bovine Serum Albumin
 548 by Tannins in the Radial Diffusion Method. *J. Sci. Food Agric.* 1997, 74, 513-523.

549 (28) Dobrev, M. A.; Frazier, R. A.; Mueller-Harvey, I.; Clifton, L. A.; Gea A.;
 550 Green, R. J. Binding of pentagalloyl glucose to two globular proteins occurs via
 551 multiple surface sites. *Biomacromolecules* **2011**, 12, 710–715.

552 (29) Hofmann, T.; Glabasnia, A.; Schwarz, B.; Wisman, K. N.; Gangwer, K. A.;
 553 Hagerman, A. E. Protein binding and astringent taste of a polymeric procyanidin,
 554 1,2,3,4,6-penta-O-galloyl-beta-D-glucopyranose, castalagin, and grandinin. *J. Agric.*
 555 *Food Chem.* **2006**, 54, 9503–9509.

- 556 (30) Li, M.; Jia, X.; Yang, J.; Deng, J.; Zhao, G. Effect of tannic acid on properties
557 of soybean (*Glycine max*) seed ferritin: A model for interaction between naturally-
558 occurring components in foodstuffs. *Food Chem.* **2012**, 133, 410–415.
- 559 (31) Stojadinovic, M.; Radosavljevic, J.; Ognjenovic, J.; Vesic, J.; Prodic, I.;
560 Stanic-Vucinic, D.; Cirkovic Velickovic T.. Binding affinity between dietary
561 polyphenols and β -lactoglobulin negatively correlates with the protein susceptibility
562 to digestion and total antioxidant activity of complexes formed. *Food Chem.* **2013**,
563 136, 1263–1271.
- 564 (32) Dufour C.; Dangles O. Flavonoid-serum albumin complexation:
565 determination of binding constants and binding sites by fluorescence spectroscopy.
566 *Biochim. Biophys. Acta* **2005**, 1721, 164–173.
- 567 (33) Skrt, M.; Benedik, E.; Podlipnik, C.; Ulrih, N. P. Interactions of different
568 polyphenols with bovine serum albumin using fluorescence quenching and molecular
569 docking. *Food Chem.* **2012**, 135, 2418–2424.
- 570 (34) Li, M.; Hagerman, A.E. Interactions between plasma proteins and naturally
571 occurring polyphenols. *Curr. Drug Metab.* **2013**, 14, 432–445.
- 572 (35) Zhang, Y.; Shi, S.; Chen, X.; Zhang, W.; Huang, K.; Peng, M. Investigation
573 on the interaction between Ilaprazole and bovine serum albumin without or with
574 different C-ring flavonoids from the viewpoint of food-drug interference. *J. Agric.*
575 *Food Chem.* **2011**, 59, 8499–8506.
- 576 (36) Huang, B.X.; Kim, H-H.; Dass, C. Probing three-dimensional structure of
577 bovine serum albumin by chemical cross-linking and mass spectrometry. *J. Am. Soc.*
578 *Mass Spectrom.* **2004**, 15, 1237–1247.

- 579 (37) Moilanen, J.; Salminen, J.-P. Ecologically neglected tannins and their
580 biologically relevant activity: chemical structures of plant ellagitannins reveal their in
581 vitro oxidative activity at high pH. *Chemoecology* **2008**, 18, 73-83.
- 582 (38) O'Brien, R.; Ladbury, J. E.; Chowdhry, B. Z. Isothermal titration calorimetry
583 of biomolecules. In: *Protein-Ligand Interactions: Hydrodynamics and Calorimetry*;
584 Harding, S. E.; Chowdhry, B. Z. (Eds.); Oxford University Press: Oxford, U.K., 2001;
585 pp. 263–286.
- 586 (39) Frazier, R. A.; Papadopoulou, A.; Mueller-Harvey, I.; Kisson, D.; Green, R. J.
587 Probing protein-tannin interactions by isothermal titration microcalorimetry. *J. Agric.*
588 *Food Chem.* **2003**, 51, 5189–5195.
- 589 (40) Freire, E.; Mayorga, O. L.; Straume, M. Isothermal titration calorimetry.
590 *Anal. Chem.* **1990**, 62, A950–A959.
- 591 (41) Chi, Z.; Liu, R.; Zhang, H. Noncovalent interaction of oxytetracycline with
592 the enzyme trypsin. *Biomacromolecules* **2010**, 11, 2454–2459.
- 593 (42) Lakowicz, J. R. *Principles of Fluorescence Spectroscopy*; 2nd ed.; Kluwer
594 Academic/Plenum Publishers: New York, 1999; p 698.
- 595 (43) Vivas, N.; Laguerre, M.; Pianet de Boissel, I.; Vivas de Gaulejac, N.; Nonier,
596 M.-F. Conformational interpretation of vescalagin and castalagin physicochemical
597 properties. *J. Agric. Food Chem.* **2004**, 52, 2073–2078.
- 598 (44) McManus, J. P.; Davis, K. G.; Beart, J. E.; Gaffney, S. H.; Lilley, T. H.;
599 Haslam, E. Polyphenol interactions. Part I. Introduction: some observations on the
600 reversible complexation of polyphenols with proteins and polysaccharides. *J. Chem.*
601 *Soc. Perkin Trans. 2*, **1985**, 1429–1438.

602 (45) Charlton, A.J.; Baxter, N.J; Khan, M.L; Moir, A.J.G.; Haslam, E.; Davies,
 603 A.P.; Williamson, M.P. Polyphenol/peptide binding and precipitation. *J. Agric. Food*
 604 *Chem.* **2002**, 50, 1593-1601.

605 (46) Mishra, B.; Barik, A.; Priyadarsini, K. I.; Mohan, H. Fluorescence
 606 spectroscopy studies on binding of a flavanoid antioxidant quercetin to serum
 607 albumins. *J. Chem. Sci.* **2005**, 117, 641–647.

608 (47) Tian, J.; Liu, J.; Tina X.; Hu, Z.; Chen, X. G. Study of the interaction of
 609 kaempferol with bovine serum albumin. *J. Mol. Struct.* **2004**, 691, 197–202.

610 (48) Tian, J.; Liu, J.; Hu, Z.; Chen, X. G. Interaction of wogonin with bovine
 611 serum albumin. *Bioorg. Med. Chem.* **2005**, 13, 4124–4129.

612 (49) Lorrain, B.; Dangles, O.; Genot, C.; Dufour, C. Chemical modeling of heme-
 613 induced lipid oxidation in gastric conditions and inhibition by dietary polyphenols. *J.*
 614 *Agric. Food Chem.* **2010**, 58, 676–683.

615 (50) Maiti, T. K.; Gosh, K. S.; Dsagupta, S. Interaction of (-)-epigallocatechin-3-
 616 gallate with human serum albumin: fluorescence, Fourier transform infrared, circular
 617 dichroism, and docking studies. *Protein Struct. Funct. Genet.* **2006**, 64, 355–362.

618 (51) Zhang, G.; Wang, L.; Pan, J. Probing the binding of the flavonoid diosmetin
 619 to human serum albumin by multispectroscopic techniques. *J. Agric. Food Chem.*
 620 **2012**, 60, 2721–2729.

621 (52) Xiao, J.; Zhao, Y.; Wang, H.; Yuan, Y.; Yang, F.; Zhang, C.; Yamamoto, K.
 622 Noncovalent interaction of dietary polyphenols with common human plasma proteins.
 623 *J. Agric. Food Chem.* **2011**, 59, 10747–10754.

624 (53) Zhu, M. J.; Phillipson, J. D.; Greengrass, P. M.; Bowery, N. E.; Cai, Y. Plant
 625 polyphenols: biologically active molecules or non-selective binders to protein?
 626 *Phytochemistry* **1997**, 44, 441–447.

627 (54) Chandler, D. Interfaces and the driving force of hydrophobic assembly.
628 *Nature* **2005**, 437, 640–647.

629 (55) Butler, L. G.; Riedl, D. J.; Lebryk, D. G.; Blytt, H. J. Interaction of proteins
630 with sorghum tannin: mechanisms, specificity and significance. *J. Am. Oil Chem. Soc.*
631 **1984**, 61, 916–920.

632

FIGURE LEGENDS

Figure 1. Structures of ellagitannins and pentagalloyl glucose.

Figure 2. ITC binding isotherms for the interaction of bovine serum albumin with the monomer-dimer pair of (a) vescalagin and (b) roburin A. Data were fitted assuming a model for two-independent binding sites and values obtained from these fits are given in Table 2.

Figure 3. ITC binding isotherms for the interaction of bovine serum albumin with the monomer-dimer pairs of (a) pendunculagin, (b) oenothien B and (c) gemin A. Data were fitted assuming a model for two-independent binding sites and values obtained from these fits are given in Table 2.

Figure 4. Fluorescence quenching of tryptophans in BSA by vescalagin, roburin A, pedunculagin, oenothien B and gemin A plotted as: (a) ratio of initial fluorescence intensity to total fluorescence intensity *versus* ellagitannin concentration; (b) ratio of initial fluorescence intensity to total fluorescence intensity *versus* ellagitannin monomer concentration; and as a (c) modified Stern-Volmer plot to reflect contributions from both static and dynamic quenching (data are shown for vescalagin, roburin A, pedunculagin and oenothien B only).

Figure 5: Relationship between the affinity constant, K_I , and the modeled accessible hydrophobic surface area (ASA_H).

Table 1: Modeling of minimized energies and associated numbers of conformers, accessible hydrophobic surface area (ASA_H) and accessible surface area (ASA).

Compound	Molecular weight (Daltons)	Energies of conformers (kcal/mol)	Number of conformers	ASA_H nm²	ASA nm²
Vescalagin	935	229 to 236	18	253	914
Roburin A	1843	406 to 412	4	407	1587
Pedunculagin	769	158 to 164	4	293	867
Oenothlein B	1569	278 to 280	2	524	1485
Pentagalloyl glucose	927	110 to 116	21	373	1041
Gemin A	1841	323	1	574	1748

Table 2: Estimated thermodynamic binding parameters for the interaction between five ellagitannins and pentagalloyl glucose^b with bovine serum albumin at pH 6. ΔH_1 and ΔH_2 are the enthalpies, and K_1 and K_2 are the equilibrium binding constants for two binding sites.

	Vescalagin	Roburin A	Pedunculagin	Oenothien B	Pentagalloyl glucose ^b	Gemin A
n_1	2.5±0.5 ^a	1.9	2.1	4.1	26	13
K_1 (M ⁻¹)	5.2 (±2.3) ×10 ⁴	2.3 × 10 ⁵	4.2 × 10 ⁴	6.5 × 10 ⁵	1.8 × 10 ⁵	1.8 × 10 ⁶
ΔH_1 (kJ mol ⁻¹)	-7.5±1.0	-15.3	-10.9	-20.6	-29	-47.4
ΔS_1 (J mol ⁻¹ K ⁻¹)	65.1	33.2	51.9	42.2	3.3	-39.3
n_2	30±4	16	30	20	26	25
K_2 (M ⁻¹)	1.1(±0.07) ×10 ³	1.8 × 10 ⁴	1.1 × 10 ³	3.3 × 10 ⁴	8 × 10 ²	6.3 × 10 ⁴
ΔH_2 (kJ mol ⁻¹)	-6.7±1.3	-8.17	-5.1	-10.3	-29	-11.4
ΔS_2 (J mol ⁻¹ K ⁻¹)	35.7	54.0	41.1	51.9	-41.7	53.6
χ^2	0.65±0.16	0.87	1.99	8.07		3.29

^a ± standard deviation, n = 3. ^b Data are included for comparison (Dobrev *et al.*²⁸).

Table 3: Estimated quenching parameters for the interaction of ellagitannins with bovine serum albumin (n=3).

Fluorescence parameters	Vescalagin	Roburin A	Pedunculagin	Oenothien B	Gemin A
$K_{SV} (M^{-1})$	$0.42 (\pm 0.07) \times 10^5$	$1.03 (\pm 0.07) \times 10^5$	$0.53 (\pm 0.1) \times 10^5$	$1.44 (\pm 0.28) \times 10^5$	$3.13 (\pm 0.06) \times 10^5$
Bathochromic (red)-shift of λ_{max} (nm)	8.7 ± 0.4	8.7 ± 0.4	4.7 ± 0.9	4.0 ± 1.0	10.3 ± 0.4

Figure 1

FIGURE 2

FIGURE 3

FIGURE 4

FIGURE 5

TOC Graphic