

Protein precipitation behaviour of condensed tannins from Lotus pedunculatus and Trifolium repens with different mean degrees of polymerization

Article

Accepted Version

Zeller, W. E., Sullivan, M. L., Mueller-Harvey, I., Grabber, J. H., Ramsay, A., Drake, C. and Brown, R. H. (2015) Protein precipitation behaviour of condensed tannins from Lotus pedunculatus and Trifolium repens with different mean degrees of polymerization. Journal of Agricultural and Food Chemistry, 63 (4). pp. 1160-1168. ISSN 0021-8561 doi: <https://doi.org/10.1021/jf504715p> Available at <https://centaur.reading.ac.uk/38988/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

To link to this article DOI: <http://dx.doi.org/10.1021/jf504715p>

Publisher: American Chemical Society

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online

Protein Precipitation Behavior of Condensed Tannins from *Lotus pedunculatus* and *Trifolium repens* with Different Mean Degrees of Polymerization

Wayne E. Zeller,^{*†} Michael L. Sullivan,[†] Irene Mueller-Harvey,[‡] John H. Grabber,[†] Aina Ramsay,[‡] Chris Drake,[‡] and Ronald H. Brown[‡]

[†]U.S. Dairy Forage Research Center, Agricultural Research Service, U.S. Department of Agriculture, 1925 Linden Drive, Madison, Wisconsin 53706, United States

[‡]Chemistry and Biochemistry Laboratory, Food Production and Quality Division, School of Agriculture, Policy and Development, University of Reading, P.O. Box 236, 1 Earley Gate, Reading RG6 6AT, United Kingdom

*Corresponding author:

Phone: 608-890-0071, Fax: 608-890-0076, e-mail: Wayne.Zeller@ars.usda.gov

Table of Contents

Figure S1. ^1H NMR (360 MHz) spectrum of condensed tannin (CT) isolated from big trefoil; Fraction 1 (BTF1) in 4:1 D_2O /acetone- d_6	S4
Figure S2. ^1H - ^{13}C HSQC NMR spectrum of condensed tannin (CT) isolated from big trefoil; Fraction 1 (BTF1) in 4:1 D_2O /acetone- d_6	S5
Figure S3. ^1H NMR (360 MHz) spectrum of condensed tannin (CT) isolated from big trefoil; Fraction 2 (BTF2) in 4:1 D_2O /acetone- d_6	S6
Figure S4. ^1H - ^{13}C HSQC NMR spectrum of condensed tannin (CT) isolated from big trefoil; Fraction 2 (BTF2) in 4:1 D_2O /acetone- d_6	S7
Figure S5. ^1H NMR (360 MHz) spectrum of condensed tannin (CT) isolated from big trefoil; Fraction 3 (BTF3) in 4:1 D_2O /acetone- d_6	S8
Figure S6. ^1H - ^{13}C HSQC NMR spectrum of condensed tannin (CT) isolated from big trefoil; Fraction 3 (BTF3) in 4:1 D_2O /acetone- d_6	S9
Figure S7. ^1H NMR (360 MHz) spectrum of condensed tannin (CT) isolated from big trefoil; Fraction 4 (BTF4) in 4:1 D_2O /acetone- d_6	S10
Figure S8. ^1H - ^{13}C HSQC NMR spectrum of condensed tannin (CT) isolated from big trefoil; Fraction 4 (BTF4) in 4:1 D_2O /acetone- d_6	S11
Figure S9. ^1H NMR (360 MHz) spectrum of condensed tannin (CT) isolated from big trefoil; Fraction 5 (BTF5) in 4:1 D_2O /acetone- d_6	S12

Figure S10. ^1H - ^{13}C HSQC NMR spectrum of condensed tannin (CT) isolated from big trefoil;

Fraction 5 (BTF5) in 4:1 D_2O /acetone- d_6S13




Figure S1. ¹H NMR (360 MHz) spectrum of condensed tannin (CT) isolated from big trefoil; Fraction 1 (BTF1) in 4:1 D₂O/acetone-*d*₆.




Figure S2. ^1H - ^{13}C HSQC NMR spectrum of condensed tannin (CT) isolated from big trefoil; Fraction 1 (BTF1) in 4:1 $\text{D}_2\text{O}/\text{acetone-}d_6$.




Figure S3. ¹H NMR (360 MHz) spectrum of condensed tannin (CT) isolated from big trefoil; Fraction 2 (BTF2) in 4:1 D₂O/acetone-*d*₆.

grad-HSQC
NEZ-7-092-5A
D2O/acetone-d6
(4: 1)




Figure S4. ¹H-¹³C HSQC NMR spectrum of condensed tannin (CT) isolated from big trefoil; Fraction 2 (BTF2) in 4:1 D₂O/acetone-*d*₆.




Figure S5. ¹H NMR (360 MHz) spectrum of condensed tannin (CT) isolated from big trefoil; Fraction 3 (BTF3) in 4:1 D₂O/acetone-*d*₆.

grad-HSQC
MEZ-7-092-5B
D2O/acetone-d6
(4: 1)




Figure S6. ^1H - ^{13}C HSQC NMR spectrum of condensed tannin (CT) isolated from big trefoil; Fraction 3 (BTF3) in 4:1 $\text{D}_2\text{O}/\text{acetone-}d_6$.




Figure S7. ¹H NMR (360 MHz) spectrum of condensed tannin (CT) isolated from big trefoil; Fraction 4 (BTF4) in 4:1 D₂O/acetone-*d*₆.




Figure S8. ¹H-¹³C HSQC NMR spectrum of condensed tannin (CT) isolated from big trefoil; Fraction 4 (BTF4) in 4:1 D₂O/acetone-*d*₆.




Figure S9. ¹H NMR (360 MHz) spectrum of condensed tannin (CT) isolated from big trefoil; Fraction 5 (BTF5) in 4:1 D₂O/acetone-*d*₆.

grad-HSQC
WEZ-7-092-7
D2O/acetone-d6
(4: 1)




Figure S10. ^1H - ^{13}C HSQC NMR spectrum of condensed tannin (CT) isolated from big trefoil; Fraction 5 (BTF5) in 4:1 $\text{D}_2\text{O}/\text{acetone-}d_6$.