

The endophytic fungus Piriformospora indica protects wheat from Fusarium crown rot disease in simulated UK autumn conditions

Article

Accepted Version

Rabiey, M., Ullah, I. ORCID: <https://orcid.org/0000-0002-9367-6741> and Shaw, M. W. (2015) The endophytic fungus *Piriformospora indica* protects wheat from *Fusarium* crown rot disease in simulated UK autumn conditions. *Plant Pathology*, 64 (5). pp. 1029-1040. ISSN 0032-0862 doi: 10.1111/ppa.12335 Available at <https://centaur.reading.ac.uk/39152/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

To link to this article DOI: <http://dx.doi.org/10.1111/ppa.12335>

Publisher: Wiley

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online

Short title: *Piriformospora indica* reduces Fusarium

The endophytic fungus *Piriformospora indica* protects wheat from Fusarium crown rot disease in simulated UK autumn conditions

M Rabiey*

I Ullah

M W Shaw

School of Agriculture, Policy and Development,

University of Reading,

Whiteknights,

Reading RG6 6AR

*Email: m.rabieyghahfarokhy@pgr.reading.ac.uk

Key words: *Piriformospora indica*, root endophytic fungus, mutualistic symbiosis, crown rot disease, *Triticum aestivum*, real-time polymerase chain reaction

19 The root endophytic fungus *Piriformospora indica* (Sebacinacea) forms mutualistic
20 symbioses with a broad range of host plants, increasing their biomass production and
21 resistance to fungal pathogens. We evaluated the effect of *P. indica* on Fusarium crown rot
22 disease of wheat, under *in vitro* and glasshouse conditions. Interaction of *P. indica* and
23 Fusarium isolates under axenic culture conditions indicated no direct antagonistic activity of
24 *P. indica* against Fusarium isolates. Seedlings of wheat were inoculated with *P. indica* and
25 pathogenic *Fusarium culmorum* or *F. graminearum* and grown in sterilised soil-free medium
26 or in a non-sterilised mix of soil and sand. Fusarium alone reduced emergence and led to
27 visible browning and reduced root growth. Roots of seedlings in pots inoculated with both
28 Fusarium isolates and *P. indica* were free of visible symptoms; seed emergence and root
29 biomass were equivalent to the uninoculated. DNA was quantified by real-time polymerase
30 chain reaction (qPCR). The ratio of Fusarium DNA to wheat DNA rose rapidly in the plants
31 inoculated with Fusarium alone; isolates and species were not significantly different. *P.*
32 *indica* inoculation reduced the ratio of Fusarium to host DNA in the root systems. The
33 reduction increased with time. The ratio of *P. indica* to wheat DNA initially rose but then
34 declined in root systems without Fusarium. With Fusarium, the ratio rose throughout the
35 experiment. The absolute amount of Fusarium DNA in root systems increased in the absence
36 of *P. indica* but was static in plants co-inoculated with *P. indica*.

37

38 **Introduction**

39 Crown rot disease of wheat, primarily caused by *Fusarium culmorum* and *F. graminearum*
 40 (Fernandez & Chen, 2005), damages wheat in most parts of the world. The disease reduces
 41 wheat grain yield and quality and wheat straw production. Infection of seedlings and basal
 42 stems leads to yield loss from damaged seedlings, pre-harvest lodging, and impaired grain
 43 filling (Schilling *et al.*, 1996). In the UK these problems are largely avoided by certified seed,
 44 seed treatment with fungicides and rotation (www.hgca.com), but *Fusarium spp.* remain a
 45 serious concern in grain because they produce a range of mycotoxins that can lead to possible
 46 human and animal health problems if they enter the food chain (Goswami & Kistler, 2004,
 47 Xu *et al.*, 2008). These *Fusarium* pathogens are soil-borne and stubble-borne and can
 48 survive in the soil and crop residues for several seasons (Leplat *et al.*, 2013). This long term
 49 survival in plant debris or grass weeds, along with the lack of commercial cultivars with
 50 resistance to *Fusarium* crown rot, makes controlling the disease difficult (Wildermuth *et al.*,
 51 1997). The effects of agronomic practices on this disease are often unpredictable (Bailey *et*
 52 *al.*, 2000) and depend on the causal species as well as the environmental conditions.

53 *Piriformospora indica* (sebacinales: basidiomycota) is a root endophytic fungus with a wide
 54 host range that was first isolated from the rhizosphere of woody shrubs in the Thar region of
 55 northwest India (Verma *et al.*, 1998). All members of the Sebacinales are involved in
 56 mycorrhizal associations (Weiss *et al.*, 2004). *P. indica*, like arbuscular mycorrhizal fungi,
 57 has plant growth promoting effects, but, in contrast to mycorrhizal fungi, can be cultured on
 58 various synthetic media (Verma *et al.*, 1998). *P. indica* can mobilise and transport phosphorus
 59 ,nitrogen and micronutrients from soil to the infected host plant via plant-fungal interfaces
 60 (Malla *et al.*, 2004, Sherameti *et al.*, 2005, Varma *et al.*, 2013, Yadav *et al.*, 2010). It has also

been reported that *P. indica* can improve growth in a range of economically important monocot and dicot hosts (Bagde et al., 2010, Varma et al., 1999, Varma et al., 2000).

Piriformospora indica has been shown to increase resistant to biotic stresses including a wheat leaf disease (caused by *Blumeria graminis* f.sp. *tritici*), a wheat stem base disease (caused by *Oculimacula* Spp.), wheat and barley root rot diseases (caused by *Fusarium culmorum*, *Gaeumannomyces graminis* var. *tritici*) (Harrach et al., 2013, Serfling et al., 2007, Deshmukh & Kogel, 2007), a maize root disease (caused by *F. verticillioides*) (Kumar et al., 2009) and a lentil vascular wilt disease (caused by *Fusarium oxysporum* f. sp. *lentis*) (Dolatabadi et al., 2012). In tomato infected with *Verticillium dahliae*, *P. indica* increased leaf and fruit biomass and decreased disease severity. Also in tomato, *P. indica* reduced the concentration of Pepino mosaic virus in shoots (Fakhro et al., 2010). *Piriformospora indica* also increased plant tolerance to abiotic stresses including salt stress in barley (Alikhani et al., 2013, Baltruschat et al., 2008), wheat (Zarea et al., 2012) and tomato (Cruz et al., 2010). The fungus conferred drought tolerance in Chinese cabbage and enhanced seed production and grain yield (Michal Johnson et al., 2013, Sun et al., 2010). Previous investigations, have been concentrated in tropical and subtropical conditions. It remains to be shown whether *P. indica* is suited to temperate climatic conditions.

In this investigation, we tested the hypothesis that *P. indica* would reduce damage to wheat seedlings by restricting growth of *F. culmorum* and *F. graminearum* on roots in controlled environmental chambers adjusted to UK autumn conditions. Pathogen progression in the presence and absence of *P. indica* colonising simultaneously with or after *Fusarium* was measured.

Materials and methods

Fungal inoculation

Piriformospora indica was obtained from Dr. Patrick Schafer, Warwick University, UK and was grown on agar containing complex modified *Aspergillus* medium (CM medium) (Pham *et al.*, 2004). To produce inoculum of *P. indica*, five plugs of 5 mm discs of 4 days old *P. indica* culture were added to 500 mL flasks of CM medium and incubated on an orbital shaker at 140 rpm at room temperature ($21 \pm 1^\circ\text{C}$) for 14 days.

Isolates of *F. culmorum* (98/11 and UK.99) and *F. graminearum* (576 and 602.1), of UK origin, were obtained from the School of Biological Science at the University of Reading and Rothamsted Research Centre, UK and cultured on potato dextrose agar (PDA). Inoculum was prepared by the methods described by Ghahfarokhy *et al.* (2011).

Test for antagonistic activity

Interactions between *P. indica* and Fusarium isolates were examined by the method described by Ghahfarokhi and Goltapeh (2010). A 5 mm mycelial disc of *P. indica* was placed on one side of a PDA plate and incubated at room temperature ($21 \pm 1^\circ\text{C}$). Single 5 mm discs of Fusarium mycelium taken from the margins of 4 day old cultures were placed on the other side of the plates, simultaneously or 3-4 days after. To see the interaction between *P. indica* and Fusarium isolates microscopically, a clean glass microscope slide was placed in the middle of Petri dishes and a thin layer of PDA poured onto it. Single 5 mm discs of 4 day old cultures of *P. indica* and Fusarium isolates were placed at opposite ends of the slide simultaneously or 3-4 days apart and incubated at room temperature ($21 \pm 1^\circ\text{C}$). After 3-4 days, when leading hyphae of each culture met, the slides were observed microscopically using a LeitzDialux 20 microscope attached to a Canon camera (EOS, 300D).

109 **Plant materials and glasshouse experiments**

110 Seeds of winter wheat cv. Battalion were surface disinfected by rinsing for 2 mins in a 20
 111 mL/L sodium hypochlorite (Fisher scientific, UK), followed by three rinses in sterilized
 112 distilled water, and germinated on damp filter paper in a Petri dish at room temperature under
 113 natural indoor light for 48 hours.

114 To determine whether *P. indica* interacted with wheat to reduce Fusarium crown rot, pre-
 115 germinated wheat seeds were planted into 4-inch pots (5 seeds per pot), filled with a 1:1
 116 mixture of vermiculite (Medium, Sinclair, UK) and sand, steam sterilised at 121°C for 60 min
 117 on two consecutive days. The pots were incubated at temperatures ranging between 15 °C
 118 and 25 °C; humidity and light were not controlled. Inoculations were performed at the time of
 119 sowing or 7 days later in a 3 × 3 factorial combination by mixing 4 g of *P. indica* and 6 g of
 120 *F. culmorum* into the surface layer of the soil, without disturbing the seedling roots. Harvest
 121 was performed at 7, 14, 21, and 30 days after inoculation (dai) and DNA concentrations of
 122 the fungi in the root system determined. Each time point was independently replicated.

123 *P. indica* and *F. culmorum* interaction during the first week after inoculation was tested in the
 124 glasshouse in conditions similar to the above experiment. Inoculations were done at the time
 125 of sowing and roots were harvested daily for one week, DNA concentrations of the fungi and
 126 wheat in the root system determined and a sample stained for microscopy. The experiment
 127 had four treatments, $\pm P. indica$ and $\pm F. culmorum$, with two replications.

128 In a confirmatory experiment inoculations were done at the time of sowing in a 2×2 factorial
 129 combination with 4 g of *P. indica* and 6 g of *F. culmorum*. Harvest was performed at 1, 2, 4,
 130 8, 16 and 32 days after inoculation and DNA concentrations of both fungi and wheat in the
 131 root system determined.

A further experiment was done to determine whether the interactions occurred under cooler, conditions more similar to UK field environments. Germinated seeds were planted in a 1:1 mixture of non-sterilised soil (John Innes Composts, BHGS Ltd, UK) and sand and pots were incubated in a controlled environment chamber. The experiment lasted 42 days. For the first 14 d, the day-length was 12 h and temperature and humidity were 15°C, 65% , respectively, during day and 10°C, 65% during night; for the second 14 d conditions were adjusted to 12°C, 70% during day and 9°C, 70% during night; and for the last 14 d the day length was reduced to 10 h with conditions set at 10°C, 75% during day and 7°C, 75% during night. Pots were arranged in two randomised blocks. The experiment had 10 treatments with two replicates and five harvests. The treatments were based on 2×5 factorial combinations of $\pm P. indica$ with one of the following: no amendment, *F. culmorum* 98/11, *F. culmorum* UK.99, *F. graminearum* 576 or *F. graminearum* 602.1. One pot of each treatment in each replicate was harvested at 7, 17, 28, 35 and 42 dai.

Each pot received 60 mL of fresh nutrient solution once a week. Nutrient solution was prepared each week using tap water with the final concentrations given: NO_3^- 10 mM, PO_4^{2-} 1 mM, K^+ 6 mM, Ca^{2+} 1.5 mM, Mg^{2+} 1 mM, SO_4^{2-} 1.5 mM, Fe 10 μM , Mn^{2+} 1 μM , Zn^{2+} 0.01 μM , Cu^{2+} 0.1 μM , MoO_4^{2-} 0.07 μM and $\text{B}_4\text{O}_7^{2-}$ 0.07 μM . Sodium metasilicate (100 mg/L) included to control powdery mildew.

Staining and microscopy

Wheat root samples inoculated with *P. indica*, Fusarium isolates, and both fungi together were stained using black ink (Pelikan Fountain Pen Ink, Niche Pens Ltd, UK) (Vierheilig et al., 1998). Roots were cleared by soaking them in 10% (w/v) KOH for one hour at 80°C, then rinsed 5 times with tap water. Cleared roots were covered with 2% HCl (v/v) for at least 30 min. Thereafter, HCl was poured off and roots were covered with 50 g/L black ink for 30

min at 80°C. Roots were de-stained by rinsing in tap water and viewed under a microscope with 10x and 40x objectives.

DNA isolation and primer development

Total genomic DNA was isolated from 100 mg of harvested roots using a Qiagen DNeasy plant mini kit (Qiagen, UK) following the manufacturer's instructions. Samples were eluted into 100 µl of elution buffer and stored at –20°C until required. Single species genomic DNA standards were obtained from roots of uninoculated plants and from mycelia of *P. indica* and *Fusarium* isolates scraped off the agar. Bulk DNA concentration was measured using a NanoDrop-lite spectrophotometer (Thermo Scientific, UK). The extent of shearing of DNA was determined by electrophoresis of an aliquot of DNA in a 1% agarose gel.

Primers were designed using the Primer BLAST tool from NCBI (<http://www.ncbi.nlm.nih.gov/tools/primer-blast>) to amplify fragments of *P. indica* *tef* gene for EF-1-alpha (accession number: AJ249911.2; Pi-forward: TCCGTCGCGCACCATT and Pi-reverse: AAATCGCCCTCTTTCCACAA, 84 bp), *Fusarium* elongation factor 1 alpha (EF1a) (accession number: JX534485; for *F. culmorum*, F1-forward: GCCCTCTTCCCACAAACCATTCC and F1-reverse: CTCGGCGGCTTCCTATTGACAG, 85 bp and for *F. graminearum*, F2-forward: AAGCCGAGCGTGAGCGTGGTA and F2-reverse: CGGGAGCGTCTGATAGTCGTGTTA, 142 bp) and wheat translation elongation factor 1 alpha-subunit (TEF1) (accession number: M90077; Wt- forward: GTGCACCAAATCTTCCTGCC, Wt-reverse: GGTTATGGAATGTAGATGCTCGG, 71 bp). The accession numbers were obtained from <http://www.ncbi.nlm.nih.gov>. All primers were supplied by Invitrogen (Life Technologies, UK). To assess specificity of the primers for the targeted species and investigate any cross reactivity, genomic DNA isolated from pure cultures of *P. indica* and *Fusarium* isolates and root tissue of wheat seedlings were subjected to PCR using all primer sets.

Quantification of *P. indica* and *Fusarium* in wheat roots

The amount of *Fusarium* and *P. indica* in wheat root samples was quantified by real-time PCR (qPCR). qPCR was performed in a 20 µl final reaction volume using 1X SYBR[®] Green Jump Start[™] TaqReady Mix[™] (Sigma Aldrich, UK), 0.25 µM forward and reverse primers, 1.5 µl of sample DNA and 7.5µl molecular grade water, in a 72 tube rotor of a Rotor-Gene 6000 System (Corbett Life Sciences, UK). Thermal cycling was set up at one cycle of 95 °C for 2 min; then 40 cycles of 95 °C for 15 s and 60 °C for 1 min followed by melt curve analysis from 65 to 95 °C at the rate of 0.5 °C per second. PCR controls in every assay included no template controls (NTC) and genomic DNA standards in duplicate for *Fusarium* isolates, *P. indica* and wheat. Serial dilutions of pure genomic wheat, *Fusarium* and *P. indica* DNA standards were initially tested in triplicate to determine a calibration curve and PCR efficiencies. Data were obtained and analysed using Rotor-Gene 6000 series software Version 1.7. After quantification, estimates of *F. culmorum*, *F. graminearum* and *P. indica* colonization of wheat tissues were obtained by dividing the concentration of fungal DNA by the concentration of wheat DNA. Absolute mass of DNA of each fungus in a root system was estimated by multiplying the concentration of fungal DNA by the ratio of root weight to the sample weight that was taken for DNA extraction.

Statistical analysis of experiments

ANOVA was used to analyse all data using GenStat 16th ed, (VSN, UK) with appropriate blocking. Where applicable, data were log and arcsine transformed to stabilize the residual variance and aid interpretation.

Results

Interaction of *P. indica* and Fusarium

Neither Fusarium isolates nor *P. indica* growth was visibly affected by the presence of the other fungus under axenic culture conditions on PDA, and there was no zone of inhibition at the contact point of two fungal colonies. There was occasional loose coiling of *P. indica* around Fusarium hyphae but no clear evidence of mycoparasitism (Fig. 1a,b).

Fusarium- inoculated root samples of both species showed extensive growth of Fusarium , with the mycelium completely covering the roots by the final observation date, when brown symptoms were clearly visible. In *P. indica*-Fusarium inoculated plants, Fusarium colonisation was visually much less, but colonisation by *P. indica* was extensive.

Piriformospora indica colonisation started on root surfaces in the differentiation zone behind the root meristem with inter- and intracellular penetration of epidermal cells, during the first two to three days after inoculation, with hyphae filling up the cells. By four days after inoculation coiled hyphae could occasionally be seen inside the cells.. Later, a little colonisation could be observed in epidermal cells of the meristematic and elongation zones of roots. *P. indica* chlamydospores were not observed until six days after inoculation (Fig. 1c,d).




Fig.1. Interaction of *Piriformospora indica* and *Fusarium* in agar plates and in wheat roots; (a). Agar plate co-cultivated with *F. culmorum* and *P. indica*; (b). Interaction of coiled hypha of *P. indica* around *F. culmorum* in agar plates at the encounter point; (c). *P. indica* chytridospores inside wheat root cells, the fungus was not detected in endodermic and central part of the root, (d). *P. indica* hyphae and chytridospores inside wheat root cells.

Effect of *P. indica* on emergence rate, root weight and pathogen DNA concentration

The emergence rates of seeds inoculated with *F. culmorum* and *F. graminearum* and *P. indica* were evaluated seven days after sowing. Seeds inoculated with *F. culmorum* and *F. graminearum* isolates emerged less often than the uninoculated ($p < 0.001$). Seeds inoculated with *P. indica* alone had the same emergence rate as the uninoculated. The emergence rate of seeds inoculated with both pathogen and *P. indica* was significantly higher than *Fusarium*-inoculated plants but slightly lower than the uninoculated ($p = 0.02$) (Fig. 2




Fig.2. Emergence rates of seeds inoculated with *Fusarium* and *Piriformospora indica* evaluated 7 days after sowing; data were arcsine transformed. (a). Roots inoculated with *F. culmorum* and *P. indica* simultaneously at sowing time (s.e.d= 0.09, d.f= 57); (b). Roots inoculated with *F. culmorum* (98/11 and UK.99), *F. graminearum* (576 and 602.1) and *P. indica* simultaneously at sowing time (s.e.d= 0.07, d.f= 89). Each bar represents mean \pm 2 SEM (P:*P. indica* and F:*Fusarium*).

243 Root weights were evaluated at the final harvest. Roots of plants inoculated with *P. indica*
244 alone at sowing or 7 days later had weights equivalent to the control. Roots inoculated with
245 *F. culmorum* or *F. graminearum* had 40% lower root weight ($p < 0.001$). Roots of plants
246 inoculated with *P. indica* prior to Fusarium or simultaneously weighed roughly the same as
247 uninoculated plants and much more than the roots inoculated with Fusarium alone ($p < 0.001$).
248 *P. indica* inoculated 7 days after *F. culmorum* was less effective (Fig. 3).

249

250




Fig.3. Root weight of samples (mg) inoculated with *Fusarium* and *Piriformospora indica* evaluated at last harvest; data were Log₁₀ transformed. (a). Roots inoculated with *F. culmorum* or *P. indica* simultaneously or 7 days after sowing, harvested at 30 dai (s.e.d= 0.07, d.f= 8); (b). Roots inoculated with *F. culmorum* (98/11 and UK.99), *F. graminearum* (576 and 602.1) and *P. indica* simultaneously at sowing time, harvested at 42 dai (s.e.d= 0.07, d.f= 9); (c). Roots inoculated with *F. culmorum* or *P. indica* simultaneously at sowing, harvested at 32 dai (s.e.d= 0.02, d.f= 3). Each bar represent mean \pm 2 SEM, (P: *P. indica*, F: *Fusarium*, P0: *P. indica* added to soil at sowing, P7: *P. indica* added to soil at 7 days after sowing, F0: *F. culmorum* added to soil at sowing and F7: *F. culmorum* added to soil at 7 days after sowing).

264

265 The absolute quantity of Fusarium DNA in the root systems without *P. indica* grew at about
266 10% day⁻¹ throughout the experiment (Fig 4 a-c,f). The rate of growth of Fusarium inoculated
267 at 7 dai was similar to that inoculated at sowing time (Fig 4 a,b). The relative rate of increase
268 was constant for *F. graminearum* but declined in *F. culmorum* particularly in the first
269 experiment (Fig 4a-c). In co-inoculated samples, the absolute amount of pathogen was static
270 or slightly declining from 7-42 d (Fig 4a, b, d, f) after an initial period of increase (Fig. 4 e,f).

271

272

273









Fig.4. The absolute growth of Fusarium in inoculated wheat roots. The absolute amount obtained by adding \log_{10} fungal DNA to \log_{10} (root weight/sample weight in mg). (a). *F. culmorum* added to soil at sowing (F0); *Piriformospora indica* added simultaneously (P0) or 7 days after sowing (P7); (b). *F. culmorum* added to soil 7 days after sowing (F7); *P. indica* added at sowing (P0) or simultaneously 7 days after sowing (P7); (c). *F. culmorum* 98/11, *F. culmorum* UK.99, *F. graminearum* 576 or *F. graminearum* 602.1 added at sowing time; (d). *F. culmorum* 98/11, *F. culmorum* UK.99, *F. graminearum* 576 or *F. graminearum* 602.1 and *P. indica* added simultaneously at sowing time; (e). *F. culmorum* added to soil at sowing (F0) and *P. indica* added simultaneously (P0), during the first week of inoculation; (f). *F. culmorum* added to soil at sowing (F0) and *P. indica* added simultaneously (P0), during the first month of inoculation. Each point represent mean ± 2 SEM (for a and b; s.e.d= 0.2 and d.f= 23), (for F. c. 98/11 and PF.c. 98/11: s.e.d= 0.14 and d.f= 9; for F. c. UK.99 and PF.c. UK.99: s.e.d= 0.12 and d.f= 9; for F.g. 576 and PF.g. 576: s.e.d= 0.2 and d.f= 9; for F.g. 602.1 and PF.g.602.1: s.e.d= 0.2 and d.f= 9), (for e, s.e.d= 0.13, d.f= 11) and (for f, s.e.d= 0.2, d.f=11).

The ratio of *F. culmorum* or *F. graminearum* DNA to plant DNA, in the absence of *P. indica*, grew approximately exponentially at about $18\% \text{ day}^{-1}$ (Fig 5 a,c,f), after the first 7 days; growth of *F. culmorum* in the first week was faster (Fig 5e,f). Despite the difference in temperatures, both glasshouse and environmental chamber experiments had similar rates of fungal growth. Increase in *F. graminearum* DNA was faster than increase in *F. culmorum* DNA (Fig 5c). The rate of growth of Fusarium inoculated at 7 dai was similar to that inoculated at sowing time (Fig 5a, b). In the presence of *P. indica*, Fusarium growth was immediately reduced to the rate of growth of the root system (Fig 5 e,f) and then declined (Fig 5 b,d). *P. indica* inoculation 7 d after the pathogen reduced the rate of Fusarium growth relative to the root similarly to the reduction when inoculated simultaneously (Fig. 5b). Because of the initial period of growth alone, the *F.culmorum* to root ratio remained consistently higher when *P. indica* inoculation was delayed until 7 d after *F. culmorum* inoculation.









Fig.5. Development of the ratio of Fusarium DNA to wheat DNA in inoculated wheat roots. The ratio obtained by subtracting \log_{10} fungal DNA from \log_{10} wheat DNA. (a). *F. culmorum* added to soil at sowing (F0); *Piriformospora indica* added simultaneously (P0) or 7 days after sowing (P7); (b). *F. culmorum* added to soil 7 days after sowing (F7); *P. indica* added at sowing (P0) or simultaneously 7 days after sowing (P7); (c). *F. culmorum* 98/11, *F. culmorum* UK.99, *F. graminearum* 576 or *F. graminearum* 602.1 added at sowing time; (d). *F. culmorum* 98/11, *F. culmorum* UK.99, *F. graminearum* 576 or *F. graminearum* 602.1 and *P. indica* added simultaneously at sowing time; (e). *F. culmorum* added to soil at sowing (F0) and *P. indica* added simultaneously (P0), during the first week after inoculation; (f). *F. culmorum* added to soil at sowing (F0) and *P. indica* added simultaneously (P0), during the first month of inoculation. Each point represents mean \pm 2 SEM (for a and b; s.e.d= 0.2 and d.f= 23), (for F.c. 98/11 and PF.c. 98/11: s.e.d= 0.15 and d.f= 9; for F.c. UK.99 and PF.c. UK.99: s.e.d= 0.08 and d.f= 9; for F.g. 576 and PF.g. 576: s.e.d= 0.2 and d.f= 9; for F.g. 602.1 and PF.g.602.1: s.e.d= 0.2 and d.f= 9), (for e; s.e.d= 0.1, d.f= 11) and (for f, s.e.d= 0.2, d.f= 11).

The absolute quantity of *P. indica* DNA in the root systems of soil free medium, in the absence of Fusarium, increased in the first 7 days after inoculation (Fig 6a), then decreased from a peak of 10^4 copies/root system to 10^3 over the 30 days of the experiment (Fig 6b,c,e); but slightly increased, under simulated autumn conditions, by 42 days into the experiment (Fig 6d). In the presence of Fusarium, *P. indica* DNA grew gradually throughout the experiment (Fig 6a-e). The rate of growth of *P. indica* was lower under the simulated autumn conditions than under temperatures ranging between 15°C and 25°C (Fig 6b-d).








Fig.6. The absolute growth of *Piriformospora indica* in inoculated wheat roots. The absolute amount obtained by adding \log_{10} fungal DNA to \log_{10} (root weight/sample weight in mg). (a). *P. indica* added to soil at sowing (P0) and *Fusarium culmorum* added simultaneously (F0), during the first week of inoculation; (b). *P. indica* added to soil at sowing (P0); *F. culmorum* added simultaneously (F0) or 7 days after sowing (F7); (c). *P. indica* added to soil 7 days after sowing (P7); *F. culmorum* added at sowing (F0) or simultaneously 7 days after sowing (F7); (d). *P. indica*, *F. culmorum* 98/11, *F. culmorum* UK.99, *F. graminearum* 576 or *F. graminearum*602.1 added at sowing time; (e). *P. indica* added to soil at sowing (P0) and *F. culmorum* added simultaneously (F0), during the first month of inoculation. Each point represent mean \pm 2 SEM (for a; s.e.d= 0.1 and d.f=11),(for b and c; s.e.d= 0.2 and d.f= 23), (for d; s.e.d= 0.3 and d.f= 24) and (for e, s.e.d= 0.1, d.f= 11).

The ratio of *P. indica* DNA to plant DNA, in the absence of *F. culmorum*, grew exponentially at about 25% day⁻¹ in the first 7 days after inoculation (Fig 7a), then declined, then stayed constant for the remainder of experiment from 14 to 30 dai (Fig 7b,c). However, this early increase was not consistent (Fig 7e).The rate of growth of *P. indica* inoculated at 7 dai was similar to that inoculated at sowing time (Fig 7b,c). In the presence of *F. culmorum*, the rate growth of *P. indica* was static throughout the experiment (Fig 7a,b,c,e). In the experiment under simulated autumn condition the ratio of *P. indica* DNA to wheat DNA, in the absence or presence of *Fusarium* isolates, grew slowly at about 2% day⁻¹ throughout the experiment (Fig 7d).








Fig.7. Development of the ratio of *Piriformospora indica* DNA to wheat DNA in inoculated wheat roots. The ratio obtained by subtracting \log_{10} fungal DNA from \log_{10} wheat DNA. (a). *P. indica* added to soil at sowing (P0) and *Fusarium culmorum* added simultaneously (F0), during the first week after inoculation; (b). *P. indica* added to soil at sowing (P0); *F. culmorum* added simultaneously (F0) or 7 days after sowing (F7); (c). *P. indica* added to soil 7 days after sowing (P7); *F. culmorum* added at sowing (F0) or simultaneously 7 days after sowing (F7); (d). *P. indica*, *F. culmorum* 98/11, *F. culmorum* UK.99, *F. graminearum* 576 or *F. graminearum* 602.1 added at sowing time; (e). *P. indica* added to soil at sowing (P0) and *F. culmorum* added simultaneously (F0), during the first month of inoculation. Each point represent mean ± 2 SEM (for a; s.e.d= 0.1 and d.f=11), (for b and c; s.e.d= 0.3 and d.f= 23), (for d; s.e.d= 0.3 and d.f= 24) and (for e, s.e.d= 0.2, d.f =11).

Discussion

In these experiments *P. indica* very effectively controlled *F. culmorum* and *F. graminearum* under simulated conditions similar to UK autumn, even though *P. indica* was found in the Thar region, India, which experiences extreme temperature conditions.

As in other *P. indica* studies, the mechanism appeared to be indirect. Dual culture tests of *P. indica* and *F. culmorum* or *F. graminearum* and microscopy showed no capability of either fungus to inhibit the other, with no inhibition zone at the interaction point and no other direct antagonistic activities. This is consistent with Kumar et al (2009) and Deshmukh and Kogel (2007) who reported that *P. indica* did not have any direct antagonistic effect on *F. graminearum* and *F. verticillioides* respectively, in-vitro. However, Ghahfarokhi and Goltapeh (2010) found a clear inhibition zone at the interaction point of *Gaeumannomyces graminis* var. *tritici* and *P. indica*. This could be a species difference or due to environmental effects, in particular the incubation temperature in Ghahfarokhi and Goltapeh was 28 °C, the most favourable temperature for *P. indica* growth.

In inoculated roots *P. indica* penetration started at the differentiation zone of the roots, with inter- and intracellular hyphae penetration during the first two to three dai. *P. indica* hyphae filled up the cortical and epidermal cells. Chlamydospores were visible from six days after

391 inoculation. Occasionally, coiled hyphae could be observed within root cells. Jacobs *et al.*
 392 (2011) proposed a colonisation model for *P. indica* in Arabidopsis root, which started with
 393 inter- and intracellular penetration of rhizodermal and cortical tissues and then root hair cells
 394 by three days after inoculation. Fungal hyphae branched and sometimes formed whorls.
 395 Finally sporulation started at seven dai; this is completely consistent with our observations.

396 Surprisingly, pathogen DNA was slightly higher than in plants inoculated with pathogen
 397 alone during the first week after inoculation, in all experiments. This effect was probably due
 398 to the slight extra supply of exogenous nutrients from the substrate of the *P. indica* inoculum.
 399 Brown symptoms on root and crown were obvious in the Fusarium-inoculated samples,
 400 which reflected the extensive invasive growth of Fusarium hyphae in the samples, which was
 401 confirmed microscopically. In the presence of *P. indica*, the ratio of pathogen DNA to wheat
 402 DNA increased much more slowly and then decreased by the end of the experiment. The
 403 results are consistent with previous work in other host-pathogen systems. Kumar *et al.* (2009)
 404 reported PCR analysis of maize samples inoculated with *P. indica* and *F. verticillioides*. They
 405 showed that *P. indica* suppressed further colonization by *F. verticillioides*. Harrach *et al.*
 406 (2013) reported preinoculation of barley roots with *P. indica* prior to *F. culmorum* resulted in
 407 reduced colonization of roots by *F. culmorum*, which is consistent with less root rot–
 408 symptom expression and a reduced loss of biomass. Deshmukh and Kogel (2007) reported a
 409 decrease in the relative amount of *F. graminearum* DNA in barley roots in the presence of *P.*
 410 *indica*, followed by a sharp decrease at 19 days after inoculation of *P. indica*.

411 Inoculation of plants with *P. indica* before pathogen had a greater effect on both the ratio
 412 between pathogen and host DNA and the actual amount of pathogen than simultaneous or
 413 delayed inoculation. In the absence of Fusarium, the absolute quantity of *P. indica* DNA and
 414 the ratio of *P. indica* DNA to plant DNA decreased to a steady level after the first 7 days in
 415 the warm environment, but increased slightly under cool conditions. These results are

consistent with a number of possible modes of action. For example *P. indica* might interfere with host signalling pathways leading to oxidative burst, which are essential to successful Fusarium establishment (Varma et al., 2012, Waller et al., 2005). Although qPCR is a precise and reliable method to quantify DNA, caution needs to be taken in interpreting the data. qPCR results must be verified by other methods and understood in the context of the sampling protocol. Fusarium causes massive plant cell death, which might result in overestimation by qPCR of the abundance of Fusarium DNA in root tissues that contain less intact plant DNA (Harrach *et al.*, 2013). Hogg et al. (2007) found that Fusarium crown rot disease severity and symptoms in wheat were often, but not always, correlated with actual Fusarium colonization. Strausbaugh *et al.* (2005) did experiments in both field and glasshouse. In their field study, they found no correlation between root-rot severity index and Fusarium DNA quantities in root samples. However, in their glasshouse study percent infected root area was correlated with Fusarium DNA quantities in both wheat and barley. This contrast in their results might have various causes. It is possible that there were sampling problems in the field study. For example rotting might be so fast in soil that they only ever sampled nearly healthy plant tissues.

Our studies show that *P. indica* can protect wheat from damage by Fusarium disease at the seedling stage, in simulated UK conditions. However, the ecological-side-effects of *P. indica* are still unclear: how *P. indica* interact with other beneficial soil microorganisms, like arbuscular mycorrhizal fungi, how it affects soil functioning, such as turnover of soil organic matter, incorporation of residues, etc, and what effects *P. indica* has on other soil-borne diseases. These must be considered in further studies.

Acknowledgment

This work was funded by the Sir Halley Stewart Trust.

References

- Alikhani M, Khatabi B, Sepehri M, Nekouei MK, Mardi M, Salekdeh GH, 2013. A proteomics approach to study the molecular basis of enhanced salt tolerance in barley (*Hordeum vulgare* L.) conferred by the root mutualistic fungus *Piriformospora indica*. *Molecular Biosystems* **9**, 1498-510. <http://www.rsc.org/molecularbiosystems>.
- Bagde US, Prasad R, Varma A, 2010. Interaction of Mycobiont: *Piriformospora indica* with Medicinal plants and plants of Economic importance. *African Journal of Biotechnology* **9**, 9214-26. <http://www.academicjournals.org/AJB/>.
- Bailey KL, Gossen BD, Derksen DA, Watson PR, 2000. Impact of agronomic practices and environment on diseases of wheat and lentil in southeastern Saskatchewan. *Canadian Journal of Plant Science* **80**, 917-27.
- Baltruschat H, Fodor J, Harrach BD, Niemczyk E, Barna B, Gullner G, Janeczko A, Kogel KH, Schafer P, Schwarczinger I, Zuccaro A, Skoczowski A, 2008. Salt tolerance of barley induced by the root endophyte *Piriformospora indica* is associated with a strong increase in antioxidants. *New Phytologist* **180**, 501-10.
- Cruz C, Martins-Loucao MA, Varma A, 2010. The influence of plant co-culture of tomato plants with *Piriformospora indica* on biomass accumulation and stress tolerance. *Vi International Symposium on Mineral Nutrition of Fruit Crops* **868**, 123-7.
- Deshmukh SD, Kogel KH, 2007. *Piriformospora indica* protects barley from root rot caused by *Fusarium graminearum*. *Journal of Plant Diseases and Protection* **114**, 263-8.
- Dolatabadi HK, Goltapeh EM, Mohammadi N, Rabiey M, Rohani N, Varma A, 2012. Biocontrol Potential of Root Endophytic Fungi and Trichoderma Species Against Fusarium Wilt of Lentil Under In vitro and Greenhouse Conditions. *Journal of Agricultural Science and Technology* **14**, 407-20.
- Fakhro A, Andrade-Linares DR, Von Bargen S, Bandte M, Buttner C, Grosch R, Schwarz D, Franken P, 2010. Impact of *Piriformospora indica* on tomato growth and on interaction with fungal and viral pathogens. *Mycorrhiza* **20**, 191-200.
- Fernandez MR, Chen Y, 2005. Pathogenicity of Fusarium species on different plant parts of spring wheat under controlled conditions. *Plant Disease* **89**, 164-9.
- Ghahfarokhi RM, Goltapeh ME, 2010. Potential of the root endophytic fungus *Piriformospora indica*, *Sebacina vermifera* and *Trichoderma* species in biocontrol of take-all disease of wheat *Gaeumannomyces graminis* var. *tritici* in vitro, in Iran. *Journal of Agricultural Technology* **6**, 11-8. <http://www.ijat-aatsea.com/>.
- Ghahfarokhy MR, Goltapeh EM, Purjam E, Pakdaman, B.S., Modarres Sanavy SaM, Varma A, 2011. Potential of mycorrhiza-like fungi and Trichoderma species in biocontrol of Take-all Disease of wheat under greenhouse condition *Journal of Agricultural Technology* **7(1)**, 185-95. <http://www.ijat-aatsea.com/>.
- Goswami RS, Kistler HC, 2004. Heading for disaster: *Fusarium graminearum* on cereal crops. *Molecular Plant Pathology* **5**, 515-25.

- 479 Harrach BD, Baltruschat H, Barna B, Fodor J, Kogel K-H, 2013. The Mutualistic Fungus
480 *Piriformospora indica* Protects Barley Roots from a Loss of Antioxidant Capacity Caused by
481 the Necrotrophic Pathogen *Fusarium culmorum*. *Molecular Plant-Microbe Interactions* **26**,
482 599-605.
- 483 Hogg AC, Johnston RH, Dyer AT, 2007. Applying real-time quantitative PCR to fusarium
484 crown rot of wheat. *Plant Disease* **91**, 1021-8.
- 485 Jacobs S, Zechmann B, Molitor A, Trujillo M, Petutschnig E, Lipka V, Kogel KH, Schafer P,
486 2011. Broad-spectrum suppression of innate immunity is required for colonization of
487 Arabidopsis roots by the fungus *Piriformospora indica*. *Plant Physiology* **156**, 726-40.
- 488 Kumar M, Yadav V, Tuteja N, Johri AK, 2009. Antioxidant enzyme activities in maize plants
489 colonized with *Piriformospora indica*. *Microbiology-Sgm* **155**, 780-90.
- 490 Leplat J, Friberg H, Abid M, Steinberg C, 2013. Survival of *Fusarium graminearum*, the
491 causal agent of Fusarium head blight. A review. *Agronomy for Sustainable Development* **33**,
492 97-111.
- 493 Malla R, Prasad R, Kumari R, Giang PH, Pokharel U, Oelueeller R, Varma A, 2004.
494 Phosphorus solubilizing symbiotic fungus: *Piriformospra indica*. *Endocytobiosis Cell*
495 *Research* **15**, 579-600.
- 496 Michal Johnson J, Lee Y-C, Camehl I, Sun C, Yeh K-W, Oelmüller R, 2013. *Piriformospora*
497 *indica* promotes growth of chinese cabbage by manipulating auxin homeostasis - Role of
498 auxin in symbiosis. In. A. Varma (Ed.), *Piriformospora indica*, *Soil Biology* 33. Berlin-
499 Heidelberg: Springer-Verlag.
- 500 Pham GH, Singh A, Kumari R, Malla R, Prasad R, Sachdev M, Rexer KH, Kost G, Luis P,
501 Kaldorf M, Buscot F, Herrmann S, Peskan T, Oelmuller R, Sexena AK, Declerck S, Mittag
502 M, Stabentheiner E, Hehl S, Varma A, 2004. Interaction of *Piriformospora indica* with
503 diverse microorganisms and plants In. A. varma, L. Abbot, D. Warner, and R. Hampp (ed.)
504 Plant Surface Microbiology. pp. 237-265 Springer, Heidelberg.
- 505 Schilling AG, Moller EM, Geiger HH, 1996. Polymerase chain reaction-based assays for
506 species-specific detection of *Fusarium culmorum*, *F-graminearum*, and *F-avenaceum*.
507 *Phytopathology* **86**, 515-22.
- 508 Serfling A, Wirsal SGR, Lind V, Deising HB, 2007. Performance of the biocontrol fungus
509 *Piriformospora indica* on wheat under greenhouse and field conditions. *Phytopathology* **97**,
510 523-31.
- 511 Sherameti I, Shahollari B, Venus Y, Altschmied L, Varma A, Oelmuller R, 2005. The
512 endophytic fungus *Piriformospora indica* stimulates the expression of nitrate reductase and
513 the starch-degrading enzyme glucan-water dikinase in tobacco and Arabidopsis roots through
514 a homeodomain transcription factor that binds to a conserved motif in their promoters.
515 *Journal of Biological Chemistry* **280**, 26241-7.
- 516 Strausbaugh CA, Overturf K, Koehn AC, 2005. Pathogenicity and real-time PCR detection of
517 *Fusarium* spp. in wheat and barley roots. *Canadian Journal of Plant Pathology-Revue*
518 *Canadienne De Phytopathologie* **27**, 430-8.

- 519 Sun C, Johnson JM, Cai D, Sherameti I, Oelmüller R, Lou B, 2010. *Piriformospora indica*
 520 confers drought tolerance in Chinese cabbage leaves by stimulating antioxidant enzymes, the
 521 expression of drought-related genes and the plastid-localized CAS protein. *Journal of Plant*
 522 *Physiology* **167**, 1009-17.
- 523 Varma A, Kost G, Oelmüller R, Eds., 2013. *Piriformospora indica; Sebaciniales and their*
 524 *biotechnological applications*. Soil Biology 33. Berlin-Heidelberg: Springer-Verlag. pp 17.
- 525 Varma A, Rai MK, Sahay NS, 2000. Microbial- biotechnology: New paradigms and role in
 526 sustainable agriculture. In. *Microbial- biotechnology for sustainable development and*
 527 *productivity* (Ed. R.C. Rajak). Scientific Publishers, India. pp. 22-37.
- 528 Varma A, Sherameti I, Tripathi S, Prasad, R., , Das A, Sharma, M., , Bakshi M, Michal
 529 Johnson, J., , Bhardwaj S, Arora M, Rastogi K, Agarwal A, Kharkwal A, Talukdar S, Bagde
 530 U, Bisaria V, Upadhyaya C, Won P, Chen Y, Ma J, Lou B, Adya A, Zhong L, Meghvanshi
 531 M, Gosal S, Srivastava R, Johri A, Cruz C, Oelmüller R, 2012. The symbiotic fungus
 532 *Piriformospora indica*: Review. In. B. Hock (Ed.), *Fungal Associations*, 2nd Edition The
 533 *Mycota IX*. Berlin Heidelberg: Springer-Verlag.
- 534 Varma A, Verma S, Sudha SN, Butehorn B, Franken P, 1999. *Piriformospora indica*, a
 535 cultivable plant-growth-promoting root endophyte with similarities to arbuscular mycorrhizal
 536 fungi. *Applied and Environmental Microbiology* **65**, 2741-4.
- 537 Verma S, Varma A, Rexer KH, Hassel A, Kost G, Sarbhoy A, Bisen P, Butehorn B, Franken
 538 P, 1998. *Piriformospora indica*, gen. et sp. nov., a new root-colonizing fungus. *Mycologia* **90**,
 539 896-903.
- 540 Vierheilig H, Coughlan AP, Wyss U, Piche Y, 1998. Ink and vinegar, a simple staining
 541 technique for arbuscular-mycorrhizal fungi. *Applied and Environmental Microbiology* **64**,
 542 5004-7.
- 543 Waller F, Achatz B, Baltruschat H, Fodor J, Becker K, Fischer M, Heier T, Huckelhoven R,
 544 Neumann C, Von Wettstein D, Franken P, Kogel KH, 2005. The endophytic fungus
 545 *Piriformospora indica* reprograms barley to salt-stress tolerance, disease resistance, and
 546 higher yield. *Proceedings of the National Academy of Sciences of the United States of*
 547 *America* **102**, 13386-91.
- 548 Weiss M, Selosse MA, Rexer KH, Urban A, Oberwinkler F, 2004. Sebaciniales: a hitherto
 549 overlooked cosm of heterobasidiomycetes with a broad mycorrhizal potential. *Mycological*
 550 *Research* **108**, 1003-10.
- 551 Wildermuth GB, Thomas GA, Radford BJ, Mcnamara RB, Kelly A, 1997. Crown rot and
 552 common root rot in wheat grown under different tillage and stubble treatments in southern
 553 Queensland, Australia. *Soil & Tillage Research* **44**, 211-24.
- 554 Xu XM, Parry DW, Nicholson P, Thomsett MA, Simpson D, Edwards SG, Cooke BM,
 555 Doohan FM, Monaghan S, Moretti A, Tocco G, Mule G, Hornok L, Beki E, Tatnell J, Ritieni
 556 A, 2008. Within-field variability of Fusarium head blight pathogens and their associated
 557 mycotoxins. *European Journal of Plant Pathology* **120**, 21-34.
- 558 Yadav V, Kumar M, Deep DK, Kumar H, Sharma R, Tripathi T, Tuteja N, Saxena AK, Johri
 559 AK, 2010. A Phosphate Transporter from the Root Endophytic Fungus *Piriformospora indica*

560 Plays a Role in Phosphate Transport to the Host Plant. *Journal of Biological Chemistry* **285**,
561 26532-44.

562 Zarea MJ, Hajinia S, Karimi N, Goltapeh EM, Rejali F, Varma A, 2012. Effect of
563 *Piriformospora indica* and *Azospirillum* strains from saline or non-saline soil on mitigation
564 of the effects of NaCl. *Soil Biology & Biochemistry* **45**, 139-46.

565

566

567

568