

Staphylococcus aureus MnhF mediates cholate efflux and facilitates survival under human colonic conditions

Article

Accepted Version

Sannasiddappa, T., Hood, G., Hanson, K., Costabile, A., Gibson, G. and Clarke, S. (2015) *Staphylococcus aureus* MnhF mediates cholate efflux and facilitates survival under human colonic conditions. *Infection and Immunity*, 83 (6). pp. 2350-2357. ISSN 0019-9567 doi:
<https://doi.org/10.1128/IAI.00238-15> Available at
<https://centaur.reading.ac.uk/39959/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

To link to this article DOI: <http://dx.doi.org/10.1128/IAI.00238-15>

Publisher: American Society for Microbiology

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online

Staphylococcus aureus MnhF mediates cholate efflux
and facilitates survival under human colonic conditions

Thippeswamy H. Sannasiddappa^{1†}, Graham A. Hood¹, Kevan J. Hanson¹,
Adele Costabile², Glenn R. Gibson² and Simon R. Clarke^{1*}

¹School of Biological Sciences, University of Reading,
Whiteknights, Reading, RG6 6AJ, United Kingdom

²Food Microbial Sciences Unit, Department of Food and Nutritional Sciences,
University of Reading, Reading, RG6 6AP, United Kingdom

*Corresponding author:

s.r.clarke@reading.ac.uk
Tel: +44 118 378 8895
Fax: +44 118 378 6537

[†] Present address: Institute of Microbiology and Infection,
University of Birmingham, Birmingham, B15 2TT, United Kingdom

1 **Abstract**

2 Resistance to the innate defences of the intestine is crucial for the survival and
3 carriage of *Staphylococcus aureus*, a common coloniser of the human gut. Bile salts
4 produced by the liver and secreted into the intestines are one such group of molecules
5 with potent anti-microbial activity. The mechanisms by which *S. aureus* is able to
6 resist such defences in order to colonize and survive in the human gut are unknown.
7 Here we show that *mnhF* confers resistance to bile salts, which can be abrogated by
8 efflux pump inhibitors. MnhF mediates efflux of radiolabelled cholic acid in both *S.*
9 *aureus* and when heterologously expressed in *Escherichia coli*, rendering them
10 resistant. Deletion of *mnhF* attenuated survival of *S. aureus* in an anaerobic three-
11 stage continuous culture model of the human colon (gut model), which represent
12 different anatomical areas of the large intestine.

13 **Introduction**

14 *Staphylococcus aureus* is a ubiquitous and highly adaptable human pathogen
15 responsible for a significant global burden of morbidity and mortality. The bacterium
16 lives as a commensal in the nares of 20-25% of the population at any one time (1, 2).
17 While nasal colonisation is a well-established risk factor for most types of *S. aureus*
18 infections, several recent studies have suggested that colonisation of the intestine,
19 which occurs in *c.* 20% of individuals and which by and large has been overlooked,
20 could have important clinical implications (3). Patients with *S. aureus* intestinal
21 colonisation can serve as an important source of transmission, as they often
22 contaminate the adjacent environment (4). Similarly, such patients display an
23 increased frequency of skin colonisation (5). A study in intensive care and liver
24 transplant units showed that patients with both rectal and nares colonisation by MRSA
25 had a significantly higher risk of disease (40%) than did patients with nasal
26 colonisation alone (18%) (6). Furthermore, a study of hospitalised patients in the
27 United States reported co-colonisation by *S. aureus* and vancomycin-resistant
28 enterococci in >50% of the individuals studied (7). Thus it is likely that intestinal
29 colonisation by *S. aureus* provides the pathogen with a potential opportunity to
30 acquire new antibiotic resistance genes.

31

32 While the clinical implications of intestinal colonisation by *S. aureus* are still
33 relatively ill-defined, it is assumed that carriage is a risk for intestinal infection; *S.*
34 *aureus* can induce pseudomembranous colitis that is histologically distinct from that
35 caused by *Clostridium difficile* (8). Multiple studies have demonstrated frequent
36 intestinal colonisation in infants, particularly in those that were breast-fed and that a
37 positive correlation exists with development of allergies (9-13). While a role for *S.*

38 *aureus* intestinal carriage in development of systemic *S. aureus* disease has not been
39 established, colonisation of the intestinal lumen of mice can lead to the pathogen
40 crossing the intestinal epithelial barrier and subsequent spread to the mesenteric
41 lymph nodes (14, 15).

42

43 As a common commensal and pathogen, *S. aureus* must resist the human host's innate
44 defences that have evolved to limit its *in vivo* growth and spread. In particular, bile
45 represents a major challenge to bacteria that survive transit through the stomach and
46 enter the intestines. Bile is a digestive secretion that plays an essential role in
47 emulsification and solubilisation of lipids. We have previously demonstrated survival
48 of *S. aureus* in a human colonic model fed with physiological levels of bile (16).
49 Resistance to bile salts has been demonstrated to be important for intestinal survival
50 of several enteric pathogens, but in *S. aureus* such an understanding is lacking. The
51 role of the *S. aureus* *mnhABCDEFG* locus in bile resistance was identified using a
52 Tn917 library screened for bile-sensitive mutants. MnhF is homologous to
53 mammalian bile salt transporters, thus we hypothesized that it was involved in bile
54 resistance and therefore survival of *S. aureus* in conditions modeling the human
55 colon.

56

57 Here we provide molecular proof that a cause of bile salt resistance in *S. aureus* is
58 efflux, catalysed by MnhF. This represents the first description of an intestinal
59 colonisation factor in this pathogen.

60

61

62

63 **Materials and Methods**

64 **Bacteria, plasmids and growth conditions**

65 The strains and plasmids used in this work are listed in Tables 1 and 2, respectively.
66 *Escherichia coli* strains were grown on Luria–Bertani medium, using selection with
67 the antibiotic ampicillin (100 µg/mL) where appropriate. *S. aureus* was grown on
68 Brain Heart Infusion (BHI) (Oxoid) at 37°C. Where appropriate, antibiotics were
69 added at the following concentrations: erythromycin 5 µg/mL, lincomycin 25 µg/mL.
70 Phage transductions were as described previously (23).

71

72 **Determination of minimum inhibitory concentration (MIC)**

73 The MICs of selected bile salts, sodium cholate (CA), sodium chenodeoxycholate
74 (CDCA), sodium deoxycholate (DCA), sodium glycocholate (GCA), and sodium
75 taurocholate (TCA) were determined by broth dilution. MICs were determined by
76 doubling dilutions and MICs were reproduced in 3 independent experiments.

77

78 **Time-course measurement of bacterial viability upon exposure to bile salts.**

79 Overnight cultures were grown to mid-exponential phase in BHI broth at 37°C with
80 shaking. After harvesting, cells were washed twice with sterile 5 mM HEPES buffer
81 (pH 7.2) containing 10 mM glucose, then resuspended in the same buffer to an OD₆₀₀
82 0.5. Cells were incubated with various concentrations of bile salt for 30 minutes at
83 37°C. At 10 minute intervals, dilutions from each of the bile salt treated groups were
84 made with sterile peptone saline diluent. Dilutions were plated onto tryptic soy agar
85 plates and incubated for overnight at 37°C. Colonies were counted, and percentage
86 viabilities calculated based on the initial untreated cell suspension.

87

88 **Generation of an in-frame *mnhF* mutant.**

89 For the $\Delta mnhF$, DNA fragments corresponding to *c.* 0.7 kb upstream and downstream of
90 *mnhF* were amplified using Pwo polymerase (Roche) with primers
91 $\Delta mnhFLFor/\Delta mnhFLRev$ and $\Delta mnhFRFor/\Delta mnhFRRev$ (Table 3). Following
92 purification, PCR products were digested with *Bam*HI/*Eco*RI and cloned into pMAD.
93 The resulting plasmid was used to transform electrocompetent *S. aureus* RN4220 (24).
94 Plasmids were transduced into SH1000 using $\phi 11$ phage. The temperature sensitive
95 nature of plasmid replication was exploited to integrate the plasmid into the bacterial
96 chromosome, by plating cells on media containing erythromycin and lincomycin at 42°C.
97 After further rounds of plating, erythromycin and lincomycin sensitive colonies were
98 isolated and the loss of *mnhF* confirmed by PCR.

99

100 **Cloning and expression of *mnhF*.**

101 The *mnhF* gene was amplified by PCR with *S. aureus* SH1000 DNA. For cloning into *S.*
102 *aureus*, *mnhFFor2* and *mnhFRev* (Table 3) were used. PCR products were digested with
103 *Eco*RI and *Bam*HI and ligated into similarly digested pRMC2. This created pMnhF2,
104 where *mnhF* is fused to P_{xyI/tetO}, which is under the control of TetR and induced with
105 anydrotetracycline. For cloning into *E. coli*, oligonucleotides *mnhFFor1* and *mnhFRev*
106 (Table 3) were used. PCR products were digested with *Eco*RI and *Bsp*HI and ligated into
107 similarly digested pBAD/His A. This created pMnhF1, where *mnhF* is fused to P_{BAD},
108 which is under tight control of AraC.

109

110 **Bile salt accumulation assay.**

111 Accumulation of cholic acid in *S. aureus* was quantified using a previously described
112 method (25). Briefly, *S. aureus* and *E. coli* were grown in BHI and LB broth
113 respectively, at 37°C to an OD₆₀₀ *c.* 0.6. Cells were centrifuged (5 mins, 16,000g),

114 washed twice in 25mM potassium phosphate buffer (pH 7.0) containing 1mM MgSO₄
115 and resuspended in same buffer to a concentration of 100 OD units/mL. One μCi of
116 ¹⁴C labelled cholic acid (American Radiolabelled Chemicals) with specific
117 radioactivity of 55 mCi/mmol was added, to a final concentration of 18 μM, cells
118 were incubated at 37°C for 2 h. Cells were then diluted to 10 OD units/ml in 25 mM
119 potassium phosphate buffer (pH 7.0) containing 1 mM MgSO₄, 20 mM glucose and
120 0.2 mM non-radiolabelled cholic acid, and incubated at 37°C. Incorporation of
121 radiolabelled cholic acid was measured by scintillation counting. At the indicated
122 time, 250 μl cells were centrifuged at 16,000g for 2 min, and the pellets resuspended
123 in 500 μl of sterile water and 3 ml of Ulitma Gold scintillation cocktail (Perkin
124 Elmer). CPM were counted in a Beckman LS 6500 Coulter liquid scintillation
125 counter.

126

127 **Quantitative real-time PCR.**

128 mRNAs from mutant and wild type strains were quantified using quantitative real-
129 time PCR (qRT-PCR). Cells were grown in triplicate as described above, then treated
130 with RNA protect (Qiagen) and RNA was isolated using the Qiagen RNeasy Mini kit.
131 DNA was removed using Turbo DNase-free (Life Technologies). Purified RNA was
132 quantified using a nonodrop ND-1000 spectrophotometer (Thermo Scientific). 0.5 μg
133 of RNA was reverse transcribed using the Tetro cDNA synthesis kit (Bioline). qRT-
134 PCR was performed using the Aligent qPCR System and iQ SYBR Green Supermix
135 (Biorad). The relative amounts of RNAlII mRNA in parental wild type and mutant
136 cells was determined by relative quantification using *gyrB*, based on consistent levels
137 observed in previous studies (26, 27, 28, 29). The oligonucleotides used for qRT-
138 PCR are listed in Table 3.

139

140 **Three-stage continuous culture colonic model system (human gut model).**

141 The three-stage continuous culture model of the human colon has been described
142 previously (16, 30). The experiment was carried out in triplicate using faecal samples
143 from three different volunteers. After obtaining verbal informed consent, a standard
144 questionnaire to collect information regarding the health status, drugs use, clinical
145 anamnesis, and lifestyle was administrated before the donor was ask to provide a
146 faecal sample. No volunteers had received antibiotics, probiotics, steroids or other
147 drugs with a proven impact on gut microbiota for at least 3 months before sampling.
148 None of them had any history of gastrointestinal disorder. All healthy faecal donors
149 had the experimental procedure explained to them and were given the opportunity to
150 ask questions. The University of Reading research Ethics Committee exempted this
151 study from review because no donors were involved in any intervention and waived
152 the need for written consent due to the fact the samples received were not collected by
153 means of intervention. All faecal samples were collected on site, kept in an anaerobic
154 cabinet (10 % H₂, 10 % CO₂ and 80 % N₂) and used within a maximum of 15 minutes
155 after collection. Samples were diluted 1/10 w/v in anaerobic PBS (0.1 mol/l phosphate
156 buffer solution, pH 7.4) and homogenized (Stomacher 400, Seward, West Sussex,
157 UK) for 2 minutes at 460 paddle-beats.

158

159 Samples were plated onto BHI agar containing 0.01% (w/v) potassium tellurite as a
160 selective agent at different dilutions in PBS (from 10² to 10⁹ CFU/ml) in triplicate for
161 each time point to measure bacterial counts.

162

163 **Statistical analysis**

164 All experiments were repeated three times and data were presented as \pm standard error
165 of mean. Analysis was performed using GraphPad Prism 5 software. Experimental
166 data were analysed by One-Way Anova and Two-Way Anova method, using
167 Bonferroni post-test analysis.

168

169 **Results**

170 **Identification of a bile salt resistance locus.**

171 Genes conferring resistance to bile were identified by replica plating *S. aureus*
172 SH1000 Tn917 insertion libraries on BHI agar and onto BHI agar containing 18%
173 (w/v) bile salts (Oxoid), which represented $0.8 \times$ MIC. Six colonies were unable to
174 grow in the presence of bile salts, but exhibited no growth defect on BHI agar in the
175 absence of bile. Sequencing of the genomic DNA flanking the transposon insertion
176 site of bile sensitive strains was carried out in order to identify the DNA insertion
177 sites of Tn917, revealing that all six strains were siblings containing the transposon
178 inserted in the same gene, namely the previously described *mnhA*, the first gene in the
179 polycistronic *mnhABCDEFG* operon which encodes a Na^+/H^+ antiporter (31).
180 *Bacillus subtilis* contains the orthologous *mrpABCDEFG* operon that has an identical
181 function, however *mrpF* and by extension *mnhF*, are homologous to mammalian bile
182 transporters and *mrpF* mediates cholic acid efflux (32, 33).

183

184 **MnhF mediates resistance to bile salts.**

185 We hypothesized that MnhF was responsible for the observed bile salt resistance
186 phenotype. To test this, an in-frame Δ *mnhF* strain was created in *S. aureus* SH1000.
187 The mutant strain had no growth defect when grown on BHI solid or liquid media in

188 the absence of bile salts (results not shown). Compared to the parental wild type, the
189 $\Delta mnhF$ strain had a reduced MIC for unconjugated bile salts and, in particular, cholic
190 acid (Table 4). Complementation of mutation with *mnhF* under the control of an
191 inducible promoter restored the bile resistance phenotype to that observed in the
192 parent strain in the presence of anhydrotetracycline as an inducer (Table 4), whereas
193 there was no such resistance in the absence of the inducer (results not shown). In
194 killing assays, the $\Delta mnhF$ strain was significantly more sensitive than the parent. In
195 the presence of 1 $\mu\text{g/mL}$ anhydrotetracycline, the complemented strain exhibited a
196 similar rate of cell death as the parental wild type (Fig 1). The increased sensitivity of
197 the mutant strain was only observed with unconjugated bile salts. However it should
198 be noted that we were unable to determine the MIC of conjugated bile salts for *S.*
199 *aureus*, as they were insoluble at concentrations greater than 200 mM.

200

201 To confirm the role of *mnhF* in bile salt resistance, it was cloned under the control of
202 the arabinose-inducible P_{BAD} promoter of plasmid pBAD/HisA, which enabled
203 arabinose-dose dependent expression of MnhF in *E. coli* TG1 and TOP10 strains.
204 Expression of MnhF increased the MICs to both conjugated and unconjugated bile
205 salts in both background strains and in the case of cholic acid, the increased resistance
206 was arabinose-dose dependent (Table 5). Similarly, expression of MnhF in *E. coli*
207 decreased the bacteriostatic effects of bile salts on that bacterium (Fig 2). Thus MnhF
208 was sufficient to enable bile salt resistance in the absence of the rest of the
209 *mnhABCDEFG* operon.

210

211 **The effect of efflux pump inhibitors on bile salt resistance.**

212 Given the ability of MnhF to confer bile salt resistance and its similarity to other
213 known and putative bile efflux systems, its ability to mediate removal of cholic acid
214 from bacteria was tested. Both Phe-Arg- β -naphthylamide (PA β N), a synthetic
215 dipeptide that inhibits bacterial efflux pumps, including bile salt efflux pumps of
216 Gram negative bacteria, and reserpine, a plant alkaloid which can inhibit multidrug
217 efflux pumps in Gram positive bacteria, were tested for their ability to reduce bile salt
218 MICs in *S. aureus*. Both inhibitors caused reductions in the *S. aureus* MIC for cholic
219 acid and PA β N reduced the MIC for all three unconjugated bile salts (Table 6A),
220 however the reduction was much smaller in the $\Delta mnhF$ strain than the parental wild-
221 type, possibly indicating the presence of other bile salt efflux systems in the pathogen.
222 Similarly, in *E. coli* (pMnhF1), PA β N reduced bile salt MICs to levels lower than that
223 for untreated *E. coli* (pBAD His A) (Table 6B). Thus in both *S. aureus* and *E. coli*,
224 inhibitors of efflux pumps abrogated bile salt resistance in an MnhF dependent
225 manner.

226

227 **MnhF transports cholic acid.**

228 Given the ability of efflux pump inhibitors to reduce the MICs of certain bile salts in
229 *S. aureus*, the capacity of the MnhF to transport cholic acid was determined *in*
230 *vitro* using a ^{14}C -radiolabelled cholic acid substrate, similar to previous efflux assays
231 (25, 34, 35). *S. aureus* SH1000 and $\Delta mnhF$ strains were incubated with ^{14}C -cholic
232 acid (uptake period) and then diluted in buffer containing excess of non-radiolabelled
233 cholic acid (efflux period). Initial ^{14}C -cholic acid uptake was the same for both
234 strains (10962 ± 550 cpm for *S. aureus* SH1000 and 10278 ± 278 cpm for *S. aureus*
235 $\Delta mnhF$), but throughout the efflux period *S. aureus* $\Delta mnhF$ retained significantly
236 more of the radiolabel than the parental wild-type (Fig 3A). To further corroborate

237 these findings, efflux assays were also carried out on *E. coli* expressing MnhF. *E. coli*
238 TG1, *E. coli* TG1 (pBAD) and *E. coli* TG1 (pMnhF1) were grown overnight in LB
239 supplemented with 1% arabinose at 37°C, then incubated with ¹⁴C- cholic acid. All
240 the *E. coli* TG1 strains incorporated similar levels of ¹⁴C-cholic acid during uptake
241 period (20774 ± 363 for TG1, 23274 ± 386 for TG1: pBAD and 22435 ± 460 CPM
242 for TG1: pMnhF1). At various points after the initial incorporation of radiolabelled
243 cholic acid, cells were centrifuged and cell-associated radioactivity was determined
244 by liquid scintillation method. *E. coli* TG1 cells expressing MnhF retained
245 significantly (P<0.05) lower levels of ¹⁴C-radiolabelled cholic acid than parental TG1
246 and TG1 cells with the empty pBAD vector (TG1: pBAD) (Fig 3B). In both sets of
247 experiments the reason for increasing cell-associated radiolabel during the efflux
248 period, after which cells have been diluted in excess non-labelled cholic acid, is
249 unclear, but has also been observed in previous studies on *Listeria monocytogenes*
250 and may reflect continued incorporation of ¹⁴C-cholic acid during the efflux period
251 after dilution (25).

252

253 **Bile salt resistance is not affected by *agr*.**

254 To examine whether *agr* quorum sensing system is involved in bile salt resistance, the
255 MICs for CA, DCA and CDCA in *S. aureus* SH1001 (*agr*) were determined and
256 found to be indistinguishable from those of the wild type (results not shown).
257 Furthermore, the *agr* system is not inhibited by the *mnhF* mutation as the RNAIII
258 effector molecule is still produced (Fig 4). Thus we were unable to demonstrate a role
259 for *agr* in bile resistance.

260

261 **MnhF is required for survival of *S. aureus* in a human gut model.**

262 To examine the role of MnhF in survival of *S. aureus* in conditions found in the
263 human colon, we used a three-stage continuous culture gut model system, designed to
264 reproduce the spatial, temporal, nutritional and physicochemical characteristics of the
265 microbiota in the human colon. *In vivo* studies of colonic bacteria are hampered by
266 the lack of suitable animal models, as these do not correctly simulate the microbiota
267 and physicochemical conditions of the human colon (36). We have previously used
268 this *in vitro* model to study survival of *S. aureus* and the impact of infection on the
269 host's intestinal microflora (16).

270

271 Mutational inactivation of the whole *mnhABCDEFG* operon does not affect the ability
272 of *S. aureus* to grow at a range of pH levels (37). In order to exclude the possibility
273 that the normal pH range (5.5 to 7.5) found in the colon, influenced survival of the
274 $\Delta mnhF$ mutant, we corroborated the previous observation at pH 5.5 to 8.5 using this
275 strain (results not shown).

276

277 After inoculating vessel 1 (which models the proximal colon) of the colonic models
278 with *S. aureus* to a concentration of *c.* 2×10^{10} cfu/mL, as a single dose, the *S. aureus*
279 populations stabilised at 6 to 7 Log₁₀ units over a period of up to 8 hours. Survival of
280 *S. aureus* $\Delta mnhF$ was significantly attenuated compared to its parental strain in all
281 three vessels (Fig 5A-C).

282

283 **Discussion**

284 A complex set of interactions exists between *S. aureus* and its human host as the
285 bacterium is able to colonize several niches, both as an opportunist pathogen of great
286 medical importance and as a common commensal. In order to defend against

287 colonization by microorganisms, the host produces a range of antimicrobials such as
288 peptides, fatty acids and bile. Bile represents one significant challenge to the gut
289 microflora; in humans the liver secretes up to one liter of bile per day into the
290 intestines (38). Furthermore molecules secreted by bacteria, including *S. aureus*,
291 during infection are an important cause of metabolic cholestasis; an inability of
292 hepatocytes to produce bile (39). Bile is a complex cocktail composed principally of
293 bile salts, phospholipids, cholesterol, proteins and bilirubin (40). Originally
294 characterised as digestive molecules, bile salts have antimicrobial activity, which has
295 been attributed to their ability to damage cell membranes (41). Additionally, they
296 cause intracellular acidification, induce formation of secondary structures in RNA,
297 DNA damage and misfolding and denaturation of proteins. Thus bile salts represent a
298 serious challenge to bacterial cells in the gastrointestinal tract and bacteria that are
299 able to colonise the gut should therefore be able to overcome their toxicity.

300

301 Bile salts which pass into the large intestine undergo modification by the normal
302 microbiota (42). The major modifications include deconjugation, oxidation of
303 hydroxyl groups at C-3, C-7 and C-12, and $7\alpha/\beta$ -dehydroxylation (43, 44). Thus the
304 normal commensal inhabitants of the human gastrointestinal tract such as
305 *Lactobacillus*, *Propionibacterium* and *Bifidobacterium*, are required by the host for
306 maintenance of gut health and the ecological balance by influencing the composition
307 of the bile acids in the large intestine and by extension, the gut microbiome (45, 46).
308 Their ability to survive in the presence of bile salts indicates the existence of inherent
309 bile resistance mechanisms. Indeed, colonic commensals deploy various different
310 strategies for resisting bile. *Lactobacillus plantarum* produces a bile salt hydrolase,
311 which detoxifies bile salts by deconjugating bile salts inside the cell, turning them into

312 weaker acids, thus negating the drop in pH that they cause (47). Bifidobacteria
313 possess a number of characterised bile salt resistance mechanisms. In addition to
314 multiple efflux pumps, exposure to bile salts results in a modification of the cell
315 envelope. Increased concentrations of membrane fatty acids and altered
316 phospholipids increase membrane rigidity and reduce the permeability to lipophilic
317 bile salts (48). Similarly, exposure of *Bifidobacterium animalis* ssp *lactis* to bile salts
318 induces increased expression of exopolysaccharides, which are proposed to form a
319 protective layer around the bacterium (49).

320

321 Bile salts represent a physiological challenge for bacteria and an environmental cue;
322 *Salmonella enterica* and *Vibrio cholera* regulate intestinal colonisation and virulence
323 in response to bile (50, 51). However pathogens that inhabit the human intestines are
324 also exposed to the bactericidal nature of bile salts and hence must also exhibit
325 resistance in order to survive. Generally, Gram-negative bacteria are more innately
326 resistant than Gram positives, due to the presence of an outer membrane, which acts
327 as a barrier (38). Indeed maintenance of membrane integrity by lipopolysaccharide
328 (LPS) in the cellular envelope of Gram-negative bacteria imparts protection against
329 the actions of bile salts (52, 53). *Salmonella typhi* and *Salmonella typhimurium* are
330 able to grow at bile concentrations that are much higher than those encountered *in*
331 *vivo*. This is due, at least in part, to the presence of outer membrane efflux pumps
332 such as AcrAB (54). Similarly, HefC is an AcrB homologue that confers bile salt
333 resistance in *Helicobacter pylori* (55). The multidrug efflux pump CmeABC, of
334 *Campylobacter jejuni* mediates bile salt resistance and is required for colonisation of
335 chickens (56). Gram-positive pathogens such as *Enterococcus faecalis* and *L.*
336 *monocytogenes* also exhibit bile resistance. In addition to bile salt hydrolase

337 activities, both bacteria possess multiple bile efflux systems. Exposure of *E. faecalis*
338 to bile results in up-regulation of two open reading frames EF0420 and EF1814,
339 which are homologous to the QacA family of efflux pumps (57). *L. monocytogenes*
340 OpuC, an osmolyte transporter, as well as specialist bile transporters BilE and MdrT,
341 all confer bile salt resistance to the pathogen (58).

342

343 We demonstrated that the *mnhABCDEFG* operon in *S. aureus* confers bile salt
344 resistance to the pathogen. Previous studies have shown this operon to encode a
345 multi-subunit hetero-oligomeric antiporter system involved in efflux of monovalent
346 cations such as Na⁺, K⁺ and Li⁺ in exchange for H⁺ (59). Transposon insertion into
347 *mnhD* (also called *snoD*) resulted in reduced susceptibility to platelet microbicidal
348 protein 1 (37), thus the operon also has the ability to sensitize the pathogen to other
349 host innate antimicrobials. The function of individual components remains to be
350 determined, however *mnhF* is homologous to a hamster ileal bile salt transporter (60)
351 and rat liver organic anion transporter that was shown to efflux cholic acid (61). A
352 transposon insertion at *mnhA*, which presumably had a polar effect on the rest of the
353 operon and in-frame deletion of *mnhF*, rendered the bacterium equally susceptible to
354 bile salts. Together with our observation that cloning of *mnhF* in *E. coli* increased the
355 bile salt MIC, demonstrated that MnhF alone is sufficient to confer bile salt
356 resistance. Furthermore, MnhF acted to exclude cholic acid from both *S. aureus* and
357 *E. coli*.

358

359 In order to confirm that this increase sensitivity of *S. aureus* translated into a
360 decreased ability of *S. aureus* to survive under conditions found in the human colon,
361 we studied survival of the mutant in a well characterised *in vitro* three-stage system

362 which models the microbial and physicochemical conditions of the in the proximal,
363 transverse and distal colon (30). The $\Delta mnhF$ strain was attenuated in its ability to
364 survive in the model, compared to the parental wild type. To date, no suitable *in*
365 *vivo* models have been developed to study carriage and survival of *S. aureus* in the
366 human intestine. Laboratory mouse models of infection do not reproduce the
367 complex microbial ecosystem or the human gut's physicochemical defences (36).

368

369 The physiology of *S. aureus* in the human gut is very poorly understood, relative to
370 other niches. A recent study to determine *S. aureus* genetic traits associated with
371 observed higher rectal carriage rates was inconclusive (62), thus this is the first report
372 of an *S. aureus* intestinal colonisation factor. Given the complex nature of the gut as a
373 niche, it seems highly likely that other loci are similarly required. Indeed it would
374 appear from our data that other bile resistance factors also exist. As such much
375 remains to be discovered about the behaviour and survival of *S. aureus* in the human
376 gut.

377

378 **Acknowledgements**

379 This work was funded by a PhD Felix Scholarship to THS.

380

381 **References**

382 1. Peacock SJ, de Silva I, Lowy FD. 2001. What determines nasal carriage of
383 *Staphylococcus aureus*? Trends Microbiol. **9**:605-610.

384

385 2. Wertheim HF, Vos MC, Ott A, van Belkum A, Voss A, Kluytmans JA, van Keulen
386 PH, Vandenbroucke-Grauls CM, Meester MH, Verbrugh HA. 2004. Risk and

387 outcome of nosocomial *Staphylococcus aureus* bacteraemia in nasal carriers versus
388 non-carriers. Lancet **364**:703-705.

389

390 3. Acton DS, Plat-Sinnige MJ, van Wamel W, de Groot N, van Belkum A. 2009.
391 Intestinal carriage of *Staphylococcus aureus*: how does its frequency compare with
392 that of nasal carriage and what is its clinical impact? Eur. J. Clin. Microbiol. Infect.
393 Dis. **28**:115-127.

394

395 4. Masaki H, Asoh N, Watanabe H, Tao M, Watanabe K, Ikeda H, Matsumoto K,
396 Oishi K, Nagatake T. 2003. Possible relationship between *Staphylococcus aureus*
397 colonizing the respiratory tract and rectum and *S. aureus* isolated in a geriatric
398 hospital environment. Intern. Med. **42**:281-282.

399

400 5. Bhalla A, Aron DC, Donskey CJ. 2007. *Staphylococcus aureus* intestinal
401 colonization is associated with increased frequency of *S. aureus* on skin of
402 hospitalized patients. BMC Infect. Dis. **7**:105.

403

404 6. Squier C, Rihs JD, Risa KJ, Sagnimeni A, Wagener MM, Stout J, Muder RR,
405 Singh N. 2002. *Staphylococcus aureus* rectal carriage and its association with
406 infections in patients in a surgical intensive care unit and a liver transplant unit.
407 Infect. Control. Hosp. Epidemiol. **23**:495-501.

408

409 7. Ray AJ, Pultz NJ, Bhalla A, Aron DC, Donskey CJ. 2003. Coexistence of
410 vancomycin-resistant enterococci and *Staphylococcus aureus* in the intestinal tracts of
411 hospitalized patients. Clin. Infect. Dis. **37**:875-881.

412

413 8. Froberg MK, Palavecino E, Dykoski R, Gerding DN, Peterson LR, Johnson S.
414 2004. *Staphylococcus aureus* and *Clostridium difficile* cause distinct
415 pseudomembranous intestinal diseases. Clin. Infect. Dis. **39**:747-750.

416

417 9. Björkstén B, Naaber P, Sepp E, Mikelsaar M. 1999. The intestinal microflora in
418 allergic Estonian and Swedish 2-year-old children. Clin. Exp. Allergy **29**:342-346.

419

420 10. Lindberg E, Nowrouzian F, Adlerberth I, Wold AE. 2000. Long-time
421 persistence of superantigen-producing *Staphylococcus aureus* strains in the intestinal
422 microflora of healthy infants. Pediatr. Res. **48**:741-747.

423

424 11. Lindberg E, Adlerberth I, Hesselmar B, Saalman R, Strannegård I-L, Åberg N,
425 Wold AE. 2004. High rate of transfer of *Staphylococcus aureus* from parental skin to
426 infant gut flora. J. Clin. Microbiol. **42**:530-534.

427

428 12. Adlerberth I, Strachan DP, Matricardi PM, Ahrné S, Orfei L, Aberg N, Perkin
429 MR, Tripodi S, Hasselmar B, Saalman R, Coates AR, Bonanno CL, Panetta V, Wold
430 AE. 2007. Gut microbiota and development of atopic eczema in 3 European
431 birth cohorts. J. Allergy Clin. Immunol. **120**:343-350.

432

433 13. Lundell AC, Adlerberth I, Lindberg E, Karlsson H, Ekberg S, Aberg N, Saalman
434 R, Hock B, Steinkasserer A, Hesselmar B, Wold AE, Rudin A. 2007. Increased
435 levels of circulating soluble CD14 but not CD83 in infants are associated with early
436 intestinal colonization with *Staphylococcus aureus*. Clin. Exp. Allergy. **37**:62-71.

437

438 14. Nakamura Y, Aramaki Y, Kakiuchi T. 2001. A mouse model for postoperative
439 fetal enteritis due to *Staphylococcus* infection. J. Surg. Res. **96**:35-43.

440

441 15. Hess DJ, Garni RM, Henry-Stanley MJ, Wells CL. 2005. *Escherichia coli*
442 modulates extraintestinal spread of *Staphylococcus aureus*. Shock **24**:376-381.

443

444 16. Sannasiddappa TH, Costabile A, Gibson GR, Clarke SR. 2011. The influence of
445 *Staphylococcus aureus* on gut microbial ecology in an *in vitro* continuous culture
446 human colonic model system. PLoS One **6**:e23227.

447

448 17. Horsburgh MJ, Aish JL, White IJ, Shaw L, Lithgow JK, Foster SJ. 2002.
449 sigmaB modulates virulence determinant expression and stress resistance:
450 characterization of a functional rsbU strain derived from *Staphylococcus aureus* 8325-
451 4. J. Bacteriol. **184**:5457-5467.

452

453 18. Kreiswirth BN, Löfdahl S, Betley MJ, O'Reilly M, Schlievert PM, Bergdoll MS,
454 Novick RP. 1983. The toxic shock syndrome exotoxin structural gene is not
455 detectably transmitted by a prophage. Nature **305**:709-712.

456

457 19. Camilli A, Portnoy A, Youngman P. 1990. Insertional mutagenesis of *Listeria*
458 *monocytogenes* with a novel Tn917 derivative that allows direct cloning of DNA
459 flanking transposon insertions. J Bacteriol. **172**:3738-3744.

460

- 461 20. Arnaud M, Chastanet A, Débarbouillé M. New vector for efficient allelic
462 replacement in naturally nontransformable, low-GC-content, gram-positive bacteria.
463 2004. *Appl Environ. Microbiol.* **70**:6887-6891.
- 464
465
- 466 21. Guzman LM, Belin D, Carson MJ, Beckwith J. 1995. Tight regulation,
467 modulation, and high-level expression by vectors containing the arabinose PBAD
468 promoter. *J. Bacteriol.* **177**:4121-4130.
- 469
- 470 22. Corrigan, RM, Foster TJ. 2009. An improved tetracycline inducible expression
471 vector for *Staphylococcus aureus*. *Plasmid* **61**:126–129.
- 472
- 473 23. Novick R. 1967. Properties of a cryptic high-frequency transducing phage in
474 *Staphylococcus aureus*. *Virology* **33**:155-166.
- 475
- 476 24. Schenk S, Laddaga RA. 1992. Improved method for electroporation of
477 *Staphylococcus aureus*. *FEMS Microbiol. Lett.* **73**:133-138.
- 478
- 479 25. Quillin SJ, Schwartz KT, Leber JH. 2011. The novel *Listeria monocytogenes*
480 bile sensor BrtA controls expression of the cholic acid efflux pump MdrT. *Mol.*
481 *Microbiol.* **81**:129-142.
- 482
- 483 26. Wolz C, Goerke C, Landmann R, Zimmerli W, Fluckiger U. 2002. Transcription
484 of clumping factor A in attached and unattached *Staphylococcus aureus* in vitro and
485 during device-related infection. *Infect. Immun.* **70**:2758–2762.

486

487 26. Valle J, Toledo-Arana A, Berasain C, Ghigo J-M, Amorena B, Penadés J, Lasa I.
488 2003. SarA and not σ^B is essential for biofilm development by *Staphylococcus*
489 *aureus*. Mol. Microbiol. **48**:1075-1087.

490

491 28. Kenny JG, Ward D, Josefsson E, Jonsson I-M, Hinds J, Rees HH, Lindsay JA,
492 Tarkowski A, Horsburgh MJ. 2009. The *Staphylococcus aureus* response to
493 unsaturated long chain free fatty acids: survival mechanisms and virulence
494 implications. PLOS One **4**:e4344.

495

496 29. Chen L, Shopsin B, Zhao Y, Smyth D, Wasserman GA, Fang C, Liu L,
497 Kreiswirth BN. 2012. Real-time nucleic acid sequence-based amplification assay for
498 rapid detection and quantification of *agr* functionality in clinical *Staphylococcus*
499 *aureus* isolates. J. Clin. Micro. **50**:657-661.

500

501 30. Macfarlane GT, Macfarlane S, Gibson GR. 1998. Validation of a three-stage
502 compound continuous culture system for investigating the effect of retention time on
503 the ecology and metabolism of bacteria in the human colon. Microb. Ecol. **35**:180-
504 187.

505

506 31. Hiramatsu T, Kodama K, Kuroda T, Mizushima T, Tsuchiya T. 1998. A putative
507 multisubunit Na^+/H^+ antiporter from *Staphylococcus aureus*. J. Bacteriol. **180**:6642-
508 6648.

509

- 510 32. Ito M, Guffanti AA, Oudega B, Krulwich TA. 1999. *mnh*, a multigene,
511 multifunctional locus in *Bacillus subtilis* with roles in resistance to cholate and to Na⁺
512 and in pH homeostasis. J Bacteriol. **181**:2394-2402.
- 513
- 514 33. Ito M, Guffanti AA, Wang W, Krulwich TA. 2000. Effects of nonpolar
515 mutations in each of the seven *Bacillus subtilis mnh* genes suggest complex
516 interactions among the gene products in support of Na⁺ and alkali but not cholate
517 resistance. J. Bacteriol. **182**:5663-5670.
- 518
- 519 34. Thanassi DG, Cheng LW, Nikaido H. 1997. Active efflux of bile salts by
520 *Escherichia coli*. J. Bacteriol. **179**:2512-2518.
- 521
- 522 35. Sleator RD, Wemekamp-Kamphuis HH, Gahan CG, Abee T, Hill C. 2005. A
523 PrfA-regulated bile exclusion system (BilE) is a novel virulence factor in *Listeria*
524 *monocytogenes*. Mol. Microbiol. **55**:1183-1195.
- 525
- 526 36. Hapfelmeier S, Hardt WD. 2005. A mouse model for *S. typhimurium*-induced
527 enterocolitis. Trends Microbiol. **13**:497-503.
- 528
- 529 37. Bayer AS, McNamara P, Yeaman MR, Lucindo N, Jones T, Cheung AL, Sahl
530 HG, Proctor RA. 2006. Transposon disruption of the complex I NADH
531 oxidoreductase gene (*snoD*) in *Staphylococcus aureus* is associated with reduced
532 susceptibility to the microbicidal activity of thrombin-induced platelet microbicidal
533 protein 1. J. Bacteriol. **188**:211-222.
- 534

- 535 38. Begley M, Gahan CG, Hill C. 2005. The interaction between bacteria and bile.
536 FEMS Microbiol. Rev. **29**:625-651.
537
- 538 39. Minuk GY, Rascanin N, Sarjeant ES, Pai CH. 1986. Sepsis and cholestasis: the
539 *in vitro* effects of bacterial products on ¹⁴C-taurocholate uptake by isolated rat
540 hepatocytes. Liver **6**:199-204.
541
- 542 40. Esteller A. 2008. Physiology of bile secretion. World J. Gastroenterol.
543 **14**:5641-5649.
544
- 545 41. Scholmerich J, Becher MS, Schmidt K, Schubert R, Kremer B, Feldhaus S,
546 Gerok W. 1984. Influence of hydroxylation and conjugation of bile salts on their
547 membrane-damaging properties--studies on isolated hepatocytes and lipid membrane
548 vesicles. Hepatology **4**:661-666.
549
- 550 42. Hoffman AF. 1999. The continuing importance of bile acids in liver and
551 intestinal disease. Arch. Intern. Med. **159**:2647-2658.
552
- 553 43. Ridlon J, Kang D-J, Hylemon PB. 2006. Bile salt biotransformations by human
554 intestinal bacteria. J. Lipid Res. **47**:241-259.
555
- 556 44. Buffie CG, Bucci V, Stein RR, McKenny PT, Ling L, Gobourne A, No D, Liu H,
557 Kinnebrew M, Viale A, Littmann E, van der Brink MRM, Jenq RR, Taur Y, Sander
558 C, Cross JR, Toussaing NC, Xavier JB, Pamer EG. 2015. Precision microbiome

559 reconstitution restores bile acid mediated resistance to *Clostridium difficile*. Nature
560 **517**:205-208.

561

562 42. Guarner F, Malagelada JR. 2003. Gut flora in health and disease. Lancet
563 **361**:512-519.

564

565 43. Pfeiler EA, Klaenhammer TR. 2009. Role of transporter proteins in bile
566 tolerance of *Lactobacillus acidophilus*. Appl. Environ. Microbiol. **75**:6013-6016.

567

568 44. De Smet I, Van Hoorde L, Vande Woestyne M, Christiaens H, Verstraete W.
569 1995. Significance of bile salt hydrolytic activities of lactobacilli. J. Appl. Bacteriol.
570 **79**:292-301.

571

572 45. Ruiz L, Sanchez B, Ruas-Madiedo P, de Los Reyes-Gavilan CG, Margolles A.
573 2007. Cell envelope changes in *Bifidobacterium animalis* ssp. *lactis* as a response to
574 bile. FEMS Microbiol. Lett. **274**:316-322.

575

576 46. Ruas-Madiedo P, Gueimonde M, Arigoni F, de los Reyes-Gavilan CG, Margolles
577 A. 2009. Bile affects the synthesis of exopolysaccharides by *Bifidobacterium*
578 *animalis*. Appl. Environ. Microbiol. **75**:1204-1207.

579

580 47. Prouty AM, Gunn JS. 2000. *Salmonella enterica* serovar *typhimurium* invasion
581 is repressed in the presence of bile. Infect. Immun. **68**:6763-6769.

582

- 583 48. Peterson KM. 2002. Expression of *Vibrio cholerae* virulence genes in response
584 to environmental signals. *Curr. Issues Intest. Microbiol.* **3**:29-38.
585
- 586 49. Nesper J, Schild S, Lauriano CM, Kraiss A, Klose KE, Reidl J. 2002. Role of
587 *Vibrio cholerae* O139 surface polysaccharides in intestinal colonization. *Infect.*
588 *Immun.* **70**:5990-5996.
589
- 590 50. Crawford RW, Kestra AM, Winter SE, Xavier MN, Tsolis RM, Tolstikov V,
591 Bäumlér AJ. 2012. Very long O-antigen chains enhance fitness during *Salmonella*-
592 induced colitis by increasing bile resistance. *PLoS Pathog.* **8**:e1002918.
593
- 594 51. Prouty AM, Brodsky IE, Falkow S, Gunn JS. 2004. Bile-salt-mediated induction
595 of antimicrobial and bile resistance in *Salmonella typhimurium*. *Microbiology*;
596 **150**:775-783.
597
- 598 52. Trainor EA, Horton KE, Savage PB, Testerman TL, McGee DJ. 2011. Role of
599 the HefC efflux pump in *Helicobacter pylori* cholesterol-dependent resistance to
600 ceragenins and bile salts. *Infect. Immun.* **79**:88-97.
601
- 602 53. Lin J, Sahin O, Michel LO, Zhang Q. 2003. Critical role of multidrug efflux
603 pump CmeABC in bile resistance and in vivo colonization of *Campylobacter jejuni*.
604 *Infect. Immun.* **71**:4250-4259.
605

- 606 54. Solheim M, Aakra A, Vebo H, Snipen L, Nes IF. 2007. Transcriptional responses
607 of *Enterococcus faecalis* V583 to bovine bile and sodium dodecyl sulfate. Appl.
608 Environ. Microbiol. **73**:5767-5774.
- 609
- 610 55. Begley M, Sleaton RD, Gahan GC, Hill C. 2005. Contribution of three bile-
611 associated loci *bsh*, *pva*, and *bltB*, to gastrointestinal persistence and bile tolerance of
612 *Listeria monocytogenes*. Infect. Immun. **73**:894-904.
- 613
- 614 56. Swartz TH, Ito M, Ohira T, Natsui S, Hicks DB, Krulwich TA. 2007. Catalytic
615 properties of *Staphylococcus aureus* and *Bacillus* members of the secondary
616 cation/proton antiporter-3 (Mrp) family are revealed by an optimized assay in an
617 *Escherichia coli* host. J. Bacteriol. **189**:3081-3090.
- 618
- 619 57. Wong MH, Oelkers P, Craddock AL, Dawson PA. 1994. Expression cloning
620 and characterization of the hamster ileal sodium-dependent bile acid transporter. J.
621 Biol. Chem. **269**:1340-1347.
- 622
- 623 58. Jacquemin E, Hagenbuch B, Stieger B, Wolkoff AW, Meier PJ. 1994.
624 Expression cloning of a rat liver Na(+)-independent organic anion transporter. Proc.
625 Natl. Acad. Sci. USA. **91**:133-137.
- 626
- 627 59. Lemmens N, van Wamel W, Snijders S, Lesse AJ, Faden H, van Belkum A.
628 2011. Genomic comparisons of USA300 *Staphylococcus aureus* colonizing the
629 nose and rectum of children with skin abscesses. Microb. Pathog. **50**:192-199.
- 630

631 **Table 1. Bacterial strains**

Strain	Description/Genotype	Source or Reference
<i>S. aureus</i> SH1000	Wild type	(17)
<i>S. aureus</i> SH1001	<i>agr</i> mutation in SH1000	(17)
<i>S. aureus</i> RN4220	Accepts <i>E. coli</i> DNA	(18)
<i>S. aureus</i> <i>mnhA::Tn917</i>	Tn917 inserted into <i>mnhA</i> in SH1000	This study
<i>S. aureus</i> Δ <i>mnhF</i>	Δ <i>mnhF</i> mutation in SH1000	This study
<i>E. coli</i> Top10	<i>F- mcrA</i> Δ (<i>mrr-hsdRMS-mcrBC</i>) ϕ 80 <i>lacZ</i> Δ <i>M15</i> Δ <i>lacX74</i> <i>nupG</i> <i>recA1</i> <i>araD139</i> Δ (<i>ara-leu</i>)7697 <i>galE15</i> <i>galK16</i> <i>rpsL</i> (<i>Str^R</i>) <i>endA1</i> λ^-	Invitrogen
<i>E. coli</i> TG1	<i>F'</i> [<i>traD36 proAB⁺ lacI^q lacZ</i> Δ <i>M15</i>] <i>supE</i> <i>thi-1</i> Δ (<i>lac-proAB</i>) Δ (<i>mcrB-</i> <i>hsdSM</i>)5, (<i>r_K⁻ m_K⁻</i>)	Lucigen

632

633

634 **Table 2. Plasmids**

Plasmid name	Description	Antibiotic resistance	Source or Reference
pLTV1	Carries Tn917	Em ^R /Tc ^R	(19)
pMAD	Temperature sensitive (30°C) <i>E. coli</i> – <i>S. aureus</i> shuttle vector. pE194 ^{ts} ::pBR322	Em ^R	(20)
pBAD His A	Expression vector containing <i>araBAD</i> promoter	Ap ^R	(21)
pRMC2	<i>S. aureus</i> expression vector	Ap ^R /Cm ^R	(22)
pΔ <i>mnhF</i>	Vector for Δ <i>mnhF</i> mutation	Em ^R	This study
pMnhF1	pBAD His A containing <i>mnhF</i> internal fragment	Ap ^R	This study
pMnhF2	pRMC2 containing <i>mnhF</i> internal fragment	Ap ^R /Cm ^R	This study

635

636

637

638 **Table 3. Oligonucleotides.** Restriction endonuclease sites are underlined

Name	Sequence 5'-3'
<i>ΔmnhFLFor</i>	CCAAAAGGATCCGATCTTAATAAC
<i>ΔmnhFLRev</i>	CATTAGAATTCATTATATTTTCGCCACC
<i>ΔmnhFRFor</i>	TATGGAATTCGGTAAGGTGATTGAAC
<i>ΔmnhFRRev</i>	GCGATTGCGGATCCCTGTATGCC
<i>mnhFFor1</i>	GGGCGAAATATCATGAATCATAATG
<i>mnhFFor2</i>	GGGCGAAATAGGATCCATCATAATG
<i>mnhFRev</i>	TGATGAATTCGATAAGTGCAAGACTAATC
RNAIIIFor	ACATGGTTATTAAGTTGGGATGG
RNAIIIRev	TAAAATGGATTATCGACACAGTGA
<i>gyrBFor</i>	ATCGACTTCAGAGAGAGGTTTG
<i>gyrBRev</i>	CCGTTATCCGTTACTTTAATCCA

639

640

641

642

643

644 **Table 4. MICs of bile salts for *S. aureus* SH1000 and $\Delta mnhF$**

Bile salt	Wild type (mM)	$\Delta mnhF$ (mM)	$\Delta mnhF$ [pMnhF2] (mM)	$\Delta mnhF$ [pRMC2] (mM)
CA	22	5	22	5
DCA	1.2	0.6	1.2	0.6
CDCA	1.2	0.6	1.2	0.6
GCA	>200	>200	ND	ND
TCA	>200	>200	ND	ND

645
646 **NOTE.** CA, sodium cholate; DCA, sodium deoxycholate; CDCA, sodium
647 chenodeoxycholate; GCA, sodium glycocholate; TCA, sodium taurocholate. ND, not
648 determined.

649
650
651
652
653
654
655
656
657
658
659
660
661

662 **Table 5. MICs of bile salts for wild type and recombinant *E. coli* strains**
 663 **expressing MnhF at different levels of arabinose induction**

Bile salt	Wild type		Vector control		Recombinants					
	TG1	TOP10	TG1 pBAD	TOP10 pBAD	TG1 pMnhF1			TOP10 pMnhF1		
					0%	0.02%	2%	0%	0.02%	2%
					Arabinose	Arabinose	Arabinose	Arabinose	Arabinose	Arabinose
CA	30	30	30	30	30	60	90	30	60	90
DCA	4	4	4	4	4	>4	>4	4	>4	>4
CDCA	4	4	4	4	4	>4	>4	4	>4	>4
GCA	50	50	50	50	50	100	100	50	100	100
TCA	50	50	50	50	50	100	100	50	100	100

664
 665 **NOTE.** CA, sodium cholate; DCA, sodium deoxycholate; CDCA, sodium
 666 chenodeoxycholate; GCA, sodium glycocholate; TCA, sodium taurocholate.
 667
 668
 669
 670
 671
 672
 673

674 **Table 6.** Effect of efflux pump inhibitors on MICs of bile salts to (A) *S. aureus* and
 675 (B) *E. coli*.

676
 677
 678
 679

A.

Bile salt	<i>S. aureus</i> SH1000 (mM)			<i>S. aureus</i> Δ <i>mnhF</i> (mM)		
	Control	^a PA β N	^a Reserpine	Control	^a PA β N	^a Reserpine
CA	22	2.5	10	5	2.5	2.5
DCA	1.2	0.3	1.2	0.6	0.3	0.3
CDCA	1.2	0.3	1.2	0.6	0.3	0.3
GCA	>200	200	>200	>200	200	>200
TCA	>200	200	>200	>200	200	>200

680
 681
 682
 683

B.

Bile salt	<i>E. coli</i> TG1 (mM)			<i>E. coli</i> TG1 p <i>MnhF1</i> (mM)		
	Control	^a PA β N	^a Reserpine	Control	^a PA β N	^a Reserpine
CA	30	2.5	30	90	2.5	90
DCA	4	0.6	>4	>4	0.6	>4
CDCA	4	0.6	>4	>4	0.6	>4
GCA	50	10	50	100	10	100
TCA	50	10	50	100	10	100

684
 685
 686
 687
 688
 689

NOTE. CA, sodium cholate; DCA, sodium deoxycholate; CDCA, sodium chenodeoxycholate; GCA, sodium glycocholate; TCA, sodium taurocholate; PA β N, Phe-Arg- β -naphthylamide. ^aPA β N at 20 μ g/ml and Reserpine at 40 μ g/ml.

690 **Figure Legends**

691 **Figure 1. MnhF protects *S. aureus* against the bactericidal activity of bile salts.**

692 Viability of *S. aureus* SH1000 [■], $\Delta mnhF$ [▲], $\Delta mnhF$ (pMnhF2) [◆], $\Delta mnhF$
693 (pRMC2) [×] treated with (A) 2 mM CA, (B) 0.25 mM DCA, and (C) 20 mM GCA.

694 Data represents mean \pm standard error of mean from three independent experiments.

695 *P<0.01, †P>0.05.

696

697 **Figure 2. Heterologous expression of MnhF in *E. coli* protects against the**

698 **bacteriostatic effects of bile salts.** Viability of wild type *E. coli* TG1 and *E. coli*

699 TG1 pMnhF1 cells in LB medium containing CA (10 and 20 mM), DCA (2 and 4

700 mM) and GCA (25 and 50 mM) and then grown for overnight at 37°C. Cell counts

701 were then determined by viable plate counting. Data represents mean \pm standard error

702 of mean from three independent experiments. * P<0.001

703

704 **Figure 3. MnhF exports cholic acid.** (A) *S. aureus* SH1000 wild type [■] and

705 $\Delta mnhF$ [▲] cells were loaded with 1 μ Ci of ¹⁴C-cholic acid, and then diluted into a

706 buffer containing excess of non-radiolabelled cholic acid (0.2 mM). (B) *E. coli* TG1

707 parental type (TG1) [▲], *E. coli* TG1 expressing pBAD (TG1: pBAD) [◆] and *E.*

708 *coli* TG1 expressing pMnhF1 (TG1: pMnhF1) [■] cells grown overnight in LB under

709 1% arabinose induction, were loaded with 1 μ Ci of ¹⁴C- cholic acid, and then diluted

710 into a buffer containing excess of non-radiolabelled cholic acid (0.2 mM) and 1%

711 arabinose. At indicated times, the amount of retained ¹⁴C-cholic acid in cell pellets

712 were determined by liquid scintillation counting. Data represents mean \pm standard

713 error of mean of three independent experiments. *P<0.05

714

715 **Figure 4. Mutation of *mnhF* does not affect *agr*.** qRT-PCR was performed in
716 order to quantify amounts of RNAIII in *S. aureus* strains during exponential and
717 stationary phases of growth. Data represents mean \pm standard error of mean of three
718 independent experiments. *P>0.05.

719

720 **Figure 5. MnhF is required for *S. aureus* survival in the human colonic model.**
721 Survival of *S. aureus* SH1000 [■] and $\Delta mnhF$ [▲] in the human colonic model. (A)
722 V1, models the ascending colon, (B) V2 model the transverse colon and (C) V3
723 models the descending colon. Samples were taken at inoculation (0 h) and 4, 8, 24,
724 48, 72, and 96 hours post infection. Results are reported as means (Log₁₀ CFU/mL)
725 of the data of three colonic models \pm standard error of mean. *P<0.05; **P<0.001.






Figure 1

□ TG1: pMnhF ▨ TG1: pMnhF@0.02% Arabinose ■ TG1: pMnhF@2% Arabinose




Figure 2





Figure 3




Figure 4




Figure 5