University of Reading

Evaluation of optical techniques for characterising soil organic matter quality in agricultural soils

Article

Accepted Version

Fernández-Romero, M.L., Clark, J.M. ORCID: https://orcid.org/0000-0002-0412-8824, Collins, C.D., Parras-Alcántara, L. and Lozano-García, B. (2016) Evaluation of optical techniques for characterising soil organic matter quality in agricultural soils. Soil and Tillage Research, 155. pp. 450-460. ISSN 0167-1987 doi: 10.1016/j.still.2015.05.004 Available at https://centaur.reading.ac.uk/40551/

It is advisable to refer to the publisher's version if you intend to cite from the work. See Guidance on citing.
Published version at: http://dx.doi.org/10.1016/j.still.2015.05.004
To link to this article DOI: http://dx.doi.org/10.1016/j.still.2015.05.004

Publisher: Elsevier

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the End User Agreement.

www.reading.ac.uk/centaur

University of Reading

CentAUR

Central Archive at the University of Reading

Reading's research outputs online

EVALUATION OF OPTICAL TECHNIQUES FOR CHARACTERISING SOIL ORGANIC MATTER QUALITY IN AGRICULTURAL SOILS

M.L. Fernández-Romero [a *], J.M. Clark [b], C.D. Collins [b], L. Parras-Alcántara [a], B. Lozano-García [a]

[a] Department of Agricultural Chemistry and Soil Science, Faculty of Science, Agrifood Campus of International Excellence (ceiA3), Universidad de Córdoba, 14071 Cordoba, Spain

[b] Soil Research Centre, Department of Geography and Environmental Sciences, School of Human and Environmental Sciences, University of Reading, Whiteknights, RG6 6AB, Reading, UK

*Corresponding author: María Luisa Fernández-Romero. Tel.: +34 667964310; +44 7790399784. E-mail addresses: a52ferom@uco.es; fdezmarialuisa@gmail.com

ABSTRACT

Soil organic matter (SOM) is one of the main global carbon pools. It is a measure of soil quality as its presence increases carbon sequestration and improves physical and chemical soil properties. The determination and characterisation of humic substances gives essential information of the maturity and stresses of soils as well as of their health. However, the determination of the exact nature and molecular structure of these substances has been proven difficult. Several complex techniques exist to characterise SOM and mineralisation and humification processes. One of the more widely accepted for its accuracy is Nuclear Magnetic Resonance (NMR) spectroscopy. Despite its efficacy, NMR needs significant economic resources, equipment, material and time. Proxy measures like the fluorescence index (FI), cold and hot-water extractable carbon (CWC and HWC) and SUVA-$_{254}$ have the potential to characterise SOM and, in combination, provide qualitative and quantitative data of SOM and its processes. Spanish and British agricultural cambisols were used to measure SOM quality and determine whether similarities were found between optical techniques and ^{1}H-NMR results in these two regions with contrasting climatic conditions. High correlations ($p<0.001$) were found between the specific aromatic fraction measured with ^{1}H-NMR and SUVA-$_{254}$ (Rs=0.95) and HWC (Rs=0.90), which could be described using a linear model. A high correlation between FI and the aromatics fraction measured with ^{1}H-

Abbreviations: (SOM) Soil organic matter; (NMR) Nuclear magnetic resonance; (FI) Fluorescence index; (CWC) Cold-water extractable carbon; (HWC) Hot-water extractable carbon; (LOI) Loss of ignition; (DOM) Dissolved organic matter; (DOC) Dissolved organic carbon; (EEM) Excitation-emission matrix; (GS-UK) Soil with a grass cover, United Kingdom; (CC-UK) Cereal crops from United Kingdom; (CC-ES) Cereal crops from Spain; (OG-ES) Olive grove from Spain; (HIX) Humification index; (BIX) Biological/autochthonous index; (RU) Raman units; (SUVA-254) Specific absorbance at 254 nm

NMR (Rs=-0.976) was also observed. In view of our results, optical measures have a potential, in combination, to predict the aromatic fraction of SOM without the need of expensive and time consuming techniques.

KEYWORDS**:** SOM quality; hot-water extractable carbon; cold-water extractable carbon; fluorescence index; EEM; SUVA-$_{254}$; ^{1}H-NMR; aromatic fraction.

1. Introduction:

1.1. The importance of organic matter

Soil organic matter (SOM) is composed of organic residues that are originated from plant and animal remains and microbial products at different stages of decomposition or humification (Hur *et al.*, 2013). Additionally, it is one of the main global carbon pools, storing three times more carbon than living organisms or the atmosphere (Fischlin *et al.*, 2007; Brevik, 2012). Aside from carbon sequestration, SOM is also a measure of soil quality because of the beneficial function it has on a variety of soil processes. For instance, it reduces erosion and, therefore, increases crop production by increasing the elasticity and resistance to deformation and compactability as well as porosity and water retention (Sellami *et al.*, 2008; Paradelo and Barral, 2013). Increased water retention decreases potential runoffs by improving water infiltration in to soils and provides a store of water for plant uptake, buffering against moisture and rainfall fluctuations (Lal, 2004). This is of importance considering that the lack of water retention leads to a change in the hydrological patterns of agricultural areas and promotes the quantity and severity of floods and water-led erosion. Also, SOM leads to an increased vegetative cover, which ultimately reduces soil erosion (Cerdà, 1998, 2000; Novara *et al.*, 2011; Zhao *et al.*, 2013). Carbon mineralisation is crucial in SOM dynamics and along with carbon input, determines how much carbon accumulates in soil and releases nutrients that are essential for plant growth. Factors that affect mineralisation are the size of

67 labile carbon, environmental conditions and the local microbial community (Zhao *et al.*,
68 2008; Li *et al.*, 2013). SOM and soil assemblage; SOM decomposition and transport by
69 organisms contribute to soil stabilisation and the improvement of soil structure (Brevik
70 *et al.*, 2015). Moreover, SOM quantity has been directly related to the preservation of
71 soil aggregates, which in turn reduces soil erodibility (Novara *et al.*, 2011). Also, the
72 direct processing of SOM along with its decomposition contribute to the improvement
73 of soil chemical properties and stability (Brevik *et al.*, 2015). Therefore, optimal
74 quantities of SOM improve structure, water retention, and nutrient holding capacity of
75 soils, which has an effect in multiple aspects of the soil system. These are essential
76 considering the wider context of Earth System, as SOM conservation techniques have
77 been proven to improve the fertility of degraded soils of a wide variety of ecosystems
78 that are the main resource of large communities of developing areas of our planet, as
79 well as have an influence on biogeochemical cycles and climate change mitigation
80 (Batjes, 2014; Saha *et al.*, 2014; Srinivasarao *et al.*, 2014).

81 *1.2. SOM carbon fractions and their importance*

82 SOM has been conceptualised as containing three pools, with different
83 residence/turnover times (Trumbore, 2000). These pools are the active SOM (living
84 biomass of microorganisms and partially decomposed residues; associated with 1 year
85 turnover); the slow SOM (resistant plant material; associated to a turnover from years to
86 centuries); and passive SOM (humic substances and inert organic matter), which has
87 been traditionally associated with longer residence time (thousands of years) and more
88 stability. Some authors consider that the inert organic matter should not be considered
89 as part of the passive pool, but as a fourth pool (Trumbore, 1997; Ohno, 2002; Agren
90 and Bosatta, 2002; Sparks, 2003; Bell and Lawrence, 2009; Dungait *et al.*, 2012).
91 Although the traditional view has been that decomposition led to complex molecules

that were very stable as a result of their structure; it has recently been known that environmental conditions, organo-mineral associations and other processes influence more in SOM stability than structure, which only plays a secondary role. As a result of this new view, recent research has found that humic substances, which have always been considered high molecular mass polymers, could be simpler than originally thought (Kleber and Johnson, 2010; Schmidt *et al.*, 2011). Still, their structure is on discussion and the separation of SOM into fractions with different turnovers remains a major challenge (Kleber, 2010; Schmidt *et al.*, 2011; Schrumpf and Kaiser, 2015). Non-humic substances are composed by microbial biomass, decomposable plant material (active SOM); and resistant plant material, mainly waxes, lignified tissues and polyphenols (slow SOM) (Dungait *et al.*, 2012). Microbial biomass has been used for comparing natural and degraded ecosystems and as an early indicator of soil processes, fertility and health (García-Gil *et al.*, 2000; Brevik, 2009; Chen *et al.*, 2013).

1.3. Current SOM quality measurements. Challenges

Measures such as % Organic Matter measured by Loss of Ignition (LOI) are useful, popular and inexpensive methods to determine bulk SOM (Luke *et al.*, 2009; Salehi *et al.*, 2011). However, some studies have concluded that bulk SOM measurements cannot be used, on their own, as a representative indicator of carbon in soil due to their limitations (Koarashi *et al.*, 2005; Salehi *et al.*, 2011).

Humic substances have also been measured to determine soil quality, as their presence has been associated with a higher quality of soils as stated in section 1.2. Their study is relevant in agricultural soils, as they increase crop yield and root dry weight, although this response is not fully understood (Rose *et al.*, 2014). Therefore, the determination and characterisation of humic substances gives essential information of the maturity and stresses of soils as well as of their health.

Traditionally, alkali and acid abstraction methods have been used, to later interpret the chemistry of the extracted functional groups (Olk and Gregorich, 2006). Afterwards, these were combined with other complex techniques that enabled scientists to obtain new information on the structure and dynamic associations of humic substances (Sutton and Sposito, 2005; Schmidt *et al.*, 2011). Despite these advances, SOM dynamics and cycling still have many questions to answer, with models differing in SOM fluxes results for the future, due to their sensitivity to SOM turnover time assumptions (Schmidt *et al.*, 2011). There are a number of powerful but complex and expensive techniques that have been used for the study of soil fluxes (Helal *et al.*, 2011). The economic resources needed, along with the time required to prepare the samples and conduct the analyses, make its use with a large number of samples difficult and delays experiments, while more work is still needed to accurately determine and define the molecular structures and linkages between the SOM components (Weishar *et al.* 2003; Helal *et al.*, 2011).

Nuclear Magnetic Resonance (NMR) spectroscopy is a non-destructive technique that is valuable for the characterisation of SOM and humification processes, providing information on static and dynamic properties of molecules. This is due to its high performance to assess intermolecular interactions. The relationship between SOM, contaminants and metals can also be studied with NMR (Cardoza *et al.*, 2004). Of the various variants that exist, ^{1}H-NMR spectroscopy was used in this study. This technique analyses humic and fulvic acids dissolved in neutral or alkaline solutions to characterise the components of the substance, and gives a semi-quantitative notion of aromatic, aliphatic and carboxylic groups (Hemminga and Buurman, 1997). One of the main drawbacks of this technique is the quantity of economic resources that are necessary for its regular application in research laboratories/centres. This is due to the expensive

142 deuterated solvents and NMR tubes, as well as the expensive equipment and significant
143 sample preparation that are required (Weishar *et al.*, 2003; Cardoza *et al.*, 2004; MIT,
144 2008). Also, the technique is time consuming not only when measuring, but when
145 interpreting 2-D or 3-D data resulting from it (Cardoza *et al.*, 2004). Simpler methods
146 for the characterisation of SOM are required.

147 *1.4. Proxy measures. Opportunities to improve the ability to characterise SOM*
148 *quality*

149 Water extractable carbon is the most active component in the carbon cycle. Its quantity
150 and biological nature is affected by the extraction temperature (Bu *et al.*, 2010). Hot-
151 water extractable carbon (HWC) contains simple compounds such as microorganisms,
152 soluble carbohydrates and other compounds that account for the labile fraction of SOM
153 (Ghani *et al.*, 2003). HWC responds to land use changes in the short term and has been
154 used to detect the effects of different land management practices and for determining
155 the effects of soil amendments such as biochar or agricultural residues (Leifeld and
156 Kögel-Knabner, 2005; Uchida *et al.*, 2012; Alburquerque *et al.*, 2014; Fernández-
157 Romero *et al.*, 2014). For these reasons, it has been proven useful to obtain information
158 about soil quality (Ghani *et al.*, 2003; Xue *et al.*, 2013).

159 Fluorescence has become popular because of its potential to characterise SOM and
160 study humic substances, as it is non-destructive, simple, non-separative and accurate. As
161 a result, it has been used for determining the compositional and structural properties of
162 SOM (Chen *et al.*, 2003; Senesi and D'Orazio, 2005; Sun *et al.*, 2007; Kwiatkowska *et*
163 *al.*, 2008; Henderson *et al.*, 2009; Hur and Kim, 2009; Tang *et al.*, 2011). The intensity
164 and position of the peaks detected in the spectra are unique to each substance structural
165 and functional characteristics. For instance, higher fluorescence intensities are related
166 to a higher humification (Martins *et al.*, 2011).

The Fluorescence Index (FI) was developed to assess different properties of dissolved organic matter (DOM). It was defined by McKnight *et al.* (2001) as the ratio of emission intensities at 450-500 nm excited at 370 nm. The 450 nm point was chosen for specific characteristics of the experiment. Later, Cory *et al.* (2010) modified the ratio to 470-520 nm to reflect corrections specific to the instruments used. This index has been correlated to the aromaticity of DOM (Korak *et al.*, 2014).

Fluorescence spectroscopy can be used in combination with UV-Visible spectroscopy to characterise humic substances, as absorbance measures transitions from the ground state to the excited state, as opposed to fluorescence spectroscopy (Skoog *et al.*, 2007). Its spectra are usually uniform and provide with qualitative data when a specific wavelength is selected (Hassouna, *et al.*, 2012). Also the specific absorbance at 254 nm (SUVA-$_{254}$) has been recognised as a method to determine SOM aromaticity (Fuentes *et al.*, 2006; Chow 2006). This parameter is very useful for assessing the nature of the general composition of dissolved organic carbon (DOC), due to its high correlation with it (Weishar *et al.*, 2003).

Considering what has been described in sections 1.3 and 1.4 above, both the fluorescence and NMR techniques can be used in combination to determine the humic substances properties and the degree of aromaticity; while HWC could contribute further to the understanding of soil quality, given its usefulness to detect the biodegradation of soil biochemical properties (Ghani *et al.*, 2003; Saab and Martin-Neto, 2007; González-Pérez *et al.*, 2007).

As an illustrative and additional way to characterise and represent some of the analyses conducted and the results obtained, Excitation-Emission Matrix (EEM) spectra have been plotted. These provide information on the relative intensity of fluorescence at different excitation and emission wavelengths regions in a fast manner that is also easy

192 to interpret (Coble, 1996). Several peaks have been identified that are used to describe
193 EEM fluorescence spectra. Peak A and C refer to humic peaks. Peak A is referred to as
194 UVC-excited and is located at an excitation wavelength (λ_{Ex}) between 240-260 nm and
195 an emission wavelength (λ_{Em}) between 400-460 nm. Peak C, also referred to as UVA-
196 excited; is located at a λ_{Ex} between 320-360 nm and a λ_{Em} between 420-460 nm. There
197 are also peaks that indicate biological activity material (peaks B and T, which are
198 defined as tyrosine-like and tryptophan-like peaks respectively). B has λ_{Ex} of 270-280
199 nm and λ_{Em} of 300-315 nm whereas T has λ_{Ex} between 270-280 nm and λ_{Em} of 345-
200 360 nm (Birdwell and Engel, 2010).

201 *1.5. Aim/objective of this study*

202 The aim of this study was to evaluate the use of fluorescence spectroscopy to measure
203 SOM quality (specifically the grade of humification). The specific objectives were: (1)
204 characterise water extractable SOM quality using liquid state ^{1}H-NMR; (2) characterise
205 the quality of water extractable organic matter using fluorescence spectroscopy and UV-
206 VIS; (3) compare measures of quantity and quality of water extractable organic matter
207 with the specific organic matter fractions measured by ^{1}H-NMR like aromaticity. If
208 robust relationships and similarities between optical measures and ^{1}H-NMR are found,
209 there may be potential for fluorescence spectroscopy as a fast and more cost-effective
210 method of organic matter characterisation.

211 **2. Materials and methods**

212 *2.1. Field sites description*

213 Cambisols were sampled in two regions with contrasting climatic conditions;
214 Andalusia (South Spain) and Berkshire (South East England) (Table 1). Berkshire has a
215 temperate climate, characterised generally by relatively mild winters and summers and
216 rainfall throughout the year. The annual mean temperature ranges from 6.7 ºC and

14.5 ºC (30 years, annual mean temperature: 10.5 ºC) and the average annual rainfall is 635.4 mm (UK Met Office, 2014).

Andalusia has a Mediterranean climate, which characteristics are hot and dry summers contrasted with cool and wet winters. In the province of Cordoba (Hinojosa del Duque and Pozoblanco), the annual mean temperature is 17.6 ºC and the average annual rainfall is 536 mm (Aemet, 2014). As for the province of Jaen (Torredelcampo), its annual mean temperature is 16.2 ºC and the average annual rainfall is 646.3 mm (REDIAM, 2007).

In Berkshire, samples were collected from University of Reading Farms at Sonning, Arborfield and Shinfield. The site sampled at Sonning (GS-UK-1) has been covered with grass for over 15 years. Arborfield (CC-UK-1) was permanent pasture until autumn 2012, and has been subsequently drilled with wheat or winter barley. Shinfield (CC-UK-2) has been in an arable rotation for over 20 years and sown with either winter wheat, maize or spring barley.

In Andalusia, soil samples were collected between the provinces of Jaen (Torredelcampo) and Cordoba (Hinojosa del Duque, Pozoblanco) that were managed with conventional tillage of cereals and olive crops. CC-ES-3 was managed with a wheat-barley-fallow cycle, whereas the other sites (CC-ES-1 and CC-ES-2) were covered by wheat crops throughout the year. OG-ES-1, OG-ES-2, and OG-ES-3 were covered by olive grove.

2.2. Sample collection and preparation

Soil samples were collected from each horizon. Total soil depths are included in Table 2. Only the first horizon from each soil (specified in section 2.3) was used for the analyses. Samples were air dried and sieved with a 2 mm sieve.

2.3. Analytical methods

Cold and hot water extractable carbon were determined following Ghani *et al.* (2003). This consists in a cold and a hot extraction of the supernatant of the samples (room temperature and 80 ºC respectively). To do this, 30 ml of ultrapure water was added to 3 g of soil. Then, there was a 30-minute extraction in a shaker at 20ºC. After this, the sample was centrifuged during 20 minutes at 3500 rpm. Once this was done, the supernatant was extracted (cold extraction) and analysed (DOC). The resulting pellet was used for the rest of the steps. 30 ml of ultrapure water were added to the pellet, which was then shaken on a Vortex to mix the ultrapure water with the pellet. The sample was then left in a bath at 80 ºC during 16 hours. The sample was centrifuged at 3500 rpm during 20 minutes and was filtered with a 0.45 µm cellulose nitrate filter (hot extraction). The filtered supernatant was analysed for DOC, fluorescence, absorbance and NMR.

The techniques that were compared in this study were applied to the first horizon of each soil sample, as this was considered sufficient for the purposes of comparing these techniques. This horizon was at different depths depending on the soil sample (at 12 cm in GS-UK-1; 10.5 cm in CC-UK-1; 5.7 cm in CC-UK-2; 15 cm in CC-ES-1; 20 cm in CC-ES-2; 30 cm in CC-ES-3; 10 cm in OG-ES-1; 10 cm in OG-ES-2 and 18 cm in OG-ES-3).

Fluorescence of all the hot water extracts was measured in a Varian Eclipse Fluorescence spectrophotometer (Agilent Technologies, Santa Clara, CA, USA) at an emission wavelength from 300 to 600 nm at 5-nm increments and an excitation of 240-450 nm at 5-nm increments. All samples were run in 1 cm quartz cuvettes and in triplicate.

FI was calculated as the ratio of intensities at 450 over 500 nm with an excitation of 370 nm, as described by Cory *et al.* (2010). McKnight *et al.* (2001) introduced this index approach for the characterization of the fulvic and fraction of DOM.

The Humification Index (HIX) was calculated with the following formula:

$$(\sum I_{435\rightarrow480}) / (\sum I_{300\rightarrow345}) + (\sum I_{435\rightarrow480})$$

where I is the fluorescence intensity at each wavelength (modified of Zsolnay *et al.*, 1999 by Ohno, 2002).

The Biological/Autochthonous Index (BIX) has been calculated as the ratio of intensities at 380 nm over 430 nm with an excitation of 310 nm, as described by Huguet *et al.* (2009). This index assesses the relative contribution of autochthonous DOM in water and soil samples.

A subsample of the HWC extract was frozen and subsequently freeze-dried to remove all the water present. Deuterium oxide was added as a solvent to avoid disruptions in the spectrum, as other solvents have a proton signal that causes disruptions, as demonstrated by Cardoza *et al.* (2004), prior to a second freezing and a second freeze-drying. This second stage of freezing-freeze-drying was used to avoid that H_2O peaks interfered in the spectra. After that, deuterium oxide (100%, density of 1.107 g/ml at 25 °C) was added again as a solvent to conduct the NMR tests. NMR was measured in 1.5-1.7 ml of sample with a Bruker Avance III 700 MHz NMR spectrometer (Bruker Corporation, Billerica, MA, USA). The deuterium oxide peak was used as a calibration reference and placed in 4.75 ppm.

Absorbance of all the hot water extracts was measured with a Varian Cary 300 UV-Visible spectrometer (Agilent Technologies, Santa Clara, CA, USA), from 200-800 nm with 1 nm intervals using a 1 cm cuvette.

290 SUVA-$_{254}$ has been used for this study. This is the absorbance at 254 nm divided by the
291 DOC concentration (of the hot water extracts). Values are expressed in $l \cdot mg^{-1} \cdot m^{-1}$.
292 DOC of the cold and hot water extracts was calculated with a Shimadzu TOC 5000 total
293 organic carbon analyser (Shimadzu Corporation, Kyoto, Japan). Standards were
294 calculated using Stock solutions of 1,000 ppm.
295 % Organic matter measured by LOI was calculated using a modified version of Hieri *et*
296 *al.* (2001). A soil sample of 10 g was heated at 105 ºC for 24 hours. Then, the sample
297 was weighted again (w_1) and heated in a Muffle furnace at 550 ºC for 16 hours to ignite
298 the organic matter. The sample was weighted after this (w_2) and the % of organic matter
299 was obtained by weight difference between w_1 and w_2.
300 Soil pH was determined in 1:2.5 soil to water ratio. Texture was determined by laser
301 granulometry, using a Coulter LS 230 (Beckman/Coulter Inc. Brea, CA, USA). This
302 technique uses polarised light at three different wavelengths (450 nm, 600 nm and
303 900 nm) to analyse the particle size distribution specifically in the 0.04 µm to 0.4 µm
304 range.
305 EEM fluorescence spectra were obtained by collecting a series of emission scans of λ_{Ex}
306 240-450 nm at 5 nm intervals and λ_{Em} 300-600 nm, also at 5 nm intervals. EEM spectra
307 were plotted using RStudio v0.98.1091 (Rstudio Inc., Boston, MA, USA).
308 Intensities are reported in Raman units (RU). Raman scattering was mitigated
309 subtracting blanks that had been collected on ultrapure water from each spectrum.

310 *2.4. Statistical methods*

311 Data were tested for normality to verify the model assumptions. As the data failed the
312 normality test, non-parametric tests were used. The effect of the hot water extraction in
313 the extractability of carbon (compared to the cold water extraction) was analysed using
314 a Paired Sample Wilcoxon Signed Rank Test. The correlation between the different

315 analytical methods was tested using the Spearman Rank and assessing the significance
316 of the resulted Spearman Correlation Coefficient (Minitab 16 for Windows. Minitab
317 Inc., State College, PA, USA). Linear regressions of the Spearman Correlations were
318 also plotted (SigmaPlot 12.0 for Windows. Systat Software Inc., San Jose, CA, USA).
319 Differences of $p<0.05$ were considered statistically significant.

320 **3. Results and discussion**

321 *3.1. Soil characteristics*

322 The majority of CC-UK soils had sandy texture, with a relatively high proportion of silt
323 (Table 2). The only exception was CC-UK-2, although the sand proportion was quite
324 close to that of silt. This texture was similar to that of CC-ES-1 and CC-ES-2, although
325 CC-ES-3 had a higher proportion of silt and higher proportion of clay than of sand. All
326 the OG soils presented a texture of a majority of silt (56.7-59.5%), followed by clay. It
327 is worth considering that silt is the most erodible fraction (Table 2).
328 pH in the UK soils was generally acid, although GS-UK-1 and CC-UK-1 had values
329 between 3.58-5.66 and CC-UK-2 had values close to 7. ES soils had generally higher
330 values, ranging from 5.53-6.98 (Table 2). Some authors have related pH and
331 measurement of aromatics (Weishar *et al.*, 2003). GS-UK-1 has the lowest pH (3.58)
332 and is the one with the highest aromatics content and highest HWC.

333 *3.2. Total extractable carbon by cold and hot extractions*

334 CWC and HWC measurement results are in Table 3. HWC extracted significantly more
335 carbon ($P<0.01$) than CWC, thus proved to be a more exhaustive extraction method.
336 The increases in the values ranged from 161-605%. This was equivalent to higher
337 values by a factor between 3 and 7 respectively. The most significant increase was
338 obtained in GS-UK-1. Gregorich *et al.* (2003) obtained a similar trend in maize-cropped
339 soils of Ottawa, Canada, and found HWC exceeding CWC by a factor of two.

340 Moreover, Landgraf *et al.* (2006) found that HWC had higher carbon concentrations
341 than CWC by a factor that varied from 4 to 6 in surface horizons of forest soils in SE
342 Germany.
343 A correlation analysis was run to assess whether the C extractability differed between
344 the different soils and locations. The correlation resulted to be high ($Rs=0.91$, $p<0.01$).
345 Wang and Wang (2007) found a similar correlation ($r=0.93$, $p=<0.01$) in forest oxisols
346 of Southern China. Klose and Makeschin (2003) also found that HWC and CWC
347 increased with the same proportion in forest soils of NE Germany. On the other hand,
348 Ghani *et al.* (2003) found a positive but poor correlation.
349 HWC and CWC data were compared with SOM (measured by LOI) to assess which of
350 the two extraction methods would imply a higher correlation with the organic matter
351 from the samples (Table 4). HWC implied a higher correlation with SOM measured
352 with LOI ($Rs= 0.70$, $p<0.05$) than CWC ($Rs= 0.55$). Despite the fact that the correlation
353 between CWC and SOM was lower than that of HWC; other authors have found even
354 poorer correlations. For instance, Van Migroet *et al.* (2005) found a r^2 of 0.2 in forest
355 soils in Utah, USA.
356 *3.3. Quality of cold-water extractable carbon using UV-Vis and Fluorescence*
357 Cold-water extractable carbon (CWC) data was compared with UV-Vis and
358 fluorescence spectroscopy, using SUVA-$_{254}$ and FI respectively. The Spearman Rank
359 Correlation was calculated to evaluate the quality and reproducibility of CWC using
360 optical techniques (Table 4).
361 There were high correlation patterns between CWC and SUVA-$_{254}$ ($Rs=0.82$, $p<0.05$)
362 although not between CWC and FI ($Rs=-0.29$), where the slight correlation that could
363 be appreciated was inverse. Chow (2006) obtained a worse correlation between CWC
364 and SUVA-$_{254}$ ($R^2=0.38$), whereas that of Van Migroet *et al.* (2005) was even lower

365 (R^2=0.01). As Weishar *et al.* (2003) point out; some authors have found conflicting
366 conclusions when using SUVA-$_{254}$ to determine the aromaticity of DOC.

367 *3.4. Quality of hot-water extractable carbon using UV-Vis and Fluorescence*

368 HWC data was compared with SUVA-$_{254}$ and FI to assess whether it had higher
369 correlation than CWC. As it can be seen in Table 4, there is a high correlation (Rs=0.88,
370 $p<0.001$) between HWC and SUVA-$_{254}$ whereas that of HWC with FI is not as
371 significant (Rs=-0.53). In both cases, HWC was more significantly correlated to the
372 results obtained with fluorescence and SUVA-$_{254}$ than CWC.

373 *3.5. EEM spectra general characteristics*

374 A set of reference fluorescence spectra has been represented in Figure 1. As it can be
375 seen in Figure 1.a and 1.b, the two soils with the highest % aromatics measured with
376 ^{1}H-NMR as well as high SUVA-$_{254}$ and HWC values (GS-UK-1 and CC-UK-1) showed
377 strong peaks C (10-12 RUs), which indicate UVA-excited humic peaks as described by
378 Coble (1996). They also showed weak peaks A.
379 On the other hand, Figure 1.c. represents the soil with the highest FI, but lowest %
380 aromatic measured with ^{1}H-NMR as well as comparatively low values of SUVA-$_{254}$
381 (CC-ES-1). The EEM spectrum of this soil is very different from the ones represented in
382 Figures 1.a and 1.b, with a strong peak A or UVC-excited (8 RUs), and weak peaks C
383 and T. The presence of the latter indicates biological activity.

384 *3.6. Relationship between optical measures and NMR*

385 The aromaticity of isolated fulvic acid samples was calculated as the ratio of the area of
386 aromatic hydrogen region to the total area of the ^{1}H-NMR spectrum (% aromaticity). An
387 analogous method was used to calculate the ratio of the area of aliphatic hydrogen and
388 carbohydrate hydrogen regions.

As it can be seen in Figure 2, the aromatic hydrogen region is situated in a chemical shift of 6-8ppm by frequency. On the other hand, the carbohydrate hydrogen and aliphatic hydrogen regions are situated in a chemical shift of 3-4.2 ppm and 0.5-3 ppm respectively. For Figure 2, the two soils with the highest % aromatics were chosen for representation purposes (GS-UK-1 and CC-UK-1).

Results of these measurements can be found in Table 3, along with the results from the optical techniques. Correlation analyses were conducted comparing the results from all of them (Table 4).

The largest carbon fraction of the UK soils was carbohydrates. The same trend was found in the CC-ES soils, but not in the OG-ES soils, where the largest carbon fraction was aliphatics.

The proportion of aromatics of the UK soils ranged from 3.91%-5.98%, whereas for the CC-ES and OG-ES soils, it ranged from 2.40-3.47% and 3.90%-4.33% respectively. As these were taken from samples in the first horizon of each soil, the low level of aromaticity can be explained by the fact that plant residues may have accumulated in the surface, while microorganisms did not have enough capacity to decompose them. As Cardoza *et al.* (2004) point out, NMR is a powerful tool for investigating humic substances interactions at the molecular level.

Aromatics have positive correlations for $SUVA\text{-}_{254}$, CWC and HWC but are inversely correlated to FI. Correlations are high for aromatics compared with $SUVA\text{-}_{254}$ and HWC (Figures 3 and 4. Rs of 0.95 and 0.90 respectively; $p<0.001$), but not as good for FI (Figure 5). The correlation between aromatics and CWC, although significant, is not as strong as the one with HWC ($Rs=0.74$, $p<0.05$). Weishaar *et al.* (2003) found that $SUVA\text{-}_{254}$ is a good predictor of the humic fraction (higher $SUVA\text{-}_{254}$ indicates higher humic acid content and molecular weight in DOM solutions) and the general chemical

properties of HWC, although it does not predict the reactivity of HWC from different types of source materials nor information of the individual molecules of samples unless these are humic substances (Chin *et al.*, 1997; Weishar *et al.*, 2003). They obtained a good correlation between SUVA-$_{254}$ and aromaticity determined by ^{13}C-NMR (R=0.97), which was slightly higher than ours. Yeh *et al.* (2014) also found a high correlation between SUVA-$_{254}$ measurements and the aromatic fraction of samples of organic matter of river sediments of Taiwan. Jamieson *et al.* (2014) found that higher SUVA-$_{254}$ implied higher aromaticity in biochar from sugar maple, thus demonstrating that this effect is currently studied when characterising organic material with a high recalcitrance and that is able to retain carbon for thousands of years, due to its stability (Lehmann, 2007).

According to McKnight (2001) and other authors (Rodríguez *et al.*, 2014; Wei *et al.*, 2014), there is a correlation between the FI values and those from methods that indicate the aromaticity of humic substances such as the ratio of the aromatic carbon region area in the total NMR spectra and SUVA-$_{254}$. This pattern was observed in our soils, with negative correlations between FI and these 2 parameters (Rs=-0.63 and Rs=-0.52 respectively), although only the trend with the aromatic carbon region area in the total NMR spectra was statistically significant ($p<0.05$).

Kim *et al.* (2006) and Rodríguez *et al.* (2014) indicated that differences in the FI higher than 0.1 would imply significant differences in the aromaticity of the samples. If this assumption is applied to our case, CC-UK-2 has differences in its aromaticity with GS-UK-1 but not with CC-UK-1. CC-ES-2 has a different aromaticity than CC-ES-1 and CC-ES-3. In the same way, OG-ES-1 has differences with OG-ES-2 and OG-ES-3. When all the types are compared, there are combinations of differences in the aromaticity but a single pattern cannot be obtained.

Given the results above, the regressions between the aromatic carbon region measured with ^{1}H-NMR and SUVA-$_{254}$ and HWC are powerful. The equations we obtained can explain very significantly the relationship between these variables ($p<0.001$ for the results of ^{1}H-NMR predicted with SUVA-$_{254}$ and $p<0.01$ if they are predicted using HWC). These equations (Figure 3 and Figure 4) could therefore be used to predict the aromaticity of water-extractable carbon.

The regression of the aromaticity measured with ^{1}H-NMR with FI, although significant ($p<0.05$), did not have the same level of confidence as the one with HWC and SUVA-$_{254}$. As it can be seen in Figure 5, an outlier was identified in the Regression analysis (FI value of CC-ES-3). Figure 5 showed how the regression model improved when the outlier was removed from the analysis (Rs=-0.98). The linear regression equation that resulted after the elimination of the outlier is able to explain the ^{1}H-NMR values using FI with a very high significance ($p<0.001$).

However, according to McKnight (2001), the FI would not be sufficient to estimate aromaticity and other techniques are necessary (e.g. SUVA-$_{254}$), as geological processes can alter aromaticity without changing the FI.

SUVA-$_{254}$ did not demonstrate a particularly strong correlation with FI (Rs=-0.52, Table 4). The correlation was negative (higher SUVA-$_{254}$ led to lower FI) which coincides with authors like Kothawala *et al.* (2012) with minerals soils collected across Canada although their trend is less clear than in the study of other authors such as Williams *et al.* (2010), where the slope is more pronounced. The correlation of Williams *et al.* (2010) was similar to ours (r=-0.57 in their study and Rs=-0.52 in ours), although the statistical analysis in their case was the Pearson correlation, as their data followed a normal distribution. Hassouna *et al.* (2012) found a negative correlation as well, with slightly higher correlation in Mediterranean calcareous soils (r=-0.69). As in the case

464 described in Figure 5, if the same outlier was removed, the correlation improved
465 significantly (Rs=-0.98) and the equation of the linear regression explained with a very
466 high degree of significance ($p<0.001$) the relation between SUVA-$_{254}$ and FI.
467 Analogously, the same outlier could be removed in the analysis of HWC vs. FI,
468 obtaining a significant correlation between them (Rs=-0.88). A linear regression model
469 explains significantly ($p<0.01$) the relationship between HWC and FI and therefore
470 similar conclusions can be stated as in the case of the correlation of SUVA-$_{254}$ to FI.
471 Given the strong intrinsic relationships between the results of SUVA-$_{254}$, HWC and FI;
472 and how they are able to predict the proportion of aromaticity measured with ^{1}H-NMR,
473 we can conclude that UV-Vis absorbance and fluorescence spectroscopy can be used to
474 characterise the aromaticity of carbon and may be a plausible substitute for ^{1}H-NMR,
475 given that they are more cost-effective. Other authors such as Zornoza *et al.* (2008,
476 2015) demonstrated the effectiveness of spectroscopy (near infrared reflectance
477 spectroscopy in their study) as an accurate, cost- and time-effective method for
478 predicting and/or estimating soil biogeochemical properties and other soil parameters.
479 Correlations between aliphatics and the data from the optical measures and CWC/HWC
480 were poor (Table 4), although slight trends can be observed for SUVA-$_{254}$, FI and CWC.
481 In these cases, the correlation was positive for FI and CWC but negative for SUVA-$_{254}$.
482 Correlations were also poor between carbohydrates and the data from the optical
483 measures and HWC/CWC, with only one trend detected. This was the inverse
484 correlation (Rs=-0.35) between HWC and carbohydrates.
485 BIX and HIX did not correlate well with any of the other parameters (Table 4), with
486 very few trends detected. Of the trends detected, HIX was positively correlated with FI
487 (Rs=0.52). It is worth noting that BIX was positively correlated with carbohydrates but
488 inversely correlated to aliphatics (Rs=0.31 and -0.31 respectively), whereas the trend

was the contrary for HIX (Rs=-0.28 and 0.22). A similar trend than ours was also found by Kalbitz *et al.* (2003) between HIX and carbohydrates in forest soils, arable soils and a fen area; although theirs was stronger (r^2=0.81). Williams *et al.* (2010) studied the correlation between HIX and SUVA-$_{254}$ in watershed from mixed land use in Canada. They obtained a positive correlation (r=0.74), whereas our correlation was negative. On the other hand, Yeh *et al.* (2014) did not find a clear trend between SUVA-$_{254}$ and HIX. The significance of our comparison between SUVA-$_{254}$ and HIX values is extremely low and therefore robust conclusions on this issue cannot be stated. Bu *et al.* (2010) compared HWC and HIX in various soils with different vegetation types in Wuyi Mountain (SE China), obtaining a poor negative correlation, just like ours. However, Kalbitz *et al.* (2003) found a strong correlation between the aromatic fraction of DOM and HIX (r^2=0.80).

Yeh *et al.* (2014) calculated BIX as well, obtaining that higher SUVA-$_{254}$ values implied lower BIX values, which coincides with our comparison in this case, although its significance is not relevant enough to establish a robust conclusion.

Plotting the results from Birdwell and Engel (2010) for FI, HIX, and BIX did not led to significant correlations, which is similar to our results and, along with the authors commented above, demonstrate that a number of studies have found difficulties when correlating HIX and BIX to other parameters.

4. Conclusions

Different techniques to measure the quantity and quality of SOM were tested in cambisols from very different climatic locations and under different cropping systems. HWC extracted a higher amount and carbon than CWC, and correlated better with the %SOM (LOI) than CWC.

513 SUVA-$_{254}$ and HWC correlated significantly with the proportion of aromatics measured
514 with ^{1}H-NMR, demonstrating their complementary nature. Linear regression models
515 fitted to the data were able to explain the relationship between the specific fraction of
516 aromatics measured with ^{1}H-NMR and SUVA-$_{254}$ and HWC, and therefore allow the
517 aromatic fraction to be estimated at lower cost using SUVA-$_{254}$ and HWC.
518 A linear regression model was also able to explain the relationship between ^{1}H-NMR
519 measurements and FI, after an outlier was removed.
520 BIX and HIX were not useful indexes for our purposes, as they did not correlate well to
521 the carbon fractions measured with ^{1}H-NMR or with FI, SUVA-$_{254}$, or HWC.
522 The EEM spectra showed a difference in the peak types that appeared in the CC-UK
523 soils and in the CC-ES soil. The fact that figures 1.a and 1.b showed a stronger peak C
524 and figure 1.c. showed a stronger peak A indicated subtle qualitative differences in their
525 humic fractions. This could have been caused by the different soil management (more
526 intense agricultural activity in CC-ES-1). More work will be needed to confirm this.
527 In view of our results, optical measures have a potential, in combination, to predict the
528 aromatic fraction of SOM without the need of expensive and time consuming
529 techniques like ^{1}H-NMR; which could be very useful when the equipment is not
530 available or in instances when a high number of samples need to be analysed
531 simultaneously.

532 **5. Acknowledgments**

533 We thank J.M. Calero and M. Bell for their contribution to improve this paper.

534 **6. References**

535 Aemet, 2014. Historic series of climatic data 1971-2000. Agencia Estatal de
536 Meteorología (Spanish State Meteorological Agency). Available at:

537 http://www.aemet.es/es/serviciosclimaticos/datosclimatologicos/valoresclimatologicos?l
538 =5402&k=and [Accessed: 08th August 2014].

539 Agren, G.I., Bosatta, E., 2002. Reconciling differences in predictions of temperature
540 response of soil organic matter. Soil Biol. Biochem. 34, 129–132.

541 Alburquerque, J.A., Calero, J.M., Barrón, V., Torrent, J., Campillo, M.C., Gallardo, A.,
542 Villar, R., 2014. Effects of biochars produced from different feedstocks on soil
543 properties and sunflower growth. J. Plant Nutr. Soil Sci. 177, 16–25.

544 Batjes, N. H., 2014. Projected changes in soil organic carbon stocks upon adoption of
545 recommended soil and water conservation practices in the upper Tana River catchment,
546 Kenya. Land Degrad. Dev. 25, 278–287.

547 Bell, M., Lawrence, D., 2009. Soil carbon sequestration - Myths and mysteries.
548 Department of Primary Industries and Fisheries, Queensland, Australia. Available at:
549 http://www.futurebeef.com.au/wp-content/uploads/2011/09/Soil-Carbon-Sequestration-
550 Report.pdf [Accessed: 8th November 2014].

551 Birdwell, J.E., Engel, A.S., 2010. Characterization of dissolved organic matter in cave
552 and spring waters using UV–Vis absorbance and fluorescence spectroscopy. Org.
553 Geochem. 41, 270–280.

554 Brevik, E.C., 2009. Soil Health and Productivity, in: Verheye, W. (Ed.), Soils, Plant
555 Growth and Crop Production; Encyclopaedia of Life Support Systems (EOLSS),
556 Developed under the Auspices of the UNESCO. EOLSS Publishers, Oxford, UK.
557 Available at: http://www.eolss.net [Accessed 10th November 2014].

558 Brevik, E. C., 2012. Soils and climate change: Gas fluxes and soil processes. Soil
559 Horiz. 53, 12–23.

Brevik, E. C., Cerdà, A., Mataix-Solera, J., Pereg, L., Quinton, J. N., Six, J., Van Oost K., 2015. The interdisciplinary nature of SOIL. Soil 1, 117–129.

Bu, X., Wang, L., Ma, W., Yu, X., McDowell, W.H., Ruan, H., 2010. Spectroscopic characterization of hot-water extractable organic matter from soils under four different vegetation types along an elevation gradient in the Wuyi Mountains. Geoderma 159, 139–146.

Cardoza, L.A., Korir, A.K., Otto, W.H., Wurrey, C.J., C.K. Larive, C.K., 2004. Applications of NMR spectroscopy in environmental science. Prog. Nucl. Mag. Res. Sp. 45, 209–238.

Cerdà, A., 1998. The influence of aspect and vegetation on seasonal changes in erosion under rainfall simulation on a clay soil in Spain. Can. J. Soil. Sci. 78, 321–330.

Cerdà, A., 2000. Aggregate stability against water forces under different climates on agriculture land and scrubland in southern Bolivia. Soil Till. Res. 57, 159–166.

Chen, J., LeBoef, E.J., Dai, S., Gu, B., 2003. Fluorescence spectroscopic studies of natural organic matter fractions. Chemosphere 50, 639–647.

Chen, Y., Day, S.D., Wick, A.F., Strahm, B.D., Wiseman, P.E., Daniels, W.L., 2013. Changes in soil carbon pools and microbial biomass from urban land development and subsequent post-development soil rehabilitation. Soil Biol. Biochem. 66, 38–44.

Chin Y.P., Aiken, G.R., Danielsen, K.M., 1997. Binding of pyrene to aquatic and commercial humic substances: the role of molecular weight and aromaticity, Environ. Sci. Technol. 31, 1630–1635.

581 Chow, A.T., 2006. Disinfection byproduct reactivity of aquatic humic substances
582 derived from soils. Water Res. 40, 1426–1430.

583 Coble, P., 1996. Characterization of marine and terrestrial DOM in seawater using
584 excitation–emission matrix spectroscopy. Mar. Chem. 51, 325–346.

585 Cory, R.M., Miller, M.P., McKnight, D.M., Guerard, J.J., Miller, P.L., 2010. Effect of
586 instrument-specific response on the analysis of fulvic acid fluorescence spectra. Limnol.
587 Oceanogr. Meth. 8, 67–78.

588 Dungait, J., Hopkins, D., Gregory, A.S., Whitmore, A.P., 2012. Soil organic matter
589 turnover is governed by accessibility not recalcitrance. Glob. Chang. Biol. 18, 1781–
590 1796.

591 Fernández-Romero, M.L., Lozano-García, B., Parras-Alcántara, L., Collins, C.D.,
592 Clark, J.M., 2014. Effects of land management on different forms of soil carbon in olive
593 groves in Mediterranean areas. Land Degrad. Dev. (In press) DOI: 10.1002/ldr.2327.

594 Fischlin, A., Midgley, G.F., Price, J., Leemans, R., Gopal, B., Turley, C., Rounsevell,
595 M., Dube, P. Tarazona, J., Velichko, A., 2007. Ecosystems, their properties, goods, and
596 services, in: M. L., Canziani, O. F., Palutikof, J. P., van der Linden, P. J. & Hanson, C.
597 E. (Eds.), Climate Change: Impacts, Adaptation and Vulnerability. Contribution of
598 Working Group II to the Fourth Assessment Report of the Intergovernmental Panel on
599 Climate Change. Cambridge University Press, Cambridge, UK, pp. 211–272. ISBN:
600 978 0521 88010-7.

601 Fuentes, M., González-Gaitano, G., García-Mina, J.M., 2006. The usefulness of UV–
602 visible and fluorescence spectroscopies to study the chemical nature of humic
603 substances from soils and composts. Org. Geochem. 37, 1949–1959.

García-Gil, J.L., Plaza, C., Soler-Rovira, P., Polo, A., 2000. Long-term effects of municipal solid waste compost application on soil enzyme activities and microbial biomass. Soil Biol. Biochem. 32, 1907–1913.

Ghani, A., Dexter, M. & Perrott, K.W., 2003. Hot-water extractable carbon in soils: a sensitive measurement for determining impacts of fertilization, grazing and cultivation. Soil Biol. Biochem. 35, 1231–1243.

González-Pérez, M., Milori, D.M.B.P., Colnago, L.A., Martin-Neto, L., Melo, W.J., 2007. A laser-induced fluorescence spectroscopic study of organic matter in a Brazilian Oxisol under different tillage systems. Geoderma 138, 20–24.

Gregorich, E.G., Beare, M.H., Stoklas, U., St-Georges, P., 2003. Biodegradability of soluble organic matter in maize-cropped soils. Geoderma 113, 237– 252.

Hassouna, M., Théraulaz, F., Massiani, C., 2012. Production and elimination of water extractable organic matter in a calcareous soil as assessed by UV/Vis absorption and fluorescence spectroscopy of its fractions isolated on XAD-8/4 resins. Geoderma 189–190, 404–414.

Helal, A.A., Murad, G.A., Helal, A.A., 2011. Characterization of different humic materials by various analytical techniques. Arab. J. Chem. 4, 51–54.

Hemminga, M.A., Buurman, P., 1997. Editorial: NMR in soil science. Geoderma 80, 221–224.

Henderson, R.K., Baker, A., Murphy, K.R., Hamblya, A., Stuetz, R.M., Khan, S.J., 2009. Fluorescence as a potential monitoring tool for recycled water systems: a review. Water Res. 43, 863–881.

626 Hieri, O., Lotter, A.F., Lemke, G., 2001. Loss on Ignition as a method for estimating
627 organic and carbonate content in sediments: Reproducibility and comparability of
628 results. J. Paleolimnol. 25, 101–110.

629 Huguet, A., Vacher, L., Relexans, S., Saubusse, S., Froidefond, J.M., Parlanti, E., 2009.
630 Properties of fluorescent dissolved organic matter in the Gironde Estuary. Org.
631 Geochem. 40, 706–719.

632 Hur, J., Kim, G., 2009. Comparison of the heterogeneity within bulk sediment humic
633 substances from a stream and reservoir via selected operational descriptors.
634 Chemosphere 75, 483–490.

635 Hur, J., Park, S.W., Kim, M.C., Kim, H.S., 2013. Enhanced binding of hydrophobic
636 organic contaminants by microwave-assisted humification of soil organic matter.
637 Chemosphere 93, 2704–2710.

638 Jamieson, T., Sager, E., Guéguen, C., 2014. Characterization of biochar-derived
639 dissolved organic matter using UV–visible absorption and excitation–emission
640 fluorescence spectroscopies. Chemosphere 103, 197–204.

641 Kalbitz, K., Schmerwitz, J., Schwesig, D., Matzner, E., 2003. Biodegradation of soil
642 derived dissolved organic matter as related to its properties. Geoderma 113, 273–291.

643 Kim, H.C., Yu, M.J., Han, I., 2006, Multi-method study of the characteristic chemical
644 nature of aquatic humic substances isolated from the Han River, Korea. Appl.
645 Geochem. 21, 1226–39.

646 Kleber, M., 2010. What is recalcitrant soil organic matter? Environ. Chem. 7, 320–332.

647 Kleber, M., Johnson, M. G., 2010. Advances in understanding the molecular structure
648 of soil organic matter: implications for interactions in the environment. Adv. Agron.
649 106, 77–142.

650 Klose, S., Makeschin, F., 2003. Effects of past fly ash deposition on the forest floor
651 humus chemistry of pine stands in Northeastern Germany. For. Ecol. Manage. 183,
652 113–126.

653 Koarashi, J., Iida, T., Asano, T., 2005. Radiocarbon and stable carbon isotope
654 compositions of chemically fractionated soil organic matter in a temperate-zone forest.
655 J. Environ. Radioact. 79, 137–56.

656 Korak, J.A., Dotson, A.D., Summers, R.S., Rosario-Ortiz, F.L., 2014. Critical analysis
657 of commonly used fluorescence metrics to characterize dissolved organic matter. Water
658 Res. 49, 327–338.

659 Kothawala, D.N., Roehm, C., Blodau, C., Moore, T.R., 2012. Selective adsorption of
660 dissolved organic matter to mineral soils. Geoderma 189–190, 334–342.

661 Kwiatkowska, J., Provenzano, M., Senesi, N., 2008. Long term effects of a brown coal-
662 based amendment on the properties of soil humic acids. Geoderma 148, 200–205.

663 Lal, R., 2004. Soil carbon sequestration to mitigate climate change. Geoderma 123, 1–
664 22.

665 Landgraf, D., Leinweber, P., Makeschin, F., 2006. Cold and hot water–extractable
666 organic matter as indicators of litter decomposition in forest soils. J. Plant Nutr. Soil
667 Sci. 169, 76–82.

668 Lehmann, J., 2007. Bio-energy in the black. Front. Ecol. Environ. 5, 381–387.

669 Leifeld J., Kogel-Knabner I., 2005. Soil organic matter fractions as early indicators for
670 carbon stock changes under different land-use? Geoderma 124, 143–155.

671 Li, L.J., Han, X.Z., You, M.Y., Yuan, Y.R., Ding, X.L., Qiao, Y.F., 2013. Carbon and
672 nitrogen mineralization patterns of two contrasting crop residues in a Mollisol: Effects
673 of residue type and placement in soils. Eur. J. Soil Biol. 54, 1–6.

674 Luke, J. K., Brevik, E. C., Wood, G. L., 2009. Evaluation of loss on ignition method for
675 determining organic matter of South Georgia soils. Soil Surv. Hor. 50, 83–85.

676 Martins, T., Saab, S.C., Milori, D.M.B.P., Brinatti, A.M., Rosa, J.A., Cassaro, F.A.M.,
677 Pires, L.F., 2011. Soil organic matter humification under different tillage managements
678 evaluated by Laser Induced Fluorescence (LIF) and C/N ratio. Soil Till. Res. 111, 231–
679 235.

680 McKnight, D.M., Boyer, E.W., Westerhoff, P., Doran, P., Kulbe, T., Andersen, D.,
681 2001. Spectrofluorimetric characterization of dissolved organic matter for indication of
682 precursor organic material and aromaticity. Limnol. Oceanogr. 46, 38–48.

683 MIT, 2008. Things to know before you begin operating an NMR. Department of
684 Chemistry. Instrumentation Facility. Massachusetts Institute of Technology. Available
685 at: http://web.mit.edu/speclab/www/PDF/DCIF-Things-2-know-NMR-m08.pdf
686 [Accessed 10th June 2014].

687 Novara, A., Gristina, L., Saladino, S.S., Santoro, A., Cerdà, A., 2011. Soil erosion
688 assessment on tillage and alternative soil managements in a Sicilian vineyard. Soil Till.
689 Res. 117, 140–147.

690 Ohno, T., 2002. Fluorescence inner-filtering correction for determining the humification
691 index of dissolved organic matter. Environ. Sci. Technol. 36, 742–746.

692 Olk, D.C., Gregorich, E.G., 2006. Overview of the symposium proceedings,
693 "Meaningful pools in determining soil carbon and nitrogen dynamics". Soil Sci. Soc.
694 Am. J. 70, 967–974.

695 Paradelo R., Barral, M.T., 2013. Influence of organic matter and texture on the
696 compactability of Technosols. Catena 110, 95–99.

697 REDIAM, 2007. Historical series of meteorological data 1953-2007. Red de
698 Información Ambiental de Andalucía (Andalusian Network of Environmental
699 Information). Seville, Spain.

700 Rodríguez, F.J., Schlenger, P., García-Valverde, M., 2014. A comprehensive structural
701 evaluation of humic substances using several fluorescence techniques before and after
702 ozonation. Part I: Structural characterization of humic substances. Sci. Total
703 Environ. 476–477, 718–730.

704 Rose, M.T., Patti, A.F., Little, K.R., Brown, A.L., Jackson, W.R., Cavagnaro, T.R.,
705 2014. Chapter Two – A Meta-Analysis and Review of Plant-Growth Response to
706 Humic Substances: Practical Implications for Agriculture. Adv. Agron. 124, 37–89.

707 Saab, S.C., Martin-Neto, L., 2007. Condensed aromatic rings and E_4/E_6 ratio: humic
708 acids in gleysoils studied by NMR cp/mas ^{13}C, and dipolar dephasing. Quim. Nova 30,
709 260–263.

710 Saha, D., Kukal, S.S., Bawa, S.S., 2014. Soil organic carbon stock and fractions in
711 relation to land use and soil depth in the degraded Shiwaliks Hills of lower Himalayas.
712 Land Degrad. Dev. 25, 407–416.

713 Salehi, M.H., Beni, O.H., Harchegani, H.G., Borujeni, I.E., 2011. Refining Soil Organic
714 Matter Determination by Loss-on-Ignition. Pedosphere 21, 473–482.

715 Schmidt, M.W.I., Torn, M.S., Abiven, S., Dittmar, T., Guggenberger, G., Janssens, I.A.,
716 Kleber, M., Kögel-Knabner, I., Lehmann, J., Manning, D.A.C., Nannipieri, P., Rasse,
717 D.P., Weiner, S., Trumbore, S.E., 2011. Persistence of soil organic matter as an
718 ecosystem property. Nature 478, 49–56.

719 Schrumpf, M., Kaiser, K., 2015. Large differences in estimates of soil organic carbon
720 turnover in density fractions by using single and repeated radiocarbon inventories.
721 Geoderma 239–240, 168–178.

722 Sellami, F., Hachicha, S., Chtourou, M., Medhioub, K., Ammar, E., 2008. Maturity
723 assessment of composted olive mill wastes using UV spectra and humification
724 parameters. Bioresource Technol. 99, 6900–6907.

725 Senesi, N., D'Orazio, V., 2005. Fluorescence spectroscopy, in: Hillel, D. (Ed.),
726 Encyclopedia of Soils in the Environment. Academic Press, Elsevier Science, pp. 35–
727 52. ISBN: 978-0-12-348530-4.

728 Skoog, D.A., Holler, F.J., Crouch, S.R., 2007. Principles of Instrumental Analysis.
729 Brooks/Cole Publishing. Belmont, CA, USA. ISBN: 978-0-495-01201-6.

730 Sparks, D.L., 2003. Environmental Soil Chemistry, second ed. Academic Press,
731 Elsevier Science, San Diego, USA. ISBN 0-12-656446-9.

732 Srinivasarao, CH., Venkateswarlu, B., Lal, R., Singh, A. K., Kundu, S., Vittal, K. P.
733 R., Patel, J. J., Patel, M. M., 2014. Long-term manuring and fertilizer effects on
734 depletion of soil organic carbon stocks under pearl millet- cluster bean-castor rotation in
735 Western India. Land Degrad. Dev. 25, 173–183.

736 Sun, W.L., Ni, J.R., Xu, N., Sun, L.Y., 2007. Fluorescence of sediment humic substance
737 and its effect on the sorption of selected endocrine disruptors. Chemosphere 66, 700–
738 707.

739 Sutton, R., Sposito, G., 2005. Molecular Structure in Soil Humic Substances: The New
740 View. Environ. Sci. Technol. 39, 9009–9015.

741 Tang, Z., Yu, G., Liu, D., Xu, D., Shen, Q., 2011. Different analysis techniques for
742 fluorescence excitation–emission matrix spectroscopy to assess compost maturity.
743 Chemosphere 82, 1202–1208.

744 Trumbore, S., 1997. Potential responses of soil organic carbon to global
745 environmental change. P. Natl. Acad. Sci. USA 94, 8284-8291.

746 Trumbore, S., 2000. Age of soil organic matter and soil respiration: Radiocarbon
747 constraints on belowground C dynamics. Ecol. Appl. 10, 399–411.

748 Uchida, Y., Nishimurab, S., Akiyama, H., 2012. The relationship of water-soluble
749 carbon and hot-water-soluble carbon with soil respiration in agricultural fields. Agr.
750 Ecosyst. Environ. 156, 116–122.

751 UK Met Office, 2014. Reading 1981–2010 climatic averages. Available at:
752 http://www.metoffice.gov.uk/public/weather/climate/gcpk9wzwz [Accessed 8th
753 November 2014].

754 Van Miegroet, H., Boettinger, J.L., Baker, M.A., Nielsen, J., Evans, D., Stum, A., 2005.
755 Soil carbon distribution and quality in a montane rangeland-forest mosaic in northern
756 Utah. Forest Ecol. Manag. 220, 284–299.

757 Wang, Q.K., Wang S.L., 2007. Soil organic matter under different forest types in
758 Southern China. Geoderma 142, 349–356.

759 Wei, Z., Zhao, X., Zhu, C., Xi, B., Zhao, Y., Yu, X., 2014. Assessment of humification
760 degree of dissolved organic matter from different composts using fluorescence
761 spectroscopy technology. Chemosphere 95, 261–267.

762 Weishaar, J.L., Aiken, G.R., Bergamaschi, B.A., Fram, M.S., Fujii, R., Mopper, K.,
763 2003. Evaluation of Specific Ultraviolet Absorbance as an Indicator of the Chemical
764 Composition and Reactivity of Dissolved Organic Carbon. Environ. Sci. Technol. 37,
765 4702–4708.

766 Williams, C.J., Yamashita, Y., Wilson, H.F., Jaffé, R., Xenopoulos, M.A., 2010.
767 Unravelling the role of land use and microbial activity in shaping dissolved organic
768 matter characteristics in stream ecosystems. Limnol. Oceanogr. 55, 1159–1171.

769 Xue, S., Li, P., Liu, G., Li, Z., Zhang, C., 2013. Changes in Soil Hot-Water Extractable
770 C, N and P Fractions During Vegetative Restoration in Zhifanggou Watershed on the
771 Loess Plateau. J. Integr. Agr. 12, 2250–2259.

772 Yeh, Y.L., Yeh, K.J., Hsu, L.F., Yu, W.C, Lee, M.H., Chen, T.C., 2014. Use of
773 fluorescence quenching method to measure sorption constants of phenolic
774 xenoestrogens onto humic fractions from sediment. J. Hazard. Mater. 277, 27–33.

775 Zhao, G., Mu, X., Wen, Z., Wang, F., Gao, P., 2013. Soil erosion, conservation, and
776 eco-environment changes in the Loess Plateau of China. Land Degrad. Dev. 24, 499–
777 510.

778 Zhao, M., Zhou, J., Kalbitz, K., 2008. Carbon mineralization and properties of water-
779 extractable organic carbon in soils of the south Loess Plateau in China. Eur. J. Soil
780 Biol. 44, 158–165.

781 Zornoza, R., Guerrero, C., Mataix-Solera, J., Scow, K.M., Arcenegui, V., Mataix-
782 Beneyto, J., 2008. Near infrared spectroscopy for determination of various physical,
783 chemical and biochemical properties in Mediterranean soils. Soil Biol. Biochem. 40,
784 1923–1930.

785 Zornoza, R., Acosta, J. A., Bastida, F., Domínguez, S. G., Toledo, D. M., Faz A., 2015.
786 Identification of sensitive indicators to assess the interrelationship between soil quality,
787 management practices and human health. Soil 1, 173–185.

788 Zsolnay, A., Baigar, E., Jimenez, M., Steinweg, B., Saccomandi, F., 1999.
789 Differentiating with fluorescence spectroscopy the sources of dissolved organic matter
790 in soils subjected to drying. Chemosphere 38, 45–50.

791 **Figure captions**

792 **Figure 1.** EEM plots from: a) GS-UK-1; b) CC-UK-1; c) CC-ES-1

793 **Figure 2**. Liquid state ^{1}H-NMR spectra from: a) GS-UK-1; b) CC-UK-1

794 **Figure 3.** Correlation between % Aromatics and SUVA-$_{254}$. A linear regression model
795 was calculated to explain the relationship between the variables. The equation was y [%
796 Aromatics] = 0.250 + 0.950 x [SUVA-$_{254}$], $p<0.001$

797 **Figure 4.** Correlation between % Aromatics and HWC. A linear regression model was
798 calculated to explain the relationship between the variables. The equation was y [%
799 Aromatics] = 0.500 + 0.900 x [HWC], ($p<0.01$)

Figure 5. Correlation between % Aromatics and FI. A linear regression model was calculated to explain the relationship between the variables. The equation was y [% Aromatics] = 8.17 – 0.633 x [FI], ($p<0.05$), represented by the solid line. The removal of an outlier resulted in the equation: y [% Aromatics] = 8.89 – 0.976 x [FI], which explained the relationship with a higher significance ($p<0.001$). The second equation is represented by the dashed line