

In vitro study on the cell adhesion ability of immobilized lactobacilli on natural supports

Article

Accepted Version

Sidira, M., Kourkoutas, Y., Kanellaki, M. and Charalampopoulos, D. ORCID: <https://orcid.org/0000-0003-1269-8402> (2015) In vitro study on the cell adhesion ability of immobilized lactobacilli on natural supports. Food Research International, 76 (3). pp. 532-539. ISSN 0963-9969 doi: 10.1016/j.foodres.2015.07.036 Available at <https://centaur.reading.ac.uk/41769/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

Published version at: <http://dx.doi.org/10.1016/j.foodres.2015.07.036>

To link to this article DOI: <http://dx.doi.org/10.1016/j.foodres.2015.07.036>

Publisher: Elsevier

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online

***In vitro* study on the cell adhesion ability of immobilized lactobacilli
on natural supports**

**Marianthi Sidira^{a,b}, Yiannis Kourkoutas^b, Maria Kanellaki^a, and Dimitris
Charalampopoulos^{c*}**

a. Food Biotechnology Group, Section of Analytical Environmental and Applied
Chemistry, Department of Chemistry, University of Patras, GR-26500 Patras, Greece.

b. Applied Microbiology and Molecular Biotechnology Research Group, Department
of Molecular Biology & Genetics, Democritus University of Thrace, Alexandroupolis,
68100, Greece.

c. Department of Food and Nutritional Sciences, The University of Reading, PO Box
226, Whiteknights, Reading, RG6 6AP, UK

Corresponding author: Charalampopoulos Dimitris

Mailing address: Department of Food and Nutritional Sciences, The University of
Reading, PO Box 226, Whiteknights, Reading, RG6 6AP, UK

E-mail: d.charalampopoulos@reading.ac.uk

Abstract

The aim of the present study was to investigate the effect of probiotic immobilization onto wheat grains, both wet and freeze dried, on the adhesion properties of the probiotic cells and make comparisons with wet and freeze dried free cells. *Lactobacillus casei* ATCC 393 and *Lactobacillus plantarum* NCIMB 8826 were used as model probiotic strains. The results showed satisfactory adhesion ability of free cells to a monolayer of Caco-2 cells (>1000 CFU/100 Caco-2 cells for wet cells). Cell immobilization resulted in a significant decrease in adhesion, for both wet and freeze dried formulations, most likely because immobilized cells did not have direct access to the Caco-2 cells, but it still remained in adequate levels (>100 CFU/100 Caco-2 cells for wet cells). No clear correlation could be observed between cell adhesion and the hydrophobicity of the bacterial cells, measured by the hexadecane adhesion assay. Most notably, immobilization enhanced the monolayer integrity of Caco-2 cells, demonstrated by a more than 2-fold increase in transepithelial electrical resistance (TEER) compared to free cells. SEM micrographs ascertained the adhesion of both immobilized and free cells to the brush border microvilli. Finally, the impact of the food matrix on the adhesion properties of probiotic bacteria and on the design of novel functional products is discussed.

Keywords: Probiotics, Lactic acid bacteria, Adhesion, Caco-2 cells

1. Introduction

Nowadays, there is a growing interest in developing foods containing probiotic microorganisms, such as bifidobacteria and lactic acid bacteria (LAB). Such functional cultures may offer organoleptic, technological and nutritional advantages, but more importantly confer a health benefit to the host. Indeed, administration of probiotics has been linked to the prevention, and in some cases reduction or treatment, of various diseases, including viral or bacterial diarrhea, gastroenteritis, irritable bowel syndrome, inflammatory bowel disease (Crohn's disease and ulcerative colitis), depressed immune function, lactose intolerance, infant allergies, *Helicobacter pylori* infections, antibiotic-associated diarrhea in children and others (Deshpandea, Rao, & Patolea, 2011; Hempel et al., 2012; Ritchie, & Romanuk, 2012).

In order to deliver the health benefits, probiotic products need to contain an adequate amount of live cells (at least 10^6 - 10^7 CFU/g of product) (Boylston, Vinderola, Ghoddusi, & Reinheimer, 2004), the cells should be able to survive the acidic conditions of the upper gastrointestinal (GI) tract, adhere to mucosal-epithelial surfaces and colonize the colon, exhibit antimicrobial activity against pathogenic bacteria and bile salt hydrolase activity (Boylston et al., 2004; Kechagia et al., 2013; Nagpal et al., 2012). Adhesion of probiotics to the intestinal mucosa is an important prerequisite for transient colonization within the GI, and it is an area that attracts considerable amount of research. To evaluate the adhesion abilities, *in vitro* methods have been applied using mainly the Caco-2 cell line (Tuomola & Salminen, 1998; Hilgendorf et al., 2000). *Lactobacillus casei*, *L. plantarum* and *L. rhamnosus* are commonly used probiotic bacteria and have been shown to exert medium to strong binding towards Caco-2 cells (Segers & Lebeer, 2014; Douillard et al., 2013; Elo, Saxelin, & Salminen, 1991; Duany, Rajput, Batish, & Grover, 2011).

A variety of methods have been applied to determine the adhesive properties of bacterial cells. The microbial adhesion to hydrocarbons (MATH) method measures various forces contributing to the adhesion of bacterial cells to hydrophobic solvents, including van der Waals, electrostatic and short-range interactions, as adhesion to hydrocarbons reflects a number of physico-chemical interactions involved in adhesion and not exclusively hydrophobicity (van der Mei, van der Belt-Gritter, Pouwels, Martinez, & Busscher, 2003; Oliveira, Azeredo, Teixeira, & Fonseca, 2001). It must be noted that there is very little information on how the formulation, for example the food matrix, affects the physicochemical properties of the cells, and hence their hydrophobicity and adhesion ability.

The transepithelial electrical resistance (TEER) assay determines the attainment of permeability *in vitro* and also includes the measurement of electrical physical resistance (Klingberg, Pedersen, Cencic, & Budde, 2005). Using the TEER assay as a measure of the integrity of the tight junctions between intestinal epithelial cells, studies have shown that some bacteria can enhance the intestinal barrier function (Anderson, Cookson, McNabb, Kelly, & Roy, 2010b). Probiotic bacteria, such as *L. plantarum* WCFS1 or *L. plantarum* MB452, *Lactobacillus rhamnosus* GG, *Bifidobacterium infantis*, *Lactobacillus casei* subsp. *rhamnosus* Lcr35, have been used to enhance the intestinal barrier, which is compromised in a number of intestinal disorders, such as inflammatory bowel diseases, irritable bowel syndrome and some types of food-borne infections (Karczewski et al., 2010; Anderson et al., 2010a; Fang et al., 2010).

The incorporation of probiotic microorganisms within a food matrix presents significant challenges, not only because of the interactions of the cells with a variety of chemical components, but also because of the unfavourable conditions often

employed during food processing and storage which might lead to high losses in viability. This is in particular evident when probiotics are incorporated into complex food matrices, such as cheese and meat products (Sidira, Karapetsas, Galanis, Kanellaki, & Kourkoutas, 2014a). To overcome these adversities, immobilization of probiotic cells onto a solid support, prior to their inclusion into the targeted food matrix, can be used to protect the cells and maintain their viability, activity and functionality during processing and storage. To this end, several studies have demonstrated the successful immobilization of probiotic bacteria onto various natural food-grade material supports, such as starch (Mattila-Sandholm et al., 2002), fruit pieces (Kourkoutas et al., 2006; Kourkoutas, Xolias, Kallis, Bezirtzoglou, & Kanellaki, 2005), casein (Dimitrellou, Kourkoutas, Koutinas, & Kanellaki, 2009) and wheat grains (Bosnea et al., 2009), and their application into food products, such as sausages (Sidira et al., 2014a) and cheese (Kourkoutas et al., 2006). However, there is limited knowledge, on how immobilization affects the functional properties of the probiotic cells, and in particular their ability to adhere to epithelial cells and their action as an intestinal barrier.

The probiotic properties of free and immobilized *L. casei* ATCC 393 were previously assessed by documenting the maintenance of cell viability after transit through the GI tract, adhesion at the large intestine and regulation of the intestinal microbiota in rats (Saxami et al., 2012, Sidira et al., 2010). Although, it was suggested that adhesion to the GI tract was transient, indicating the need for daily consumption of probiotic products containing the specific strain, no information about the effect of cell immobilization on adhesion properties were available. Similarly, *in vivo* experiments showed that *L. plantarum* enhanced the intestinal barrier and induced changes in the epithelial tight junctions (Karczewski et al., 2010).

Thus, the objective of the present study was to investigate the effect of probiotic immobilization onto wheat grains (wet and freeze dried) on the adhesion properties of lactobacilli using the model probiotic strains *L. casei* ATCC 393 and *L. plantarum* NCIMB 8826 in comparison to free wet and freeze dried cells. *L. plantarum* NCIMB 8826 was selected due to the strong binding characteristics (Izquierdo et al., 2009; Kinoshita et al., 2008), resistance to acidic conditions (Charalampopoulos, Pandiella, & Webb, 2002), survival during refrigerated storage (Charalampopoulos & Pandiella, 2010) and in fruit juices (Nualkaekul & Charalampopoulos, 2011), while *L. casei* ATCC 393 due to its good survival in the GI tract (Saxami et al., 2012; Sidira et al., 2010) and excellent technological properties (Bosnea et al., 2009; Kourkoutas et al., 2006, 2005; Sidira et al., 2014a).

2. Materials and methods

2.1 Strain maintenance and growth

L. casei ATCC 393 (DSMZ, Germany) and *L. plantarum* NCIMB 8826 (National Collection of Industrial and Marine Bacteria, UK) were stored at -20 °C in 2 ml cryovials containing 20 % (v/v) glycerol. The cells were initially propagated at 37 °C in 250 ml flasks containing 100 ml of MRS Broth (Fluka, Buchs, Switzerland) and then cultivated at 37 °C in 500 ml flasks for 72 h containing 200 ml of the same culture medium. The free cells were collected by centrifugation at 3200 g and 4 °C for 15 min.

2.2 Cell immobilization

Wheat grains were used as immobilization support. Prior to use, they were boiled and sterilized at 130 °C for 15 min. Cell immobilization was carried out as described previously (Bosnea et al., 2009). In brief, 50 g of immobilization support

and ~1 g (wet weight) of *L. casei* ATCC 393 or *L. plantarum* NCIMB 8826 cells were added into 1 L flasks containing 500 ml of MRS broth. The culture was incubated at 37 °C without agitation for 48 h. The immobilized cells were collected by washing twice with sterilized ¼ Ringer's solution (Sigma-Aldrich, UK) for removal of non-immobilized free cells. The immobilized cells were either used directly or subjected to freeze-drying.

2.3 Freeze-drying of free and immobilized cells

Free and immobilized cells were freeze-dried as described previously (Siaterlis, Deepika, & Charalampopoulos, 2009). Briefly, the immobilized cells were washed three times with ¼ Ringer's solution and transferred into 250 ml sterilin polystyrene container containing 25 ml of 10 % sucrose in PBS solution used as cryoprotectant. Similarly, the free cells were re-suspended in 10 ml of 10 % sucrose in PBS solution. Both immobilized and free cell suspensions in sucrose solutions were incubated at room temperature for 1 h and then frozen at -80 °C for 24 h. The frozen cultures were then freeze-dried in a IEC Lyoprep 3000 freeze-dryer (Lyoprep, Dunstable, UK) for approximately 2-4 days. The viable cell concentration were measured both pre- and post-freeze-drying. The water activity of freeze-dried powders in all cases was ~ 0.07. Each experiment was conducted in triplicate.

2.4 Adhesion of *Lactobacillus* cells to Caco-2 cells

Adhesion of the *Lactobacillus* cells to Caco-2 cells was studied as described previously (Deepika, Green, Frazier, & Charalampopoulos, 2009). The Caco-2 cell line was cultured in Eagle's Minimum Essential Medium (EMEM, Lonza, UK) supplemented with 10 % Fetal Bovine Serum (FBS), 1 % mixture penicillin-streptomycin solution and 1 % non-essential amino acid solution at 37 °C in an atmosphere of 5 % CO₂ and 95 % relative humidity. After 21 days, the six-well tissue

culture plate was ready to use as the cells were fully differentiated. One day prior to the adhesion assay, the Caco-2 cells were fed with the above culture medium, but without the presence of antibiotics. Before adhesion, the Caco-2 monolayer was washed twice with Dulbecco's phosphate buffered saline (DPBS, Lonza) in order to remove all traces of the medium. The number of the fully differentiated cells was counted using a Nikon microscope (Kingston upon Thames, UK) and was found to be around 5×10^5 cells/ml in all cases. Then, 10 ml of free or 1 g of immobilized bacterial cells (wet or freeze dried) were washed once with DPBS and then were re-suspended in DPBS to obtain an approximate concentration of around 10^8 - 10^9 CFU/ml and 1 ml of the bacterial suspension was added to each well. The plates were incubated at 37 °C in 5 % CO₂ and 95 % air. After 1 h, the supernatant was removed from the wells and the wells were washed twice with DPBS. Subsequently, 1 ml of DPBS was added to each well and the monolayer was scalped and transferred to 9 ml of ¼ Ringer's solution. The number of probiotic cells bound to the Caco-2 cells was determined after serial dilutions and plating in MRS agar plates. Adhesion of *Lactobacillus* strains was expressed as the number of viable bacteria adhering to 100 Caco-2 cells. Each experiment was conducted in triplicate.

2.5 Adhesion to *n*-hexadecane

The microbial adhesion to *n*-hexadecane (MATH assay) was employed to study the hydrophobicity of free and immobilized cells and was carried out as described previously (Deepika et al., 2009). In brief, 10 ml of free or 1 g of immobilized bacteria (wet or freeze-dried) were washed with PBS and suspended in 10 mM KH₂PO₄ to obtain an OD₆₀₀ ~ 0.8. The pH was adjusted to 3.0 with 1 M HCl and then 2 ml of the bacterial suspension were mixed with equal volume of *n*-hexadecane. The mixture was vortexed for 1 min and was allowed undisturbed to rest

for 20 min to achieve a complete phase separation. The aqueous phase was removed and the OD₆₀₀ was measured. The level of adhesion to *n*-hexadecane was expressed as the % percentage adhesion according to the following equation:

$$\% \text{ Adhesion to hexadecane} = \left(\frac{1 - A_1}{A_0} \right) \times 100$$

where A₀: initial absorbance, A₁: absorbance after 20 min of incubation. The MATH assay was conducted in triplicate for each sample.

2.6 Trans-epithelial electrical resistance (TEER) assay

TEER was used to measure the levels of tight junction in intestinal epithelial cells according to a method described previously (Commane et al., 2005). Cell culture inserts (0.4 µm, Becton Dickinson) were coated with 0.1 % type I rat-tail collagen (Sigma-Aldrich Company Ltd., Dorset, UK) and left to dry overnight under a UV light in six-well plates (Becton Dickinson). Caco-2 cells were seeded into the inserts at 2.5 mL aliquots per well with a concentration of 5 x 10⁵ cells/ml, as described above (Section 2.4). Eagle's Minimum Essential Medium (EMEM, Lonza) supplemented with 10 % Fetal Bovine Serum (FBS), 1 % mixture penicillin-streptomycin solution and 1 % non-essential amino acid solution (2 ml of culture medium) was added to the basal compartment of each well. The cells were grown at 37 °C in an atmosphere of 5 % CO₂ and 95 % relative humidity. The culture medium was refreshed every 2 days. After 12 days of culturing, the integrity of the monolayer was evaluated by measuring the TEER using an EVOM epithelial voltohmmeter chopstick electrode (World Precision Instruments, Stevenage, UK). Readings were performed every 24 h until the TEER values stabilized (day 16). TEER at time zero was determined before bacterial samples were added to the monolayer. Then, the culture medium from both the apical and basal compartments was removed and immediately 2 ml bacterial samples were added in the apical side (10⁸-10⁹ CFU/ml),

while 2 ml medium were added in the basal compartments. The cells were maintained for 1 h at 37 °C in an atmosphere of 5 % CO₂ and 95 % relative humidity. Then, the culture medium and the bacterial sample were removed and the apical side was washed with 2 ml DPBS. Thereafter, in both the apical and basal compartments, 2 ml of culture medium were applied and TEER readings were recorded after 1, 1.5 and 17 h in order to estimate the potential increase in TEER. One insert/well in every falcon was left without bacteria as a negative control. The percentage change in TEER was expressed according to the following equation:

$$\% \text{ change in TEER} = \left[\frac{(T1 - T0)}{T0} \right] \times 100$$

where T0: TEER at time zero, T1: TEER after 1 h, 1.5 h and 17 h. Each experiment was conducted in triplicate.

2.7 Scanning electron microscopy (SEM)

Monolayers of Caco-2 cells were grown on glass coverslips as described previously (Chauviere, Coconnier, Kerneis, Fourniat, & Servin, 1992). Briefly, Caco-2 cells were prepared on glass coverslips which were placed in six-well tissue culture plates. Cells were grown in culture medium at 37 °C in 5 % CO₂ and 95 % air. After the bacterial adhesion assay, the cells were fixed with 2.5 % glutaraldehyde in 0.1 M sodium phosphate buffer (pH 7.4) for 1 h at room temperature. Then, washing with 2 ml of the phosphate buffer was performed and the cells were osmicated for 30 min with 2 % OsO₄ and washed again 3 times with the same buffer. The coverslips were gradually dehydrated with 30 % ethanol for 15 min, 50 % for 15 min, 70 % for 15 min, 80 % for 15 min, 90 % for 15 min, and 3x100 % for 15 min to achieve absolute dry samples. The cells were further dried in a critical-point dryer (Balzers CPD 030) and coated with gold. The samples were imaged using a FEI Quanta 600 FEG scanning electron microscope. Bacterial attachment was evaluated by scanning

electron microscopy using a fixed number of monolayer cells (5×10^5 cells/well) and bacteria (8.2 log CFU added per well).

2.8 Experimental design and statistical analysis

All treatments were carried out in triplicate. The experiments were designed and analyzed statistically by ANOVA. Duncan's multiple range test was used to determine significant differences among results (coefficients, ANOVA tables and significance ($P < 0.05$) were computed using Statistica v.5.0).

3. Results and Discussion

Cell immobilization techniques are usually applied in order to maintain cell viability, activity and functionality during food production, processing and storage (Deepika, Rastall, & Charalampopoulos, 2011; Lopez-Rubio, Gavara, & Lagaron, 2006), and during passage through the GI tract (Saxami et al., 2012; Sidira et al., 2010). In order to confer a health effect, high adhesion ability of the probiotic cells is required (Ouwehand & Salminen, 2003). Despite the considerable amount of research aiming at evaluating the adhesion properties of free cells and the potential interaction between probiotics and the host (Ouwehand & Salminen, 2003; Tuomola & Salminen, 1998), there is limited knowledge on how the food matrix influences the adhesion ability of probiotics (Burgain et al., 2014; Endo et al., 2014; Bove et al. 2013). This is important for the design of novel foods that are able to maintain high cell viability and functionality.

To the best of our knowledge, this is the first report concerning the investigation of the effect of the immobilization support on the adhesion properties of probiotic bacteria. The strategy adopted was to use the model probiotic strains *L. casei* ATCC 393 and *L. plantarum* NCIMB 8826, which were immobilized on wheat

grains. Wheat grains were previously used as immobilization supports of *L. casei* ATCC 393 (Bosnea et al., 2009) and the immobilized cells were incorporated in traditional foods to confer probiotic properties (Sidira et al., 2014a; Sidira, Galanis, Nikolaou, Kanellaki, & Kourkoutas, 2014b). In addition, the effect of freeze-drying was also evaluated because wet cells are incompatible with commercial and industrial needs, which require robust cultures that can maintain their availability during storage.

3.1 Immobilization and freeze-drying of lactobacilli

The results concerning the effect of freeze-drying on the viability of free and immobilised lactobacilli onto wheat grains are presented in Figure 1. It can be observed that freeze-drying had no significant ($P>0.05$) effect on cell survival for both free and immobilized cells and that high cell concentrations were obtained post freeze-drying in all cases.

3.2 Adhesion to Caco-2 cells

The results concerning the adhesion abilities of both *L. casei* ATCC 393 and *L. plantarum* NCIMB 8826 to Caco-2 cells are presented in Figure 2, and indicate that both strains showed considerable adhesion ability to Caco-2 cells (>1000 CFU/100 Caco-2 cells for free wet cells). These values were higher than those reported in previous studies for other lactobacilli, which ranged between 0.9 and 900 CFU/100 Caco-2 cells (Ren et al., 2012; Bogovic-Matijasic, Narat, & Zori, 2003; Chauviere et al., 1992; Bernet, Brassart, Neeser, & Servin., 1994; Sarem, Sarem-Damerdj, & Nicolas, 1996; Coconnier, Klaenhammer, Kerneis, Bernet, & Servin, 1992). A cell concentration of 8-9 log CFU/ml of *Lactobacillus* was added to the Caco-2 culture, as

this is the recommended concentration for probiotics in order to exert a beneficial effect in the gut.

Although cell immobilization resulted in a significant reduction of adhesion ability (>100 CFU/100 Caco-2 cells for immobilized wet cells), adhesion still remained in satisfactory levels compared to values reported in literature concerning free cells (0.9-900 CFU/100 Caco-2 cells) (Bogovic-Matijasic et al., 2003; Chauviere et al., 1992; Bernet et al., 1994; Ren et al., 2012; Sarem et al., 1996; Coconnier et al., 1992).

Freeze-drying had a negative effect on the adhesion properties of free cells, but not of immobilized cells. The adhesion to Caco-2 cells decreased significantly for free cells (86 % and 71 % reduction for *L. plantarum* and *L. casei*, respectively). According to Henriksson, Szewzyk, & Conway, (1991) the adhesion of lactobacilli is most probably mediated by a proteinaceous component on the bacterial surface. Bacterial surface layers (S-layers) are composed of protein monomers arranged in crystalline arrays (Callegari et al., 1998) and it appears that freeze-drying can induce the detachment of S-layers monomers from the cell wall of lactobacilli (Ray & Johnson, 1986). This cell surface damage that occurs during freeze-drying may provide an explanation of the results above. Nevertheless, the adhesion values still were comparable to those recorded in the literature for freeze-dried cells, ranging from 3-325 CFU/100 Caco-2 cells (Bogovic-Matijasic et al., 2003; Chauviere et al., 1992; Bernet et al., 1994).

The adoption of functional criteria (adherence to human cell lines, resistance to gastric acidity and bile acids) for the *in vitro* selection of probiotic bacteria can result in the isolation of strains capable of performing effectively in the GI tract and that may reflect certain *in vivo* effects on the host (Dunne et al., 2001). A previous

study has shown how *in vitro* methods can be used for prediction of the survival potential of lactobacilli in the human GI tract (Jacobsen et al., 1999), although Lebeer et al., (2010a) has shown that *in vitro* studies are not related with survival *in vivo*. Our results validated the suggestions of Busscher & Weerkamp (1987), according to which the role of hydrophobic cells, supposed to be associated with bacterial surface appendages, is suggested to be its dehydrating capacity, allowing the removal of the water film and yielding a small area of direct contact between the protuberant parts of the cell surface and the substrate. Noticeably, the ability of probiotics to remove vicinal water depends greatly on the strain used.

3.3 Adhesion to *n*-hexadecane

Figure 3 shows the results from the MATH assay for wet and freeze-dried cells. The results indicated that the *L. casei* strain was relatively hydrophilic compared to the *L. plantarum* strain (54.8 % and 22.7 % for free cells of *L. plantarum* and *L. casei*, respectively). Immobilization affected ($P<0.05$) positively the adhesion only for *L. casei*. On the other hand, although freeze-drying resulted in increased ($P<0.05$) adhesion ability for *L. plantarum* (both free and immobilized) and for immobilized *L. casei*, it had no effect ($P>0.05$) on free *L. casei* cells.

Cell surface hydrophobicity has been associated with bacterial adhesion to a variety of surfaces (Marin et al., 1997). The consensus is that high hydrophobicity of bacteria results in greater attractive forces and higher levels of adhesion to Caco-2 cells, whereas smaller results in lower levels of adhesion (Rijnaarts, Norde, Bouwer, Lyklema, & Zehnder, 1993). According to Reid et al. (1992), the surface hydrophobicity of *Lactobacillus* strains varies greatly. In the present study, the results indicated that *L. casei* ATCC 393 strain was relatively hydrophilic (33.3% adhesion to *n*-hexadecane), whereas *L. plantarum* NCIMB 8826 is characterized as hydrophobic

(42.86% adhesion to *n*-hexadecane). Similar results were obtained by Pelletier et al., (1997) and Harty, Patrikakis, and Knox (1993) concerning the *L. casei* cells and by Zavisic et al., (2011), Jamaly, Benjouad, and Bouksaim (2011) and Abdulla, Abed, and Saeed (2014) concerning the *L. plantarum* cells.

However, no correlation between adhesion ability to the Caco-2 monolayer and hydrophobicity was observed for immobilized *L. casei* ATCC 393. Although the strain was characterized as relatively hydrophilic, the number of adhered immobilized *L. casei* was higher than immobilized *L. plantarum* (154 to 104 CFU adhered per 100 Caco-2 cells, respectively). The opposite results were observed in free cells, as hydrophobicity correlated well with adhesion ability (1038 and 2349 CFU adhered per 100 Caco-2 cells for *L. plantarum* and *L. casei*, respectively). A possible explanation for these observations might be that the immobilized cells do not have direct access to the Caco-2 cells (Figure 4). The inability to correlate hydrophobicity with adhesion to the Caco-2 monolayer is most likely due to many factors that are involved in the adhesion of lactobacilli (Azuma & Sato, 2001). Cell surface hydrophobicity is one of the physico-chemical properties that facilitate the first contact between the microorganism and the host cells. This non-specific initial interaction is weak and reversible and precedes the subsequent adhesion process mediated by more specific mechanisms involving cell-surface proteins and lipoteichoic acids (Schillinger, Guigas, & Holzapfel, 2005). Bacterial cell surface macromolecules may interact with host pattern recognition receptors (PRRs) of the GI mucosa and elicit specific responses in the intestinal system (Bron, Baarlen, & Kleerebezem, 2012; Lebeer, Vanderleyden, & Keersmaecker, 2010b). As stated above, structural damage (S-layer) of bacteria during freeze-drying can affect the adhesion ability (Callegari et al., 1998) and reduce its cell surface hydrophobicity

significantly (Bruinsma, van der Mei, & Busscher, 2001). It therefore appears that the adhesion was not caused by hydrophobic interaction. According to Schillinger et al., (2005), hydrophobicity is not a prerequisite for a strong adherence capacity.

3.4 Trans-Epithelial Electrical Resistance Assay

The most sensitive measure of mucosal barrier function is transepithelial electrical resistance (TEER), as it reflects the degree to which ions traverse tissue (Blikslager, Moeser, Gookin, Jones, & Odle, 2007). When TEER is increased, the tight junction between the cells becomes stronger and the permeability of the monolayer is decreased (Mattar, Drongowski, Coran, & Harmon, 2001). The TEER ratio was measured before adding the bacterial inoculum (time zero), and after addition and incubation of the probiotic strains in Caco-2 monolayer. The results from the TEER assay are presented in Table 1. For all samples, the TEER values increased with incubation time, reaching the maximum after 17 h of incubation.

The results indicated that both wet and freeze-dried free cells led to an increase in TEER values (58.4 and 43.6 % for wet *L. plantarum* and *L. casei*, respectively, and 73.6 and 73.7 % for freeze-dried *L. plantarum* and *L. casei*, respectively). Similar results were also published previously reporting increases in TEER by 20-158 % for *L. plantarum* (Anderson et al., 2010a, 2010b) and 6-87 % for *L. casei* cells (Fang et al., 2010). However, significantly ($P<0.05$) higher increase was recorded for immobilized cells (151 and 170 % for wet *L. plantarum* and *L. casei*, respectively, and 151 and 121 % for freeze-dried *L. plantarum* and *L. casei*, respectively). Notably, immobilized cells resulted in TEER values at least 2-fold higher than free cells, highlighting the positive effects of cell immobilization in enhancing the persistence of probiotic cells in the Caco-2 monolayer. It is thus suggested that immobilized cells act as a protective shield against intestinal

permeability, which might be due to increased physical interactions between the immobilised bacterial cells and the Caco-2 cells, or possibly due to increased synthesis of short chain fatty acids, which are known to be important nutrients for intestinal epithelial cells (Mangell et al., 2002). Improved epithelial barrier function has been previously associated with synbiotic food products containing immobilized probiotic microorganisms on prebiotic fibres (Commane et al., 2005). The findings of this study are expected to have a significant impact in the development of novel probiotic products targeting a range of GI disorders, as they indicated that cell immobilization, which has been suggested for the production of probiotic products (Kourkoutas et al., 2006; Bosnea et al., 2009; Sidira et al., 2014a, 2014b), can potentially offer the additional benefit of increased probiotic barrier function.

3.5 Scanning electron microscopy (SEM)

Figure 4 shows images from electron microscopy demonstrating key features of the Caco-2 cells, as well as the adhesion of the bacterial cells. It was noted that both immobilized and free cells adhered to the brush border microvilli and that the starch granules of wheat grains interacted with the Caco-2 monolayer, confirming the results of adhesion and TEER assays. Importantly, the morphology of the monolayer with the expression of microvilli, the tight-junctions of the starch granules of wheat, and the immobilized cells onto wheat grains are also obvious. Similar photographs were obtained for *L. plantarum* NCIMB 8826 strain and for freeze-dried cells (data not shown).

4. Conclusions

In conclusion, the present study showed that immobilization onto wheat grains decreased significantly the adhesion ability of both *L. casei* ATCC 393 and *L. plantarum* NCIMB 8826 strains to Caco-2 monolayers most likely because

immobilized cells did not have direct access to the Caco-2 cells. However, immobilization led to a significant enhancement of the monolayer integrity, which is compromised in a number of intestinal disorders, and is an important functional attribute of probiotic strains. Overall, more *in vitro* and *in vivo* research is needed in order to understand the effect of immobilization, which can also occur naturally within a food matrix, on the functional properties of probiotics.

Acknowledgments

This research has been co-financed by the European Union (European Social Fund – ESF) and Greek national funds through the Operational Program "Education and Lifelong Learning" of the National Strategic Reference Framework (NSRF) - Research Funding Program: Heracleitus II. Investing in knowledge society through the European Social Fund. The authors wish also to acknowledge the contribution of the Electron Microscopy Laboratory at the University of Reading for their help with SEM.

References

- Abdulla, A. A., Abed, T. A., & Saeed, A. M. (2014). Adhesion, Autoaggregation and Hydrophobicity of Six *Lactobacillus* Strains. *British Microbiology Research Journal*, 4, 381-391.
- Anderson, R. C., Cookson, A. L., McNabb, W. C., Park, Z., McCann, M. J., Kelly, W. J., & Roy, N. C. (2010a). *Lactobacillus plantarum* MB452 enhances the function of the intestinal barrier by increasing the expression levels of genes involved in tight junction formation. *BMC Microbiology*, 10, 316.
- Anderson, R. C., Cookson, A. L., McNabb, W. C., Kelly, W. J., & Roy, N. C. (2010b). *Lactobacillus plantarum* DSM2648 is a potential probiotic that enhances intestinal barrier function. *FEMS Microbiology Letters*, 309, 184-192.
- Azuma, Y., & Sato, M. (2001). *Lactobacillus casei* NY1301 increases the adhesion of *Lactobacillus gasseri* NY0509 to human intestinal Caco-2 cells. *Bioscience, Biotechnology, and Biochemistry*, 65, 2326-2329.
- Bernet, M. F., Brassart, D., Neeser, J. R., & Servin, A. L. (1994). *Lactobacillus acidophilus* LA 1 binds to cultured human intestinal cell lines and inhibits cell attachment and cell invasion by enterovirulent bacteria. *Gut*, 35, 483-489.
- Blikslager, A. T., Moeser, A. J., Gookin, J. L., Jones, S. L., & Odle, J. (2007). Restoration of Barrier Function in Injured Intestinal Mucosa. *Physiological Reviews*, 87, 545-564.
- Bogovic-Matijasic, B., Narat, M., & Zori, M. (2003). Adhesion of Two *Lactobacillus gasseri* Probiotic Strains on Caco-2 Cells. *Food Technology and Biotechnology*, 41, 83-88.

461 Bosnea, L., Kourkoutas, Y., Albantaki, N., Tzia, C., Koutinas, A. A., & Kanellaki, M.
 462 (2009). Functionality of freeze-dried *L. casei* cells immobilized on wheat grains.
 463 *LWT-Food Science and Technology*, 42, 1696-1702.

464 Bove, P., Russo, P., Capozzi, V., Gallone, A. Spano, G., Fiocco D. (2013)
 465 *Lactobacillus plantarum* passage through an oro-gastro-intestinal tract simulator:
 466 Carrier matrix effect and transcriptional analysis of genes associated to stress and
 467 probiosis. *Microbiological Research*, 168, 351–359.

468 Boylston, T. D., Vinderola, C. G., Ghoddusi, H. B., & Reinheimer J. A. (2004).
 469 Incorporation of bifidobacteria into cheeses: Challenges and rewards. *International*
 470 *Dairy Journal*, 19, 315–387.

471 Bron, P. A., Baarlen, P., & Kleerebezem, M. (2012). Emerging molecular insights
 472 into the interaction between probiotics and the host intestinal mucosa. *Nature Reviews*
 473 *Microbiology*, 10, 66-78.

474 Bruinsma, G. M., van der Mei, H. C., & Busscher, H. J. (2001). Bacterial adhesion to
 475 surface hydrophilic and hydrophobic contact lenses. *Biomaterials*, 22, 3217-3224.

476 Burgain, J., Scher, J., Francius, G., Borges, F., Corgneau, M., Revol-Junelles, A. M.,
 477 Cailliez-Grimal, C., Gaiani, C. (2014). Lactic acid bacteria in dairy food: Surface
 478 characterization and interactions with food matrix components. *Advances in Colloid*
 479 *and Interface Science*, 213, 21-35.

480 Busscher, H. J., & Weerkamp, A. H. (1987). Specific and non-specific interactions in
 481 bacterial adhesion to solid substrata. *FEMS Microbiology Reviews*, 46, 165-173.

482 Charalampopoulos, D., Pandiella, S. S., & Webb, C. (2002). Growth studies of
 483 potentially probiotic lactic acid bacteria in cereal-based substrates. *Journal of Applied*
 484 *Microbiology*, 92, 851–859.

485 Charalampopoulos, D., & Pandiella, S. S. (2010). Survival of human derived
 486 *Lactobacillus plantarum* in fermented cereal extracts during refrigerated storage.
 487 *Food Science and Technology*, 43, 431–435.

488 Nualkaekul, S., & Charalampopoulos D. (2011). Survival of *Lactobacillus plantarum*
 489 in model solutions and fruit juices. *International Journal of Food Microbiology*, 146,
 490 111–117.

491 Callegari, M. L., Riboli, B., Sanders, J. W., Cocconcelli, P. S., Kok, J., Venema, G.,
 492 & Morelli, L. (1998). The Slayer gene of *Lactobacillus helveticus* CNRZ 892:
 493 cloning, sequence and heterologous Expression. *Microbiology*, 144, 719-726.

494 Chauviere, G., Coconnier, M. H., Kerneis, S., Fourniat, J., & Servin, A.L. (1992).
 495 Adhesion of human *Lactobacillus acidophilus* strain LB to human enterocyte-like
 496 Caco-2 cells. *Journal of General Microbiology*, 138, 1689-1696.

497 Coconnier, M. H., Klaenhammer, T. R., Kerneis, S., Bernet, M. F. & Servin, A. L.,
 498 (1992). Protein-Mediated Adhesion of *Lactobacillus acidophilus* BG2FO4 on Human
 499 Enterocyte and Mucus-Secreting Cell Lines in Culture. *Applied and Environmental*
 500 *Microbiology*, 58, 2034-2039.

501 Commane, D. M., Shortt, C. T., Silvi, S., Cresci, A., Hughes R. M., & Rowland I. R.
 502 (2005). Effects of Fermentation Products of Pro- and Prebiotics on Trans-Epithelial
 503 Electrical Resistance in an In Vitro Model of the Colon. *Nutrition and Cancer*, 51,
 504 102-109.

505 Deepika, G., Green, R. J., Frazier, R. A., & Charalampopoulos, D. (2009). Effect of
 506 growth time on the surface and adhesion properties of *Lactobacillus rhamnosus* GG.
 507 *Journal of Applied Microbiology*, 107, 1230-1240.

508 Deepika, G., Rastall, R. A., & Charalampopoulos D. (2011). Effect of Food Models
509 and Low-Temperature Storage on the Adhesion of *Lactobacillus rhamnosus* GG to
510 Caco-2 Cells. *Journal of Agricultural and Food Chemistry*, 59, 8661-8666.

511 Deshpandea, G., Rao, S., & Patolea S. (2011). Progress in the field of probiotics: year
512 2011. *Current Opinion in Gastroenterology*, 27, 13-18.

513 Dimitrellou, D., Kourkoutas, Y., Koutinas, A. A., & Kanellaki, M. (2009). Thermally-
514 dried immobilized kefir on casein as starter culture in dried whey cheese production.
515 *Food Microbiology*, 26, 809-820.

516 Douillard, F. P., Ribbera, A., Kant, R., Pietila, T. E., Jarvinen, H. M., Messing, M.,
517 Randazzo, C. L., Paulin, L., Laine, P., Ritari, J., Caggia, C., Lahtinen, T., Brouns, S.
518 J. J., Satokari, R., Ossowski, I., Reunanen, J., Palva, A., & Vos, W. M. (2013).
519 Comparative Genomic and Functional Analysis of 100 *Lactobacillus rhamnosus*
520 Strains and Their Comparison with Strain GG. *Plos Genetics*, 9, e1003683. doi:
521 10.1371/journal.pgen.1003683.

522 Duary, R. K., Rajput, Y. S., Batish, V. K., & Grover, S. (2011). Assessing the
523 adhesion of putative indigenous probiotic lactobacilli to human colonic epithelial
524 cells. *Indian Council of Medical Research*, 134, 664-671.

525 Dunne, C., O'Mahony, L., Murphy, L., Thornton, G., Morrissey, D., O'Halloran, S.,
526 Feeney, M., Flynn, S., Fitzgerald, G., Daly, C., Kiely, B., O'Sullivan, G. C.,
527 Shanahan, F., & Collins, J. K. (2001). *In vitro* selection criteria for probiotic bacteria
528 of human origin: correlation with *in vivo* findings. *The American Journal of Clinical*
529 *Nutrition*, 73, 386S-392S.

530 Elo, S., Saxelin, M., & Salminen, S. (1991). Attachment of *Lactobacillus casei* strain
531 GG to human colon carcinoma cell line Caco-2: comparison with other dairy strains.
532 *Letters in Applied Microbiology*, 13, 154-156.

533 Endo, A., Teräsjärvi, J., Salminen, S. (2014). Food matrices and cell conditions
534 influence survival of *Lactobacillus rhamnosus* GG under heat stresses and during
535 storage. *International Journal of Food Microbiology*, 174, 110-112.

536 Fang H. W., Fang, S. B., Chiau J. S. C., Yeung, C. Y., Chan, W. T., Jiang, C. B.,
537 Cheng, M. L., & Lee, H. C. (2010). Inhibitory effects of *Lactobacillus casei* subsp.
538 *rhamnosus* on *Salmonella* lipopolysaccharide-induced inflammation and epithelial
539 barrier dysfunction in a co-culture model using Caco-2/ peripheral blood mononuclear
540 cells. *Journal of Medical Microbiology*, 59, 573-579.

541 Harty, D. W. S., Patrikakis, M., & Knox, K. W. (1993). Identification of
542 *Lactobacillus* Strains Isolated from Patients with Infective Endocarditis and
543 Comparison of their Surface-associated Properties with those of Other Strains of the
544 Same Species. *Microbial Ecology in Health and Disease*, 6, 191-201.

545 Hempel, S., Newberry, S. J., Maher, A. R., Wang, Z., Miles, J. N. V., Shanman, R.,
546 Johnsen, B., & Shekelle, P. G. (2012). Probiotics for the Prevention and Treatment of
547 Antibiotic-Associated Diarrhea A Systematic Review and Meta-analysis. *Journal of*
548 *the American Medical Association*, 307, 1959-1969.

549 Henriksson, A., Szewzyk, R., & Conway, P. L. (1991). Characteristics of the
550 Adhesive Determinants of *Lactobacillus fermentum* 104. *Applied and Environmental*
551 *Microbiology*, 499-502.

552 Hilgendorf, C., Spahn-Langguth, H., Regardh, C. G., Lipka, E., Amidon, G.L., &
553 Langguth, P. (2000). Caco-2 versus Caco-2/HT29-MTX co-cultured cell lines:
554 Permeabilities via diffusion, inside- and outside-directed carrier-mediated transport.
555 *Journal of Pharmaceutical Sciences*, 89, 63-75.

556 Izquierdo, H., Horvatovich, P., Marchioni, E., Aoude-Werner, D., Sanz, Y., &
557 Ennahar, S. (2009). 2-DE and MS analysis of key proteins in the adhesion of

558 *Lactobacillus plantarum*, a first step toward early selection of probiotics based on
559 bacterial biomarkers. *Electrophoresis*, 30, 949–956.

560 Jacobsen, C. N., Nielsen, V. R., Hayford, A. E., Moller, P. L., Michaelsen, K. F.,
561 Pærregaard, A., Sandström, B., Tvede, M., & Jakobsen, M. (1999). Screening of
562 Probiotic Activities of Forty-Seven Strains of *Lactobacillus* spp. by *In Vitro*
563 Techniques and Evaluation of the Colonization Ability of Five Selected Strains in
564 Humans. *Applied and Environmental Microbiology*, 65, 4949-4956.

565 Jamaly, N., Benjouad A. & Bouksaim M. (2011). Probiotic Potential of *Lactobacillus*
566 strains Isolated from Known Popular Traditional Moroccan Dairy Products. *British*
567 *Microbiology Research Journal*, 1, 79-94.

568 Karczewski, J., Troost, F. J., Konings, I., Dekker, J., Kleerebezem, M., Brummer R. J.
569 M., & Wells, J. M. (2010). Regulation of human epithelial tight junction proteins by
570 *Lactobacillus plantarum* in vivo and protective effects on the epithelial barrier.
571 *American journal of physiology - Gastrointestinal and Liver Physiology*, 298, G851-
572 G859.

573 Kechagia, M., Basoulis, D., Konstantopoulou, S., Dimitriadi, D., Gyftopoulou, K.,
574 Skarmoutsou, N., & Fakiri, E. M. (2013). Health Benefits of Probiotics: A Review.
575 *ISRN Nutrition*, 1-7.

576 Kinoshita, H., Uchida, H., Kawai, Y., Kawasaki, T., Wakahara, N., Matsuo, H.,
577 Watanabe, M. Kitazawa, H., Ohnuma, S., Miura, K., Horii, A., & Saito T. (2008).
578 Cell surface *Lactobacillus plantarum* LA 318 glyceraldehyde-3-phosphate
579 dehydrogenase (GAPDH) adheres to human colonic mucin. *Journal of Applied*
580 *Microbiology*, 104, 1667–1674.

581 Klingberg, T. D., Pedersen, M. H., Cencic, A., & Budde, B. B. (2005). Application of
582 Measurements of Transepithelial Electrical Resistance of Intestinal Epithelial Cell

583 Monolayers To Evaluate Probiotic Activity. *Applied and Environmental*
584 *Microbiology*, 71, 7528-7530.

585 Kourkoutas, Y., Xolias, V., Kallis, M., Bezirtzoglou, E., & Kanellaki M. (2005).
586 *Lactobacillus casei* immobilization on fruit pieces for probiotic additive, fermented
587 milk and lactic acid production. *Process Biochemistry*, 40, 411–416.

588 Kourkoutas, Y., Bosnea, L., Taboukos, S., Baras, C., Lambrou, D., & Kanellaki, M.
589 (2006). Probiotic cheese production using *Lactobacillus casei* cells immobilized on
590 fruit pieces. *Journal of Dairy Science*, 89, 1439–1451.

591 Lebeer, S., Claes, I. J. J., Verhoeven, T. L. A., Vanderleyden, J., & Keersmaecker, S.
592 C. J. (2010a). Exopolysaccharides of *Lactobacillus rhamnosus* GG form a protective
593 shield against innate immune factors in the intestine. *Microbial Biotechnology*, 4,
594 368-374.

595 Lebeer, S., Vanderleyden, J., & Keersmaecker, S. C. J. (2010b). Host interactions of
596 probiotic bacterial surface molecules: comparison with commensals and pathogens,
597 *Nature Reviews Microbiology*, 8, 171-184.

598 Lopez-Rubio, A., Gavara, R., & Lagaron, J. M. (2006). Bioactive packaging: turning
599 foods into healthier foods through biomaterials. *Trends in Food Science and*
600 *Technology*, 17, 567-575.

601 Mangell, P., Nejdfors, P., Wang, M., Ahrne, S., Westrom, B., Thorlacius, H., &
602 Jeppsson, B. (2002). *Lactobacillus plantarum* 299v inhibits Escherichia coli induced
603 intestinal permeability. *Digestive Diseases and Sciences*, 47, 511–516,

604 Marin, M. L., Benito, Y., Pin, C., Fernández, M. F., García, M. L., Selgas, M. D., &
605 Casas, C. (1997). Lactic acid bacteria: hydrophobicity and strength of attachment to
606 meat surfaces. *Letters in Applied Microbiology*, 24, 14-18.

607 Mattar, A. F., Drongowski, R. A., Coran, A. G., & Harmon, C. M. (2001). Effect of
608 probiotics on enterocyte bacteria translocation in vitro. *Pediatric Surgery*
609 *International*, 17, 265-268.

610 Mattila-Sandholm, T., Myllarinen, P., Crittenden, R., Mogensen, G., Fonden, R., &
611 Saarela, M. (2002). Technological challenges for future probiotic foods. *International*
612 *Dairy Journal*, 12, 173–182.

613 Nagpal, R., Kumar, A., Kumar, M., Behare, P. V., Jain, S., & Yadav, H. (2012).
614 Probiotics, their health benefits and applications for developing healthier foods: a
615 review. *FEMS Microbiology Letters*, 334, 1-15.

616 Oliveira, R., Azeredo, J., Teixeira, P., & Fonseca, A. P. (2001). The role of
617 hydrophobicity in bacterial adhesion. Meeting of the Biofilm Club, *BioLine* (pp. 11-
618 22). Powys, UK.

619 Ouwehand, A. C., & Salminen, S. (2003). *In vitro* Adhesion Assays for Probiotics and
620 their in vivo NYH6 Relevance: A Review. *Microbial Ecology in Health and Disease*,
621 15, 175-184.

622 Pelletier, C., Bouley, C., Cayuela, C., Bouttier, S., Bourlioux, P., & Bellon-Fontaine,
623 M. N. (1997). Cell Surface Characteristics of *Lactobacillus casei* subsp. *casei*,
624 *Lactobacillus paracasei* subsp. *paracasei*, and *Lactobacillus rhamnosus* Strains.
625 *Applied and Environmental Microbiology*, 63, 1725–1731.

626 Ray, B., & Johnson, M. C. (1986). Freeze-drying injury of surface layer protein and
627 its protection in *Lactobacillus acidophilus*. *Cryo Letters*, 7, 210-217.

628 Reid, G., Cuperus, P. L., Bruce, A. W., van der Mei, H., Tomeczek, L., Khoury, A.
629 H., & Buscher, H. J. (1992). Comparison of Contact Angles and Adhesion to
630 Hexadecane of Urogenital, Dairy, and Poultry Lactobacilli: Effect of Serial Culture
631 Passages. *Applied and Environmental Microbiology*, 58, 1549-1553.

632 Ren, D., Li, C., Qin, Y., Yin, R., Li, X., Tian, M., Du, S., Guo, H., Liu, C., Zhu, N.,
 633 Sun, D., Li, Y., & Jin, N. (2012). Inhibition of *Staphylococcus aureus* adherence to
 634 Caco-2 cells by lactobacilli and cell surface properties that influence attachment.
 635 *Anaerobe*, 18, 508-515.

636 Rijnaarts, H. H. M., Norde, W., Bouwer, E. J., Lyklema, J., & Zehnder, A. J. B.
 637 (1993). Bacterial Adhesion under Static and Dynamic Conditions. *Applied and*
 638 *Environmental Microbiology*, 59, 3255-3265.

639 Ritchie, M. L., & Romanuk, T. N. (2012). A Meta-Analysis of Probiotic Efficacy for
 640 Gastrointestinal Diseases. *Plos One*, 7, e34938-e34938. doi:
 641 10.1371/journal.pone.0034938.

642 Sarem, F., Sarem-Damerdj, L. O., & Nicolas, J. P. (1996). Comparison of the
 643 adherence of three *Lactobacillus* strains to Caco-2 and Int-407 human intestinal cell
 644 lines. *Letters in Applied Microbiology*, 22, 439-442.

645 Saxami, G., Ypsilantis, P., Sidira, M., Simopoulos, C., Kourkoutas, Y., & Galanis, A.
 646 (2012). Distinct adhesion of probiotic strain *Lactobacillus casei* ATCC 393 to rat
 647 intestinal mucosa. *Anaerobe*, 18, 417-420.

648 Schillinger, U., Guigas, C., & Holzapfel, W. H. (2005). *In vitro* adherence and other
 649 properties of lactobacilli used in probiotic yoghurt-like products. *International Dairy*
 650 *Journal*, 15, 1289-1297.

651 Segers, M. E., & Lebeer, S. (2014). Towards a better understanding of *Lactobacillus*
 652 *rhamnosus* GG - host interactions. *Microbial Cell Factories*, 13, S1-S7.

653 Siaterlis, A., Deepika, G., & Charalampopoulos, D. (2009). Effect of culture medium
 654 and cryoprotectants on the growth and survival of probiotic lactobacilli during freeze
 655 drying. *Letters in Applied Microbiology*, 48, 295-301.

656 Sidira, M., Karapetsas, A., Galanis, A., Kanellaki, M., & Kourkoutas, Y. (2014a).
657 Effective survival of immobilized *Lactobacillus casei* during ripening and heat
658 treatment of probiotic dry-fermented sausages and investigation of the microbial
659 dynamics. *Meat Science*, 96, 948-955.

660 Sidira, M., Galanis, A., Nikolaou, A., Kanellaki, M., & Kourkoutas, Y. (2014b).
661 Evaluation of *Lactobacillus casei* ATCC 393 protective effect against spoilage of
662 probiotic dry-fermented sausages. *Food Control*, 42, 315-320.

663 Sidira, M., Saxami, G., Dimitrellou, D., Santarmaki, V., Galanis, A., & Kourkoutas,
664 Y. (2013). Monitoring survival of *Lactobacillus casei* ATCC 393 in probiotic yogurts
665 using an efficient molecular tool. *Journal of Dairy Science*, 96, 3369–3377.

666 Sidira, M., Galanis, A., Ypsilantis, P., Karapetsas, A., Progaki, Z., Simopoulou, C., &
667 Kourkoutas Y. (2010). Effect of probiotic-fermented milk administration on
668 gastrointestinal survival of *Lactobacillus casei* ATCC 393 and modulation of
669 intestinal microbial flora. *Journal of Molecular Microbiology and Biotechnology*, 19,
670 224–230.

671 Tuomola, E. M., & Salminen, S. J. (1998). Adhesion of some probiotic and dairy
672 *Lactobacillus* strains to Caco-2 cell cultures. *International Journal of Food*
673 *Microbiology*, 41, 45-51.

674 van der Mei, H. C., van der Belt- Gritter, B., Pouwels, P. H., Martinez, B., &
675 Busscher, H. J. (2003). Cell surface hydrophobicity is conveyed by S-layer proteins –
676 a study in recombinant lactobacilli. *Colloids and Surfaces B: Biointerfaces*, 28, 127-
677 134.

678 Zavisic, G., Radulovic, Z., Vranic, V., Begovic, J., Topisirovic L., & Strahinic, I.
679 (2011). Characterization and antimicrobial activity of vaginal *Lactobacillus* isolate.
680 *Archives of Biological Sciences*, 63, 29-35.

681

Table 1. Changes in Trans-Epithelial Electrical Resistance (TEER) values of Caco-2 cells in the presence of *L. plantarum* NCIMB 8826 and *L. casei* ATCC 393 cells. Changes are expressed as percentage change in TEER values. Significant differences ($P<0.05$) are indicated by different letters in superscript.

Strain	Time (h)	Free		Immobilized	
		Wet	Freeze-dried	Wet	Freeze-dried
<i>L. plantarum</i> NCIMB 8826	1.0	-1.8±0.1 ^a	6.2±0.3 ^b	32.6±1.6 ^c	8.45±0.4 ^d
	1.5	25.7±5.3 ^{c,e}	18.9±0.6 ^f	111.6±38.8 ^g	47.3±23.1 ^h
	17.0	58.4±10.3 ^h	73.6±29.9 ^{g,h,i,j,k}	150.8±54.6 ^{i,l,p}	150.7±33.1 ^{i,l,p}
<i>L. casei</i> ATCC 393	1.0	13.9±0.7 ^m	1.3±0.1 ⁿ	25.9±1.3 ^e	2.6±0.1 ^o
	1.5	10.7±2.0 ^d	41.2±8.2 ^h	86.3±3.7 ^j	35.2±9.4 ^{c,e,k}
	17.0	43.6±10.9 ^{h,k}	73.7±22.9 ^{g,h,i,j}	169.8±9.9 ^l	121.1±2.2 ^p

Legends to Figures

Figure 1. Effect of freeze-drying on viability of lactobacilli. Significant differences ($P<0.05$) are indicated by different letters.

Figure 2. Adhesion of lactobacilli to Caco-2 cells (CFU adhered per 100 Caco-2 cells). Significant differences ($P<0.05$) are indicated by different letters.

Figure 3. Adhesion of *L. plantarum* NCIMB 8826 and *L. casei* ATCC 393 to *n*-hexadecane. Significant differences ($P<0.05$) are indicated by different letters.

Figure 4. SEM micrographs showing: a) the morphology of complete Caco-2 monolayer, b) adhesion of free *L. casei* ATCC 393 bacteria to Caco-2 cells, c) adhesion of *L. plantarum* NCIMP 8826 bacteria to Caco-2 cells and the level of intercellular junctions (shown by arrows), d) starch granules of wheat interacting with the Caco-2 monolayer and e) immobilized *L. casei* ATCC 393 cells on wheat grains.




Fig. 1




Fig. 2





Fig. 3.


714



715

716 **Fig. 4a**

717



718

719 **Fig. 4b**

720

721




Fig. 4c




Fig. 4d




Fig. 4e