

Global change pressures on soils from land use and management

Article

Accepted Version

Smith, P., House, J. I., Bustamante, M., Sobocká, J., Harper, R., Pan, G., West, P., Clark, J. ORCID: <https://orcid.org/0000-0002-0412-8824>, Adhya, T., Rumpel, C., Paustian, K., Kuikman, P., Cotrufo, M. F., Elliott, J. A., McDowell, R., Griffiths, R. I., Asakawa, S., Bondeau, A., Jain, A. K., Meersmans, J. and Pugh, T. A.M. (2016) Global change pressures on soils from land use and management. *Global Change Biology*, 22 (3). pp. 1008-1028. ISSN 1365-2486 doi: 10.1111/gcb.13068 Available at <https://centaur.reading.ac.uk/42100/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

To link to this article DOI: <http://dx.doi.org/10.1111/gcb.13068>

Publisher: Wiley-Blackwell

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online

Global Change Pressures on Soils from Land Use and Management

Pete Smith¹, Jo I. House², Mercedes Bustamante³, Jaroslava Sobocká⁴, Richard Harper⁵, Genxing Pan⁶, Paul West⁷, Jo Clark⁸, Tapan Adhya⁹, Cornelia Rumpel¹⁰, Keith Paustian¹¹, Peter Kuikman¹², M. Francesca Cotrufo¹¹, Jane A. Elliott¹³, Richard McDowell¹⁴, Robert I. Griffiths¹⁵, Susumu Asakawa¹⁶, Alberte Bondeau¹⁷, Atul K. Jain¹⁸, Jeroen Meersmans¹⁹ and Thomas A.M. Pugh²⁰

¹ Institute of Biological and Environmental Sciences, Scottish Food Security Alliance-Crops & ClimateXChange, University of Aberdeen, 23 St Machar Drive, Aberdeen, AB24 3UU, UK

² Cabot Institute, School of Geographical Sciences, University of Bristol, University Road, Bristol, BS8 1SS, U

³ Departamento de Ecologia, Universidade de Brasília, I.B. C.P. 04457. Campus Universitário Darcy Ribeiro - UnB. D.F.. CEP: 70919-970 Brasília, Brazil

⁴ National Agriculture and Food Centre Lužianky, Soil Science and Conservation Research Institute Bratislava, Gagarinova 10, 827 13 Bratislava, Slovakia

⁵ School of Veterinary and Life Sciences, Murdoch University, South Street, Murdoch WA. 6150 Australia

⁶ Institute of Resources, Environment and Ecosystem of Agriculture, Nanjing Agricultural University, 1 Weigang, Nanjing 210095, China

⁷ Global Landscapes Initiative, Institute on the Environment (IonE), University of Minnesota, 325 Learning & Environmental Sciences, 1954 Buford Ave, St. Paul, MN 55108, USA

⁸ Department of Geography and Environmental Science, School of Archaeology, Geography and Environmental Science, The University of Reading, Whiteknights, PO Box 227, Reading, RG6 6AB, UK

⁹ School of Biotechnology, KIIT University, Bhubaneswar - 751024, Odisha, India

¹⁰ CNRS, Campus AgroParisTech, Bâtiment EGER, 78850 Thiverval-Grignon, France

¹¹ Department of Soil and Crop Sciences & Natural Resource Ecology Laboratory, Colorado State University, Fort Collins, Colorado 80523-1499, USA

¹² Alterra Wageningen UR, PO Box 47, 6700AA Wageningen, The Netherlands

¹³ Environment Canada, National Hydrology Research Centre, Saskatoon, Saskatchewan, S7N 3H5, Canada

¹⁴ AgResearch, Invermay Agricultural Centre, Private Bag 50034, Mosgiel, New Zealand

¹⁵ Centre for Ecology & Hydrology, Maclean Building, Benson Lane, Crowmarsh Gifford Wallingford, OX10 8BB, UK

¹⁶ Graduate School of Bioagricultural Sciences, Nagoya University, Chikusa, Nagoya 464-8601, Japan

¹⁷ Mediterranean Institute of Biodiversity and Ecology (IMBE), Aix-Marseille University, Centre National de la Recherche Scientifique (CNRS) - Institut de Recherche pour le Développement (IRD) - Université d'Avignon Pays du Vaucluse (UAPV), F-13545 Aix-en-Provence Cedex 04, France

¹⁸ Department of Atmospheric Sciences, University of Illinois @ Urbana-Champaign 105 S. Gregory Street, Urbana, IL 61801, USA

¹⁹ Department of Geography, College of Life and Environmental Sciences, University of Exeter, Armory Building, Renes Drive, Exeter EX4 4RJ, UK

²⁰ Karlsruhe Institute of Technology, Institute of Meteorology and Climate Research / Atmospheric Environmental Research (IMK-IFU), Kreuzeckbahnstrasse 19, 82467 Garmisch-Partenkirchen, Germany

50

51 ***Corresponding author:** Prof Pete Smith, Tel: +44 (0)1224 272702, Fax: +44 (0)1224
52 272703, E-mail: pete.smith@abdn.ac.uk

53 **Running head:** Global Change Pressures on Soils

54 **Keywords:** soil, land use change, land use intensity, nitrogen deposition, sulphur deposition,
55 heavy metal deposition

56 **Paper type:** Invited Review

57

58

59 **Abstract**

60
61 Soils are subject to varying degrees of direct or indirect human disturbance, constituting a
62 major global change driver. Factoring out natural from direct and indirect human influence is
63 not always straightforward, but some human activities have clear impacts. These include land
64 use change, land management, and land degradation (erosion, compaction, sealing and
65 salinization). The intensity of land use also exerts a great impact on soils, and soils are also
66 subject to indirect impacts arising from human activity, such as acid deposition (sulphur and
67 nitrogen) and heavy metal pollution. In this critical review, we report the state-of-the-art
68 understanding of these global change pressures on soils, identify knowledge gaps and
69 research challenges, and highlight actions and policies to minimise adverse environmental
70 impacts arising from these global change drivers.

71
72 Soils are central to considerations of what constitutes sustainable intensification. Therefore,
73 ensuring that vulnerable and high environmental value soils are considered when protecting
74 important habitats and ecosystems, will help to reduce the pressure on land from global
75 change drivers. To ensure that soils are protected as part of wider environmental efforts, a
76 global soil resilience programme should be considered, to monitor, recover or sustain soil
77 fertility and function, and to enhance the ecosystem services provided by soils. Soils cannot,
78 and should not, be considered in isolation of the ecosystems that they underpin and vice
79 versa. The role of soils in supporting ecosystems and natural capital needs greater
80 recognition. The lasting legacy of the International Year of Soils in 2015 should be to put
81 soils at the centre of policy supporting environmental protection and sustainable
82 development.

1. Introduction

2015 is the International Year of Soil. This represents an ideal time to take stock of scientific knowledge about the changing global pressures that humans are exerting on soils. 2015 is also the year when policy makers will adopt a new legally-binding climate agreement under the United Nations Framework Convention on Climate Change (UNFCCC), with individual countries and businesses making policies and targets on greenhouse gas emissions and removals. Soils storage and cycling of carbon and nitrogen are part of emissions and removals from the land sector. Furthermore, 2015 is the year when countries will shape and adopt a new development agenda that will build on the Millennium Development Goals (MDGs). With increasing population, issues such as food security, water security, energy security (including bioenergy production) and sustainable integrated land and resource management are central to many development research and policy agendas. Soils underpin the provision of many ecosystem services related to development.

Soils provide multiple ecosystem services, allowing sustained food and fibre production, and delivering climate regulation, flood regulation, improved air and water quality, reducing soil erosion, and provide a reservoir for biodiversity (Smith et al. 2015). All soils are subject to some degree of human disturbance, either directly through land-use and land management, or indirectly through responses to human-induced global change such as pollution and climate change. Distinguishing natural from direct and indirect human influence is not always straightforward (Smith, 2005), but some human activities and their consequences have clear impacts, and despite large heterogeneity in soil properties and responses, robust scientific knowledge exists.

Human impacts on soils largely emerge from the need to meet the food, fibre, and fuel demands of a growing population including an increase in meat consumption as developing nations become wealthier, the production of biofuels, and increasing areas of urbanization. This has led to conversion of natural land to managed land (extensification) and intensification of agricultural and other management practices on existing land such as increasing nutrient and water inputs and increasing harvest frequency to increase yields per hectare.

Land cover or land use change (e.g. from forest or natural grassland to pasture or cropland), removes biomass, changes vegetation and disturbs soils, leading to loss of soil carbon and other nutrients, changes in soil properties, and changes to above- and below-ground biodiversity. Some land cover conversions e.g. reforestation after abandonment of cropland, can increase both above- and below-ground carbon and nutrients. Land use or land management that does not result in a change of cover (e.g. forest harvest and regrowth, increased grazing intensity and intensification of crop production), can potentially result in degradation of soil properties, depending on the characteristics of the management practices.

Land use change has been accelerated by population increases and migration as food, shelter, and materials are sought and acquired. It is estimated that humans have directly modified at least 70 Mkm², or >50 percent of Earth's ice-free land area (Hooke et al. 2012). The new Global Land Cover Share-database (Latham et al., 2014) represents the major land cover classes defined by the FAO. Croplands and grasslands (including both natural grasslands and managed grazing lands) each covered 13.0 %. "Tree-covered areas" (i.e. both natural and managed forests) covered 28%, shrub-covered areas 9.5 %. Artificial surfaces (including urbanised areas) occupy 1 %. Land degradation can be found in all land cover types. Degraded land covers approximately 24% of the global land area (35 Mkm²). 23% of degrading land is under broadleaved forest, 19% under needle-leaved forests and 20-25% on rangeland (Bai *et al.*, 2008).

In this review we report the state-of-the-art understanding, and knowledge gaps concerning impacts of changes in anthropogenic land use and land management on soils, including interactions with other anthropogenic global change pressures. We also review actions and policies that limit the adverse impacts arising from these global change drivers. We make the case to put soils at the centre of research strategy and policy actions as a legacy of the International Year of Soils.

2. Land use/land cover change

Land cover change has been dominated by deforestation, but also conversion of grasslands to cropland and grazing land. Deforestation has had the greatest impact on historical soil carbon change, causing on average around 25% of soil carbon to be lost (Guo & Gifford, 2002; Murty *et al.*, 2002). Soil carbon losses largely stem from oxidation of the organic matter as

well as soil erosion.

Deforestation affected an estimated 13 million hectares per year between 2000 and 2010; net forest loss was 5.2 million hectares per year (FAO, 2010). Most of this recent deforestation has taken place in tropical countries (FAO, 2010; Hansen et al., 2013). Over 50% of tropical forest loss occurred in Brazil and Indonesia, largely driven by a few commodities: timber, soy, beef, and oil palm (West *et al.*, 2014). There has been a reduced rate of deforestation in some regions over the last decade, most notably Brazil (INPE, 2014), largely because of land use conservation policies (Soares-Filho *et al.*, 2014; Nolte *et al.*, 2013) as well as economics. Most developed countries with temperate and boreal forest ecosystems – and more recently, countries in the Near East and Asia – are experiencing stable or increasing forest areas in contrast to the large scale historic deforestation in these regions, with afforestation reported in Europe, USA, China, Vietnam and India (FAO 2013).

Changes in soil properties can vary markedly with type of land cover change, climate, and method, extent of vegetation removal (e.g. land clearing, fires, mechanical harvest) and management post-harvest. For example, West *et al.* (2010) estimated that clearing land in the tropics generally emits three times the amount of carbon per ton of annual crop production compared to clearing land in temperate areas. Emissions are particularly high when organic peatland/wetland soils are drained to enable agriculture as the initial soil carbon is higher, and drainage results in large losses of carbon as previously anaerobic soils become aerobic, allowing the organic matter to oxidise. For example, clearing forest on organic soils for palm oil production in Kalimantan emits nine-times more carbon than clearing on neighbouring mineral soils (Carlson & Curran, 2013). Impacts of deforestation can be reduced by avoiding deforestation on organic soils, and on steep slopes prone to erosion.

There is large heterogeneity in soil measurements of carbon, nitrogen, microbes etc., and still many areas of the world with poor data coverage. Models can be used to fill gaps in spatial coverage and look at past and future time periods, but these too give very variable results. Nevertheless there are some clear signals that can be obtained from meta-analyses of field data and global model results of land use/land cover change with respect to soil carbon.

2.1. Observations of impacts of land cover change

Table 1 presents the results of different meta-analysis studies across different climatic zones that compared the impacts of land use changes on SOC (Guo & Gifford 2002; Don *et al.* 2011; Poeplau *et al.* 2011; Bárcena *et al.* 2014; Murty *et al.* 2002; Wei *et al.* 2014a). Changes in SOC after the conversion of forests to croplands ranged from -24 to -52% without marked differences between climatic regions. The conversion of pastures to other uses (tree plantations and particularly, croplands) also induced decreases in SOC (-10% and -59%, respectively). On the other hand, the substitution of croplands by other land uses (forest regrowth, tree plantation, grassland, pasture) resulted in an increase of SOC (+18 to +53%). In the case of afforestation, soil C increase with time after afforestation, and C sequestration depends on prior land use, climate and the tree species planted.

Fewer meta-analysis studies are available for changes in soil N with changes in land uses. A compilation with predominance of data from tropical sites indicated that average loss of 15% of soil N after conversion of forests to croplands (Murty *et al.* 2002). In Australia, N losses after conversion of native vegetation to perennial pasture and cropland were more than 20% and 38%, respectively (Dalal *et al.* 2013) while in China N loss (0-10 cm depth) was 21% and 31% after 4 and 50 years after conversion of forests to cropland (Wei *et al.* 2014b). Similarly to what was described for SOC, afforestation in subtropical zone results in a significant increase of N stocks 50 years after conversion (Li *et al.* 2012).

[Table 1 here]

2.2. Modelled impacts of land cover change

Dynamic Global Vegetation Models (DGVMs) are used to look at the combined effects of land use change, climate, CO₂, and in some cases N deposition, on vegetation and soil properties over time. A few global models include some aspects of forest, grassland or cropland management (Bondeau *et al.* 2007; Lindeskog *et al.* 2013; Drewniak *et al.* 2013; Jain *et al.* 2005). Most DGVMs do not currently model peatland soils. In Tables 1 and 2, and Figures 1 and 2, we show impacts of past land cover and management change on soil carbon and nitrogen as calculated by three DGVMs: ISAM (Jain *et al.* 2013; El-Masri *et al.* 2013; Barman *et al.* 2014 a,b); LPJ-GUESS (Smith *et al.* 2001; Lindeskog *et al.* 2013); and LPJmL (Bondeau *et al.* 2007). The ISAM and LPJ-GUESS models were run with the HYDE historical land use change data set (History Database of the Global Environment; Klein

Goldewijk *et al.* 2011). ISAM included wood harvest following (Hurtt *et al.* 2011). The LPJmL group combined 3 land use change data sets with the geographic distribution of global agricultural lands in the year 2000. All models were run with historical climate and CO₂, and additionally N deposition in the ISAM model only as it includes a nitrogen cycle. The effects of land cover change were isolated by comparing model runs with and without land use/management (Le Quéré *et al.* 2014). Table 2 and Figure 1 show the loss of soil carbon due to historical land use change from 1860 to 2010 (note there was land use change causing soil carbon loss prior to 1860 particularly in Europe and central Asia, but results are not shown as they were not available for all three models). As with the observed data (Table 1) high carbon losses are associated with the conversion of forests to croplands. Figure 2 shows the mineral soil C and N concentration of different land cover types in different geographic ranges.

[Figure 1 & 2; Table 2 here]

Differences between the models are large for some systems and regions due to different land use change data, different land cover definitions, and different processes included in the models. For example, soil carbon losses are higher in the LPJmL model (Table 2, Figure 1) in part due to greater land cover change in their land cover reconstructions, while their boreal grassland soil carbon is high due to the inclusion of permafrost slowing soil carbon decomposition (Figure 2). Treatment of management processes turns out to be an important differentiator. ISAM shows strong decreases of soil carbon in some regions e.g. the southern Boreal zone (Figure 1) where the inclusion of wood harvest removes carbon and nutrients from the soil, while increases in soil carbon in parts of the mid.-latitudes are due to regrowth of forest following abandonment of agricultural land.

In semi-arid to arid regions, LPJ-GUESS and LPJmL show opposite signs of soil carbon change after conversion of natural land to pastures (Figure 1), primarily because LPJ-GUESS simulates a greater fraction of woody vegetation than LPJmL in these regions under potential natural vegetation. Conversion of woody vegetation to pasture slightly increases soil carbon (see the meta analysis of Guo & Gifford 2002), partly because of boosted productivity and higher turnover rates adding more C to the soil, while the change from potential natural grassland to managed pasture (for which the literature is sparse) results in a soil carbon

decrease in LPJmL. Pasture management strategies can have a large influence on the soil carbon storage (see Section 4.3), and may also be partly be responsible for differences.

Vegetation models are embedded in Earth System Models (ESMs) used to project future climates under different human activity including different land management. Some significant differences between future model climate projections stem from the differences in modeling soil carbon, in particular, the strength of the relationship between increasing temperatures and the increasing rate of soil carbon decomposition (Q_{10}) causing climate-carbon feedbacks *via* CO₂ emissions (Friedlingstein *et al.* 2006). A recent intercomparison of 11 ESMs used in the IPCC 5th Assessment Report (Todd-Brown *et al.* 2013), found the estimate of global soil carbon from ESMs ranged from 510 to 3040 PgC across 11 ESMs compared to an estimate of 890-1600 PgC (95% confidence interval) from the Harmonized World Soil Data Base (FAO/IIASA/ISRIC/ISSCAS/JRC, 2012), with all models having difficulty representing the spatial variability of soil carbon at smaller (1 degree) scales compared to empirical data. In all models NPP and temperature strongly influenced soil carbon stocks, much more so than in the observational data, and differences between models was found to be largely due to the representation of NPP and the parameterization of soil decomposition sub-models. A similar, systematic analysis of DGVMs including benchmarking with observational data, and careful testing of assumptions and process representations in these models, making use of the very large number of observations that have become available in the years since these algorithms were formulated (e.g. Medlyn *et al.* 2015), could significantly improve model performance. This, along with better representation of critical biological and geochemical mechanisms would improve model capability (Todd-Brown *et al.* 2013).

2.3 Drainage and conversion of peatlands/wetlands for agriculture

The organic soils in peatlands/wetlands store vast quantities of carbon which decomposes rapidly when they are drained for agriculture or commercial forestry, resulting in emissions of CO₂ and N₂O to the atmosphere (Hooijer *et al.*, 2010). Other services, in particular water storage and biodiversity, are negatively impacted. Drainage increases vulnerability to further losses through fire. The majority of soil carbon is concentrated in peatlands in the boreal zone and tropical peatland forests in Southeast Asia. These areas, along with wetlands along the banks of rivers, lakes and estuaries have increasingly been developed for croplands/bioenergy

production over recent decades. The FAO emissions database estimates that globally there are 250 000 km² of drained organic soils under cropland and grassland, with total GHG emissions (N₂O plus CO₂) of 0.9 Pg CO₂eq yr⁻¹ in 2010, with the largest contributions from Asia (0.44 Pg CO₂eq yr⁻¹) and Europe (0.18 Pg CO₂eq yr⁻¹; FAOSTAT, 2013; Tubiello *et al.*, 2015). Joosten (2010) estimated that there are >500 000 km² of drained peatlands in the world, including under forests, with CO₂ emissions having increased from 1.06 Pg CO₂ yr⁻¹ in 1990 to 1.30 Pg CO₂ yr⁻¹ in 2008, despite a decreasing trend in developed countries, from 0.65 to 0.49 Pg CO₂ yr⁻¹, primarily due to natural and artificial rewetting of peatlands. In Southeast Asia, CO₂ emissions from drained peatlands in 2006 were 0.61 ± 0.25 Pg CO₂ yr⁻¹ (Hooijer *et al.*, 2010). Conversion of peatlands in Southeast Asia is increasing, particularly for oil palm expansion, where cleared peatlands typically emit ~9 times more carbon than neighbouring mineral soils (Carlson & Curran 2013). In China, between 1950 and 2000, 13 000 km² of wetland soils were shifted to cultivated arable lands, which led to a SOC loss of 5.5 Pg CO₂, mostly from peatlands in Northeast China and Tibet (Zhang *et al.*, 2008).

Soil drainage also affects mineral soils. Meersmans *et al.* (2009) showed that initially poorly drained valley soils in Belgium have lost significant amount of topsoil SOC (i.e. between ~70 and 150 t CO₂ ha⁻¹ over the 1960 – 2006 period), most probably as a consequence of intensified soil drainage practices for cultivation purposes.

3. Agricultural management

To meet projected increases in food demand, crop production will need to increase by 70–110% by 2050 (World Bank, 2008; Royal Society of London, 2009; Tilman *et al.*, 2011). This can be achieved either through further expansion of agricultural land (extensification), or through intensification of production on existing land. Intensification is widely promoted as the more sustainable option because of the negative environmental consequences of land expansion through deforestation and wetland cultivation (Foley *et al.*, 2011). For example, Burney *et al.* (2010) estimate that intensification of production on croplands between 1961 and 2010 avoided the release of 590 PgCO₂eq. Increased productivity per unit land area can be achieved through a variety of management practices, such as fertilization, irrigation and increased livestock density, but these can lead to adverse consequences for the soil and wider environment (Tilman *et al.*, 2002). Intensifying land use can potentially reduce soil fertility (without additional inputs) and its ability to sustain high production, as well as soil resilience

to extreme weather under climate change, pests and biological invasion, environmental pollutants and other pressures. Some key management practices and consequences are highlighted below and summarised in Table 3.

[Table 3 here]

3.1 Nutrient management

Cultivation of soils results in a decline in soil nutrients (nutrient mining). Nutrient inputs, from both natural and synthetic sources, are needed to sustain soil fertility and supply nutrient requirements for crop production. Nutrient supply can improve plant growth which increases organic matter returns to the soil, which in turn can improve soil quality (see section 3.5), so balanced nutrient supply has a positive impact on soils (Smith *et al.*, 2015). Overuse, however, has negative environmental consequences. Annual global flows of nitrogen and phosphorus are now more than double natural levels (Matson *et al.*, 1997; Smil, 2000; Tilman *et al.*, 2002). In China, for example, N input in agriculture in the 2000's was twice that in 1980's (State Bureau of Statistics-China, 2005).

Between 50-60% of nutrient inputs remain in agricultural soils after harvest (West *et al.*, 2014) and can enter local, regional, and coastal waters becoming a major source of pollution such as eutrophication leading to algal blooms (Carpenter *et al.*, 1998). In many places around the world, over-use of synthetic nitrogen fertilizers is causing soil acidification and increased decomposition of soil organic matter, leading to loss of soil function in over-fertilized soils (Ju *et al.*, 2009; Tian *et al.*, 2012).

Use of fertilisers and manures contributes to climate change through their energy intensive production and inefficient use (Tubiello *et al.*, 2015). Globally, approximately 3-5% of nitrogen additions are released as nitrous oxide (N₂O) to the atmosphere when both direct (from soils) and indirect (e.g. downstream from nitrate leaching) emissions are considered (Galloway *et al.*, 2004), and N₂O has ~300 times the radiative forcing of carbon dioxide (IPCC, 2007). Recent research indicates that the relationship between nitrogen application and N₂O emissions is non-linear, resulting in an increasing proportion of added N being emitted, as application rate increases (Philibert *et al.*, 2013; Shcherbak *et al.*, 2014). China,

India, and the United States account for ~56% of all N₂O emissions from croplands, with 28% from China alone (West *et al.*, 2014). Overuse of nitrogen and phosphorus fertilizer can contribute to eutrophication of water bodies, adversely affecting water quality and biodiversity (Galloway *et al.*, 2003, 2004, 2008).

Nutrient use-efficiency can be significantly increased, and nitrate losses to water and N₂O emissions can be reduced, through changes in rate, timing, placement, and type of application, as well as balancing fertilization (Venterea *et al.*, 2012; Snyder *et al.*, 2014). It has been estimated that current levels of global cereal production could be maintained while decreasing global nitrogen application by 50% (Mueller *et al.*, 2014).

3.2 Carbon management: reduced disturbance and organic matter additions

Agricultural soils have the potential to store additional carbon than at present if best management practices are used (Paustian *et al.*, 1997; Smith, 2008; Smith, 2012). Soil organic matter content of soils can be increased through use of improved crop varieties or grassland species mixtures with greater root mass or deeper roots (Kell, 2012), improved crop rotations in which C inputs are increased over a rotation (Burney *et al.*, 2010), greater residue retention (Wilhelm *et al.*, 2004), and use of cover crops during fallow periods to provide year-round C inputs (Burney *et al.*, 2010; Poeplau & Don, 2015). Several studies report that soil carbon increases in croplands under no-till management (West & Post, 2002; Ogle *et al.*, 2005). However, the carbon benefits of no-till may be limited to the top 30cm of soil (Blanco-Canqui & Lal 2008; Powlson *et al.*, 2014). Baker *et al.* (2007) found that total soil carbon was similar in non-till and conventional systems, suggesting that carbon accumulation is occurring at different depths in the soil profile under different management schemes. Given the larger variability in sub-surface horizons and lack of statistical power in most studies, more research is needed on soil carbon accumulation at depth under different tillage regimes (Kravchenko & Robertson, 2010).

Adding plant-derived carbon from external sources such as composts and biochar can increase soil carbon stocks. Composts and biochars are more slowly decomposed compared to fresh plant residues, with mean residence times several (composts) to 10-100 (biochars) longer than un-composted organic materials (Ryals *et al.*, 2015; Lehmann *et al.*, 2015). Recent developments suggest that biochar, from the pyrolysis of crop residues or other

biomass, can consistently increase crop N use efficiency while greatly (over 25%) reducing direct N₂O emissions from N fertilizers (Liu *et al.*, 2012; Huang *et al.*, 2012), as well as enhancing soil fertility (Woolf *et al.*, 2010).

3.3 Water management

The amount of irrigated croplands has doubled in the last 50 years and now accounts for 70% of all water use on the planet (Gleick, 2003). While irrigated crops cover 24% of all cropland area, they account for 34% of all production (Siebert & Döll, 2010). Irrigation is concentrated in precipitation-limited areas such as India, China, Pakistan, and the USA, which account for 72% of irrigation water use (West *et al.*, 2014). Agricultural water-use competes with uses for human and natural ecosystems exacerbating water stress in dry regions. Increased irrigation has occurred in many areas of world agriculture due to the increasing frequency of drought under the climate change (West *et al.*, 2014). Where irrigation increases productivity (e.g. in drought prone areas), organic carbon inputs to the soils would be expected to increase, increasing soil organic matter content (section 3.2).

Irrigation can increase soil salinity in dry regions with high salt content in the subsoil (Ghassemi *et al.*, 1995; Setia *et al.*, 2011). Where salinization occurs, additional irrigation is needed to “flush” the salts beyond the root zone of the crops, which can further exacerbate stress on water resources, particularly when using underground water sources. Saline soils, which have a high concentration of soluble salts, occupy approximately 3.1% (397 Mha) of the world’s land area (FAO, 1995). Climate change (need for more frequent irrigation) and increases in human population (increasing demand for more production) are likely to increase the extent of saline soils (Rengasamy, 2008). The energy required by plants or soil organisms to withdraw water from the soil or retain it in cells increases with decreasing osmotic potential. As soils dry out, the salt concentration in the soil solution increases (decreasing osmotic potential), so two soils of different texture may have the same electrical conductivity, but the osmotic potential is lower in the soil with low water content (Setia *et al.*, 2011a; Chowdhury *et al.*, 2011; Ben-Gal *et al.*, 2009). The accumulation of salts in the root zone has adverse effects on plant growth activity, not only due to negative osmotic potential of the soil solution resulting in decreased availability of water to plants, but also ion imbalance and specific ion toxicity (Chowdhury *et al.*, 2011). Salinity affects microorganisms mainly by decreasing osmotic potential, which affects a wide variety of metabolic activities and alters

the composition and activity of the microbial community (Chowdhury *et al.*, 2011) and thereby soil organic matter decomposition.

In saline soils, SOC content is influenced by two opposing factors: reduced plant inputs which may decrease SOC, and reduced rates of decomposition (and associated mineralisation of organic C to CO₂) which could increase SOC content if the C input were unchanged. Using a modified Rothamsted Carbon model (RothC) with a newly-introduced salinity decomposition rate modifier and a plant input modifier (Setia *et al.*, 2011b, 2012), Setia *et al.* (2013) estimated that, historically, world soils that are currently saline have lost an average of 3.47 t SOC ha⁻¹ since they became saline. With the extent of saline soils predicted to increase under the future climate, Setia *et al.* (2013) estimated that world soils may lose 6.8 Pg SOC due to salinity by the year 2100. Soil salinization is difficult to reverse, but salt tolerant plant species could be used to rehabilitate salt affected soils (Setia *et al.*, 2013).

Water efficiency can be improved through management practices that reduce water requirement and evaporation from the soil (such as adding mulch as groundcover), more precise irrigation scheduling and rates, fixing leaks in dryland irrigation systems, improved application technology (e.g., drip irrigation) and use of intermittent irrigation in rice paddies. Given that water limitation is projected to become even more limiting in several semi-arid regions, e.g. Sub-Saharan Africa, where the human population will probably increase most in the future, and climate change impacts are projected to be severe, improved water harvesting methods, e.g. storage systems, terracing and other methods for collecting and storing runoff, are required to make best use of the limited water resource.

3.4 Harvest frequency

Approximately 9% of crop production increases from 1961-2007 was from increasing the harvest frequency (Alexandratos & Bruinsma, 2012). The global harvested area (i.e. counting each time an area is harvested) increased four times faster than total cropland area between 2000 and 2011 (Ray & Foley, 2013). The fraction of net primary production (NPP) extracted by humans is increasing (Haberl *et al.*, 2007). Global warming is increasing the total area suitable for double or even triple cropping in subtropical and warm temperate regions (Liu *et al.*, 2013). The increase results from fewer crop failures, fewer fallow years, and an increase in multi-cropping.

Increasing harvest frequency can reduce soil quality by e.g. continuously removing soil nutrients and increasing soil compaction through greater soil traffic, but if legumes are included in rotations as harvest frequency increases, soil quality could be improved. Increasing harvest frequency may require increasing pesticide and herbicide use, and increased use of fertilisers contributing to pollution (section 3.1). The net effect will depend on the effectiveness of the management practices followed.

3.5 Soil compaction

Soil compaction causes degradation of soil structure by increasing soil bulk density or decreasing porosity through externally or internally applied loads, as air is displaced from the pores between the soil grains (McCarthy, 2007; Alakukku, 2012). It is the most important subtype of physical soil deterioration, covering 68 Mha globally when first mapped in the 1990s (Oldeman et al., 1991). Compaction of agricultural soils often results from heavy machinery or from animal trampling, so is more likely to occur in intensive agricultural systems (machinery use and high stocking densities), and affects physical, chemical and biological properties of soil. Top soil compaction can be reversed and controlled, but when compaction creates impermeable layers in the subsoil, this is less easily reversed.

Subsoil compaction can disrupt nutrient water flows, which in turn can lead to reduced crop yields, poorer crop quality and can give rise to increased GHG emissions, water and nutrient run-off, erosion, reduced biodiversity and reduced groundwater recharge (Batey, 2009). Where compaction cannot be avoided, mitigation is necessary. Biological approaches to mitigation include planting deep rooted plants such as agroforestry; chemical methods include fertilization (to overcome yield penalty, though not to remedy compaction); and technical measures include machinery in which planting does not coincide with wheel tracks, wide tyres / reduced tyre pressures to reduce pressure per unit area, and precision farming to retain the same wheel tracks each year (Hamza & Anderson, 2005).

3.6 Livestock density

Livestock production is projected to increase significantly in order to meet the growing demand from a growing population and increase in per-capita meat consumption, with total

demand for meat expected to grow by more than 200 Mt by 2050 (Alexandratos & Bruinsma, 2012). The greatest increases in per-capita consumption are projected to be in developing and transition countries (Alexandratos & Bruinsma, 2012). Since the 1970s, most increased livestock production has resulted from intensification: increasing livestock density and shifting to a greater fraction of livestock raised in industrial conditions (Bouwmann *et al.*, 2006). For example, 76-79% of pork and poultry production is industrialized (Herrero & Thornton, 2013). Manure, inputs for growing feed, and soil loss from intensively managed areas can be major sources of water pollution to local and downstream freshwater ecosystems. Clearing natural ecosystems for new pastures, particularly in arid and semi-arid regions, typically occurs on low-productivity lands with a much higher risk of soil erosion and soil carbon/nutrient depletion (Alexandratos & Bruinsma, 2012), and negatively impacts water storage and biodiversity. The impacts of livestock production are particularly prevalent for beef production, which has a least an order of magnitude greater impact on land, water, GHGs, and reactive nitrogen compared to other livestock (Eshel *et al.*, 2014; Ripple *et al.*, 2014). Moreover, industrial livestock production had led to an increased use of veterinary medicines, antibiotics and hormones, posing potential risks to soil, water, ecosystems and human health. Improved grazing management (e.g. optimised stocking density) can reduce soil degradation, and thereby maintain and enhance organic matter content (McSherry & Ritchie, 2013; see sections 3.2 and 4.3), and can reduce soil compaction, thereby increasing infiltration and water storage and reduce risk of runoff and flooding downstream (Marshall *et al.*, 2009).

4. Other land management

4.1 Forest management

Logging and fire are the major causes of forest degradation in the tropics (Bryan *et al.*, 2013). Logging removes nutrients and negatively affects soil physical properties and nutrient levels (soil and litter) in tropical (e.g. Olander *et al.*, 2005; Villela *et al.*, 2006; Alexander, 2012) and temperate forests (Perez *et al.*, 2009). Forest Fires affect many physical, chemical, mineralogical, and biological soil properties, depending on fire regime (Certini, 2005). Increased frequency of fires contributes to degradation, and reduces the resilience of the biomes to natural disturbances. A meta-analysis of 57 publications (Nave *et al.*, 2011) showed that fire caused a significant decrease in soil C (-26%) and N (-22%). Fires reduced

forest floor storage (pool sizes only) by an average of 59% (C) and 50% (N), but the relative concentrations of these two elements did not change. Prescribed fires caused smaller reductions in C and N storage (-46% and -35%) than wildfires (-67% and -69%). These differences are likely because of lower fuel loads or less extreme weather conditions in prescribed fires, both factors that result in lower fire intensity. Burned forest floors recovered their C and N pools in an average of 128 and 103 years, respectively. Among mineral soil layers, there were no significant changes in C or N storage, but C and N concentrations declined significantly (-11% and -12%, respectively). Mineral soil C and N concentrations were significantly reduced in response to wildfires, but not after prescribed burning.

Forest fires produce charcoal, or black carbon, some of which can be preserved over centuries and millennia in soils. Dissolved black carbon (DBC) from burning of the Brazilian Atlantic forest continued to be mobilized from the watershed each year in the rainy season, despite the fact that widespread forest burning ceased in 1973 (Dittmar *et al.*, 2012).

A large field study in the Amazon (225 forest plots) on the effects of anthropogenic forest disturbance (selective logging, fire, and fragmentation) on soil carbon pools showed that the first 30 cm of the soil pool did not differ between disturbed primary forests and undisturbed areas of forest, suggesting a resistance to impacts from selective logging and understory fires (Berenguer *et al.*, 2014). As with deforestation, impacts of human disturbances on the soil carbon are of particular concern in tropical forests located on organic soils and on steep easily-eroded slopes.

4.2 Shifting cultivation

Shifting cultivation practices, where land is cleared through fire, have been practiced for thousands of years, but recent increasing demographic pressure has reduced the duration of the fallow period, affecting the system sustainability. Moreover, especially in Southeast Asia where urbanisation is expanding in fertile planes, shifting cultivation is practiced in sloping uplands, which are prone to soil and carbon loss by erosion (Chaplot *et al.*, 2005). A review by Ribeiro Filho *et al.* (2013) reported negative impact on SOC associated with the conversion stage, modified by the characteristics of the burning. Chop-and-mulch of enriched fallows appears to be a promising alternative to slash-and-burn, conserving soil bulk density, and significantly increasing nutrient concentrations and organic matter content compared to

burnt cropland, and a control forest in a study in the Amazon (Comtea *et al.*, 2012).

4.3 Grassland management and dryland degradation

Grasslands, including rangelands, shrublands, pastureland, and cropland sown with pasture and fodder crops, cover 26% of the global ice-free land area and 70% of the agricultural area, and contain about 20% of the world's soil organic carbon (C) stocks. Grasslands on every continent have been degraded due to human activities, with about 7.5% of grassland having been degraded because of overgrazing (Conant, 2012). A meta-analysis (McSherry & Ritchie, 2013) of grazer effects on SOC density (17 studies that include grazed and un-grazed plots) found higher grazing intensity was associated with increased SOC in grasslands dominated by C4 grasses (increase of SOC by 6–7%), but with lower SOC in grasslands dominated by C3 grasses (decrease of SOC by an average 18%). An increase in mean annual precipitation of 600 mm resulted in a 24% *decrease* in the magnitude of the grazer effect on finer textured soils, but on sandy soils the same increase in precipitation produced a 22% *increase* in the grazer effect on SOC (McSherry & Ritchie, 2013).

Land use dynamics and climate change are the major drivers of dryland degradation with important feedbacks, with changes in plant community composition (e.g. shrub encroachment and decrease in vegetation cover; D'Odorico *et al.*, 2013). A review by Ravi *et al.* (2010) indicated soil erosion as the most widespread form of land degradation in drylands, with wind and water erosion contributing to 87% of the degraded land. Grazing pressure, loss of vegetation cover, and the lack of adequate soil conservation practices increase the susceptibility of these soils to erosion. The degree of plant cover is negatively related to aridity, and an analysis of 224 dryland sites (Delgado-Baquerizo *et al.*, 2013) highlighted a negative effect of aridity on the concentration of soil organic C and total N, but a positive effect on the concentration of inorganic P, possibly indicating the dominance of physical processes such as rock weathering, a major source of P to ecosystems, over biological processes that provide more C and N through litter decomposition (Delgado-Baquerizo *et al.*, 2013).

Soil carbon dynamics in pastures strongly depend on management, with soil carbon increases or decreases observed for different combinations of animal densities and grazing frequency (Conant 2012; Machmuller *et al.* 2015). Different grazing strategies, especially in the semi-

natural dryland biomes, have large implications for vegetation and the carbon balance (Yates *et al.* 2000). Under certain conditions, grazing can lead to increased annual net primary production over un-grazed areas, particularly with moderate grazing in areas with a long evolutionary history of grazing and low primary production but this does not always lead to an increase in soil carbon (e.g. Badini *et al.* 2007); grazing, like crop harvest, fundamentally leads to the rapid oxidation of carbon that would otherwise be eventually transferred to the soil. It has long been recognised that the potential effects of management on carbon storage in grassland and dryland soils are substantially greater than that of climate change or CO₂ enhancement (Ojima *et al.* 1993), and Henderson *et al.* (2015) estimated that the optimization of grazing pressure could sequester 148 Tg CO₂ yr⁻¹.

4.4 Artificial surfaces, urbanisation and soil sealing

In 2014, 54% of the world's population was urban, and by 2050, two thirds of the global population will be urban. Many regions in the world, (such as Europe and Asia) are affected by migration of populations from rural area to large megacities. Africa and Asia have more rural populations, but are urbanizing faster than the other regions (World Urbanization Prospects, 2014). With urbanization comes land take (development of scattered settlements in rural areas, the expansion of urban areas around an urban nucleus, and densification on land within an urban area) and soil sealing. Soil sealing refers to the permanent covering of an area of land and its soil by impermeable artificial material (e.g. asphalt and concrete), for example through buildings and roads. The area actually sealed is only part of a settlement area, and gardens, urban parks and other green spaces are not covered by an impervious surface (Prokop *et al.*, 2011).

Sealing by its nature has a major effect on soil, diminishing many of its benefits (Tóth *et al.*, 2007). It is normal practice to remove the upper layer of topsoil, which delivers most of the soil-related ecosystem services, and to develop a strong foundation in the subsoil and/or underlying rock to support the building or infrastructure. Loss of ecosystem and social services (mainly on high-quality soils) include impacts on water resources (e.g. reduction of rainfall absorbed by the soil, reduction of soil water holding capacity affecting flooding), on soil biodiversity when sealing prevents recycling of dead organic material (Marfenina *et al.* 2008), on the carbon cycle due to topsoil and vegetation removal (Davies *et al.*, 2011).

Sealing through expansion of urban areas can also lead to agricultural land becoming more

marginal since the best agricultural land around settlements is lost as they expand, with agricultural land displaced to more marginal land.

Appropriate mitigation measures can be taken in order to maintain some of the soil functions. In urban planning management, objectives to reduce the impact of soil sealing include: i) preventing the conversion of green areas, ii) re-use of already built-up areas (e.g. brownfield sites Meuser, 2010; Hester & Harrison, 2001 – though remediation of contaminated sites can be costly; Maderova & Paton, 2013), iii) using (where appropriate) permeable cover materials instead of concrete or asphalt supporting green infrastructure, and iv) implementation of compensation measures. In order to deliver this mitigation a number of actions are necessary, e.g. reduction of subsidies that act as drivers for unsustainable land take and soil sealing (Prokop *et al.*, 2011), and strong collaboration between relevant public authorities and governance entities (Siebielec *et al.*, 2010). Development impacts can be reduced by inclusion of green infrastructure, a network of high-quality green spaces and other environmental features that have a positive effect on well-being (Gill *et al.*, 2007) as well as soils. In some regions, urban sprawl is exacerbated by insufficient incentives to re-use brownfield (derelict, underused or abandoned former industrial or commercial) sites, putting increasing pressure on greenfield land take.

Actions to alleviate pressures on soils driven by sealing fall into three categories: limiting, mitigating and compensating. Actions to limit soil sealing centre around reduction of land take through development of spatial urban planning and environmental protection. Mitigation of soil sealing entails use of strategic environmental assessment for plans and programmes, use of permeable materials and surfaces, green infrastructure within built and urban environments, and natural water harvesting. Compensating soil sealing entails reclamation of degraded land, re-use of extracted topsoil, de-sealing and is incentivised by land take fees and application of environmental cost calculations.

5. Anthropogenic environmental change pressures that interact with land management pressures on soils

In addition to the direct impacts of humans on soils *via* land use change and land management, anthropogenic activity has indirect impacts through human-induced environmental change, such as pollution and climate change. These interact with land

management. Soils provide a temporary but labile store for pollutants (Meuser, 2010). Natural processes can release pollutants back to the atmosphere, make them available to be taken up by plants and organisms, leached in to surface waters (Galloway *et al.*, 2008) and/or transported to other areas by soil erosion (Ravi *et al.*, 2010). Pollutants disrupt natural biogeochemical cycles by altering both soil quality and function through direct changes to the nutrient status, acidity and bioavailability of toxic substances and also by indirect changes to soil biodiversity, plant uptake and litter inputs (EEA, 2014). Soil sensitivity to atmospheric pollution varies with respect to key properties influenced by geology (cation exchange capacity, soil base saturation, aluminium), organic matter, carbon to nitrogen ratio (C:N) and water table elevation (EEA, 2014).

Atmospheric pollutant deposition impacts on soils vary with respect to soil sensitivity to a specific pollutant and the actual pollutant load. Sulphur, nitrogen and heavy metals are released in to the atmosphere by fossil fuel combustion (e.g. power generation, industry and transport) and non-combustion processes (e.g. agricultural fertilizers, waste). These pollutants are transported off-site and deposited as either dry or wet deposition, which can cross national borders. Deposition is enhanced in forests and with altitude because of reduced wind speeds and greater precipitation, respectively, impacting remote areas. Harmful effects to soil function and structure occur where deposition exceeds the ‘critical load’ that a particular soil can buffer (Nilsson & Grennfelt, 1988). Spatial differences in soil sensitivity (commonly defined by the ‘critical load’) and pollutant deposition result in an uneven global distribution of impacted soils (Figure 3). For instance, global emissions of sulphur and nitrogen have increased 3-10 fold since the pre-industrial period (van Aardenne *et al.*, 2001), yet only 17% of the global land area sensitive to acidification is in a region where deposition exceeds the critical load (Bouwman *et al.*, 2002).

Emissions of pollutants, notably sulphur, across Europe and North America have declined since the 1980s following protocols established under the 1979 Convention on Long-range Transboundary Air Pollution (LRTAP) and the 1990 US Clean Air Act Amendments (CAAA) (Greaver *et al.*, 2012; Reis *et al.*, 2012; EEA, 2014). Conversely, emissions are likely to increase in response to industrial and agricultural development in south and east Asia, sub-Saharan Africa and South America (Kuylenstierna *et al.*, 2001; Dentener *et al.*,

2006). Further emission increases are occurring in remote areas due to mining activity, such as oil sand extraction in Canada (Kelly *et al.*, 2010; Whitfield *et al.*, 2010).

5.1 Sulphur deposition

Sulphur emissions are primarily from combustion of coal and oil, typically associated with power generation and heavy industry. In 2001, regions with deposition in excess of 20 kg S ha⁻¹ yr⁻¹ where China and Republic of Korea, western Europe and eastern North America (Vet *et al.*, 2014; Figure 3a). Deposition in un-impacted areas is <1 kg S ha⁻¹ yr⁻¹ (Figure 3a). Pollution control measures have seen an 80% reduction in pollutant sulphur deposition across Europe between 1990 and 2010 (Reis *et al.*, 2012), and emissions in China have declined since 2005 (Fang *et al.*, 2013).

Soil acidification is a natural process that is altered and accelerated by sulphur and nitrogen deposition (Greaver *et al.*, 2012). Sulphur oxides (SO_x) react with water to form sulphuric acid (H₂SO₄). Excess inputs of acidity (H⁺) displace soil base cations (e.g. calcium (Ca²⁺) and magnesium (Mg²⁺)) from soil surfaces into solution, which are subsequently lost by leaching (Reuss & Johnson, 1986). Mineral soils can buffer base cation losses if inputs from rock weathering and/or atmospheric dust deposition exceed the amount lost. Therefore, the global distribution of acid sensitive soils is associated with conditions that favour development of soils with low cation exchange capacity and base saturation (Bouwman *et al.*, 2002; Figure 3c). Wetland can also buffer inputs of acidity through biological sulphate reduction, although acidity can be mobilised again following drought and drainage (Tipping *et al.*, 2003; Laudon *et al.*, 2004; Daniels *et al.*, 2008). Organic acids can also buffer mineral acidity in naturally acidic organic soils (Krug and Frink, 1983).

Decreased soil fertility or 'sterilisation' due to loss of nutrients and mobilisation of toxic metals, particularly Al, are caused by acidification. Impacts in Scandinavia over the 1960s-80s included declines in freshwater fish populations and damage to forests (EEA, 2014). Sulphur can also stimulate microbial processes that make mercury bioavailable, leading to bioaccumulation in the food chain (Greaver *et al.*, 2012). In agricultural soils in Europe, however, fertilizer inputs of sulphur have increased to combat crop sulphur deficiencies as a result of sulphur emission controls (Bender & Weigel, 2011).

Acidification is reversible, evident by increases in soil pH following decreased sulphur emissions, although the recovery time varies; some areas with organic soils where deposition has declined are showing either slow or no recovery (Greaver *et al.*, 2012; Lawrence *et al.*, 2012; RoTAP, 2012). On agricultural soils, lime can be applied to increase soil pH. However, 50-80% of sulphur deposition on land is on natural, non-agricultural land (Dentener *et al.*, 2006). Application of lime to naturally acidic forest soils can cause further acidification of deep soil layers whilst increasing decomposition in surface litter, with no improvement in tree growth (Lundström *et al.*, 2003).

Wider effects of acidification are starting to be understood through long-term monitoring. Decreased organic matter decomposition due to acidification has increased soil carbon storage in tropical forests (Lu *et al.*, 2014). However, in temperate forest soils acidification can lead to reduced C:N ratios in soil, which in turn increases nitrification (Aber *et al.*, 2003), but on already acidic soils reduces nitrification. In wetland soils, methane (CH₄) emissions have also been suppressed by sulphur deposition (Gauci *et al.*, 2004). Conversely, declining sulphur deposition has been associated with increased dissolved organic carbon fluxes from organic soils (Monteith *et al.*, 2007), and decreased soil carbon stocks in temperate forest soils (Oulehle *et al.*, 2011; Lawrence *et al.*, 2012).

5.2 Nitrogen deposition

Nitrogen deposition covers a wider geographical area than sulphur, as the sources are more varied, including extensive agriculture fertilizer application, ammonia derived from livestock operations, and biomass burning in addition to fossil fuel combustion (Figure 3b). Regions with deposition in excess of 20 kg N ha⁻¹ yr⁻¹ in 2001 were western Europe, South Asia (Pakistan, India, Bangladesh) and eastern China (Vet *et al.*, 2014); although extensive areas with 4 kg N ha⁻¹ yr⁻¹ were found across North, Central and South America, Europe and Sub-Saharan Africa. By contrast, 'natural' deposition in un-impacted areas is around 0.5 kg N ha⁻¹ yr⁻¹ (Dentener *et al.*, 2006). While emissions related to fossil fuel combustion have declined along with sulphur across Europe, agricultural sources of nitrogen are likely to stay constant in the near future across Europe (EEA, 2014), whilst overall global emissions are likely to increase (Galloway *et al.*, 2008). Nitrogen deposition in China's industrialized and intensively managed agricultural areas in the 2000s was similar to peaks in Western Europe during the 1980s before mitigation (Liu *et al.*, 2013).

Deposition of nitrogen induces a ‘cascade’ of environmental problems, including both acidification and eutrophication that can have both positive and negative effects on ecosystem services (Galloway *et al.*, 2003). Excluding agricultural areas where nitrogen is beneficial, 11% of land surface received nitrogen deposition above 10 kg N ha⁻¹ yr⁻¹ (Dentener *et al.*, 2006; Bouwman *et al.* 2002; Figure 3d). In Europe, eutrophication has and will continue to impact a larger area than acidification (EEA, 2014).

Nitrogen fertilisation can increase tree growth (Magnani *et al.*, 2007) and cause changes in plant species and diversity (Bobbink *et al.*, 2010), which in turn will alter the amount and quality of litter inputs in to soils, notably the C:N ratio and soil-root interactions (RoTAP, 2012). However, increased carbon sequestration (Reay *et al.*, 2008) may be offset by increased emissions of the greenhouse gases N₂O and CH₄ (Liu & Greaver, 2009). Long-term changes caused by nitrogen deposition are uncertain as transport times vary between environmental systems; and the only way to remove excess nitrogen is to convert it to an unreactive gas (Galloway *et al.*, 2008).

[Figure 3 here]

5.3 Heavy metal deposition

Heavy metal emissions are associated with coal combustion and heavy industry. In the UK, deposition is responsible for 25-85% of inputs to UK soils (Nicholson *et al.*, 2003). In Europe, the areas at risk from cadmium, mercury and lead deposition in 2000 were 0.34%, 77% and 42% respectively, although emissions are declining (Hettelingh *et al.*, 2006). Tighter legislation to control industrial emissions of heavy metals are helping to reduce the environmental load of heavy metals in many regions, though rapid industrial growth in some regions such as East Asia is increasing pressures on soil from heavy metal deposition. Global heavy metal emissions and deposition are poorly understood in comparison to sulphur and nitrogen; although the on-site impact of heavy metal contamination has been well studied (Guo *et al.*, 2014). Metals in bioavailable form have toxic effects on soil organisms and plants, influencing the quality and quantity of plant inputs to soils, rate of decomposition and, importantly, can bio-accumulate in the food chain. Some heavy metals will persist for centuries as they are strongly bound to soil organic matter (RoTAP, 2012), although they can

791 be mobilised to bioavailable form following drought-induced acidification, drainage and soil
792 erosion (Tipping *et al.*, 2003; Rothwell *et al.*, 2005).

793
794 Whilst the direct impacts of sulphur, nitrogen and heavy metals on inorganic soil chemical
795 processes are generally well understood, many uncertainties still exist about pollutant impacts
796 on biogeochemical cycling, particularly interactions between organic matter, plants and
797 organisms in natural and semi-natural systems (Greaver *et al.*, 2012). Process understanding
798 is dominated by research in Europe and North America (e.g. Bobbink *et al.*, 2010). Research
799 is needed across Asia, Africa and South and Central America where soil properties and
800 environmental conditions differ. Models need to be developed to examine the combined
801 effects of air pollutants and their interactions with climate change impacts and feedbacks on
802 greenhouse gas balances and carbon storage (Spranger *et al.*, 2008; RoTAP, 2012). Air
803 quality, biodiversity and climate change policies all impact on soils. A more holistic approach
804 to protecting the environment is needed, particularly as some climate change policies (e.g.
805 biomass burning, carbon capture and storage) have potential to impact air quality and,
806 therefore, soil quality (Reis *et al.*, 2012; RoTAP, 2012; Aherne & Posch, 2013).

807
808 Indirect impacts on soils can be addressed largely by preventing the pollution at source, or by
809 mitigating the adverse effects where these have already occurred. Air pollution control on
810 coal burning and increased car and fleet efficiency standards has been effective in reducing
811 sulphur deposition in many areas of the world, particularly in Europe since the 1970s.
812 Substitution of coal with bioenergy might also reduce sulphur emissions, but unless burned
813 cleanly in a controlled way, can also release pollutants to the air. In terms of nitrogen,
814 ammonia abatement techniques when fertilizers are spread (e.g. slurry injection) are helping
815 to reduce N deposition (Sutton *et al.*, 2007).

816 817 **6. Existing policies and practices that alleviate global change pressures on soils** 818 **from land use and management**

819
820 The previous text has highlighted specific anthropogenic activities that exert or alleviate
821 pressures on soils. Actions that alleviate pressures on soils driven by land use change and
822 land management can be broadly divided into three categories, those that:

- 1) Prevent conversion of natural ecosystems to other uses (e.g. protected areas, reduced deforestation, prevention of wetland drainage, intensification rather than extensification);
- 2) Prevent soil degradation (erosion control, fire management, reduced tillage / conservation agriculture, long term fallows, flood protection, use of organic amendments, intercropping, improved rotations); and
- 3) Result in soil / ecosystem restoration (e.g. peatland rewetting, afforestation, re-vegetation on degraded lands, improved grass varieties, appropriate animal stocking densities, bioremediation).

Policies to encourage such actions were recently reviewed by Bustamante *et al.* (2014) and include:

- a) Economic incentives, e.g., special credit lines for low carbon agriculture and forestry practices and projects, payment for ecosystem services (such as carbon storage) and tradable credits such as carbon,
- b) Regulatory approaches, e.g. enforcement of environmental law to protect natural areas, set-aside policies,
- c) Research, development and diffusion investments, e.g. increase of resource use-efficiency, livestock improvement,
- d) Information and certification schemes, e.g. in China, forest certification to promote sustainable forest management, state regulation for protecting mandatory arable lands, protection projects on Tibetan grasslands, a national wetland protection programme, and the “grain for green” programme.

Many of these actions and policies are not directed at soil conservation, but nevertheless have an effect on soil quality. Two of the main pieces of international policy that have served to reduce pressures on soils, directly and indirectly, are the United Nations Convention to Combat Desertification (CCD) and the United Nations Framework Convention on Climate Change (UNFCCC). In general, policies and actions are important at all scales from international conventions to local action, and local activity is encouraged by policies at regional, national and global level. Policies to sustainably increase land productivity, for example, can prevent land use change, and there are various other supporting actions that can help deliver improvements, e.g. agricultural research, technology transfer, knowledge transfer

and improved rural infrastructure. Some examples of policies that impact on land management and soil quality are given below.

6.1 United Nations Framework Convention on Climate Change (UNFCCC) and other climate specific policies

Soil carbon storage and nutrient cycling as climate services are being increasingly recognised e.g. under UNFCCC as part of national reporting and accounting, as part of life-cycle greenhouse gas assessments for biofuels, in various regional initiatives and national efforts. The UNFCCC is an international treaty, which came into force in 1994, setting an overall framework for intergovernmental efforts to tackle the challenge posed by climate change. The requirements for the 196 country Signatories (or 'Parties') to the UNFCCC include adopting national mitigation policies and publishing national inventories of anthropogenic emissions and sinks of greenhouse gases including activities on the land such as afforestation, deforestation, agricultural management and wetland drainage and rewetting. Developed country signatories have legally binding targets under the Kyoto Protocol and can count land based emissions or sinks towards meeting these targets, thus incentivising activities that protect soil carbon. Developing countries currently have voluntary targets and several countries have made pledges that include reduced deforestation (e.g. Brazil and Indonesia) or afforestation (e.g. 400000 km² in China). Under the Clean Development Mechanism (CDM) developed countries can fund projects in developing countries that generate certified emission reduction credits (CERs). China, for example, has the largest number of CERs in the world (IFPRI, 2011). Brazil also has 180 CDM projects, the third largest number of CERs after China and India (Cole & Liverman, 2011). A number of projects in Africa, North America and South Asia have a significant component for soil greenhouse gas emission reduction of soil carbon sequestration, financed through the Verified Carbon Standard or the American Carbon Registry.

As part of negotiations leading to the new climate treaty in Paris in December 2015, all parties will be required to submit INDCs (Intended Nationally determined Contributions). The new treaty will also include provision for REDD+ (reduced Emissions from Deforestation and Degradation, including management of forests and enhancement of forest carbon stocks). This could go some way to protecting forest soils, and negotiations have been intense around methods for monitoring reporting and verification, with key issues such

as permanence (the risk the forest may be lost at a later date due to management or environmental change) and leakage (displacement of land use change to other areas), and how to finance such activities.

6.2 United Nations Convention to Combat Desertification (CCD)

The CCD entered into force in December 1996; today 179 countries acknowledge it as a legally binding framework to tackle land degradation and promote sustainable development in fragile ecosystems. The Global Mechanism was established under the convention to "promote actions leading to the mobilization and channelling of substantial financial resources, including for the transfer of technology, on a grant basis, and/or on concessional or other terms, to affected developing country Parties". In September 2011 the United Nations General Assembly declared a goal of building a world with no land degradation. In October 2011 parties to the CCD issued a declaration calling for zero land degradation and for adopting sustainable land management as a way to achieve sustainable development.

6.3 Millennium Development Goals (MDGs)

Of the eight MDGs (UNDP, 2014a), soil protection is most relevant to the goal to ensure environmental sustainability, since soils are critical in underpinning environmental sustainability (Smith *et al.*, 2015). A complementary MDG, to develop a global partnership for development, will improve the governance structure to deliver soil security. The other MDG to which soils plays a critical contribution is the goal to eradicate extreme poverty and hunger, with the role of soils in supporting food provision critical for the latter part of this MDG (Smith *et al.*, 2015). The MDGs are currently being revisited to set a post-2015 development agenda (UNDP, 2014b), with discussion around the themes of localising the post-2015 development agenda, helping to strengthen capacities and build effective institutions, participatory monitoring for accountability, partnerships with civil society, engaging with the private sector, and culture and development. The key emerging principles from the dialogue are participation, inclusion, and the need for strengthened capacities and partnerships (UNDP, 2014b). It is important that soils play their role in delivering this post-2015 agenda.

6.4 Protected areas and the Convention on Biological Diversity (CBD)

Many measures to protect biodiversity and vulnerable habitats also protect the soils underpinning them, so numerous conservation actions around the world serve to protect soils, even if this was not the primary aim (Smith *et al.*, 2013). Between 1990 and 2010, the amount of forest land designated primarily for the conservation of biological diversity increased by 35 percent, indicating a political commitment to conserve forests. These forests now account for 12 percent of the world’s forests (FAO, 2010). In India, a Supreme Court ruling in 2011 on effective self-governance of “common” or communal land by local communities may help to protect these valuable resources, and thereby the soils that underpin them. Soil biodiversity is known to be important for soil function (Bodelier, 2011), yet it rarely receives the attention enjoyed by larger flora and fauna within the ecosystem.

6.5 Reduced deforestation and forest management

Various actions have been implemented to reduce deforestation (Bustamante *et al.*, 2014), and to reduce the impact of forestry activities, such as reduced impact logging. UNFCCC, carbon markets and other international environmental programs have contributed to global efforts to reduce deforestation in addition to other sustainable natural resource management programs in countries and by industry. For example, zero deforestation commitments made by several companies (many made in 2014), and activities from bodies such as the Roundtable for Sustainable Palm Oil (RSPO) and the Forest Stewardship Council (FCO) certification scheme. Land improvement has increased in East Asia between 1981 and 2006 despite population increase, attributed largely to policies promoting tree planting and forest plantation programs in China and Korea. In Brazil, deforestation was rapidly reduced after national laws and regulations were enacted to protect forests in the 1990s and early 2000s (including the soy moratorium and the forest code), followed up by state and municipal governments setting further by-laws enforcing the deforestation moratorium (Bustamante *et al.*, 2014).

6.6 Agricultural policies and practices

The pressures on soils imposed by land use intensity change can be mitigated by regulation of over-grazing and reduction of over-stocking on grazed grasslands, return of crop residues to the soil, reduced tillage, best management practices, targeted nutrient management and

precision farming on croplands, and wetland / floodplain restoration. These actions have been encouraged by various policies. Some examples include: The EU set-aside programme of the 1990s encouraged less intensive use of agricultural land where production is low and environmental impacts are high. The EU Common Agricultural Policy ties agricultural subsidies to implementation of best management practices and environmental protection, for example through pillar 2 (rural development programmes) providing crop insurance for lower fertilizer application rates; in Africa, policies for integrated land management to help protect vulnerable soils; China's conservation tillage program (2012-2030); the USA Conservation Reserve Program (set aside marginal lands, steep slopes).

7. Conclusion: Keeping soils central to the science and policy agendas

The International Year of Soils in 2015 is an excellent opportunity to raise the profile of soils in the minds of national and international policy makers, land managers, timber and agro-industries, and the public. Ensuring that vulnerable and high environmental value soils (e.g. peatlands) are considered when making policies and decisions about which habitats and ecosystems to convert or to protect, will help to reduce the pressure on soils particularly vulnerable to global change drivers such as land use and land management, and maintain important ecosystem services. This is in part happening with agendas around valuation of ecosystem services and life-cycle assessments of impacts of land use change that include soil carbon. At a time when governments are negotiating a legally binding climate change treaty and making national targets for greenhouse gas reduction, and revisiting the Millennium Development Goals, keeping soil carbon and nitrogen central to land based greenhouse gas monitoring and reporting will maintain awareness with policy makers and industries with emissions reduction targets. Both science and policy agendas are increasingly concerned with long-term food security, ensuring that soils are central to considerations of how to achieve on-going increases in production will enable those increases to be more sustainable into the future.

Research and policy regarding soil quality and sustainability is abundant, but patchy and disjointed. To ensure that soils are protected as part of on-going wider environmental and sustainable production efforts, soils cannot, and should not, be considered in isolation of the ecosystems that they underpin, but the role of soils in supporting ecosystems and natural capital needs greater recognition (Robinson *et al.*, 2013, 2014). This can, in part, be enhanced

through education and awareness-raising which has started during the International Year of the Soils in 2015. The time is ripe to consider a global soil resilience programme, under the auspices of a global body such as the UN or one of its delivery agencies such as the FAO to monitor, recover or sustain soil fertility and function, and to enhance the ecosystem services provided by soils. The lasting legacy of the International Year of Soils in 2015 should be to bring together robust scientific knowledge on the role of soils, and to put soils at the centre of policy supporting environmental protection and sustainable development.

Acknowledgements

The input of PS and PCW contributes to the Belmont Forum/FACCE-JPI funded DEVIL project (NE/M021327/1) and for PS also contributes to the EU FP7 SmartSoil project (Project number: 289694). TAMP acknowledges funding from European Commission's 7th Framework Programme, under Grant Agreement numbers 282672 (EMBRACE) and 603542 (LUC4C). AKJ was supported by NSF (AGS 12-43071), DOE (DE-SC0006706), and NASA (NNX14AD94G).

References

- Aber, J.D., Goodale, C.L., Ollinger, S.V., Smith, M.-L., Magill, A.H., Martin, M.E., Hallett, R.A., Stoddard, J.L. (2003) Is Nitrogen Deposition Altering the Nitrogen Status of Northeastern Forests? *BioScience*, **53**, 375-375.
- Aherne, J., Posch, M. (2013) Impacts of nitrogen and sulphur deposition on forest ecosystem services in Canada. *Current Opinion in Environmental Sustainability*, **5**, 108–115.
- Alakukku, L. (2012) Soil Compaction. In: Jakobsson, C. (Ed.): *Ecosystem Health and Sustainable Agriculture I: Sustainable Agriculture*. Uppsala University. URL: www.balticuniv.uu.se/index.php/component/docman/doc_download/1256-chapter-28-soil-compaction. (Accessed 4th June 2015).
- Alexander, A.B. (2012) Soil compaction on skid trails after selective logging in moist evergreen forest of Ghana. *Agriculture and Biology Journal of North America* doi:10.5251/abjna.2012.3.6.262.264.
- Alexandratos, J., Bruinsma, J. (2012) *World agriculture towards 2030/2050: the 2012 revision*. FAO Report, Rome.
- Badini, O., Stockle, C.O., Jones, J.W., Nelson, R., Kodio, A., Keita, M. (2007) A simulation-based analysis of productivity and soil carbon in response to time-controlled rotational grazing in the West African Sahel region. *Agricultural Systems*, **94**, 87-96

- 1029 Bai, Z.G., Dent, D.L., Olsson, L., Schaepman, M.E. (2008) Global assessment of land
1030 degradation and improvement. 1. Identification by remote sensing. Report 2008/01, ISRIC –
1031 World Soil Information, Wageningen.
- 1032 Baker, J.M., Ochsner, T.E., Venterea, R.T., Griffis, T.J. (2007) Tillage and soil carbon
1033 sequestration - what do we really know? *Agriculture, Ecosystems & Environment*, **118**, 1-5.]
- 1034 Bárcena, T.G., Kiær, L.P., Vesterdal, L., Stefánsdóttir, H.M., Gundersen, P., Sigurdsson,
1035 B.D. (2014) Soil carbon stock change following afforestation in Northern Europe: a meta -
1036 analysis. *Global Change Biology*, **20**, 2393-2405.
- 1037 Barman, R., Jain, A.K., Liang, M. (2014a) Climate-driven uncertainties in terrestrial gross
1038 primary production: a site-level to global scale analysis, *Global Change Biology*, doi:
1039 10.1111/gcb.12474.
- 1040 Barman, R., Jain, A.K., Liang, M. (2014b) Climate-driven uncertainties in terrestrial energy
1041 and water fluxes: a site-level to global scale analysis, *Global Change Biology*, doi:
1042 10.1111/gcb.12473.
- 1043 Batey, T. (2009) Soil compaction and soil management – a review. *Soil Use and*
1044 *Management*, **12**, 335-345.
- 1045 Batjes, N.H. (2012) *ISRIC-WISE derived soil properties on a 5 by 5 arc-minutes global grid*
1046 *(ver. 1.2)*. Wageningen, ISRIC - World Soil Information (www.isric.org). 52pp.
- 1047 Bender, J., Weigel, H.-J. (2011) Changes in atmospheric chemistry and crop health: A
1048 review. *Agronomy for Sustainable Development*, **31**, 81–89.
- 1049 Ben-Gal, A., Borochoy-Neori, H., Yermiyahu, U., Shani, U. (2009) Is osmotic potential a
1050 more appropriate property than electrical conductivity for evaluating whole-plant response to
1051 salinity? *Environmental and experimental Botany*, **65**, 232-237.
- 1052 Berenguer, E., Ferreira, J., Gardner, T.A., Aragão, L.E.O.C., Camargo, P.B., Cerri, C.E.,
1053 Durigan, M., Oliveira Jr., R.C., Vieira, I.C.G., Barlow, J. (2014) A large-scale field
1054 assessment of carbon stocks in human-modified tropical forests. *Global Change Biology*, doi:
1055 10.1111/gcb.12627.
- 1056 Blanco-Canqui, H., Lal, R. (2008) No-tillage and soil-profile carbon sequestration: An on-
1057 farm assessment. *Soil Science Society of America Journal* **72**, 693-701.
- 1058 Bobbink, R., Hicks, K., Galloway, J., Spranger, T., Alkemade, R., Ashmore, M., Bustamante,
1059 M., Cinderby, S., Davidson, E., Dentener, F., Emmett, B., Erisman, J.-W., Fenn, M., Gilliam,
1060 F., Nordin, A., Pardo, L., De Vries, W. (2010) Global assessment of nitrogen deposition
1061 effects on terrestrial plant diversity: a synthesis. *Ecological Applications*, **20**, 30–59.
- 1062 Bodelier, P.L.E. (2011) Toward understanding, managing, and protecting microbial
1063 ecosystems. *Frontiers in Microbiology*, **2**, 80.
- 1064 Bondeau, A., Smith, P.C., Zaehle, S., Schaphoff, S., Lucht, W., Cramer, W., Gerten, D.
1065 (2007) Modelling the role of agriculture for the 20th century global terrestrial carbon balance.
1066 *Global Change Biology* **13**, 679-706.
- 1067 Bouwman, A.F., Vuuren, D.P. Van, Derwent, R.G., Posch, M. (2002) A Global Analysis of
1068 Acidification and Eutrophication of Terrestrial Ecosystems. *Water, Air, and Soil Pollution*,
1069 **141**, 349–382.

- 1070 Bouwman, L., van der Hoek, K., van Drecht, G., & Eickhout, B. (2006). World livestock and
 1071 crop production systems, land use and environment between 1970 and 2030. In: Brouwer, F.
 1072 & McCarl, B.A. (Eds.), *Agriculture and Climate Beyond 2030*, pp. 75–89, Springer,
 1073 Netherlands.
- 1074 Burney, J.A., Davis, S.J., Lobell, D.B. (2010) Greenhouse gas mitigation by agricultural
 1075 intensification. *Proceedings of the National Academy of Sciences* **107**, 12052–12057.
- 1076 Bustamante, M., Robledo-Abad, C., Harper, R., Mbow, C., Ravindranath, N.H., Sperling, F.,
 1077 Haberl, H., de Siqueira Pinto, A., Smith, P. 2014. Co-benefits, trade-offs, barriers and
 1078 policies for greenhouse gas mitigation in the Agriculture, Forestry and Other Land Use
 1079 (AFOLU) sector. *Global Change Biology* **20**, 3270–3290.
- 1080 Carlson, K.M. & Curran, L.M. (2013) Refined carbon accounting for oil palm agriculture:
 1081 disentangling potential contributions of indirect emissions and smallholder farmers. *Carbon*
 1082 *Management*, **4**, 347–349.
- 1083 Carpenter, S.R., Caraco, N.F., Correll, D.L., Howarth, R.W., Sharpley, A.N., Smith, V.H.
 1084 (1998) Nonpoint pollution of surface waters with phosphorus and nitrogen. *Ecological*
 1085 *Applications*, **8**, 559–568.
- 1086 Certini, G. (2005) Effects of fire on properties of forest soils: a review. *Oecologia*, **143**, 1–10.
- 1087 Chaplot, V., Rumpel, C., Valentin, C. (2005) Water erosion impact on soil and carbon
 1088 redistributions within uplands of South-East Asia. *Global Biogeochemical Cycles*, **19**,
 1089 GB4004, doi:10.1029/2005GB002493.
- 1090 Chowdhury N, Marschner P, Burns R. (2011) Response of microbial activity and community
 1091 structure to decreasing soil osmotic and matric potential. *Plant and Soil*, **344**, 241–254.
- 1092 Cole, J.C, Liverman, D.M. (2011) Brazil's Clean Development Mechanism governance in the
 1093 context of Brazil's historical environment–development discourses. *Carbon Management*, **2**,
 1094 145–160.
- 1095 Comtea, I., Davidson, R., Lucotte M., Carvalho, C.J.R., Oliveira, F.A., Silva, B.P.,
 1096 Rousseau, G. (2012) Physicochemical properties of soils in the Brazilian Amazon following
 1097 fire-free land preparation and slash-and-burn practices. *Agriculture, Ecosystems and*
 1098 *Environment*, **156**, 108–115.
- 1099 Conant, R.T. (2012) Grassland soil organic carbon stocks: status, opportunities, vulnerability.
 1100 In: Lal, R., Lorenz, K., Hüttl, R.F., Schneider, B.U., von Braun, J. (Eds), *Recarbonization of*
 1101 *the Biosphere*, pp. 275–302, Springer, Dordrecht.
- 1102 D'Odorico, P., Bhattachan, A., Davis, K.F., Ravi, S., Runyan, C.W. (2013) Global
 1103 desertification: drivers and feedbacks. *Advances in Water Resources*, **51**, 326–344.
- 1104 Dalal, R.C., Thornton, C.M., Cowie, B.A. (2013) Turnover of organic carbon and nitrogen in
 1105 soil assessed from $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ changes under pasture and cropping practices and
 1106 estimates of greenhouse gas emissions. *Science of the Total Environment*, **465**, 26–35.
- 1107 Daniels, S.M., Evans, M.G., Agnew, C.T., Allott, T.E.H. (2008) Sulphur leaching from
 1108 headwater catchments in an eroded peatland, South Pennines, U.K. *The Science of the Total*
 1109 *Environment*, **407**, 481–96.
- 1110 Davies, Z.G., Edmondson, J.L., Heinemeyer, A., Leake, J.R. & Gaston, K.J. (2011) Mapping
 1111 and urban ecosystem service: quantifying above-ground carbon storage at a city-wide scale.
 1112 *Journal of Applied Ecology*, **48**, 1125–1134.

- 1113 Delgado-Baquerizo, M., Maestre, F.T. Gallardo, A. Bowker, M.A., Wallenstein, M.D.,
 1114 Quero, J.L. *et al.* (2013) Decoupling of soil nutrient cycles as a function of aridity in global
 1115 drylands. *Nature*, **502**, 672–676.
- 1116 Dentener, F., Drevet, J., Lamarque, J.F., Bey, I., Eickhout, B., Fiore, A.M. *et al.* (2006)
 1117 Nitrogen and sulfur deposition on regional and global scales: A multimodel evaluation.
 1118 *Global Biogeochemical Cycles*, **20**, GB4003.
- 1119 Dittmar, T., Rezende, C.E., Manecki, M., Niggemann, J., Ovalle, A.R.C., Stubbins, A.,
 1120 Bernardes, M.C. (2012) Continuous flux of dissolved black carbon from a vanished tropical
 1121 forest biome. *Nature Geoscience*, **5**, 618–622.
- 1122 Don, A., Schumacher, J., Freibauer, A. (2011) Impact of tropical land-use change on soil
 1123 organic carbon stocks – a meta-analysis. *Global Change Biology*, **17**, 1658–1670.
- 1124 Drewniak, B., Song, J., Prell, J., Kotamarthi, V.R., Jacob, R. (2013) Modeling agriculture in
 1125 the Community Land Model. *Geoscientific Model Development*, **6**, 495–515. Available at:
 1126 <http://www.geosci-model-dev.net/6/495/2013/> (accessed 14th June 2015).
- 1127 EEA (2014) *Effects of air pollution on European ecosystems*, Copenhagen, European
 1128 Environment Agency.
- 1129 El-Masri, B., Barman, R., Meiyappan, P., Song, Y., Liang, M., Jain, A. (2013) Carbon
 1130 dynamics in the Amazonian basin: integration of eddy covariance and ecophysiological data
 1131 with a land surface model. *Agricultural & Forest Meteorology*, **182**, 156–167.
- 1132 Eshel, G., Shepon, A., Makov, T., Milo, R. (2014) Land, irrigation water, greenhouse gas,
 1133 and reactive nitrogen burdens of meat, eggs, and dairy production in the United States.
 1134 *Proceedings of the National Academy of Sciences*, **111**, 11996–12001.
- 1135 Fang, Y., Wang, X., Zhu, F., Wu, Z., Li, J., Zhong, L., Chen, D., Yoh, M. (2013) Three-
 1136 decade changes in chemical composition of precipitation in Guangzhou city, southern China:
 1137 has precipitation recovered from acidification following sulphur dioxide emission control?
 1138 *Tellus B*, **65**, Article Number 20213.
- 1139 FAO (1995) Global network on integrated soil management for sustainable use of salt-
 1140 affected soils. FAO Land and Plant Nutrition Managment Service, Rome, Italy.
- 1141 FAO (2010) Global Forest Resources Assessment (FRA) 2010. FAO, Rome. Available at:
 1142 <http://www.fao.org/forestry/fra/fra2010/en/> (accessed 14 February 2015).
- 1143 FAO (2013) *FAO statistical yearbook - World Food and Agriculture*. ISBN 978-92-5-
 1144 107396-4.
- 1145 FAO/IIASA/ISRIC/ISSCAS/JRC (2012) Harmonized World Soil Database (version 1.10),
 1146 FAO, Rome, Italy and IIASA, Laxenburg, Austria, 2012.
- 1147 Foley, J.A., Ramankutty, N., Brauman, K.A., Cassidy, E.S., Gerber, J.S., Johnston, M. *et al.*
 1148 (2011) Solutions for a cultivated planet. *Nature*, **478**, 337–342.
- 1149 Friedlingstein, P., Cox, P., Betts, R., Bopp, L., von Bloh, W., Brovkin, V., Cadule, P., Doney,
 1150 S., Eby, M. & Fung, I. (2006) Climate-Carbon Cycle Feedback Analysis: Results from the
 1151 C⁴MIP Model Intercomparison. *Journal of Climate*, **19**, 3337–3353.
- 1152 Galloway, J.N., Aber, J.D., Erisman, J.W., Seitzinger, S.P., Howarth, R.W., Cowling, E.B.,
 1153 Cosby, B.J. (2003) The Nitrogen Cascade. *BioScience*, **53**, 341.
- 1154 Galloway, J.N., Dentener, F.J., Capone, D.G., Boyer, E.W., Howarth, R.W., Seitzinger, S.P.
 1155 *et al.* (2004) Nitrogen cycles: Past, present, and future. *Biogeochemistry*, **70**, 153–226.

- 1156 Galloway, J.N., Townsend, A.R., Erisman, J.W., Bekunda, M., Cai, Z., Freney, J.R. *et al.*,
 1157 2008. Transformation of the nitrogen cycle: recent trends, questions, and potential solutions.
 1158 *Science*, **320**, 889–92.
- 1159 Gauci, V., Matthews, E., Dise, N., Walter, B., Koch, D., Granberg, G., Vile, M. (2004) Sulfur
 1160 pollution suppression of the wetland methane source in the 20th and 21st centuries.
 1161 *Proceedings of the National Academy of Sciences*, **101**, 12583–12587.
- 1162 Ghassemi, F., Jakeman, A.J., Nix, H.A. (1995) *Salinisation of land and water resources:*
 1163 *Human causes, management and case studies*. Canberra, Australia: Centre for Resource and
 1164 Environmental Studies.
- 1165 Gill, S.E., Handley J.F., Ennos A.R., Pauleits S. (2007) Adapting cities for climate change:
 1166 the role of the green infrastructure. *Built Environment*, **33**, 115–133.
- 1167 Gleick, P.H. (2003) Global freshwater resources: Soft-path solutions for the 21st century.
 1168 *Science*, **302**, 1524–1528.
- 1169 Greaver, T.L., Sullivan, T.J., Herrick, J.D., Barber, M.C., Baron, J.S., Cosby, B.J. *et al.*
 1170 (2012) Ecological effects of nitrogen and sulfur air pollution in the US: what do we know?
 1171 *Frontiers in Ecology and the Environment*, **10**, 365–372.
- 1172 Guo L.B., Gifford R.M. (2002) Soil carbon stocks and land use change: a meta-analysis
 1173 *Global Change Biology*, **8**, 345–360.
- 1174 Guo, K., Liu, Y.F., Zeng, C., Chen, Y.Y., Wei, X.J. (2014) Global research on soil
 1175 contamination from 1999 to 2012: A bibliometric analysis. *Acta Agriculturae Scandinavica*,
 1176 *Section B — Soil & Plant Science*, **64**, 377–391.
- 1177 Haberl, H., Erb, K.H., Krausmann, F., Gaube, V., Bondeau, A., Plutzar, C. *et al.* (2007)
 1178 Quantifying and mapping the human appropriation of net primary production in Earth's
 1179 terrestrial ecosystems. *Proceedings of the National Academy of Sciences, USA* **104**, 12942–
 1180 12947.
- 1181 Hamza, M., Anderson, W. (2005) Soil compaction in cropping systems: A review of the
 1182 nature, causes and possible solutions. *Soil and Tillage Research* **82**, 121 - 145.
- 1183 Hansen, M.C., Potapov, P.V., Moore, R., Hancher, M., Turubanova, S.A., Tyukavina, A. *et*
 1184 *al.* (2013). High-Resolution Global Maps of 21st-Century Forest Cover Change. *Science*,
 1185 **342**, 850–853.
- 1186 Henderson, B.B., Gerber, P.J., Hilinski, T.E., Falcucci, A., Ojima, D.S., Salvatore, M.,
 1187 Conant, R.T. (2015) Greenhouse gas mitigation potential of the world's grazing lands:
 1188 Modeling soil carbon and nitrogen fluxes of mitigation practices. *Agriculture, Ecosystems &*
 1189 *Environment*, **207**, 91–100.
- 1190 Herrero, M., & Thornton, P. K. (2013) Livestock and global change: Emerging issues for
 1191 sustainable food systems. *Proceedings of the National Academy of Sciences*, **110**, 20878–
 1192 20881.
- 1193 Hester, R.E., Harrison R.M. (2001) *Assessment and contamination of contaminated land*.
 1194 Royal Society of Chemistry, 164pp.

- 1195 Hettelingh, J.P., Sliggers, J., van het Bolcher, M., Denier van der Gon, H., Groenenberg, B.J.,
 1196 Ilyin, I. *et al.* (2006) *Heavy Metal Emissions, Depositions, Critical Loads and Exceedances*
 1197 *in Europe*, Den Haag, Netherlands.
- 1198 Hooijer A., Page, S., Canadell, J.G., Silvius, M., Kwadijk, J., Wosten, H., Jauhiainen, J.
 1199 (2010) Current and future CO₂ emissions from drained peatlands in Southeast Asia.
 1200 *Biogeosciences*, **7**, 1505–1514.
- 1201 Huang, M., Yang, L., Qin, H., Jiang, L., Zou, Y. (2013) Quantifying the effect of biochar
 1202 amendment on soil quality and crop productivity in Chinese rice paddies. *Field Crops*
 1203 *Research*, **11**, 172–177.
- 1204 Hurtt, G.C., Chini, L.P., Froking, S., Betts, R.A., Feddema, J., Fischer, G. *et al.* (2011)
 1205 Harmonization of Land-Use Scenarios for the Period 1500–2100: 600 Years of Global
 1206 Gridded Annual Land-Use Transitions, Wood Harvest, and Resulting Secondary Lands,
 1207 *Climatic Change*, **109**, 117–161.
- 1208 IFPRI (2011) Global Food Policy Report 2011. Available at: <http://www.ifpri.org/gfpr/2011>.
- 1209 INPE (2014) Description of the PRODES project. Available at:
 1210 <http://www.obt.inpe.br/prodes/index.php>.
- 1211 IPCC (2007) *Climate Change 2007. The Physical Science Basis*. Cambridge University Press,
 1212 Cambridge, UK.
- 1213 Jain, A.K., West, T., Yang, X., Post, W. (2005) Assessing the impact of changes in climate
 1214 and CO₂ on potential carbon sequestration in agricultural soils. *Geophysical Research*
 1215 *Letters*, **32**, L19711, doi:10.1029/2005GL023922.
- 1216 Jain, A.K., Meiyappan, P., Song, Y., House, J. (2013) CO₂ emissions from land-use change
 1217 affected more by nitrogen cycle, than by the choice of land-cover data. *Global Change*
 1218 *Biology*, doi: 10.1111/gcb.12207.
- 1219 Joosten, H. (2010) *The global peatland CO₂ picture - peatland status and drainage related*
 1220 *emissions in all countries of the world*. Wetlands International, Wageningen, The
 1221 Netherlands.
- 1222 Ju, X., Xing, G., Chen, X., Zhang, S., Zhang, L., Liu, X. *et al.* (2009) Reducing
 1223 environmental risk by improving N management in intensive Chinese agricultural systems.
 1224 *Proceedings of the National Academy of Sciences* **106**, 3041–3046.
- 1225 Kell, D. (2012) Large-scale sequestration of atmospheric carbon via plant roots in natural and
 1226 agricultural ecosystems: why and how. *Philosophical Transactions of the Royal Society, B*.
 1227 **367**, 1589–1597, 2012.
- 1228 Kelly, E.N., Schindler, D.W., Hodson, P. V., Short, J.W., Radmanovich, R., Nielsen, C.C.
 1229 (2010) Oil sands development contributes elements toxic at low concentrations to the
 1230 Athabasca River and its tributaries. *Proceedings of the National Academy of Sciences* **107**,
 1231 16178–16183.
- 1232 Klein Goldewijk, K., Beusen, A., Van Dreht, G., De Vos, M. (2011) The HYDE 3.1
 1233 spatially explicit database of human-induced global land-use change over the past 12,000
 1234 years. *Global Ecology & Biogeography*, **20**, 73–86.
- 1235 Kravchenko, A.N., Robertson, G.P. (2010) Whole-profile soil carbon stocks: The danger of
 1236 assuming too much from analyses of too little. *Soil Science Society of America Journal*, **75**,
 1237 235–240.

- 1238 Krug, E.C., Frink, C.R. (1983) Acid Rain on Acid Soil: A New Perspective. *Science*, **221**,
1239 520–525.
- 1240 Kuylensstierna, J.C., Rodhe, H., Cinderby, S., Hicks, K. (2001) Acidification in developing
1241 countries: ecosystem sensitivity and the critical load approach on a global scale. *Ambio*, **30**,
1242 20–28.
- 1243 Le Quéré, C., Peters, G.P., Andres, R.J., Andrew, R.M., Boden, T.A., Ciais, P. *et al.* 2014.
1244 Global carbon budget 2013. *Earth System Science Data*, **6**, 235–263.
- 1245 Laudon, H., Dillon, P.J., Eimers, M.C., Semkin, R.G., Jeffries, D.S. (2004) Climate-induced
1246 episodic acidification of streams in central ontario. *Environmental Science & Technology*, **38**,
1247 6009–6015.
- 1248 Lawrence, G.B., Shortle, W.C., David, M.B., Smith, K.T., Warby, R.A.F. & Lapenis, A.G.
1249 (2012) Early Indications of Soil Recovery from Acidic Deposition in U.S. Red Spruce
1250 Forests. *Soil Science Society of America Journal*, **76**, 1407.
- 1251 Lehmann J., Czimczik, C., Laird, D., Sohi, S. (2015) Stability of biochar in soil. In:
1252 Lehmann, J. & Joseph, S. (Eds.), *Biochar for Environmental Management: Science*,
1253 *Technology and Implementation*, pp. 235-282, Taylor and Francis, London, UK.
- 1254 Li, D., Niu, S., Luo, Y. (2012) Global patterns of the dynamics of soil carbon and nitrogen
1255 stocks following afforestation: a meta - analysis. *New Phytologist*, **195**, 172-181.
- 1256 Lindeskog, M., Arneth, A., Bondeau, A., Waha, K., Seaquist, J., Olin, S. *et al.* (2013)
1257 Implications of accounting for land use in simulations of ecosystem carbon cycling in Africa.
1258 *Earth System Dynamics*, **4**, 385–407.
- 1259 Liu, L., Greaver, T.L. (2009) A review of nitrogen enrichment effects on three biogenic
1260 GHGs: the CO₂ sink may be largely offset by stimulated N₂O and CH₄ emission. *Ecology*
1261 *Letters*, **12**, 1103–1117.
- 1262 Liu, L., Xu, X., Zhuang, D., Chen, X., Li, S. (2013) Changes in the potential multiple
1263 cropping system in response to climate change in China from 1960–2010. *PLoS ONE* **8**,
1264 e80990. doi:10.1371/journal.pone.0080990.
- 1265 Liu, X., Zhang, Y., Han, W., Tang, A., Shen, J., Cui, Z. *et al.* (2013) Enhanced nitrogen
1266 deposition over China. *Nature*, **494**, 459–462.
- 1267 Liu, X.Y., Qu, J.J., Li, L.Q., *et al.* (2012) Can biochar amendment be an ecological
1268 engineering technology to depress N₂O emission in rice paddies? - A cross site field
1269 experiment from South China. *Ecological Engineering*, **42**, 168-173.
- 1270 Liu, Z.H., Jiang, L.H., Zhang, W.J., Zheng, F.L., Wang, M., Lin, H.T. (2008) Evolution of
1271 fertilization rate and variation of soil nutrient contents in greenhouse vegetable cultivation in
1272 Shandong. *Pedologica Sinica*, **45**, 296-303. (in Chinese with English abstract).
- 1273 Lu, X., Mao, Q., Gilliam, F.S., Luo, Y., Mo, J. (2014) Nitrogen deposition contributes to soil
1274 acidification in tropical ecosystems. *Global Change Biology*, doi:10.1111/gcb.12665.
- 1275 Lundström, U.S., Bain, D.C., Taylor, A.F.S. & van Hees, P.A.W. (2003) Effects of
1276 acidification and its mitigation with lime and wood ash on forest soil processes: a review.
1277 *Water, Air and Soil Pollution* **3**, 5–28.
- 1278 Machmüller, M.B., Kramer, M.G., Cyle, T.K., Hill, N., Hancock, D., Thompson, A. (2015)
1279 Emerging land use practices rapidly increase soil organic matter. *Nature Communications*, **6**,
1280 Article Number: 6995. doi: 10.1038/ncomms7995.

- 1281 Maderova, L., Paton, G.I. (2013) Deployment of microbial sensors to assess zinc
1282 bioavailability and toxicity in soils. *Soil Biology and Biochemistry* **66**, 222–228.
- 1283 Magnani, F., Mencuccini, M., Borghetti, M., Berbigier, P., Berninger, F., Delzon, S. *et al.*
1284 (2007) The human footprint in the carbon cycle of temperate and boreal forests. *Nature*, **447**,
1285 848–50.
- 1286 Marfenina, O.E., Ivanova, A.E. Kislova E.E., Sacharov, D.S. (2008) The mycological
1287 properties of medieval culture layers as a form of soil “biological memory” about
1288 urbanization. *Journal of Soils and Sediments*, **8**, 340–348.
- 1289 Marshall, M. R., Francis, O. J., Frogbrook, Z. L., Jackson, B. M., McIntyre, N. *et al.* (2009)
1290 The impact of upland land management on flooding: results from an improved pasture
1291 hillslope. *Hydrological Processes*, **23**, 464–475.
- 1292 Matson, P.A., Parton, W.J., Power, A.G., Swift, M.J. (1997). Agricultural intensification and
1293 ecosystem properties. *Science*, **277**, 504–509.
- 1294 McCarthy, D.F. (2007). *Essentials of Soil Mechanics and Foundations*. Upper Saddle River,
1295 NJ: Pearson Prentice Hall.
- 1296 McSherry, M.E., Ritchie, M.E. (2013) Effects of grazing on grassland soil carbon: a global
1297 review. *Global Change Biology*, **19**, 1347–1357.
- 1298 Medlyn, B.E., Zaehle, S., De Kauwe, M.G., Walker, A.P., Dietze, M.C., Hanson, P.J. *et al.*
1299 2015. Using ecosystem experiments to improve vegetation models. *Nature Climate Change*,
1300 **5**, 528–534.
- 1301 Meersmans, J., Van Wesemael, B., De Ridder, F. Dotti, M.F., De Baets, S. Van Molle, M.
1302 (2009) Changes in organic carbon distribution with depth in agricultural soils in northern
1303 Belgium, 1960–1990. *Global Change Biology* **15**, 2739–2750.
- 1304 Meuser, H. (2010) *Contaminated Urban Soils*. Springer Science & Business Media, 340pp.
- 1305 Monteith, D.T., Stoddard, J.L., Evans, C.D., de Wit, H.A., Forsius, M., Høgåsen, T. *et al.*
1306 (2007) Dissolved organic carbon trends resulting from changes in atmospheric deposition
1307 chemistry. *Nature*, **450**, 537–540.
- 1308 Mueller, N.D, West, P.C., Gerber, J.S., MacDonald, G.K., Polasky, S., Foley, J.A. (2014) A
1309 tradeoff frontier for global nitrogen use and cereal production. *Environmental Research*
1310 *Letters* **9**, 054002, doi:10.1088/1748-9326/9/5/054002.
- 1311 Murty, D., Kirschbaum, M.U.F., McMurtrie, R.E., McGilvray, H. (2002) Does conversion of
1312 forest to agricultural land change soil carbon and nitrogen? A review of the literature. *Global*
1313 *Change Biology*, **8**, 105–123.
- 1314 Nave, L.E, Vance, E.D., Swanston, C.W., Curtis, P.S. (2011) Fire effects on temperate forest
1315 soil C and N storage. *Ecological Applications* **21**, 1189–1201.
- 1316 Nicholson, F.A., Smith, S.R., Alloway, B.J., Carlton-Smith, C., Chambers, B.J. (2003) An
1317 inventory of heavy metals inputs to agricultural soils in England and Wales. *The Science of*
1318 *the Total Environment*, **311**, 205–219.
- 1319 Nilsson, J., Grennfelt, P. (1988) *Critical Loads for Sulphur and Nitrogen*, Copenhagen,
1320 Nordic Council of Ministers.

- 1321 Nolte, C., Agrawal, A., Silvius, K.M., Soares-Filho, B.S. (2013) Governance regime and
 1322 location influence avoided deforestation success of protected areas in the Brazilian Amazon.
 1323 *Proceedings of the National Academy of Sciences*, **110**, 4956-4961.
- 1324 Ogle, S., Breidt, F.J., Paustian, K. (2005) Agricultural management impacts on soil organic
 1325 carbon storage under moist and dry climatic conditions of temperate and tropical regions.
 1326 *Biogeochemistry*, **72**, 87-121.
- 1327 Ojima, D.S., Dirks, B., Glenn, E.P., Owensby, C.E., Scurlock, J.O. (1993) Assessment of C
 1328 budget for grasslands and drylands of the world. *Water Air and Soil Pollution*, **70**, 95-109.
- 1329 Olander, L.O., Bustamante, M.C.C., Asner, G.P., Telles, E., do Prado, Z.A. (2005) Surface
 1330 soil changes following selective logging in an Eastern Amazon forest. *Earth Interactions* **9**,
 1331 1-19.
- 1332 Oldeman, L.R., Hakkeling, R.T.A. & Sombroek, W.G. (1991) Global Assessment of Soil
 1333 Degradation GLASOD, second revised edition October 1991. Wageningen: International Soil
 1334 Reference and Information Centre; Nairobi: United Nations Environment Programme.
- 1335 Oulehle, F., Evans, C.D., Hofmeister, J., Krejci, R., Tahovska, K., Persson, T. *et al.* (2011)
 1336 Major changes in forest carbon and nitrogen cycling caused by declining sulphur deposition.
 1337 *Global Change Biology*, **17**, 3115–3129.
- 1338 Paustian, K., Andrén, O., Janzen, H.H., Lal, R., Smith, P., Tian, G. *et al.* (1997) Agricultural
 1339 soils as a sink to mitigate CO₂ emissions. *Soil Use and Management*, **13**, 230–244.
- 1340 Perez, C.A., Carmona, M.R., Fariña, J.M., Armesto, J.J. (2009) Selective logging of lowland
 1341 evergreen rainforests in Chiloe Island, Chile: Effects of changing tree species composition on
 1342 soil nitrogen transformations. *Forest Ecology and Management* **258**, 1660–1668.
- 1343 Philibert, A., Loyce, C. Makowski, D., Bernacchi, C.J. (2012) Quantifying uncertainties in
 1344 N₂O emission due to N fertilizer application in cultivated areas. *PloS One* **7**, e50950. doi:
 1345 10.1371/journal.pone.0050950.
- 1346 Poeplau, C., Don, A. (2015) Carbon sequestration in agricultural soils via cultivation of cover
 1347 crops – A meta-analysis. *Agriculture, Ecosystems & Environment*, **200**, 33-41.
- 1348 Poeplau, C., Don, A., Vesterdal, L., Leifeld, J., Wesemael, B., Schumacher, J., Gensior, A.
 1349 (2011) Temporal dynamics of soil organic carbon after land-use change in the temperate zone
 1350 – carbon response functions as a model approach. *Global Change Biology* **17**, 2415–2427.
- 1351 Powlson, D.S., Stirling, C.M., Jat, M.L., Gerard, B.G., Palm, C.A., Sanchez, P.A., Cassman,
 1352 K.G. (2014) Limited potential of no-till agriculture for climate change mitigation. *Nature*
 1353 *Climate Change* **4**, 678–683.
- 1354 Prokop G., Jobstmann H., Schöbauer A. (2011) Overview on best practices for limiting soil
 1355 sealing and mitigating its effects in EU-27 (Environment Agency Austria), technical Report –
 1356 2011-50, ISBN: 978-92-79-20669-6. <http://ec.europa.eu/environment/soil/sealing.htm>
- 1357 Ravi, S., Breshears, D.D., Huxman, T.E., D'Odorico, P. (2010) Land degradation in drylands:
 1358 Interactions among hydrologic–aeolian erosion and vegetation dynamics. *Geomorphology*
 1359 **116**, 236–245.
- 1360 Ray, D.K., Foley, J.A. (2013) Increasing global crop harvest frequency: recent trends and
 1361 future directions. *Environmental Research Letters*, **8**, 044041. doi:10.1088/1748-
 1362 9326/8/4/044041

- 1363 Reay, D.S., Dentener, F., Smith, P., Grace, J., Feely, R. (2008) Global nitrogen deposition
1364 and carbon sinks. *Nature Geoscience*, **1**, 430-437. doi: 10.1038/ngeo230.
- 1365 Reis, S., Grennfelt, P., Klimont, Z., Amann, M., Apsimon, H., Hettelingh, J.-P. *et al.* (2012)
1366 From acid rain to climate change. *Science*, **338**, 1153-1154.
- 1367 Rengasamy P. (2008) Salinity in the landscape: A growing problem in Australia. *Geotimes*
1368 **53**, 34-39.
- 1369 Reuss, J.O., Johnson, D.W. (1986) *Acid Deposition and the Acidification of Soils and Waters*,
1370 Ecological. New York, Springer Verlag.
- 1371 Ribeiro-Filho, A.A., Adams, C., Sereni Murrieta, R.S. (2013) The impacts of shifting
1372 cultivation on tropical forest soil: a review. *Bol. Mus. Para. Emilio Goeldi. Cienc. Hum.*,
1373 *Belém*, **8**, 693-727.
- 1374 Ripple, W.J., Smith, P., Haberl, H., Montzka, S.A., McAlpine, C., Boucher, D.H. (2014)
1375 Ruminants, climate change and climate policy. *Nature Climate Change*, **4**, 2-5.
- 1376 Robinson, D.A., Fraser, I., Dominati, E.J., Davíðsdóttir, B., Jónsson, J.O.G., Jones, L. *et al.*
1377 2014. On the value of soil resources in the context of natural capital and ecosystem service
1378 delivery. *Soil Science Society of America Journal* (in press).
- 1379 Robinson, D.A., Hockley, N., Cooper, D.M., Emmett, B.A., Keith, A.M., Lebron, I. *et al.*
1380 (2013) Natural capital and ecosystem services, developing an appropriate soils framework as
1381 a basis for valuation. *Soil Biology and Biochemistry*, **57**, 1023-1033.
- 1382 RoTAP (2012) *Review of Transboundary Air Pollution (RoTAP): Acidification*,
1383 *Eutrophication, Ground Level Ozone and Heavy Metals in the UK*, Edinburgh, Contract
1384 Report to the Department for Environment, Food and Rural Affairs. Centre for Ecology &
1385 Hydrology.
- 1386 Rothwell, J.J., Robinson, S.G., Evans, M.G., Yang, J., Allott, T.E.H. (2005) Heavy metal
1387 release by peat erosion in the Peak District, southern Pennines, UK. *Hydrological Processes*,
1388 **19**, 2973-2989.
- 1389 Royal Society of London. (2009). *Reaping the benefits: science and the sustainable*
1390 *intensification of global agriculture*. London, UK: London.
- 1391 Ryals, R., Hartman, M.D., Parton, W.J., DeLonge, M., Silver W.L. (2015) Long-term climate
1392 change mitigation potential with organic matter management on grasslands. *Ecological*
1393 *Applications*, **25**, 531-545.
- 1394 Setia, R., Gottschalk, P., Smith, P., Marschner, P., Baldock, J. & Smith, J. (2013) Soil salinity
1395 decreases global soil organic carbon stocks. *Science of the Total Environment*, **465**, 267-272.
- 1396 Setia, R., Marschner, P., Baldock, J., Chittleborough, D., Smith, P., Smith, J. (2011a) Salinity
1397 effects on carbon mineralization in soils of varying texture. *Soil Biology and Biochemistry*,
1398 **43**, 1908-1916.
- 1399 Setia, R., Smith, P., Marschner, P., Baldock, J., Chittleborough, D.J., Smith, J. (2011b)
1400 Introducing a decomposition rate modifier in the Rothamsted carbon model to predict soil
1401 organic carbon stocks in saline soils. *Environmental Science & Technology*, **45**, 6396-6403.

- 1402 Setia, R., Smith, P., Marschner, P., Gottschalk, P., Baldock, J., Verma, V. *et al.* (2012)
 1403 Simulation of salinity effects on past, present and future soil organic carbon stocks.
 1404 *Environmental Science & Technology*, **46**, 1624-1631.
- 1405 Shcherbak, I., Millar, N., Robertson, G.P. (2014) Global meta-analysis of the nonlinear
 1406 response of soil nitrous oxide (N₂O) emissions to fertilizer nitrogen. *Proceedings of the*
 1407 *National Academy of Sciences*. doi: 10.1073/pnas.1322434111.
- 1408 Shi, S., Zhang W., Zhang P., Yu Y., Ding, F.A. (2013) Synthesis of change in deep soil
 1409 organic carbon stores with afforestation of agricultural soils. *Forest Ecology and*
 1410 *Management*, **296**, 53–63.
- 1411 Siebert, S. & Döll, P. (2010) Quantifying blue and green virtual water contents in global crop
 1412 production as well as potential production losses without irrigation. *Journal of Hydrology*
 1413 **384**, 198-217.
- 1414 Siebielec G., Lazar S., Kaufmann C., Jaensch, S. (2010) *Handbook for measures enhancing*
 1415 *soil function performance and compensating soil loss during urbanization process*. Urban
 1416 SMS – Soil Management Strategy project, 37pp. www.urban-sms.eu
- 1417 Sitch, S., Smith, B., Prentice, I., Arneth, A., Bondeau, A., Cramer, W., *et al.* (2003)
 1418 Evaluation of ecosystem dynamics, plant geography and terrestrial carbon cycling in the LPJ
 1419 dynamic global vegetation model. *Global Change Biology* **9**, 161-185.
- 1420 Smil, V. (2000) Phosphorus in the environment: natural flows and human interferences.
 1421 *Annual Review of Energy and the Environment*, **25**, 53–88.
- 1422 Smith, B., Prentice, I., Sykes, M (2001) Representation of vegetation dynamics in the
 1423 modelling of terrestrial ecosystems: comparing two contrasting approaches within European
 1424 climate space. *Global Ecology and Biogeography* **10**, 621-637.
- 1425 Smith, J.U., Gottschalk, P., Bellarby, J., Chapman, S., Lilly, A., Towers, W. *et al.* (2010)
 1426 Estimating changes in national soil carbon stocks using ECOSSE – a new model that includes
 1427 upland organic soils. Part I. Model description and uncertainty in national scale simulations
 1428 of Scotland. *Climate Research* **45**, 179-192.
- 1429 Smith, P. (2005) An overview of the permanence of soil organic carbon stocks: influence of
 1430 direct human-induced, indirect and natural effects. *European Journal of Soil Science*, **56**,
 1431 673-680.
- 1432 Smith, P. (2008) Land use change and soil organic carbon dynamics. *Nutrient Cycling in*
 1433 *Agroecosystems* **81**, 169-178.
- 1434 Smith, P. (2012) Soils and climate change. *Current Opinion in Environmental Sustainability*
 1435 **4**, 539–544.
- 1436 Smith, P., Ashmore, M., Black, H., Burgess, P.J., Evans, C., Quine, T. *et al.* (2013a) The role
 1437 of ecosystems and their management in regulating climate, and soil, water and air quality.
 1438 *Journal of Applied Ecology*, **50**, 812–829.
- 1439 Smith, P., Cotrufo, M.F., Rumpel, C., Paustian, K., Kuikman, P.J., Elliott, J. A. *et al.* (2015)
 1440 Biogeochemical cycles and biodiversity as key drivers of ecosystem services provided by
 1441 soils. *SOIL Discussions* **2**, 537-586, 2015.
- 1442 Smith, P., Davies, C.A., Ogle, S., Zanchi, G., Bellarby, J., Bird, N. *et al.* (2012) Towards an
 1443 integrated global framework to assess the impacts of land use and management change on
 1444 soil carbon: current capability and future vision. *Global Change Biology* **18**, 2089–2101.

- 1445 Snyder, C.S., Davidson, E.A., Smith, P., Venterea, R.T. (2014) Agriculture: sustainable crop
1446 and animal production to help mitigate nitrous oxide emissions. *Current Opinion in*
1447 *Environmental Sustainability* **9-10**, 46-54.
- 1448 Soares-Filho, B., Rajao, R. Macedo, M. Carneiro, A., Costa, W. Coe, M. *et al.* (2014) LAND
1449 USE Cracking Brazil's Forest Code. *Science*, **344**, 363-364.
- 1450 Song, G.H., Li, L.Q., Pan, G.X., Zhang, Q. (2005) Topsoil organic carbon storage of China
1451 and its loss by cultivation. *Biogeochemistry* **74**, 47-62.
- 1452 Spranger, T., Hettelingh, J.-P., Slootweg, J., Posch, M. (2008) Modelling and mapping long-
1453 term risks due to reactive nitrogen effects: an overview of LRTAP convention activities.
1454 *Environmental Pollution*, **154**, 482-487.
- 1455 State Bureau of Statistics-China (2005) *50 Years Rural Statistics of New China*. China
1456 Statistics Press, Beijing, China.
- 1457 Sutton M.A., E. Nemitz, J.W. Erisman, C. Beier, K. Butterbach Bahl, P. Cellier *et al.* (2007)
1458 Challenges in quantifying biosphere-atmosphere exchange of nitrogen species.
1459 *Environmental Pollution*, **150**, 125-139.
- 1460 Tian, H.Q., Lu, C.Q., Melillo, J., Ren, R., Huang, Y., Xu, X.F. *et al.* (2012) Food benefit and
1461 climate warming potential of nitrogen fertilizer uses in China. *Environmental Research*
1462 *Letters*, **7**, doi:10.1088/1748-9326/7/4/044020.
- 1463 Tilman, D., Balzer, C., Hill, J., Befort, B.L. (2011) Global food demand and the sustainable
1464 intensification of agriculture. *Proceedings of the National Academy of Sciences*, **108**, 20260-
1465 20264.
- 1466 Tilman, D., Cassman, K. G., Matson, P. A., Naylor, R., Polasky, S. (2002) Agricultural
1467 sustainability and intensive production practices. *Nature*, **418**, 671-677.
- 1468 Tipping, E., Smith, E., Lawlor, A., Hughes, S., Stevens, P. (2003) Predicting the release of
1469 metals from ombrotrophic peat due to drought-induced acidification. *Environmental*
1470 *Pollution*, **123**, 239-253.
- 1471 Todd-Brown, K.E.O., Randerson, J.T., Post, W.M., Hoffman, F.M., Tarnocai, C., Schuur,
1472 E.A.G. & Allison, S.D. (2013) Causes of variation in soil carbon simulations from CMIP5
1473 Earth system models and comparison with observations. *Biogeosciences*, **10**, 1717-1736.
- 1474 Tóth, G., Stolbovoy, V., Montanarella, L. (2007) *Soil Quality and Sustainability Evaluation –*
1475 *An Integrated approach to support soil-related policies of the European Union*. EUR 22721
1476 EN. Available at: <http://ec.europa.eu/environment/soil/biodiversity.htm> (accessed 14th
1477 February 2015).
- 1478 Tubiello, F.N., Salvatore, M., Condor Golec, R., Rossi, S., Ferrara, A., Biancalani, R. *et al.*
1479 (2015) The contribution of agriculture, forestry and other land use activities to global
1480 warming, 1990–2012. *Global Change Biology*, doi: 10.1111/gcb.12865.
- 1481 UNDP (2014a) *The Millennium Development Goals 2014*. United Nations Development
1482 Group 2014. 59pp. Available at:
1483 [http://www.undp.org/content/dam/undp/library/MDG/english/UNDP_MDGReport_EN_2014](http://www.undp.org/content/dam/undp/library/MDG/english/UNDP_MDGReport_EN_2014_Final1.pdf)
1484 [Final1.pdf](http://www.undp.org/content/dam/undp/library/MDG/english/UNDP_MDGReport_EN_2014_Final1.pdf). (Accessed 4th June 2015)
- 1485 UNDP (2014b) *Delivering the post-2015 development agenda. Opportunities at the national*
1486 *and local levels*. United Nations Development Group 2014. 44pp. Available at:
1487 [http://www.undp.org/content/dam/undp/library/MDG/Post2015/UNDP-MDG-Delivering-](http://www.undp.org/content/dam/undp/library/MDG/Post2015/UNDP-MDG-Delivering-Post-2015-Report-2014.pdf)
1488 [Post-2015-Report-2014.pdf](http://www.undp.org/content/dam/undp/library/MDG/Post2015/UNDP-MDG-Delivering-Post-2015-Report-2014.pdf). (Accessed 4th June 2015)

- 1489 Van Aardenne, J.A., Dentener, F.J., Olivier, J.G.J., Goldewijk, C.G.M.K. & Lelieveld, J.
1490 (2001) A 1°×1° resolution data set of historical anthropogenic trace gas emissions for the
1491 period 1890-1990. *Global Biogeochemical Cycles*, **15**, 909–928.
- 1492 Venterea, R.T., Maharjan, B., Dolan, M.S. (2011) Fertilizer source and tillage effects on
1493 yield-scaled nitrous oxide emissions in a corn cropping system. *Journal of Environmental*
1494 *Quality*, **40**, 1521-1531.
- 1495 Vet, R., Artz, R.S., Carou, S., Shaw, M., Ro, C.-U., Aas, W. *et al.* (2014) A global
1496 assessment of precipitation chemistry and deposition of sulfur, nitrogen, sea salt, base
1497 cations, organic acids, acidity and pH, and phosphorus. *Atmospheric Environment*, **93**, 3–100.
- 1498 Villela, D.M., Nascimento, M.T., Aragão, L.E.O.C., Gama, D.M. (2006) Effect of selective
1499 logging on forest structure and nutrient cycling in a seasonally dry Brazilian Atlantic forest.
1500 *Journal of Biogeography*, **33**, 506–516.
- 1501 Wei, X., Shao, M., Gale, W., Li, L. (2014a) Global pattern of soil carbon losses due to the
1502 conversion of forests to agricultural land. *Scientific Reports* **4**, 4062. doi: 10.1038/srep040.
- 1503 Wei, X., Huang, L., Xiang, Y., Shao, M., Zhang, X., Gale, W. (2014b) The dynamics of soil
1504 OC and N after conversion of forest to cropland. *Agricultural and Forest Meteorology*, **194**,
1505 188-196.
- 1506 West, P.C., Gerber, J.S., Engstrom, P.M., Mueller, N.D., Brauman, K.A., Carlson, K.M. *et al.*
1507 (2014) Leverage points for improving global food security and the environment. *Science*,
1508 **345**, 325–328.
- 1509 West, P.C., Gibbs, H.K., Monfreda, C., Wagner, J., Barford, C.C., Carpenter, S.R., Foley,
1510 J.A. (2010) Trading carbon for food: Global comparison of carbon stocks vs. crop yields on
1511 agricultural land. *Proceedings of the National Academy of Sciences* **107**, 19645-19645.
- 1512 West, T.O., Post, W.M. (2002) Soil organic carbon sequestration rates by tillage and crop
1513 rotation. *Soil Science Society of America Journal*, **66**, 1930-1940.
- 1514 Whitfield, C.J., Aherne, J., Watmough, S.A., McDonald, M. (2010) Estimating the sensitivity
1515 of forest soils to acid deposition in the Athabasca Oil Sands Region, Alberta. *Journal of*
1516 *Limnology*, **69**, 201–208.
- 1517 Wilhelm, W.W., Johnson, J.M.F., Hatfield, J.L., Voorhees, W.B., Linden, D.R. (2004) Crop
1518 and soil productivity response to corn residue removal: A literature review. *Agronomy*
1519 *Journal* **96**, 1-17, (2004).
- 1520 Woolf, D., Amonette, J.E., Street-Perrott, F.A. Lehmann, J., Joseph, S. (2010) Sustainable
1521 biochar to mitigate global climate change. *Nature Communications*, **1**, Article 56. doi:
1522 10.1038/ncomms1053.
- 1523 World Bank (2008) *World Development Report 2008: Agriculture for Development*. World
1524 Bank, Washington, DC.
- 1525 World Urbanization Prospects (2014) World's population increasingly urban with more than
1526 half living in urban areas. Available at:
1527 [http://www.un.org/en/development/desa/news/population/world-urbanization-prospects-](http://www.un.org/en/development/desa/news/population/world-urbanization-prospects-2014.html)
1528 [2014.html](http://www.un.org/en/development/desa/news/population/world-urbanization-prospects-2014.html) (accessed on 14th February 2015).
- 1529 Yates, D.N., Kittel, T.G.F., Cannon, R.F. (2000) Comparing the correlative holdridge model
1530 to mechanistic biogeographical models for assessing vegetation distribution response to
1531 climatic change. *Climatic Change*, **44**, 59-87.

- 1532 Zhang, X.H., Li, D.Y., Pan, G., Li, L.Q., Lin, F., Xu, X.W. (2008) Conservation of wetland
1533 soil c stock and climate change of China. *Adv. Climate Change Research*, **4**, 202-208.

Tables

Table 1. Observed and modelled soil carbon change (%) when converting from land cover classes in the left hand column to land cover classes listed across the top. Results are from meta-analysis of observations from the sources listed below. Model results (range across three models) are shown for comparison in square brackets, range across the ISAM, LPJml, and LPJ_GUESS models (see text), although note this calculated as difference in soil carbon under the different land classes in 2010 and is thus not modelled loss/gain after a conversion. Negative numbers represent loss of soil carbon.

		Regrowth Forest	Tree plantation	Grassland	Pasture	Cropland
Forest	Global	-9% (2)	-13% (3) ^a		+8% (3)	-42% (3)
	Trop.				-12% (2)	-41% (1)
						-25% (2) ^b
						-30% (2) ^c
	Temp.				[-40 to -63%]	-24% (5)
						[-51 to -62%]
	Boreal				[-52% to +17]	-52% (1)
						-36% (4)
						[-24 to -60%]
						-31% (1)
					[-14 to -49%]	[-63 to -65%]
Grassland	Global				[-1 to +15%]	[-2 to -6%]
	Trop					-32% (4)
	Temp					[-15 to -53%]
	Boreal				[-28 to +3%]	[-70 to -79%]
					[-26 to -71%]	
Pasture	Global		-10% (3)			-59% (3)
	Trop					[-19 to +0.5%]
	Temp					[-17 to -35%]
	Boreal					[-28 to -59%]
Cropland	Global	+53% (3)	+18% (3)	+28% (4)	+19% (3)	
	Trop		+29% (2)		+26% (2)	
	Temp	+16% (4)	+20% (6)			
	Boreal					

Footnotes: ^a Broadleaf tree plantations onto prior native forest or pasture did not affect soil C stocks whereas pine plantations reduced soil C stocks by -12 to -15%; ^b Annual crops; ^c Perennial crops; 1 Wei *et al.* (2014a); 2 Don *et al.* (2011); 3 Guo & Gifford (2002; tropical and temperate zones compiled); 4 Poeplau *et al.* (2011); 5 Murty *et al.* (2014); 6 Barcena *et al.* (2014).

Table 2. Soil carbon loss due to land use change 1860 to 2010 (PgCO₂)

Model	Tropical	Temperate	Boreal	Global
LPJ-GUESS	46	55	1	109
LPJmL	128	95	0	227
ISAM	63	139	19	221
Mean	79	96	7	186

Table 3. Threats to soil resource quality and functioning under increasing intensity of agricultural management


Agricultural management practice	Specific issue	Distribution	Major environmental consequence	Knowledge gap
Cropping practice	Harvest frequency	Global	Soil quality and resilience	Impact on total C and nutrient cycles
	Monoculture	Global but particularly in developing and transition countries	Soil health, pesticide residue in intensively managed monocultures	Biological resilience
Use of agrochemicals	Over fertilization	Particularly in some developing countries	Soil acidification, water pollution, N ₂ O emission and nitrate accumulation	Rate reducing versus balancing
Irrigation	Submerged Rice	Developing countries, Asian	Water scarcity, methane emission	Trade-offs C and water,
	Arid/semi-arid regions	Arid/semi-arid regions	Secondary salinization, water scarcity	Competition use of water
Livestock management	Over-grazing	Largely in developing countries	Soil degradation, water storage, C loss	Forage versus feed crops?
	Industrial breeding	Largely in industrialized and transition countries	Waste pressure, water pollution, residue of veterinary medicine and antibiotics	Safe waste treatment and recycling
Agriculture in wetlands	Wetland drainage	Developing and transition countries	C loss	Agro-benefit versus natural value

Figure Legends


Figure 1. Maps of change in soil carbon due to land use change and land management from 1860 to 2010 from three vegetation models. Pink indicates loss of soil carbon, blue indicates carbon gain.

Figure 2. Soil carbon and nitrogen under different land cover types in three different vegetation models (values are the annual average over the period 2001 to 2010).


Figure 3. Uneven global distribution of soils sensitive to pollution by (a) acidification and (b) eutrophication (measured by soil C:N) compared to uneven distribution of atmospheric (c) sulphur and (d) nitrogen pollution. Soils most sensitive to acidification have low base saturation and cation exchange capacity, as defined by (Kuylenstierna *et al.*, 2001). Acidification is caused by both sulphur and nitrogen. Eutrophication is caused by nitrogen. Soil data in (a) and (b) were produced using the ISRIC-WISE derived soil properties (ver 1.2) (Batjes, 2012) and the FAO Digital Soil Map of the World. Atmospheric deposition data in (c) and (d) were provided by the World Data Centre for Precipitation Chemistry (<http://wdcpc.org>, 2014) and are also available in Vet *et al.* (2014). Data show the ensemble-mean values from the 21 global chemical transport models used by the Task Force on Hemispheric Transport of Air Pollution (HTAP) (Dentener *et al.*, 2006). Total wet and dry deposition values are presented for sulphur, oxidized and reduced nitrogen.



254x190mm (96 x 96 DPI)




254x190mm (96 x 96 DPI)




254x190mm (96 x 96 DPI)


(a) Soil sensitivity to acidification




(c) Atmospheric sulphur deposition (2001)



(b) Soil carbon to nitrogen ratio (C:N)



(d) Atmospheric nitrogen deposition (2001)



168x101mm (300 x 300 DPI)