

The effects of diesel exhaust pollution on floral volatiles and the consequences for honey bee olfaction

Article

Accepted Version

Lusebrink, I., Girling, R. D. ORCID: <https://orcid.org/0000-0001-8816-8075>, Farthing, E., Newman, T. A., Jackson, C. W. and Poppy, G. M. (2015) The effects of diesel exhaust pollution on floral volatiles and the consequences for honey bee olfaction. *Journal of Chemical Ecology*, 41 (10). pp. 904-912. ISSN 0098-0331 doi: <https://doi.org/10.1007/s10886-015-0624-4> Available at <https://centaur.reading.ac.uk/43963/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

Published version at: <http://dx.doi.org/10.1007/s10886-015-0624-4>

To link to this article DOI: <http://dx.doi.org/10.1007/s10886-015-0624-4>

Publisher: Springer Verlag

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online

THE EFFECTS OF DIESEL EXHAUST POLLUTION ON FLORAL VOLATILES AND
THE CONSEQUENCES FOR HONEY BEE OLFACTION

INKA LUSEBRINK ^{1,*}

¹*Centre for Biological Sciences*

Life Sciences Building 85

University of Southampton

Southampton

United Kingdom SO17 1BJ

e-mail: I.Lusebrink@soton.ac.uk

phone: +44 (023) 8059 4258

ROBBIE D. GIRLING ^{1,3}

¹*Centre for Biological Sciences*

Life Sciences Building 85

University of Southampton

Southampton

United Kingdom SO17 1BJ

³*Current address:*

School of Agriculture, Policy and Development

University of Reading

Reading

United Kingdom RG6 6AR

EMILY FARTHING ¹

¹*Centre for Biological Sciences*

Life Sciences Building 85

University of Southampton

Southampton

United Kingdom SO17 1BJ

TRACEY A. NEWMAN ²

² CES

Faculty of Medicine

Institute for Life Sciences

University of Southampton

Southampton

United Kingdom SO17 1BJ

CHRIS W. JACKSON ¹

¹*Centre for Biological Sciences*

Life Sciences Building 85

University of Southampton

Southampton

United Kingdom SO17 1BJ

GUY M. POPPY ¹

¹*Centre for Biological Sciences*

Life Sciences Building 85

University of Southampton

Southampton

United Kingdom SO17 1BJ

1 **Abstract** – There is growing evidence of a substantial decline in pollinators within Europe
2 and North America, most likely caused by multiple factors such as diseases, poor nutrition,
3 habitat loss, insecticides and environmental pollution. Diesel exhaust could be a contributing
4 factor to this decline, since we found that diesel exhaust rapidly degrades floral volatiles,
5 which honey bees require for flower recognition. In this study we exposed eight of the most
6 common floral volatiles to diesel exhaust in order to investigate, whether it can affect volatile
7 mediated plant-pollinator interaction. Exposure to diesel exhaust altered the blend of common
8 flower volatiles significantly: myrcene was considerably reduced, β -ocimene became
9 undetectable, and β -caryophyllene was transformed into its *cis*-isomer isocaryophyllene.
10 Proboscis extension response (PER) assays showed that the alterations of the blend reduced
11 the ability of honey bees to recognize it. The chemically reactive nitrogen oxides fraction of
12 diesel exhaust gas was identified as capable of causing degradation of floral volatiles.

13

14 **Key Words** - Floral scent compounds, diesel exhaust, nitrogen oxides, scent degradation,
15 scent recognition, proboscis extension response.

INTRODUCTION

16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41

Since the winter and spring of 2006-7 there have been reports of elevated colony losses of managed European honey bees (*Apis mellifera*) in the U.S.A., Europe, the Middle East, and Japan (Abrol 2012; Oldroyd 2007). Furthermore, there is growing evidence of substantial losses of many wild pollinator species worldwide, mainly within Europe and North America (Biesmeijer et al. 2006; Potts et al. 2010). Declines of managed honey bees and wild pollinators may have serious implications, particularly because global food security is considered to be dependent on animal pollination (Abrol 2012). Even though most staple food crops do not require insect pollination (Ghazoul 2005), 35% of the world crop production for human food depends on pollinators (Klein et al. 2007). Those pollinated crops are particularly important for our food diversity and add nutritional value to our diet (Steffan-Dewenter et al. 2005). Some crops, such as oilseed rape (*Brassica napus*), are considered self-fertile (Free 1993), but insect pollination can strongly contribute to increased yield and market value (Bommarco et al. 2012). About 73% of cultivated crop varieties are pollinated by some type of bee, with the European honey bee dominating crop pollination worldwide (Abrol 2012). The drivers behind the global pollinator decline are likely to be multifactorial and include fragmentation and loss of habitat, increased pesticide use, decreased resource diversity, alien species, spread of pathogens, and climate change (Epstein et al. 2013; Kerr et al. 2015; Oldroyd 2007; Potts et al. 2010, Vanbergen et al. 2013). However, it is unlikely that this is a definitive list, and it is, therefore, important to investigate additional potential stressors that could result in negative effects on bee fitness.

Bees use mixtures of olfactory and visual stimuli to find suitable host plants (Dötterl and Vereecken 2010). They rely mainly on olfactory cues during their initial foraging bouts, and visual cues become more important in host-plant location as bees gain more experience (Dobson 1994). However, floral scent remains an important stimulus for experienced bees,

42 since it allows them to discriminate between rewarding and non-rewarding flowers (Dobson
43 2006; Wright and Schiestel 2009).

44 Ozone or reactive species, such as hydroxyl and nitroxyl or nitrate radicals, readily
45 react with volatile organic compounds (VOC) (Atkinson and Arey 2003; Calogirou et al.
46 1999), which contribute to floral scent and could, therefore, impact upon VOC-mediated
47 plant-insect interactions. McFrederick et al. (2008) modelled the dispersion of three common
48 floral scent compounds (linalool, myrcene, and β -ocimene) under different air pollution
49 scenarios and concluded that increasing pollution levels may impair the recognition of floral
50 scents by pollinators.

51 In an earlier study we were able to show that diesel exhaust alters the VOC
52 composition of a synthetic floral odour blend (Girling et al. 2013) designed to mimic the
53 oilseed rape (OSR) cultivar *Brassica napus*, cv. Topas. It consisted of eight compounds,
54 which are behaviourally active in honey bees, mixed in their naturally occurring ratios (Blight
55 et al. 1997). The exposure of the OSR blend to diesel exhaust rendered two compounds
56 undetectable, which significantly reduced the ability of honey bees to recognize the altered
57 floral scent (Girling et al. 2013). Whether such changes deleteriously affect honey bee
58 foraging or foraging by other pollinators that utilise these compounds is currently unknown.
59 Three of the volatile compounds in OSR are floral scent compounds that occur in more than
60 50% of seed plants (reviewed by Knudsen et al. 2006).

61 In order to investigate further the effects of diesel exhaust emissions on floral VOCs
62 and to study whether such effects could be a more widespread phenomenon, we exposed eight
63 of the 12 most common floral scent compounds (Knudsen et al. 2006) to diesel exhaust, none
64 of which are found in OSR, mixed in equal amounts. To investigate if bees that were
65 conditioned to our floral volatile blend were able to still recognize it after they underwent the
66 alterations caused by diesel exhaust, we used the classical conditioning of the proboscis
67 extension reflex (PER; Bitterman et al. 1983; Kuwabara 1957). Additionally, we exposed all

68 eight common floral scent compounds individually to diesel exhaust with the intention of
69 determining their possible fate in a diesel exhaust polluted environment. In order to confirm
70 the general mechanism behind the alteration of floral scent components in such an
71 environment, we also exposed all compounds to various concentrations of NO_x, since that
72 was the observed mechanism behind the floral scent degradation in our earlier study (Girling
73 et al 2013).

74

75

76

METHODS AND MATERIALS

77

78 *Floral Volatile Blend.* Based on a review by Knudsen et al. (2006) on the diversity and
79 distribution of floral scent, we selected eight common floral compounds that occur in more
80 than half of all the families of seed plants: β -Pinene, myrcene, limonene, β -ocimene,
81 benzaldehyde, β -caryophyllene, methyl salicylate, and benzyl alcohol (Table 1). We mixed all
82 eight compounds in equal amounts to produce a common floral volatile (CFV) blend.

83

84 *Floral Volatile Exposures.* For the exposures the general procedure was to place 1 μ L of CFV
85 blend on a filter paper (2.1cm circle, grade 3MM; Whatman plc, Maidstone, UK) and a 4.5cm
86 stir bar in a 1000mL glass bottle (VWR International Ltd., West Sussex, UK). To assess the
87 influence of UV light the tests were replicated in both clear and amber glass bottles. The CFV
88 blend was exposed to either ambient air or diesel exhaust at room temperature. The ambient
89 air treatment consisted of a bottle of air collected from the laboratory. Diesel exhaust was
90 collected into the bottle from the exhaust pipe of a Suntom SDE 6500 E diesel generator
91 (Fuzhou Suntom Power Machinery Co., Ltd. Fuzhou, China) for 3min at a flow rate of
92 1L/min. Bottles were sealed with two layers of Parafilm M[®] (Bemis Flexible Packaging,
93 Oshkosh, WI, USA) and a lid. The lid contained a 1mm bore hole in its centre to allow

94 insertion of the solid-phase microextraction (SPME) fibre holder for the collection of
95 volatiles. The bottle content was mixed with a stir bar at 300rpm. The content of the bottle
96 was sampled after 1min, 30min, 1h and 2h. At each sample time point the SPME fibre was
97 extended into the bottle and exposed to the volatiles for 5min. The SPME fibre was then
98 inserted into the injector of the gas chromatograph linked to the mass spectrometer (GC/MS)
99 and allowed to desorb for 10min. This 10min desorption period was sufficient to clean the
100 SPME fibre so that the same fibre could be used to collect VOCs from the same bottle and
101 filter paper at a later time point. After the collection of VOC at all four time points the bottle
102 was cleaned with hexane, dried under nitrogen gas, and left in the fume hood overnight to
103 remove the hexane residue. Each of the air treatment and bottle type combinations were
104 repeated five times at room temperature for each of the two floral blends.

105

106 *Floral Blend GC/MS Analyses.* After each VOC adsorption period the SPME fibre assembly
107 (65µm PDMS/DVB, fused silica, 24Ga, manual holder; Supelco, Bellefonte, PA, USA) was
108 injected into a Hewlett Packard HP6890/5972A GC/MS at an injection port temperature of
109 250°C and a split ratio of 10:1. The GC/MS was equipped with a SPME injection sleeve (I.D.
110 0.75mm, Supelco) and HP-Innowax column (I.D. 0.25mm, length 30m; film thickness
111 0.25µm; Agilent Technologies, Santa Clara, CA, USA). The helium carrier gas flow was set
112 at 1.0mL/min, and the oven temperature was held for 2min at 50°C, increased to 70°C by 5°C
113 per min and then ramped up to 240°C by 10°C/min.

114 Peaks were identified using the same standards used to produce the floral volatile
115 blends (Table 1). Isocaryophyllene the rearrangement product of β-caryophyllene, was
116 identified based on catalytic conversion of β-caryophyllene in clove oil following US patent
117 3621070 (Rachlin 1971).

118

119 *Preparation of Bees.* Honey bees were kept on the University of Southampton campus in an
120 apiary (50° 56' 10"N, 1° 23' 39"W). During the summers of 2012 and 2013, on days when
121 bees showed high activity, 30-35 returning forager bees were caught at their hive entrance
122 between 14:00-16:00 GMT. Bees were caught in Sterilin 30mL universal containers (Sterilin
123 Limited, Cambridge, UK) and kept individually. They were immobilized on ice and
124 transferred into blue 1mL pipette tips that were cut to resemble the PER tubes used by
125 Bitterman et al. (1983) and harnessed to it with two strips of cloth tape (tesa[®] extra Power
126 Perfect). Harnessed bees could freely move their antennae, mouth parts and forelegs (Fig. 1).
127 The PER tubes containing bees were kept in Eppendorf centrifuge tube racks placed inside
128 plastic boxes with perforated lids. The bottom of the box was lined with wet tissue to provide
129 humidity and prevent the bees from desiccating. The bees were fed with a 30% sucrose
130 solution between 16:00-18:00 and kept in an environmentally controlled room at 20°C
131 overnight. PER assays were conducted the next morning for which bees were equally divided
132 into groups of 7-10 bees (depending on overnight survival).

133

134 *Proboscis Extension Response Assay.* One harnessed bee at a time was placed in the
135 experimental arena (W×D×H= 60cm×45cm×55cm) 3cm in front of an odour delivery system.
136 Behind the bee, an extraction fan removed the odour from the arena in order to avoid a build-
137 up of any stimuli. The odour delivery system was custom-built. It allowed for a constant
138 stream of fresh air in the arena through Teflon tubing which flowed into a glass tube. The
139 conditioning stimuli were delivered through a three channel system that ended in the same
140 glass tube as the fresh air, before it discharged into the arena. All three channels consisted of
141 electronic valves and Teflon tubing connected to 10mL glass tubes, in which the stimuli were
142 placed on a piece of filter paper. One channel served as the control (clean air), which was
143 always open when no volatiles were delivered. In order to deliver scented air, the control
144 channel was switched. The airflow passing through the odour delivery part of the system was

145 set at 300mL/min and the fresh air at 200mL/min air. Hence, the bees were constantly
146 exposed to an airflow of 500mL/min. The following timeline applied to the PER assay: A bee
147 was allowed to adapt to the experimental arena for 10s, then was presented with the
148 conditioning stimulus (CS) for 10s. Five seconds after the onset of CS, the antennae of the bee
149 was touched with a cocktail stick that was dipped in 30% sucrose solution (unconditioning
150 stimulus (US)). Upon extension of her proboscis, the bee was allowed to feed on the sucrose
151 solution for the remaining part of the 10s US period. The bee was then removed from the
152 arena. Thirty seconds was allowed between bees to give the extraction fan additional time to
153 clear the experimental area from any remaining odours.

154 The bees were conditioned six times to the CFV blend. During these conditioning
155 trials most bees learned to associate the sugar reward with the odour stimuli. Bees which had
156 learned to extend their proboscis in response to the odour stimuli by the sixth conditioning
157 trial were used in the subsequent recognition trials. The recognition trial resembled the
158 conditioning trial but without the use of the US. Its purpose was to test the proboscis response
159 of the bees to the original flower blends and to three different test stimuli for each blend
160 (Table 2). The test stimuli were based on the original blend but some chemical compounds
161 were omitted based on our results from the floral volatile exposure experiment. Extension of
162 the proboscis in response to the test stimulus was rated as a positive recognition. The PER
163 assay was repeated until $n \geq 25$ was achieved for each test group of bees. The results show the
164 percentage of bees that had successfully learned to recognize the floral odour blends and then
165 responded to either of the test stimuli.

166
167 *Fate of Individual Floral Volatiles.* The floral volatile exposure method described above was
168 altered slightly when investigating the fate of all individual compounds as these analyses were
169 conducted on a replacement Agilent Technologies 7890B/5977A GC/MS. Because of its
170 higher sensitivity only 0.2 μ L of each compound was placed on a filter paper and transferred

171 into an amber bottle with either ambient air or diesel exhaust. The content of the bottle was
172 sampled after 30min, and the SPME fibre was exposed to the volatile for 1min. The SPME
173 fibre was then inserted into the injection port of the GC/MS at 250°C and a split ratio of 10:1.
174 The GC/MS was equipped with a SPME injection sleeve, a HP-Innowax column (I.D.
175 0.25mm, length 30m; film thickness 0.25µm) and a retention gap (I.D. 0.18mm, length 4.1m,
176 film thickness 0µm; both Agilent Technologies, Santa Clara, CA, USA). The helium carrier
177 gas flow and oven temperature programme remained the same as for the floral blend analyses
178 described above.

179

180 *Diesel Exhaust Measurements.* In order to determine the concentration of toxic and reactive
181 gases in our diesel exhaust we used a Wolfsense TG501 probe equipped with O₂, NO, NO₂,
182 SO₂, and CO photochemical sensors (GrayWolf Sensing Solutions, Shelton, CT, USA).
183 Diesel exhaust was pumped through the probe at a flow rate of 1L/min by attaching a
184 calibration hood. Measured values were logged every 30s. Five 3min measurements were
185 taken to mimic the diesel exhaust collection for floral volatile exposures.

186

187 *Influence of Nitrogen Oxides.* For the exposures to nitric oxides (NO_x) a filter paper with 1µL
188 of the CFV blend and a stir bar were added to an amber 1000mL bottle. Nitrogen oxides were
189 produced by reducing nitric acid with elemental copper, which resulted in a 1:1 ratio of NO
190 and NO₂, which was confirmed by the Wolfsense TG501 probe. Concentrations of 20ppm,
191 2ppm, 0.2ppm per bottle were achieved by using gas tight syringes and volumetric
192 calculations. The bottle was closed with two layers of Parafilm M[®] and a lid. The content was
193 allowed to stir for 30min at 300rpm before the SPME fibre was extended into the bottle.
194 SPME exposure and the GC/MS method were the same as for floral blend volatile exposures.

195

196 *Statistical Analysis.* As we found that UV light had no statistically significant effect on the
197 composition of volatiles in our experiments, we pooled the data gained from exposures in
198 amber and clear glass bottles. Due to the large number of samples for the floral volatile
199 exposures, it was necessary to use more than one SPME fibre, which did not allow
200 comparative statistical analyses based on peak area according to the SPME guidelines of the
201 journal. However, the SPME fibres used were from the same lot, and the peak area data of the
202 floral volatile exposures had less variance than that of the NO_x exposure data that was
203 collected using a single SPME fibre. Therefore, we plotted the peak areas as bar graphs with
204 95% confidence interval error bars, which provides a graphical display of the significance of
205 differences of the CFV blend when exposed to either ambient air or diesel exhaust (Fig. 6,
206 online supplementary material).

207 The PER assay data was analysed by comparing the recognition of each test stimuli to
208 the response of bees to the complete CFV blend. Pairwise comparisons were made using χ^2 -
209 tests. The criterion for significance was corrected after Bonferroni: $\alpha' = \alpha/k$, where k is the
210 number of comparisons.

211 The influence of NO_x on the CFV blend was measured with the same SPME fibre, so
212 we were able to directly compare the effect of the different NO_x concentrations on the
213 abundance of compounds. The peak area data was analysed with a MANOVA followed by
214 univariate ANOVAs for each compound and Tukey post-hoc tests for comparing the
215 influence of different NO_x concentrations on the single compounds. All statistical analyses
216 were conducted with SPSS 20.0 for Windows (IBM Corporation, Armonk, New York, USA).

217

218

219

RESULTS

220

221 *Floral Volatile Exposures.* At the one minute time point of diesel exhaust exposure of the
222 CFV blend the compounds myrcene, *Z*- and *E*- β -ocimene were drastically reduced, and
223 β -caryophyllene started to be transformed into its *cis*-isomer, isocaryophyllene (Fig. 2). After
224 30min, when all compounds in the mixture had reached equilibrium in the glass bottle,
225 myrcene remained strongly reduced, whereas *Z*- and *E*- β -ocimene (assignment of geometrical
226 isomers followed Babushok et al. 2011) became undetectable, and β -caryophyllene was
227 completely transformed into isocaryophyllene.

228

229 *Proboscis Extension Response Assays.* The recognition assay for the CFV blend revealed that
230 the PER response of the honey bees was significantly influenced by the test stimuli
231 composition. The original CFV blend was recognized by 93% of all bees in the absence of a
232 sugar reward (Fig. 3). The CFV blend without β -ocimene (CFV-1) was still recognized by
233 76% of bees compared to the original blend, which is non-significant ($\chi^2_{(1)} = 2.928$, $P =$
234 0.092). Omission of myrcene (CFV-2) had a significant effect on the bees' PER response, and
235 only 37% still recognized the test stimuli ($\chi^2_{(1)} = 18.941$, $P < 0.001$). The test stimuli which
236 lacked β -ocimene and myrcene (CFV-3) was recognized by 39% of the bees, which is
237 significantly lower than the recognition of the full CFV blend ($\chi^2_{(1)} = 17.923$, $P < 0.001$).

238

239 *Fate of Individual Floral Volatiles.* Investigation on the effect of diesel exhaust on each single
240 floral volatile showed that β -caryophyllene was transformed into its *cis*-isomer, confirming
241 the results that we had seen when we exposed the complete CFV blend to diesel exhaust.
242 These further investigations revealed that the β -caryophyllene standard sample also contained
243 α -copaene and α -humulene as impurities (verified by standard injection). The amount of both
244 of them was also drastically reduced after being exposed to diesel exhaust (see supplementary
245 material, Fig. 6), however, we were unable to identify any degradation products. Similarly, we

246 could not trace the fate of myrcene and β -ocimene, the two acyclic monoterpenes in the CFV
247 blend. Benzyl alcohol was partly oxidized to benzaldehyde.

248
249 *Diesel Exhaust Measurement.* The averages of all measurements of all toxic and reactive
250 gases \pm S.E. acquired with the Wolfsense TG501 probe are given in Table 3. Carbon
251 monoxide represented the biggest fraction of the emission gases. Within the NO_x fraction, NO
252 and NO_2 were produced approximately at a ratio of 1:1. No sulphur dioxide was detected
253 which is most likely due to the fact that we used low sulphur diesel.

254
255 *Influence of Nitrogen Oxides.* The abundances of myrcene, *Z*- β -ocimene, *E*- β -ocimene, β -
256 caryophyllene, and benzyl alcohol were altered significantly by NO_x (Table 4). In particular,
257 the abundances of myrcene, *Z*- β -ocimene, *E*- β -ocimene, β -caryophyllene, and benzyl alcohol
258 were significantly reduced when exposed to 20ppm NO_x (Fig. 4). The conversion of β -
259 caryophyllene into isocaryophyllene began at a concentration of 0.2ppm NO_x .

260

261

262

DISCUSSION

263

264 Exposure to diesel exhaust led to a reduction of myrcene in the CFV blend, β -ocimene
265 disappeared, and β -caryophyllene was converted into its *cis*-isomer, isocaryophyllene.

266 Investigations on nitration mechanisms have shown that the reaction of unsaturated
267 compounds with nitrogen dioxide, which is abundant in diesel exhaust, can lead to *cis/trans*-
268 isomerization (Augusto et al. 2002; Titov 1963). In the blend used for the behavioural
269 recognition assays we omitted myrcene and β -ocimene. The lack of β -ocimene did not
270 influence the recognition of the test stimulus significantly. When myrcene was absent,
271 recognition decreased significantly to only 37%. Similarly, when both compounds were

272 missing 39% of honey bees recognized the test stimuli, suggesting that the absence of
273 myrcene predominantly contributes to the lack of recognition. Similarly, Reinhard et al.
274 (2010) identified myrcene as a key odorant and ocimene as a non-key odorant in the
275 recognition of complex scent mixtures by honey bees. Isocaryophyllene was not available for
276 the behavioural assays, however, we hypothesise that the change from β -caryophyllene to
277 isocaryophyllene could also have changed the insects' odour perception of the CFV blend.
278 Geometric isomers may differ in their quality of odour (Roderick 1966), e.g. to humans, the
279 cyclic sesquiterpene (*E*)- γ -bisabolene smells soapy and spicy, whereas the odour of (*Z*)- γ -
280 bisabolene is described as fatty and woody (Kjeldsen et al. 2003). The discrimination of
281 geometric isomers by insects has been mainly studied in the context of straight chain sex-
282 pheromones of moths (e.g. Klun et al. 1973; Naka et al. 2013; Silverstein and Young 1976).

283 Investigations into the fates of individual flower volatiles could not detect substantial
284 amounts (if any) of degradation products of the acyclic terpenes myrcene and β -ocimene. On
285 their oxidative way to carbon dioxide, most biogenetic volatiles pass acetone and
286 formaldehyde (Atkinson and Arey 2003), which, due to their low boiling points, would have
287 been difficult to find by our analytical method. However, primary oxidation products of
288 myrcene such as 4-methylene-5-hexenal and of β -ocimene such as 4-methyl-3,5-hexadienal
289 (Lee et al. 2006, Reissell et al. 2002) should have been reliably registered.

290 Each of the floral scent compounds used in our study that degraded when exposed to
291 diesel exhaust also occur in numerous other plant families: β -caryophyllene in 52% (46
292 families), β -ocimene in 71% (*E*- β -ocimene in 64 families and *Z*- β -ocimene in 44 families),
293 and myrcene in 70% (63 families) of all investigated seed plant families (Knudsen et al.
294 2006). Half of the highly ubiquitous flower scent compounds that we tested experienced
295 significant reduction as a result of exposure to diesel exhaust, indicating that it has the
296 potential to disrupt and modify plant floral volatile signalling at a broad scale.

297 Diesel exhaust emissions have received considerable attention due to their health risks
298 and environmental concerns, the main focus has been on particulate matter and NO_x
299 (Setiabudi et al. 2004), which are considered to be the main air pollutants of diesel exhaust
300 together with carbon monoxide and hydrocarbons. The NO_x concentrations of our diesel
301 exhaust were approximately 20ppm for both NO and NO₂. When we exposed the floral
302 volatiles to 20ppm NO_x we observed that it had a similar effect as diesel exhaust: β-
303 caryophyllene was turned into isocaryophyllene, the amount of myrcene was reduced, β-
304 ocimene was reduced as well, but still detectable, whereas it had completely disappeared
305 when exposed to diesel exhaust. In contrast to oxygenation involving oxygen species, radical
306 reactions induced by NO_x appear to be responsible for the majority of the changes observed in
307 our CFV blend when exposed to diesel exhaust. In addition, the particulate matter in diesel
308 exhaust may have yet unknown adsorptive and catalytic properties. Because of their isoprene
309 (butadiene) substructure, myrcene and β-ocimene are particularly labile, and radical reactions
310 may easily cause polymerisation, leading to products that would escape our analytical system.

311 Ambient air quality standards in the EU (ec.europa.eu) and the US (www.epa.gov)
312 suggest a maximum hourly average NO₂ concentration of 0.1ppm, a concentration at which β-
313 caryophyllene started turning into isocaryophyllene in this study, and α-terpinene and
314 phenylacetaldehyde were already significantly degraded, in our previous study. Furthermore,
315 in 2013 the UK Supreme Court declared that the nitrogen dioxide limits are regularly
316 exceeded in 16 zones across the UK (European Commission, press release).

317 Ambient NO_x levels can exhibit strong diurnal patterns, with maximum values in the
318 early morning and another increase in the late afternoon (Fuentes et al. 2007), following
319 periods of high traffic. Fluctuating NO_x levels will change the ratio of compounds in floral
320 scent throughout the course of the day (McFrederick et al. 2008). Additionally, high NO_x
321 levels lead to increased ozone levels (Fehsenfeld et al. 1992), which may further contribute to
322 the degradation of floral scent. Changing floral scent of a flower can cause floral-constant

323 pollinators, like honey bees, to reject the flower (Wright and Schiestl 2009), which may
324 impact on pollination efficiency. To date, there is only little research on how air pollution
325 directly impedes on plant-insect interactions. Fuentes et al. (2013) found that ozone pollution
326 obstructs host finding in the cucumber beetle (*Acalymma vittatum*). Pinto et al. (2007a; 2007b;
327 2008) investigated the impacts of ozone on tritrophic interactions under both laboratory and
328 field conditions. They showed that parasitoids are able to use herbivore-induced plant VOCs
329 to find their hosts in the presence of ozone, but preferred the intact signal over the ozone-
330 degraded signal. There is also a possibility that the reaction products of floral scent
331 compounds with ozone and other air pollutants might form new compounds that may serve as
332 novel signals (McFrederick et al. 2009).

333 Our study demonstrates that NO_x and/or other components of diesel exhaust are
334 capable of degrading floral signals and have the potential to alter floral recognition. How this
335 impacts upon the fitness of pollinators requires further investigations, however it has been
336 proposed that pollinators in polluted areas may need to spend more time searching for
337 adequate host plants, which if true could lead to decreased fitness and decreased pollination
338 rates (McFrederick et al. 2008).

339

340

341 **Acknowledgements**

342 We acknowledge Wittko Francke for his valuable comments on the manuscript, Christine M.
343 Reitmayer for her help with beekeeping. We would also like to thank the Erbilgin lab at the
344 University of Alberta for providing us with a *trans*-verbenol mass spectrum. Funding for this
345 research has been provided through the Leverhulme Trust Research Project Grant RPG-089.

346

347

348

REFERENCES

349 Abrol DP (2012) Decline in pollinators. In: Pollination Biology: Biodiversity Conservation
350 and Agricultural Production. Springer, Netherlands, pp 545-601 doi:10.1007/978-94-007-
351 1942-2_17

352 Atkinson R, Arey J (2003) Gas-phase tropospheric chemistry of biogenic volatile organic
353 compounds: a review. Atmospheric Environment 37, Supplement 2:197-219
354 doi:10.1016/s1352-2310(03)00391-1

355 Augusto O, Bonini MG, Amanso AM, Linares E, Santos CCX, De Menezes SL (2002)
356 Nitrogen dioxide and carbonate radical anion: two emerging radicals in biology. *Free Radical*
357 *Bio Med* 32:841-859 doi:10.1016/S0891-5849(02)00786-4

358 Babushok VI, Linstrom PJ, Zenkevich IG (2011) Retention indices for frequently reported
359 compounds of plant essential oils. *J Phys Chem Ref Data* 40:043101 doi:10.1063/1.3653552

360 Biesmeijer JC, Roberts SPM, Reemer M, Ohlemüller R, Edwards M, Peeters T, Schaffers AP,
361 Potts SG, Kleukers R, Thomas CD, Settele J, Kunin WE (2006) Parallel declines in
362 pollinators and insect-pollinated plants in Britain and the Netherlands. *Science* 313:351-354
363 doi:10.1126/science.1127863

364 Bitterman ME, Menzel R, Fietz A, Schafer S (1983) Classical conditioning of proboscis
365 extension in honeybees (*Apis mellifera*). *J Comp Psychol* 97:107-119

366 Blight MM, Métayer ML, Pham-Delègue MH, Pickett JA, Marion-Poll F, Wadhams LJ
367 (1997) Identification of floral volatiles involved in recognition of oilseed rape flowers,
368 *Brassica napus* by honeybees, *Apis mellifera*. *J Chem Ecol* 23:1715-1727
369 doi:10.1023/B:JOEC.0000006446.21160.c1

370 Bommarco R, Marini L, Vaissière B (2012) Insect pollination enhances seed yield, quality,
371 and market value in oilseed rape. *Oecologia* 169:1025-1032 doi:10.1007/s00442-012-2271-6

372 Calogirou A, Larsen BR, Kotzias D (1999) Gas-phase terpene oxidation products: a review.
373 *Atmos Environ* 33:1423-1439 doi:10.1016/s1352-2310(98)00277-5

374 Dobson HEM (1994). Floral volatiles in insect biology. In: Bernays EA (ed) *Insect-plant*
375 *interactions*. CRC Press, London. pp. 47–81.

376 Dobson HEM (2006) Relationship between floral fragrance composition and type of
377 pollinator. In: Dudareva N and Pichersky E (eds) *Biology of Floral Scent*. CRC Press, pp 147-
378 198 doi:10.1201/9781420004007.sec4

379 Dötterl S, Vereecken NJ (2010) The chemical ecology and evolution of bee-flower
380 interactions: a review and perspectives. *Can J Zool* 88:668-697 doi:10.1139/Z10-031

381 [ec.europa.eu] European Commission press release: Environment: Commission takes action
382 against UK for persistent air pollution problems. [http://europa.eu/rapid/press-release_IP-14-](http://europa.eu/rapid/press-release_IP-14-154_en.htm)
383 [154_en.htm](http://europa.eu/rapid/press-release_IP-14-154_en.htm)

384 Epstein D, Frazier JL, Purcell-Miramontes M, Hackett K, Rose R, Erickson T, Moriarty T,
385 Steeger T (2013) USDA: Report on the national stakeholders conference on honey bee health.
386 <http://www.usda.gov/documents/ReportHoneyBeeHealth.pdf>. Accessed 27 August 2015

387 Fehsenfeld F, Calvert J, Fall R, Goldan P, Guenther AB, Hewitt CN, Lamb B, Liu S, Trainer
388 M, Westberg H, Zimmerman P (1992) Emissions of volatile organic compounds from
389 vegetation and the implications for atmospheric chemistry. *Global Biogeochemical Cycles*
390 6:389-430 doi:10.1029/92GB02125

391 Fuentes J, Wang D, Bowling D, Potosnak M, Monson R, Goliff W, Stockwell W (2007)
392 Biogenic hydrocarbon chemistry within and above a mixed deciduous forest. *J Atmos Chem*
393 56 (2):165-185 doi:10.1007/s10874-006-9048-4

394 Fuentes JD, Roulston TaH, Zenker J (2013) Ozone impedes the ability of a herbivore to find
395 its host. *Environ Res Letters* 8 (1):014048 doi:10.1088/1748-9326/8/1/014048

396 Free JB (1993) *Insect pollination of crops*. 2nd edn. Academic Press, London; New York

397 Ghazoul J (2005) Buzziness as usual? Questioning the global pollination crisis. *Trends Ecol*
398 *Evol* 20:367-373 doi:10.1016/j.tree.2005.04.026

399 Girling RD, Lusebrink I, Farthing E, Newman TA, Poppy GM (2013) Diesel exhaust rapidly
400 degrades floral odours used by honeybees. *Sci Rep* doi:10.1038/srep02779

401 Kerr JT, Pindar A, Galpern P, Packer L, Potts SG, Roberts SM, Rasmont P, Schweiger O,
402 Colla SR, Richardson LL, Wagner DL, Gall LF, Sikes DS, Pantoja A (2015) Climate change
403 impacts on bumblebees converge across continents *Science* 349:177-180
404 doi:10.1126/science.aaa7031

405 Kjeldsen F, Christensen LP, Edelenbos M (2003) Changes in Volatile Compounds of Carrots
406 (*Daucus carota* L.) During Refrigerated and Frozen Storage *J Agr Food Chem* 51:5400-5407
407 doi:10.1021/jf030212q

408 Klein A-M, Vaissière BE, Cane JH, Steffan-Dewenter I, Cunningham SA, Kremen C,
409 Tscharntke T (2007) Importance of pollinators in changing landscapes for world crops. *P Roy*
410 *Soc Lond B Bio* 274:303-313 doi:10.1098/rspb.2006.3721

411 Klun JA, Chapman OL, Mattes KC, Wojtkowski PW, Beroza M, Sonnet PE (1973) Insect Sex
412 Pheromones: Minor Amount of Opposite Geometrical Isomer Critical to Attraction *Science*
413 181:661-663 doi:10.1126/science.181.4100.661

414 Knudsen J, Eriksson R, Gershenzon J, Ståhl B (2006) Diversity and distribution of floral
415 scent. *Bot Rev* 72:1-120 doi:10.1663/0006-8101(2006)72[1:dadofs]2.0.co;2

416 Kuwabara M (1957) Bildung des bedingten Reflexes von Pavlovs Typus bei der Honigbiene,
417 *Apis mellifica*. *Journal of the Faculty of Science, Hokkaido University, Series 6, Zoology,*
418 13:458-464

419 Lee A, Goldstein AH, Keywood MD, Gao S, Varutbangkul V, Bahreini R, Mg NL, Flagan
420 RC, Seinfeld JH (2006) Gas-phase products and secondary aerosol yields from the ozonolysis
421 of ten different terpenes. *J Geophysical Res Atmospheres* 111 doi: 10.1029/2005JD006437

422 McFrederick QS, Fuentes JD, Roulston T, Kathilankal JC, Lerdau M (2009) Effects of air
423 pollution on biogenic volatiles and ecological interactions. *Oecologia* 160:411-420
424 doi:10.1007/s00442-009-1318-9

425 McFrederick QS, Kathilankal JC, Fuentes JD (2008) Air pollution modifies floral scent trails.
426 *Atmos Environ* 42:2336-2348 doi:10.1016/j.atmosenv.2007.12.033

427 Naka H, Suzuki T, Watarai T, Horie Y, Mochizuki F, Mochizuki A, Tsuchida K, Arita Y,
428 Ando T (2013) Identification of the sex pheromone secreted by *Synanthedon tenuis*
429 (Lepidoptera: Sesiidae) *Appl Entomol Zool* 48:27-33 doi:10.1007/s13355-012-0148-0

430 Oldroyd BP (2007) What's killing American honey bees? *PLoS Biol* 5:e168
431 doi:10.1371/journal.pbio.0050168

432 Pinto D, Blande J, Nykänen R, Dong W-X, Nerg A-M, Holopainen J (2007a) Ozone degrades
433 common herbivore-induced plant volatiles: Does this affect herbivore prey location by
434 predators and parasitoids? *J Chem Ecol* 33:683-694 doi:10.1007/s10886-007-9255-8

435 Pinto D, Nerg A-M, Holopainen J (2007b) The role of ozone-reactive compounds, terpenes,
436 and green leaf volatiles (GLVs), in the orientation of *Cotesia plutellae*. *J Chem Ecol* 33:2218-
437 2228 doi:10.1007/s10886-007-9376-0

438 Pinto DM, Himanen SJ, Nissinen A, Nerg AM, Holopainen JK (2008) Host location behavior
439 of *Cotesia plutellae* Kurdjumov (Hymenoptera: Braconidae) in ambient and moderately
440 elevated ozone in field conditions. *Environ Pollut* 156:227-231
441 doi:10.1016/j.envpol.2007.12.009

442 Potts SG, Biesmeijer JC, Kremen C, Neumann P, Schweiger O, Kunin WE (2010) Global
443 pollinator declines: trends, impacts and drivers. *Trends Ecol Evol* 25:345-353
444 doi:10.1016/j.tree.2010.01.007

445 Rachlin S (1971) Process for preparing isocaryophyllene. US Patent 3621070, 192.14, R448

446 Reinhard J, Sinclair M, Srinivasan MV, Claudianos C (2010) Honeybees learn odour mixtures
447 via a selection of key odorants. *PLoS ONE* 5:e9110 doi:10.1371/journal.pone.0009110

448 Reissell A, Aschmann SM, Atkinson R, Arey J (2002) Products of the OH radical- and O₃-
449 initiated reactions of myrcene and ocimene. *J Geophysical Res* 107:4138 doi:
450 10.1029/2001JD001234

451 Roderick WR (1966) Current ideas on the chemical basis of olfaction. *J Chem Education*
452 43:510 doi:10.1021/ed043p510

453 Setiabudi A, Makkee M, Moulijn JA (2004) The role of NO₂ and O₂ in the accelerated
454 combustion of soot in diesel exhaust gases. *Appl Catal B Environ* 50:185-194
455 doi:10.1016/j.apcatb.2004.01.004

456 Silverstein RM, Young JC (1976) Insects generally use multicomponent pheromones. In:
457 Beroza, M (ed) *Pest Management with Insect Sex Attractants*; ACS Symposium Series;
458 American Chemical Society: Washington, DC, 1976:1-29

459 Steffan-Dewenter I, Potts SG, Packer L (2005) Pollinator diversity and crop pollination
460 services are at risk. *Trends Ecol Evol* 20:651-652 doi:10.1016/j.tree.2005.09.004

461 Titov AI (1963) The free radical mechanism of nitration. *Tetrahedron* 19:557-580
462 doi:10.1016/S0040-4020(01)98544-8

463 Vanbergen AJ, the Insect Pollinators Initiative (2013) Threats to an ecosystem service:
464 pressures on pollinators. *Front Ecol Environ* 11 (5):251-259 doi:10.1890/120126

465 Wright GA, Schiestl FP (2009) The evolution of floral scent: the influence of olfactory
466 learning by insect pollinators on the honest signalling of floral rewards. *Funct Ecol* 23:841-
467 851 doi:10.1111/j.1365-2435.2009.01627.x

468 [www.epa.gov] United States Environmental Protection Agency. National Ambient Air
469 Quality Standards. <http://www.epa.gov/air/criteria.html>

470 **Table 1** Information about compound purity, vapour pressure, supplier information and their occurrence in seed
 471 plant families as reported in Knudsen et al. 2006

Common flower volatile blend				
compound	purity (%)	vapour pressure (mm Hg⁻¹ at 25°C)	supplier	occurrence
β-Pinene	98	2.9300	Sigma Aldrich ^a	59%
Myrcene	90	2.0900	Sigma Aldrich ^a	70%
Limonene	97	1.9800	Sigma Aldrich ^a	71%
β-Ocimene	≥ 90	1.5590	Sigma Aldrich ^a	71%
Benzaldehyde	99	0.1270	Sigma Aldrich ^a	64%
β-Caryophyllene	≥ 80	0.0130	Sigma Aldrich ^a	52%
Methyl salicylate	≥ 99	0.0343	Fluka ^b	57%
Benzyl alcohol	99	0.0940	Sigma Aldrich ^a	56%

472 ^a Sigma Aldrich (St. Louis, MO, USA), ^b Fluka (Sigma-Aldrich, Buchs, Switzerland)

473 **Table 2** Compound composition of the common floral volatile and the oilseed rape blend and their according test
 474 stimuli for the proboscis extension response assays

Common flower volatile blend				
compound	CFV	CFV-1	CFV-2	CFV-3
β-Pinene	1 μL	1 μL	1 μL	1 μL
Myrcene ^a	1 μL	1 μL	-	-
Limonene	1 μL	1 μL	1 μL	1 μL
β-Ocimene ^b	1 μL	-	1 μL	-
Benzaldehyde	1 μL	1 μL	1 μL	1 μL
β-Caryophyllene ^c	1 μL	1 μL	1 μL	1 μL
Methyl salicylate	1 μL	1 μL	1 μL	1 μL
Benzyl alcohol	1 μL	1 μL	1 μL	1 μL
dose	8 μL	7 μL	7 μL	6 μL

475 ^a compound reduced by diesel exhaust, ^b compounds erased by diesel exhaust, ^c compound turns into its *cis*-isomer

476 **Table 3** Diesel exhaust gas composition

gas (unit)	mean	S.E.
Oxygen (%)	19.78 ±	0.10
Nitric Oxide (ppm)	19.84 ±	2.27
Nitrogen Dioxide (ppm)	17.54 ±	1.25
Sulphur Dioxide (ppm)	0.00 ±	0.00
Carbon Monoxide (ppm)	226.84 ±	20.47

477

478 **Table 4** ANOVA results for single compounds which abundances were significantly decreased by NO_x exposure
479 (also see Fig. 4)

compound	<i>F</i> -value	P-value
Myrcene	$F_{(3,12)} = 11.803$	0.001
<i>Z</i> - β -Ocimene	$F_{(3,12)} = 89.226$	< 0.001
<i>E</i> - β -Ocimene	$F_{(3,12)} = 96.951$	< 0.001
β -Caryophyllene	$F_{(3,12)} = 568.459$	< 0.001
Benzyl alcohol	$F_{(3,12)} = 9.184$	0.002

480

481 **Fig. 1** Harnessed honey bee ready for PER assay

482

483 **Fig. 2** Total ion current (TIC) chromatogram of the common flower volatile under 30min
484 ambient air and diesel exhaust exposure. Consecutive numbers represent the following
485 chemical compounds: β -Pinene (1), myrcene (2), limonene (3), *Z*- β -ocimene (4a),
486 *E*- β -ocimene (4b), benzaldehyde (5), β -caryophyllene (6a), isocaryophyllene (6b) methyl
487 salicylate (7), benzyl alcohol (8).


488

489 **Fig. 3** Proboscis extension response assay results representing the ability of honey bees to
490 recognize the common floral volatile (CFV) blend, CFV blend without β -ocimene (CFV-1),
491 CFV blend without myrcene (CFV-2), and CFV blend without both (CFV-3). The response to
492 the test stimuli was compared to the response to the full blend. A Bonferroni correction was
493 applied, and all significant differences are indicated with an asterisk at a 0.016 level of
494 significance


495

496 **Fig. 4** The effect of 30min exposures of different nitrogen oxides ($\text{NO}_x = \text{NO}$ and NO_2 at a
497 1:1 ratio) concentrations on the abundance of all floral compounds used in this study. Bars
498 represent the average peak area (\pm S.D). Different lowercase letters indicate a statistically
499 significant difference.


500




501
502 Fig. 1



503
504 Fig. 2
505



506
507 Fig. 3
508



509 Fig. 4
510