

Extractability and characteristics of proteins deriving from wheat DDGS

Article

Accepted Version

Chatzifragkou, A. ORCID: <https://orcid.org/0000-0002-9255-7871>, Prabhakumari, P. C., Kosik, O., Lovegrove, A., Shewry, P. R. and Charalampopoulos, D. ORCID: <https://orcid.org/0000-0003-1269-8402> (2016) Extractability and characteristics of proteins deriving from wheat DDGS. Food Chemistry, 198. pp. 12-19. ISSN 0308-8146 doi: 10.1016/j.foodchem.2015.11.036 Available at <https://centaur.reading.ac.uk/46416/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

To link to this article DOI: <http://dx.doi.org/10.1016/j.foodchem.2015.11.036>

Publisher: Elsevier

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online

1 **Extractability and characteristics of proteins deriving from wheat**
2 **DDGS**

3 Afroditi Chatzifragkou¹, Parvathy Chandran Prabhakumari¹, Ondrej Kosik², Alison
4 Lovegrove², Peter R. Shewry² and Dimitrios Charalampopoulos^{1*}

5 ¹Department of Food and Nutritional Sciences, University of Reading, Whiteknights,
6 PO Box 226, Reading RG6 6AP, UK

7 ²Department of Plant Science, Rothamsted Research, Harpenden, AL5 2JQ,
8 Hertfordshire, UK

9

10 *Corresponding author: d.charalampopoulos@reading.ac.uk

11

Abstract

Wheat Distillers' Dried Grains with Solubles (DDGS) and in-process samples were used for protein extraction. Prolamins were the predominant protein components in the samples. The absence of extractable α - and γ -gliadins in DDGS indicated protein aggregation during the drum drying processing stage. Prolamin extraction was performed using 70% (v/v) ethanol or alkaline-ethanol solution in the presence of reducing agent. DDGS extracts had relatively low protein contents (14-44.9%, w/w), regardless of the condition applied. The wet solids were the most suitable raw material for protein extraction, with recovery yields of ~ 55% (w/w) and protein content of ~58% (w/w) in 70% (v/v) ethanol. Protein extracts from wet solids were significantly rich in glutamic acid and proline. Mass balance calculations demonstrated the high carbohydrate content (~ 50%, w/w) of solid residues. Overall, the feasibility of utilising in-process samples of DDGS for protein extraction with commercial potential was demonstrated.

Keywords: DDGS, protein, wheat, distillery, extraction, in-process samples, amino acids

1. Introduction

Distillers' Dried Grains with Solubles (DDGS) is the principal by-product of the dry-grind distillation process, generated mainly from beverage alcohol plants (e.g. whisky and neutral spirits distilleries) or from grain-based fuel-ethanol plants. In the case of distilleries, single or blended grains including wheat, barley, maize and rye can be utilised as feedstock, whereas fuel-ethanol plants use either corn (maize) (US) or wheat (Europe) as starting materials.

During the dry-grind process, in the case of bioethanol production, whole grains are milled and liquefied, followed by the addition of amylolytic enzymes for starch conversion into fermentable glucose. In distillery plants, saccharification of the milled grain is carried out using malted barley instead of external enzymes and a food-grade process is followed, as the end-product (potable ethanol) is intended for human consumption. For both bioethanol and potable ethanol production, yeast is added to ferment the sugars into ethanol and carbon dioxide. At the end of the fermentation, the whole stillage undergoes distillation by direct steam injection. Ethanol is further purified via dehydration, whereas the non-volatile components (spent solids) are centrifuged to produce a liquid fraction (thin stillage) and a solid fraction (wet solids). Around 15% or more of the thin stillage is recycled to the liquefaction process of the ground grain, whereas the remaining is concentrated in a series of steam driven evaporators, mixed with wet solids and drum dried to produce the final DDGS (Kim et al. 2008; Liu, 2011). The drying process applied at the last stage is intensive, as the air temperature can be over 500 °C at the dryer inlet and over 100 °C at the dryer outlet. Partial recycling of DDGS to the drum dryer can also occur in order to increase the drying efficiency of the equipment and improve the consistency of the produced

53 DDGS (Kingsly et al. 2010). Overall, for 100 kg of grain approximately 40 litres of
54 ethanol, 32 kg of DDGS and 32 kg of CO₂ are generated (Schingoethe, 2006).

55 Because it is enriched in protein, as well as in water-soluble vitamins and minerals,
56 DDGS has been long marketed as feed for livestock (including poultry) (Klopfenstein,
57 Erickson & Bremer, 2008; Schingoethe, Kalscheur, Hippen & Garcia, 2009). DDGS
58 derived from wheat contains around 28-38% (w/w) of protein, whereas for maize
59 DDGS the protein levels range within 28-31% (w/w) (Kim et al. 2010). The major
60 parameters influencing the cost-effectiveness of bioethanol production from cereal
61 grains include the cost of raw materials, as well as the revenue derived from DDGS.
62 In Europe, bioethanol production is currently driven by the EU mandates on biofuel
63 framework (Directive 2009/28/EC), thus the increased bioethanol demand is likely to
64 result in increased DDGS availability. As a result, current research is focused on
65 identifying alternative uses of DDGS, other than animal feed. To this end, existing
66 bioethanol or distillery companies could implement a biorefinery approach, where
67 DDGS is fractionated into several added value compounds including proteins,
68 carbohydrates and phytochemicals.

69 In contrast to the literature on maize DDGS, a limited number of studies have
70 investigated the extraction of protein from wheat-based DDGS (Bandara, Che & Wu,
71 2011; Hong, Avramenko, Stone, Abbott, Classen & Nickerson, 2012; Xu, Reddy
72 & Yang, 2007). Wheat grains contain gluten proteins that account for 80% of the total
73 wheat protein, with the remaining 20% corresponding to a heterogeneous group of
74 structural and metabolic proteins, including a major group of water soluble
75 components with molecular weight (MW) lower than 25 kDa (Veraverbeke &
76 Delcour, 2002). By contrast, gluten proteins are largely insoluble in water due to their

77 high non-polar amino acids content (in particular proline and glutamine) and serve as
78 storage reserves in the wheat grain (prolamins) (Shewry, 1999). Prolamins comprise
79 both alcohol-soluble monomers (gliadins) and alcohol-insoluble polymers (glutenins)
80 with the individual glutenin subunits being alcohol-soluble in their reduced state.
81 Prolamin monomers and subunits show considerable diversity in MW, ranging from
82 10 to 100 kDa (Shewry & Halford, 2002). The extraction of proteins from DDGS at
83 high yield and purity remains a challenge; DDGS proteins often show low
84 extractability possibly due to the intensive heating applied at the final stage of the
85 production process. Looking towards potential applications, DDGS proteins can be
86 exploited for the production of biodegradable films, coatings and biodegradable
87 plastics, which can be used for food, agricultural and industrial applications (Day,
88 Augustin, Batey & Wrigley, 2006). Wheat protein (gluten) has therefore been
89 extensively studied as a natural starting material for the development of biodegradable
90 films, due to its remarkable cohesive and elastic properties, as well as its susceptibility
91 to chemical modifications (Irissin-Mangata, Bauduin, Boutevin & Gontard, 2001;
92 Kuktaine et al. 2011). Further applications of gluten include in aquaculture feed and in
93 pet food, as an adhesive material in tapes and medical bandages, or as a biodegradable
94 polymer material for the slow release of pesticides or fertilising agents (Day et al.
95 2006; Majeed, Ramli, Mansor & Man, 2015).

96 The aim of this study was to investigate the extractability of proteins from various
97 samples originating from a distillery plant, i.e. wheat DDGS, wet and spent solids (the
98 latter also known as whole stillage). The composition of the extracted proteins and
99 their amino acid content were determined and are discussed in order to evaluate the

effect of the multi-step DDGS production process on the properties of the proteins at each stage of production.

2. Materials and methods

2.1 Raw materials

Distillers' Dried Grains with Solubles (DDGS) and in-process samples of wet solids and spent solids were kindly provided by a distillery plant in UK. The distillery plant uses a mixture of 95% (w/w) wheat and 5% (w/w) barley as starting material for potable ethanol manufacture. After being received, samples were frozen at -80 °C. After determination of their moisture content (Section 2.2), samples were lyophilised in a VirTis Bench Top (USA) freeze-drier, initially set at -55 °C for 48 h, packed in polyethylene bags and subsequently stored at -20 °C, until further analysis.

2.2 Compositional analysis of samples

All samples were milled using a conventional coffee grinder in order to reduce their particle size to less than 0.5 mm. The moisture content was determined by drying at 105 °C until a constant weight was reached (at least 18 h of drying needed). Ash was determined after drying the samples in a muffle furnace at 550±10 °C for at least 6 h until a constant weight was reached. Kjeldahl analysis was used to determine total protein using N×5.7 as the conversion factor. Starch content was measured using the Megazyme total starch assay kit (Megazyme International, Ireland). The lipid content was measured gravimetrically after extraction with a Soxhlet apparatus using petroleum ether (Merck, Germany) as solvent.

The composition of the carbohydrates in the samples was determined after a two-step acid hydrolysis procedure according to the National Renewable Energy Laboratory

protocol (NREL/TP-510-42618). The material (300 mg) was first hydrolysed with 72% v/v of sulphuric acid at 30 °C for 1 h and then in diluted acid (4%, v/v) at 121 °C for 30 min. During hydrolysis the polysaccharides are hydrolysed into monosaccharides (glucose derived from cellulose and β -glucan, and xylose and arabinose derived from hemicellulose) which were quantified by HPLC (Agilent, 1100 series) with an Aminex HPX-87H column (300 mm \times 7.8 mm, Bio-Rad, California, USA) and a refractive index detector. The operating conditions were: sample volume 20 μ L; mobile phase 0.005 M H₂SO₄; flow rate 0.6 mL/min; column temperature 65 °C. According to the NREL protocol, during acid hydrolysis lignin is fractionated into acid soluble and acid insoluble material. Acid-soluble lignin was measured with a UV-Vis spectrometer at 320 nm and acid-insoluble lignin gravimetrically after subtracting the ash and protein contents of the samples. The lignin content of samples is presented as the sum of acid soluble lignin and acid insoluble residue.

2.3 Osborne fractionation of DDGS and in-process samples

DDGS, wet and spent solid samples were subjected to Osborne fractionation according to the method of Lookhart and Bean (1995). Briefly, 100 mg of sample were sequentially extracted with deionised water, 0.5 M NaCl (Sigma, UK), 70% (v/v) aqueous ethanol (Sigma, UK) and 50% (v/v) 1-propanol (Merck, Germany) with 1% (w/v) dithiothreitol (DTT) (Sigma, UK), in order to extract the water-soluble albumins, salt-soluble globulins, ethanol-soluble prolamins and ethanol-insoluble prolamins (as reduced subunits), respectively. A 1:10 (w/v) solids-to-liquid ratio was used for the extractions, which were performed in a thermomixer (Eppendorf, UK) with constant mixing (1400 rpm), at 60 °C for 30 min. Extractions for each sample

were done in duplicate and the supernatants were collected by centrifugation (8,000×g for 5 min). In the case of sodium chloride, an additional wash with deionised water was performed in order to remove the residual salt. The protein contents of the Osborne fractionated supernatants were determined using the Bradford reagent assay (Sigma, UK) (Bradford, 1976).

2.4 Protein extraction

2.4.1 Aqueous-ethanol extraction of proteins

Lyophilised and milled samples were subjected to protein extraction using different extraction conditions. Initially, all samples were treated with hexane at a 1:10 (w/v) solid-to-hexane ratio at room temperature for 8 h in order to remove the oil content. Hexane was removed by filtration through a Whatman No 1 paper and the solids were placed in an oven at 45 °C overnight to remove any residual hexane. A two-stage process was subsequently applied to the de-fatted samples to extract the water insoluble proteins. Specifically, 10 g of each sample were mixed with 70% (v/v) aqueous ethanol in a 1:10 (v/w) ratio and incubated under constant shaking for 30 min at different temperatures (50, 70 and 90 °C). Supernatants were removed by centrifugation (8,000×g, 15 min) and the residues mixed with 70% (v/v) of aqueous ethanol in a 1:10 (v/w) ratio containing varying concentrations of sodium metabisulfite (Fluka, UK) (0.5, 1.0 or 1.5% w/v) as reducing agent. After mixing the samples for 30 min at different temperatures, again at 50, 70 and 90 °C, they were centrifuged (10,000×g, for 10 min at 25 °C), and the second step of the extraction was repeated. Deionised water was added to the collected supernatants in order to dilute the ethanol concentration to below 20% (v/v) and the samples were placed at -20 °C for 4 h to precipitate the proteins. The precipitated proteins were collected by

centrifugation (15,000×g, for 20 min at 2 °C), washed with distilled water, lyophilized in a VirTis Bench Top (USA) freeze-drier for 48 h, and stored at -20°C.

2.4.2 Alkaline-ethanol extraction of proteins

Alkaline conditions were also investigated for the extraction of the proteins in DDGS, wet and spent solid samples. These were incorporated in the second stage of the 2-step extraction process described in 2.4.1, in which aqueous ethanol (45 or 70%, v/v) was mixed with 0.05 or 0.1 M of NaOH (Fluka, UK) and 1.0% (w/v) sodium metabisulfite, in a solid-to-liquid ratio of 1:10. This extraction step was carried out twice at 70 °C for 30 min and the supernatants were collected following centrifugation (10,000×g, for 10 min at 25°C). Extracted proteins were then precipitated with 2 M HCl at pH 5.5 and collected by centrifugation (15,000×g, for 20 min at 10°C), washed with distilled water, lyophilized (VirTis Bench Top, USA) and stored at -20 °C. For both aqueous-ethanol and alkaline-ethanol extractions, the protein contents of the dried extracted samples were determined by Kjeldahl analysis. The protein content and protein yield of dried extracts were calculated as follows:

$$\text{Protein content of extract (\%)} = \frac{\text{Total Kjeldahl Nitrogen} \times 5.7}{\text{Dry weight of extract}} \times 100$$

$$\text{Protein yield (\%)} = \frac{\text{Protein content of extract}}{\text{Protein concentration in original sample}} \times 100$$

2.5 SDS-PAGE of samples and protein isolates

To identify the sub-units of water-insoluble proteins present in the original samples, they were extracted sequentially according to Singh, Shepherd and Cornish (1991). Briefly, gliadins were extracted three times from 20 mg samples with 0.1 mL 50% (v/v) 1-propanol for 30 min at 65 °C and the supernatants from the three extractions containing the gliadin fraction were collected by centrifugation (3,000×g for 2 min)

and pooled together. The solid residues, free of gliadins, were incubated with 50% (v/v) 1-propanol in 0.08M Tris-HCl (pH 8.0) with 1% (v/v) p-mercaptoethanol (Sigma, UK) as reducing agent and 1.4% (v/v) 4-vinylpyridine (Sigma, UK) as alkylating agent of sulfhydryl groups, in order to extract glutenin subunits; the supernatant containing the glutenin fraction was collected by centrifugation (3,000×g for 2 min). The supernatants containing the gliadin and glutenin fractions, respectively, were diluted in sample buffer [2% (v/v) SDS, 40% (w/v) glycerol, 0.02% (w/v) bromoethyl-blue in 0.08M Tris-HCl (pH=8.0)] and loaded onto a 1.0 mm 4-12% Bis-Tris pre-casted gel (NuPAGE Novex, UK). Proteins were separated in an XCell Surelock™ unit (Invitrogen, UK) at constant voltage (200 V) for 35 min. Gels were washed three times with purified water, stained with SimplyBlue SafeStain buffer (Life Technologies, UK) for 1 h at room temperature and washed with distilled water to obtain a clear background. The molecular weights of the visualised bands were estimated using Novex Sharp pre-stained protein standards (Invitrogen, UK).

The protein fractions extracted after aqueous-ethanol and alkaline-ethanol treatments of the samples were also separated based on their molecular weights using an XCell Surelock™ unit (Invitrogen, UK) according to the protocol provided by the supplier. Specifically, protein samples were reduced by treatment with NuPAGE LDS buffer and reducing agent (dithiothreitol) at 70 °C for 10 min. Electrophoresis was performed as described above.

2.6 Amino acid analysis

The original solid samples as well as lyophilized protein extracts (10 mg) were hydrolysed using 6M HCl (Fluka, UK) and 1% (w/v) phenol (Sigma, UK) at 110 °C for 24 h, in oxygen-free pressure tubes. After hydrolysis, aliquots (100 µL) were

neutralised and derivatised using the EZ-Faast amino acid derivatisation kit (Phenomenex, UK). The kit is based on a solid-phase extraction that binds amino acids and enables the derivatisation in aqueous solution of both the amine and carboxylic groups of amino acids at room temperature. Amino acid profiles were determined using a Gas Chromatography-Mass Spectrometry instrument (Agilent 6890/5975) as described by Elmore, Koutsidis, Dodson, Mottram & Wedzicha (2005). Norvaline was used as internal standard and detected amino acids were quantified according to standard solutions supplied by the manufacturer. Methionine, cysteine and tryptophan were not detected as they were degraded by acid hydrolysis.

2.7 Thermogravimetric analysis (TGA)

TGA analysis was carried out on the protein extracts using a Thermogravimetric Analyzer (TA-Q600SDT TGA). 10 mg samples were heated in an aluminium open pan (Perkin-Elmer) from 30 to 800 °C, with a heating rate of 20 °C/min under nitrogen flow (20 mL/min).

2.8 Statistical analysis

Data are presented as mean values and their respective standard deviations from three replicates. One-way ANOVA was used to calculate the significance between the means of the samples treated under different extraction conditions at $p < 0.05$.

3. Results and discussion

3.1 Composition of DDGS, wet solids and spent solids

The compositions of DDGS, wet solids and spent solids are shown in Table 1 as percentage concentration per dry weight basis (db). As expected, the dry matter was

higher in the case of DDGS, due to the thermal drying process carried out at the final stage of DDGS production. By contrast, wet and spent solids contained significant amounts of moisture (66.8% and 77.9%, respectively). In terms of protein, DDGS contained around 30% (db) of protein, whereas lower concentrations were present in wet (20%, db) and spent solids (25%, db). Similar values for wheat DDGS have been previously reported (Pedersen et al. 2014; Cozannet et al. 2010; Ortín & Yu, 2009), with the small differences probably resulting from differences in the processes used between different plants, seasonal variation in the harvested wheat, and a different N-to-protein conversion factor (6.25 over 5.7). The lipid content was similar in DDGS (3.4% db) and wet solids (2.9% db) but significantly higher ($P<0.05$) in spent solids (5.4%, db). Low concentrations of starch were detected in all samples (1.4-2.6%, db). In terms of the non-starch carbohydrate content, the values for cellulose and hemicellulose (Table 1) did not vary significantly between DDGS and the in-process samples. The ash content was slightly higher in spent solids (4.4%, db) than in DDGS (3.9%, db). Thin stillage, and consequently spent solids, as also shown in this study, typically contain the highest contents of ash among the different in-process samples (Liu, 2011; Hong et al. 2012). Blending of wet solids with condensed thin stillage and subsequent drying to give DDGS resulted in a lower ash content of DDGS compared to spent solids. Finally, DDGS had a higher lignin content (5.3%, db) compared to wet and spent solids. Pedersen et al. (2014) determined the composition of DDGS of various origins, including wheat, maize and mixed cereals, and found differences in the Klason lignin content among the DDGS samples. These were attributed to an extent to the presence of non-lignin sources in the Klason lignin fraction, such as Maillard-reaction products. The latter are formed during the mixing and drying of wet

solids as a result of the reaction between reducing sugars and lysine residues, and are condensed in thin stillage (Pahm, Pedersen & Stein, 2009).

3.2 Osborne fractionation of DDGS and in-process samples

A modified Osborne protocol was carried out for DDGS and in-process samples in order to identify the nature and solubility in different solvents of the various protein fractions present in the samples (Fig 1). Salt-soluble globulins were the least abundant group in all samples, accounting for 14% (w/w) of the total extracted protein in spent solids and about 10% (w/w) or less in wet solids and DDGS, respectively. The albumin content varied significantly among samples; it was the major protein fraction of spent solids accounting for ~ 41% (w/w), followed by ~18% (w/w) in DDGS and ~10% (w/w) in wet solids. Taking into account the fact that spent solids are a mixture of fermentation liquid and grain residues, it is expected that a substantial amount of the protein content in spent solids could be attributed to non-gluten proteins. These are mainly water soluble, metabolic or structural proteins and also include the amylolytic enzymes used in the fermentation process. Moreover, from a process point of view, the mixing of wet solids and the concentration of thin stillage taking place during the production of DDGS, contributed considerably to the presence of about 20% of albumins in DDGS. Alcohol soluble gliadins were the second most abundant protein fraction in all samples. Small differences in their concentration occurred between spent and wet solids (Fig. 1), whereas in DDGS they accounted for 33% (w/w) of the total extracted protein. Despite the fact that gliadins are readily soluble in aqueous alcohol, it is unlikely that they are solubilised during the fermentation process, as the ethanol concentration is only around 18% (v/v) at the end point of the fermentation. Glutenins were the major fraction in wet solids and DDGS, accounting

for 55% (w/w) and 42% (w/w) of the total extracted protein, respectively. Glutenins and gliadins contain high levels of proline and glutamine and serve as storage proteins in the starchy endosperm cells of the wheat grain (Shewry et al., 2002). Glutenins comprise a heterogeneous mixture of high and low molecular weight subunits assembled into polymers stabilised by inter-chain disulphide bonds (Veraverbeke & Delcour, 2002). However, they are only extractable in aqueous alcohol as reduced subunits in the presence of a reducing agent. It should be noted that the yield of total extracted proteins according to the Osborne fractionation method (measured by Bradford and compared to the initial protein content of the samples), was 20.3% for DDGS, 27.9% for wet solids and 28.4% for spent solids, respectively. Although the Osborne method has been widely used to extract proteins based on solubility, quantification can be problematic due to the fact that the different protein groups can overlap in their solubility in the different solvents, leading to partial cross-contamination of the fractions (Shewry, 1999; DuPont, Chan, Lopez & Vensel, 2005). However, in our study, Osborne fractionation proved to be a useful tool for identifying key differences between the protein contents of the samples, and demonstrated the influence of certain process steps on specific protein groups, such as albumins.

In order to further characterise the protein content of DDGS and in-process samples, prolamins were sequentially extracted according to the protocol of Singh, Shepherd and Cornish (1991). SDS PAGE analysis (Fig 2a) of the gliadins and glutenins present in spent solids, wet solids and DDGS was conducted. Based on the molecular weight ladder (lane 1), distinctive bands with molecular weight of around 50 kDa were present in spent and wet solids, corresponding to ω -gliadins, as well as bands with molecular weights of 40 and 30 kDa, corresponding α - and γ -gliadins, respectively. In

DDGS, only bands corresponding to ω -gliadins were distinctively present. It has been reported that high temperatures (above 100°C) can result in re-arrangements leading to the formation of new disulphide bonds among the sulphur-rich α , β - and γ -gliadins, whereas ω -gliadins do not contain cysteine residues and thus cannot form disulphide bonds (Schofield et al. 1983). This could explain the presence of only ω -gliadins in the DDGS sample, as the latter is subjected to an intensive thermal treatment at the last stage of its production (drum drying). All samples demonstrated intense bands at the top of the gel, suggesting the presence of high molecular weight aggregates of glutenin or gliadin subunits or even polymerised gliadins that were not able to enter the gel. Distinctive bands corresponding to low molecular weight glutenin subunits were present in spent and wet solids (30-60 kDa), whereas these bands were only present in traces in DDGS. The intensities of the glutenin bands on the SDS-PAGE gels confirmed the results obtained from the Osborne fractionation which indicated that glutenins were the most abundant proteins in the samples. In most of the gliadin and glutenin protein fractions, a clear band was obtained around 20 kDa. This could correspond to albumins, resulting from partial cross-contamination during Osborne fractionation, or partially hydrolysed proteins.

3.3 Extraction of proteins from DDGS and in-process samples

One of the major goals of this study was to investigate the methodology for the extraction of proteins from DDGS and in-process samples. Water-insoluble proteins (i.e. gliadins and glutenins) were mainly targeted, as these could serve as suitable starting materials for the development of biodegradable polymers for food and non-food applications, as previously shown for gliadins and glutenins derived from wheat grains (Kuktaite et al. 2011; Lagrain et al. 2010). Gliadins and reduced glutenin

340 subunits are both soluble in aqueous (60-70% v/v) ethanol (Shewry, 1999). Reducing
341 agents are typically used to improve protein extraction, as they reduce the disulphide
342 bonds present both within (intra-chain) and between (inter-chain) gluten protein
343 subunits (Shewry & Tatham, 1997). Dithiothreitol (DDT) and β -mercaptoethanol (β -
344 ME) are most widely utilised for this purpose. However, these chemicals are not
345 suitable for commercial production because of their toxicity. Alternatively, sodium
346 metabisulfite is a preferable reducing agent, as it is food grade and has lower toxicity
347 and odour compared to other reducing agents (Park, Bean, Wilson & Schober, 2006).

348 The first set of extraction experiments was carried out using 70% (v/v) aqueous
349 ethanol, in order to determine the effects of temperature and reducing agent
350 concentration on protein extractability. The protein content of the dried extracts as
351 determined by Kjeldahl analysis is presented in Table 2. Extraction at 50 °C resulted
352 in low protein content, ranging between 14-32 %, in all extracts depending on the
353 reducing agent concentration. The greatest amount of protein was present in the
354 extracts from the wet solids (~32%) followed by spent solids (~23%) and then DDGS
355 (~14%). Extraction at 70 °C improved significantly ($P<0.05$) the protein content of all
356 samples compared to 50 °C, with the highest being ~45% for DDGS, ~58% for wet
357 solids and ~62% for spent solids; the optimum reducing agent concentration was in
358 most cases 1%. At 90 °C the protein content of the extracts decreased significantly
359 ($P<0.05$) compared to that at 70 °C for all samples. In terms of the protein extraction
360 yield (% of protein per total protein of original sample), the best extraction conditions
361 were identified as 70 °C and 1% reducing agent, resulting in protein extraction yields
362 of 30.1% (w/w) for DDGS, 55.3% (w/w) for wet solids and 52.1% (w/w) for spent
363 solids.

In the presence of ethanol, only the hydrophobic fraction of wheat protein is solubilised, as a result of the disruption of low-energy hydrogen bonds in the decreased dielectric constant of the medium. Reduction of the disulphide bonds is responsible for the solubilisation of small amounts of ω -gliadins that are present in glutenin (the D type low molecular weight subunits) and some low molecular weight glutenin subunits, which in turn renders the remaining gluten proteins (comprising high molecular weight glutenins as well as α -, β - and γ -gliadins) soluble in hot ethanol solution (Mimouni, Robin, Azanza & Raymond, 1998). In this study, the use of reducing agent and 70 °C led to the extraction of water-insoluble prolamins, with the extraction efficiency being dependent on the starting material. However, at elevated temperatures (around 100 °C), the rich-sulphur α -, β - and γ -gliadins undergo disulphide bond rearrangements which reduces their solubility; this could be the reason for the lower protein extraction seen in the case of DDGS compared to the other samples in all extraction temperatures, and particularly at 90 °C (Table 2). Moreover, under such conditions, glutenin polymerisation can occur via sulphhydryl-disulphide inter-chain exchange reactions between polymers (Lagrain et al., 2008). The latter may be further facilitated by a temperature-dependent unfolding of the tertiary structure of the proteins. Recently, Hong and co-workers (2012) stated that the protein extraction efficiency of samples post-distillation (i.e. spent solids) is higher compared to DDGS, as a result of heat-induced protein denaturation and increased disulphide bonding within and among proteins, which occurs during the final drum drying step of the process.

The second set of extraction experiments was carried out using an alkaline-aqueous ethanol solution at 70 °C and a reducing agent concentration of 1.0%, as these were

shown from the previous experiments to be the optimal conditions for extraction. Alkalis and acids can partially hydrolyse protein molecules into smaller peptide fragments, which typically increases their solubility and extractability. In these experiments, the proteins were extracted with 45% or 70% (v/v) aqueous ethanol in the presence of 0.05M or 0.1M of NaOH. As shown in Table 2, 45% ethanol combined with 0.05M NaOH resulted in low protein contents in the extracts derived from wet and spent solids (~ 13% and 20%, respectively) and DDGS (~21%). The extractability of proteins was significantly ($P<0.05$) increased with 70% (v/v) ethanol, in particular in the presence of relatively high concentrations of alkali (0.1M); the protein content of the dried extracts was ~39% for DDGS, ~49% for wet solids and ~52% for spent solids, whereas the extraction yields were 27.1% for DDGS, 33.4% for wet solids and 31.2% for spent solids, respectively.

SDS-PAGE analyses of the proteins in the aqueous ethanol and alkaline-aqueous ethanol extracts are shown in Fig 2b. For both extraction methods, distinctive bands were obtained for all samples in the range of 35-50 kDa, most likely corresponding to a mixture of α -, γ - and ω -gliadins and low molecular weight glutenin subunits. Visualisation of the gels suggests that the profile of the extracted proteins was not considerably affected by the extraction conditions, and that the main differences in the protein content of the extracts obtained under the different conditions were primarily quantitative rather than qualitative.

Taking the above results into account, it can be deduced that aqueous ethanol extraction (pH~10) was a more efficient method for the extraction of proteins from the wheat DDGS and in-process samples compared to alkaline-aqueous ethanol extraction (pH~12). Utilising aqueous ethanol for extraction of proteins from DDGS

or in-process samples would be particularly attractive for distilleries and bioethanol plants. Moreover, the presence of alkali in the extraction process could result in corrosion of equipment in the long-term.

Although the literature on the extraction of proteins from wheat DDGS is limited, a few studies have studied the extraction of proteins from DDGS from other cereals. Xu et al. (2007) reported an extraction yield of 44% with 90% protein content for corn DDGS using 70% ethanol and 0.25% sodium sulfite at acidic pH. More recently, in a two fraction extraction process with 70% (v/v) aqueous 2-propanol and 70% (v/v) aqueous ethanol, Anderson, Ilankovan & Lamsai (2012) achieved an extraction yield of 70% of α -zein from maize DDGS. In another study, Wang, Tilley, Bean, Sun & Wang (2009) investigated the extraction efficiency of kafirin proteins (prolamins) from sorghum DDGS and reported an extraction yield of 44% with a kafirin content of 98.8% using acetic acid under reducing conditions. In the same context, Bandara et al. (2011) studied the efficiency of protein extraction from triticale DDGS and demonstrated that alkaline-ethanol conditions gave extraction yields between 21-30% (w/w) and a maximum protein content of ~66% (w/w). The present study is the first to investigate the extraction of proteins from in-process samples produced during the wheat DDGS production. Comparison of the extractability of proteins within samples shows that wet solids are the most appropriate starting material for protein extraction. Under optimal extraction conditions, 55.3% of the total protein was recovered from wet solids, with a protein content of 58% (w/w). From an industrial perspective, protein recovery and purity are very important for the translation of the process to large scale extraction. Commercial gluten products extracted from wheat contain around 75% protein. Therefore, efficient extraction using DDGS or in-process

samples as starting material should ideally result in a protein-rich extract with a similar purity. To this end, the addition of an ultrafiltration step post-reduction would reduce the amounts of carbohydrates and other non-protein components in the protein extracts and increase their purity.

3.4 Composition of protein extracts and solid residues

Table 3 shows the compositions of the protein extracts and their respective solid residues after ethanol extraction of DDGS, spent and wet solids samples. Very small amounts of water-soluble carbohydrates were detected in all aqueous ethanol extracts (2.4-5.1%, w/w), with the spent solids containing the smallest amount. Glucose was the major monosaccharide determined after hydrolysis, indicating the presence of starch followed by xylose and arabinose, the latter indicating the presence of soluble arabinoxylans, which are the major non-starch polysaccharides in wheat grain (Saulnier, Peneau & Thibault, 1995). On the other hand, the solid residue after extraction had a high content of water unextractable polysaccharides which was around 49% (w/w) for all samples. The monosaccharides composition (Table 3) indicated the presence of insoluble β -glucan, cellulose and water unextractable arabinoxylan. The protein content of the solid residues was ~4.7% for spent solids, ~7.4% for wet solids and ~11.6% for DDGS, i.e. the reverse ranking of that obtained for protein extractability.

Moreover, the mass balances for the principal components (i.e. protein and carbohydrates) were calculated. It should be noted that because the current study focused on the extraction of water-insoluble proteins, the contents of gliadins and glutenins determined by Osborne analysis (Fig 1) were taken into account for calculating the protein mass balance. Based on the data in Table 1, only ~69% of the

initial protein (gliadins and glutenins) content was recovered from DDGS. This could be attributed to only partial precipitation of the low molecular weight proteins in the extraction liquid, as well as to the thermal denaturation of DDGS proteins during the drying stage. At the drying stage DDGS is subjected to intense and prolonged thermal treatment (higher than 100°C). Under these conditions, the disulphide bonds present in all wheat gluten proteins (except ω -gliadins) may undergo rearrangements to form cross-links in highly insoluble denatured aggregates (Wang, Wei, Li, Bian & Zhao, 2009; Hong et al. 2012). On the other hand, protein recoveries from wet and spent solids were noticeably higher (89-93%), reflecting the higher protein extractability of these samples. In terms of the carbohydrate mass balances, the recoveries ranged from 86 to 94% for all samples. These values are very good considering that approximately 5% of the material could be lost during the intermediate washing steps. The high carbohydrate contents of the solid residues after protein extraction indicate that these materials could be a potential source of non-starch polysaccharides, and if processed to oligosaccharides could provide functional ingredients (prebiotics, stabilisers, emulsifiers) for food and non-food applications.

Further information on the proteins present in the extracts of DDGS and wet solids was provided by thermogravimetric analysis (TGA). Degradation of the samples was carried out under nitrogen and the observed peaks are presented as derivatives of the weight loss as a function of temperature. As depicted in Fig 4, a peak was identified for both samples at around 53-60°C. This was more intense in the case of wet solids and corresponded to the loss of free and bound water. Extracts of both DDGS and wet solids exhibited a prominent broad peak in the range of 230-370°C, which was attributed to the breakage of the covalent peptide bonds of amino acids, as well as to

the cleavage of disulphide, O-N and O-O bonds in protein molecules (Sun, Song & Zheng, 2007). Moreover, the analysis showed an additional peak for the DDGS extract at about 730°C, which is probably associated with the degradation of lignin components (Sahoo, Seydibeyoğlu, Mohanty & Misra, 2011).

3.5 Amino acid compositions of solid samples and protein extracts

Fig 5a shows the relative concentrations of amino acids in hydrolysates of the DDGS, wet and spent solids samples. Glutamic acid (which is mostly derived from the deamidation of glutamate), proline, leucine and phenylalanine were the major amino acids in the samples and are representative of wheat gluten proteins (Wieser, 2007). It is worth noting the reduced concentration of lysine in the DDGS sample, as lysine is the limiting essential amino acid in wheat grain proteins for the nutrition of humans and monogastric livestock (Shewry, 2007) but is labile to heating (Almeida, Htoo, Thomson & Stein, 2013). Fig 5b shows the relative concentration of individual amino acids in the wet solid and DDGS protein samples, obtained after aqueous ethanol and alkaline-aqueous ethanol extraction. The increased amino acid content of the wet solid extracts reflects the increased protein extractability of wet solids compared to the intensively thermally treated DDGS sample. Comparing the amino acid profile obtained between the two extraction methods, aqueous ethanol conditions showed increased specificity towards glutamic acid, phenylalanine and proline. These amino acids are present in α -, γ -, ω -gliadins and low molecular subunits of glutenin (Shewry, Tatham, Forde, Kreis & Mifflin, 1986) and as shown by SDS-PAGE, these were the major protein groups in the extracts. Apart from the potential utilisation of protein extracts as starting material for biodegradable plastics, the high content of glutamic acid could justify its extraction and utilisation as building block for chemical

compounds such as succinonitrile or acrylonitrile (Lammens, Franseen, Scott & Sanders, 2012). To this end, glutamic acid is a non-essential amino acid and its extraction would not compromise the nutritional value of DDGS used as livestock feed.

4. Conclusions

Aqueous ethanol extraction was more effective than alkaline-aqueous ethanol for extracting water-insoluble proteins from DDGS and in-process samples. The extractability of the proteins and their compositional characteristics were highly influenced by the starting raw material, i.e. wet solids, spent solids or DDGS. Protein was less efficiently extracted from DDGS, probably due to the decreased solubility of protein aggregates formed during the intensive thermal treatment during the drum drying stage. This is also indicated by the low recovery of α - and γ -gliadins. The wet solids exhibited the highest protein extractability (gliadins and glutenins), with a maximum recovery yield of 55% (w/w) (on the basis of total protein) and a protein content of 58% (w/w), and hydrolysates were particularly rich in glutamic acid and proline. The solid residues after extraction had a high carbohydrate content, which renders them amenable to enzymatic processing for the production of bioactive carbohydrates, such as prebiotic oligosaccharides, or for use as fibre-rich livestock feed. Overall, the research demonstrated the feasibility of utilising in-process samples from the DDGS production process for the extraction of proteins with good commercial potential.

Acknowledgements

The authors would like to acknowledge the Integrated Biorefining Research and Technology Club (IBTI) of the UK Biotechnology and Biological Sciences Research Council (BBSRC) for their financial support on a collaborative research project entitled “Development of a process scheme for the production of high value functional products from DDGS” (BB/J019429/1–University of Reading; BB/J019380/1 – Rothamsted Research).

References

- Almeida, F.N., Htoo, J.K., Thomson, J., Stein, H.H. (2013). Amino acid digestibility of heat damaged distillers dried grains with soluble fed to pigs. *Journal of Animal Science and Biotechnology*, 4, 44-54.
- Anderson, T.J., Ilankovan, P., Lamsai, B.P. (2012). Two fraction extraction of α -zein from DDGS and its characterization. *Industrial Crops and Products*, 37, 466-472.
- Bandara N, Chen L, Wu J. (2011). Protein extraction from triticale distillers grains. *Cereal Chemistry*, 88, 553-559.
- Bradford, MM. (1976) A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Analytical Biochemistry*, 72, 248-254.
- Cozannet, P., Primot, Y., Gady, C., Métayer, J.P., Callu, P., Lessire, M., Skiba, F., Noblet, J. (2010). Ileal digestibility of amino acids in wheat distillers dried grains with solubles for pigs. *Animal Feed Science and Technology*, 158, 177-186.
- Day, L., Augustin, M.A., Batey, I.L., Wrigley, C.W. (2006). Wheat-gluten uses and industry needs. *Trends in Food Science and Technology*, 17, 82-90.
- Directive 2009/28/EC of the European Parliament and of the Council on the promotion of the use of energy from renewable sources and amending and

556 subsequently repealing Directives 2001/77/EC and 2003/30/EC. 2009, OJ
 557 L140/16.

558 DuPont, F.M., Chan, R., Lopez, R., Vensel, W.H. (2005). Sequential extraction and
 559 quantitative recovery o gliadins, glutenins and other proteins from small samples
 560 of wheat flour. *Journal of Agriculture and Food Chemistry*, 53, 1575-1584.

561 Elmore, J.S., Koutsidis, G., Dodson, A.T., Mottram, D., Wedzicha, B. (2005).
 562 Measurement of acrylamide and its precursors in potato, wheat and rye model
 563 systems. *Journal of Agricultural and Food Chemistry*, 53, 1286-1293.

564 Hong, G.P., Avramenko, N., Stone, A., Abbott, D., Classen, H., Nickerson M. (2012).
 565 Extractability and molecular modification of gliadin and glutenin proteins
 566 withdrawn from different stages of a commercial ethanol fuel/distillers dried
 567 grains with solubles process using a wheat feedstock. *Cereal Chemistry*, 89, 276-
 568 283.

569 Irissin-Mangata, J., Bauduin, G., Boutevin, B., Gontard, N. (2001). New plasticizer
 570 for wheat gluten films. *European Polymer Journal*, 37, 1533-1541.

571 Kim, Y., Mosier, N.S., Hendrickson, R., Ezeji, T., Blaschek, H., Dien, B., Cotta, M.,
 572 Dale, B., Ladisch, M.R. (2008). Composition of corn dry-grind ethanol by-
 573 products: DDGS, wet cake and thin stillage. *Bioresource Technology*, 99, 5165-
 574 5176.

575 Kim, Y., Hendrickson, R., Mosier, N.S., Ladisch, M.R., Bals, B., Balan, V., Dale,
 576 B.E., Dien, B.S., Cotta, M.A. (2010). Effect of compositional variability of
 577 distillers' grains on cellulosic ethanol production. *Bioresource Technology*, 101,
 578 5385-5393.

579 Kingsly, A.R.P., Ileleji, K.E., Clementson, C.L., Garcia, A., Maier, D.E., Stroshine,
 580 R.L., Radcliff, S. (2010). The effect of process variables during drying on the

581 physical and chemical characteristics of corn dried distillers grains with soluble
 582 (DDGS)-Plant scale experiments. *Bioresource Technology*, 101, 193-199.

583 Klopfenstein, T.J., Erickson, G.E., Bremer, V.R. (2008). Use of distillers by-products
 584 in the beef cattle feeding industry. *Journal of Animal Science*, 86, 1223-1231.

585 Kuktaine, R., Privelic T.S, Cerenius, Y., Hedenqvist, M.S, Gällstedt, M.,Marttila, S.,
 586 Ignell, R., Popineau, Y., Tranquet, O., Shewry, P.R., Johansson, E. (2011).
 587 Structure and morphology of wheat gluten films: From polymeric protein
 588 aggregates toward superstructure arrangements. *Biomacromolecules*, 12, 1438-
 589 1448.

590 Lagrain, B., Goderis, B., Brijs, K., Delcour, J.A. (2010). Molecular basis of
 591 processing wheat gluten toward biobased materials. *Biomacromolecules*, 11, 533-
 592 541.

593 Lammens, T.M., Franssen, M.C.R., Scott, E.L., Sanders, J.P.M. (2012). Availability
 594 of protein-derived amino acids as feedstock for the production of bio-based
 595 chemicals. *Biomass and Bioenergy*, 44, 168-181.

596 Liu, K. (2011). Chemical composition of distillers grains, a review. *Journal of*
 597 *Agricultural and Food Chemistry*, 59, 1508-1526.

598 Lookhart, G.L., Bean, S.R. (1995). Separation and characterization of wheat protein
 599 fractions by high-performance capillary electrophoresis. *Cereal Chemistry*, 72(6),
 600 527-532.

601 Majeed, Z., Ramli, N.K, Mansor, N., Man, Z. (2015) A comprehensive review on
 602 biodegradable polymers and their blends used in controlled-release fertilizer
 603 processes. *Reviews in Chemical Engineering*, 31, 69-95.

604 Mimouni, B., Robin, J.M., Azanza, J.L., Raymond, J. (1998). Wheat flour proteins:
 605 Isolation and functionality of gliadin and HMW-glutenin enriched fractions.
 606 *Journal of the Science of Food and Agriculture*, 78, 423-428.

607 Ortín, W.G.N., Yu, P. (2009). Nutrient variation and availability of wheat DDGS,
 608 corn DDGS and blend DDGS from bioethanol plants. *Journal of the Science of*
 609 *Food and Agriculture*, 89, 1754-1761.

610 Pahm, A.A., Pedersen, C., Stein, H.H. (2009). Standardized ileal digestibility of
 611 reactive lysine in distillers dried grains with soluble fed to growing pigs. *Cereal*
 612 *Chemistry*, 57, 535-539.

613 Park, S.H., Bean, S.R., Wilson, J.D., Schober, T.J. (2006). Rapid isolation of sorghum
 614 and other cereal starches using sonication. *Cereal Chemistry*, 83, 611-616.

615 Pedersen, M.B., Dalsgaard, S., Knudsen, K.E.B., Yu, S., Lærke, H.N. (2014).
 616 Compositional profile and variation of distillers dried grains with soluble from
 617 various origins with focus on non-starch polysaccharides. *Animal Feed Science*
 618 *and Technology*, 197, 130-141.

619 Sahoo, S., Seydibeyoğlu, M.Ö., Mohanty, A.K., Misra, A. (2011). Characterization of
 620 industrial lignins for their utilisation in future value added applications. *Biomass*
 621 *and Bioenergy*, 32, 4230-4237.

622 Saulnier, L., Peneau, N., Thibault, J.F. (1995). Variability in grain extract viscosity
 623 and water-soluble arabinoxylan content in wheat. *Journal of Cereal Science*, 22,
 624 259-264.

625 Schingoethe, D.J., Kalscheur, K.F., Hippen, A.R., Garcia, A.D. (2009). Invited
 626 review: The use of distillers products in dairy cattle diets. *Journal of Dairy*
 627 *Science*, 92, 5802-5813.

628 Schingoethe, D.J. (2006). Utilization of DDGS by cattle. *Proc. 27th Western Nutrition*
629 *Conf, Winnipeg, Manitoba, Canada*, 19-20.

630 Schofield, J. D., Bottomley, R. C., Timms, M. F., and Booth, M. R. (1983). The effect
631 of heat on wheat gluten and the involvement of sulphhydryl disulphide interchange
632 reactions. *Journal of Cereal Science*, 1, 241-253.

633 Shewry, P.R. (2007). Improving the protein content and composition of cereal grain.
634 *Journal of Cereal Science*, 46, 239-250.

635 Shewry, P.R., Halford, N.G. (2002). Cereal seed storage proteins: structures,
636 properties and role in grain utilisation. *Journal of Experimental Botany*, 53, 947-
637 958.

638 Shewry, P.R. (1999). The synthesis, processing and deposition of gluten proteins in
639 the developing wheat grain. *Cereal Food World*, 44, 587-589.

640 Shewry, P.R., Tatham, A.S. (1997). Disulphide bonds in wheat gluten proteins.
641 *Journal of Cereal Science*, 25, 207-227.

642 Shewry, P.R., Tatham, A.R., Forde, J., Kreis, M., Mifflin, B.J. (1986). The
643 classification and nomenclature of wheat gluten proteins: A reassessment.
644 *Journal of Cereal Science*, 4, 97-106.

645 Singh, N.K., Shepherd, K.W., Cornish, G.B. (1991). A simplified SDS-PAGE
646 procedure for separating LMW subunits of glutenin. *Journal of Cereal Science*,
647 14, 203-208.

648 Sun, S., Song, Y., Zheng, Q. (2007). Morphologies and properties of thermo-molded
649 biodegradable plastics based on glycerol-plasticized wheat gluten. *Food*
650 *Hydrocolloids*, 21, 1005-1013.

651 Veraverbeke, W.S., Delcour, J.A. (2002). Wheat protein composition and properties
 652 of wheat glutenin in relation to breadmaking functionality. *Critical Reviews in*
 653 *Food Science and Nutrition*, 42, 179-208.

654 Wang, Y., Tilley, M., Bean, S., Sun, X.S., Wang, D. (2009). Comparison of methods
 655 for extraction of kafirin proteins from sorghum distillers dried grains with
 656 solubles. *Journal of Agricultural and Food Chemistry*, 57, 8366-8372.

657 Wang, J., Wei, Z., Li, L., Bian, K., Zhao M. (2009). Characteristics of enzymatic
 658 hydrolysis of thermal-treated wheat gluten. *Journal of Cereal Science*, 50, 205-
 659 209.

660 Wieser, H. (2007). Chemistry of gluten proteins. *Food Microbiology*, 24, 115-119.

661 Xu, W., Reddy, N., Yang, Y. (2007). An acidic method of zein extraction from
 662 DDGS. *Journal of Agricultural and Food Chemistry*, 55, 6279-6284.

663

664

Table 1. Chemical composition of DDGS and in-process samples


(in %, db)	DDGS	Wet solids	Spent solids
Dry matter	96.6±0.7	33.2±1.1	22.1±1.2
Crude protein	29.1±1.7	19.8±1.2	25.6±1.1
Crude fat	3.4±0.1	2.9±0.9	5.4±0.4
Cellulose and β-glucan	14.9±0.4	15.1±0.6	16.3±1.1
Starch	2.6±0.10	2.0±0.3	1.4±0.19
Hemicellulose	25.1±1.6	28.0±1.3	25.5±0.9
Xylose	(16.7±0.9)	(18.4±0.7)	(16.7±0.6)
Arabinose	(8.3±0.8)	(9.6±0.6)	(8.9±0.3)
Lignin	5.3±0.7	4.1±0.5	3.8±0.3
Ash	3.9±0.5	2.1±0.3	4.4±0.6

665

666

667

668



669

670 Fig. 1 Protein composition of fractionated spent solids (grey), wet solids (dark grey)
 671 and DDGS (black), based on Osborne protocol

672

673

674

675

Table 2. Protein content of isolates (% w/w) derived from DDGS and in-process samples during different extraction conditions, as determined by Kjeldahl analysis

Protein content of isolates (% w/w)						
Extraction Temperature (°C)	SMB (% w/w)	EtOH (% v/v)	NaOH (M)	DDGS	Wet solids	Spent solids
50	0.5	70	-	14.0 (0.8)	27.0 (1.3)	23.5 (1.2)
	1.0			14.5 (0.7)	31.7 (1.7)	22.9 (0.8)
	1.5			14.9 (0.5)	29.9 (1.7)	23.5 (0.9)
70	0.5	70	-	34.1 (3.2)	47.8 (2.9)	53.6 (2.8)
	1.0			42.7 (2.1)	55.6 (2.9)	62.4 (0.5)
	1.5			44.9 (1.5)	58.2 (0.5)	54.7 (0.3)
90	0.5	70	-	38.1 (1.3)	43.7 (0.3)	42.4 (0.3)
	1.0			29.6 (0.1)	42.6 (0.1)	45.8 (0.3)
	1.5			27.2 (0.2)	39.1 (0.2)	40.2 (0.3)
70	1.0	45	0.10	22.1 (0.6)	19.7 (1.7)	13.5 (1.4)
		70	0.05	27.5 (1.4)	25.9 (0.8)	36.6 (1.8)
		70	0.10	39.5 (2.1)	49.1 (0.7)	51.8 (2.2)

Data in parenthesis represent standard deviation values





Fig 2. (a) SDS-PAGE according to the protocol by Singh et al. (1991) of original samples: Lane 1, Molecular weight marker; lanes 2-4, Gliadin proteins of: spent solids (lane 2), wet solids (lane 3) and DDGS (lane 4), respectively; Lanes 4-6, Glutenin

687 proteins of: spent solids (lane 4), wet solids (lane 5) and DDGS (lane 6). (b) SDS-
688 PAGE of proteins extracted in aqueous-ethanol or alkaline-ethanol solutions: Lane 1,
689 Molecular weight marker; lanes 2-4, Ethanol extracted proteins of wet solids (lane 2),
690 spent solids (lane 3) and DDGS (lane 4); lanes 5-7, Alkaline (0.1M)-ethanol (70%,
691 v/v) extraction of wet solids (lane 5), spent solids (lane 6) and DDGS (lane 7).
692

693 Table 3 Protein and carbohydrate content of ethanol extracted proteins and their solid
694 residues and mass balance calculations compared to the starting raw materials
695 (DDGS, wet solids, spent solids)

	Protein extracts (% db)			Solid residues (% db)			Mass balance ^b (%, per 100 g)		
	DDGS	Wet solids	Spent solids	DDGS	Wet solids	Spent solids	DDGS	Wet solids	Spent solids
Protein^a	44.7	55.6	62.4	11.6	7.4	4.7	68.7	89.2	92.8
Carbohydrates	4.2	2.4	5.1	49.7	49.0	49.1	94.2	85.8	92.8
Glucose	2.1	1.2	2.8	20.1	18.6	19.8	87.3	83.1	89.2
Xylose	1.5	0.9	1.1	17.3	19.1	18.4	90.7	78.8	86.8
Arabinose	0.6	0.3	1.1	12.3	11.3	10.9	101.8	92.6	108.9

696 Data presented as mean values

697 ^a: Protein content measured by Kjeldahl

698 ^b: Mass balance for protein calculated by taking into account Osborne analysis results for gliadin and
699 glutenin content (45% in spent solids, 78% in wet solids and 73% in DDGS).

700




Fig 4. TGA analysis of ethanol extracted proteins from DDGS (black line) and wet solids (grey line)





Fig 5. Amino acid analysis of samples: (a) Relative amino acid concentration of spent solids (grey), wet solids (dark grey) and DDGS (black) after acid hydrolysis; (b) Relative amino acid concentration in wet solid protein extracted with ethanol, (light grey), wet solid protein extracted with alkaline-ethanol (grey), DDGS protein

714 extracted with ethanol (dark grey) and DDGS protein extracted with alkaline-ethanol
715 (black).