

Still scratching the surface: how much of the 'black box' of soil ectomycorrhizal communities remains in the dark?

Article

Accepted Version

Pickles, B. J. ORCID: <https://orcid.org/0000-0002-9809-6455> and Pither, J. (2014) Still scratching the surface: how much of the 'black box' of soil ectomycorrhizal communities remains in the dark? *New Phytologist*, 201 (4). pp. 1101-1105. ISSN 0028-646X doi: 10.1111/nph.12616 Available at <https://centaur.reading.ac.uk/48103/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

Published version at: <http://dx.doi.org/10.1111/nph.12616>

To link to this article DOI: <http://dx.doi.org/10.1111/nph.12616>

Publisher: Wiley

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online

1 **Still scratching the surface: how much of the “black box” of soil**
2 **ectomycorrhizal communities remains in the dark?**

3

4 Brian J. Pickles, Jason Pither

5

6 Irving K. Barber School of Arts and Sciences, Department of Biology, University of British
7 Columbia, Okanagan campus, 3333 University Way, Kelowna, BC V1V 1V7, Canada.

8

9 Author for correspondence:

10 *Brian J Pickles*

11 *Tel: +1 250 807 9965*

12 *Fax: +1 250 807 8005*

13 *Email: brian.pickles@ubc.ca*

14

Pickles & Pither 2013

15 **Article type: Letter**

16

17 **Key words:** Community ecology; diversity; ectomycorrhizas; meta-analysis; rooting depth;

18 soil depth.

19

Pickles & Pither 2013

Article

Symbiotic soil organisms such as ectomycorrhizal fungi (EMF) were long thought of as an inscrutable “black-box”, yet the advent of molecular technologies has driven rapid advances in identification and enumeration of their diversity (Horton & Bruns, 2001; Buée *et al.*, 2009). For instance, one 20 cm soil core can impressively yield 100’s of fungal OTUs (Taylor *et al.*, 2013). Importantly root symbionts play functional roles in sequestration or breakdown of soil carbon pools (Trumbore & Czimczik, 2008; Harrison *et al.*, 2011a,b; Clemmensen *et al.*, 2013; Kramer *et al.*, 2013), nutrient and water cycling (Virginia *et al.*, 1986; Read & Perez-Moreno, 2003), alteration of soil porosity (Perry *et al.*, 1990), and provision of sustenance for different trophic levels (Coleman & Whitman, 2005).

Yet root symbionts occur and contribute to function far deeper in the soil than is usually sampled (Jenkins *et al.*, 1988; Dalpé *et al.*, 2000; Bornyasz *et al.*, 2005). Soil properties vary considerably among ecosystems (Schenk, 2005; Dickie *et al.*, 2013), hence so too does rooting depth (see below) – even within a single species (Stone & Kalisz, 1991; Canadell *et al.*, 1996). However, in practice, we are (understandably) encouraged to employ uniform sampling strategies, even if these are known to only scratch the surface of potential symbiont habitat in some ecosystems (see below). Although the issue of limited-depth sampling has been raised before (Taylor, 2002), we are unaware of any efforts to quantify how much of the “black box” typically remains out of reach of standard sampling techniques. Such information would be extremely timely, due to the growing interest in accurately characterizing global patterns of EMF diversity and distribution (Dickie & Moyerson 2008; Vellinga *et al.*, 2009; Tedersoo *et al.*, 2012).

To begin addressing this question, we gathered sampling depth data from recent field studies of EMF, and analysed these in relation to published data compiled by ecosystem ecologists regarding (i) maximum rooting depths of trees and shrubs, including 137 EMF host species distributed among 29 host genera, and (ii) estimates for 8 ecosystems of the mean depth above

Pickles & Pither 2013

which 95% of all roots are located. Rooting depth data were derived from the following sources: (i) EMF host species/genera from Stone & Kalisz (1991) and Canadell *et al.* (1996), (ii) ecosystem data from Schenk & Jackson (2002). Sampling depth data were obtained from EMF studies published in the last 5 years of *New Phytologist* (Table S1). While the concepts discussed here are equally applicable to all soil-borne root symbionts, for the sake of brevity we focus our attention specifically on EMF and their hosts.

Based on 27 articles that reported sampling depth, the average was 13.4 cm (± 1.59 s.e.m.), with a median value of 10 cm. This sampling depth was approximately doubled in boreal and semi-arid ecosystems, and halved in semi-arid and tropical evergreen ecosystems. In comparison, none of the 29 ectomycorrhizal host genera for which data was available exhibited maximum rooting depths shallower than 50 cm (Fig. 1a), and on average maximum rooting depth among the 137 host species is 530 cm (± 44 cm s.e.m.) (Fig. 1b). Correspondingly, the average proportion of maximum rooting depth assessed is estimated to be 0.068 (± 0.0071 s.e.m.) across all host genera. If we consider maximum rooting depth as a proxy for the amount of habitat available to symbionts, then an enormous amount of potential habitat remains under-sampled, even within the Pinaceae (Fig. 1), which, according to a 2008 literature survey (Dickie & Moyerson, 2008), represented the focal family in 62% of all EMF studies.

Although maximum rooting depth is a crucial variable in research examining ecosystem function (Canadell *et al.*, 1996; Jackson *et al.*, 1996; Schenk, 2005), it could be argued that for our purposes it provides an overly pessimistic outlook on the completeness of current sampling efforts. We therefore also considered the EMF sampling depth data in relation to estimates of ecosystem-specific mean rooting depths calculated using 16 to 59 observations per ecosystem type, spanning all tree and shrub species for which rooting depth data existed (Schenk & Jackson, 2002). Using an average sampling depth of 13.4 cm, the proportion of the mean ecosystem rooting depth sampled varied from a high of 0.47 for tundra, to a low of 0.08 for

Pickles & Pither 2013

Mediterranean ecosystems (Fig. 2a), with a mean of 0.178 (\pm 0.0442 s.e.m.). Thus, even in tundra ecosystems where rooting depths are comparatively shallow (Fig. 2b), typical sampling methods are likely to access less than 50% of the mean depth of host roots.

Although striking, our findings do not necessarily mean that standard sampling methods are always doomed to miss an important or sizeable component of the symbiont assemblage associated with any given host. Indeed, it is likely that some studies - especially those occurring in shallow rooting regions (e.g. tundra ecosystems) - could yield reasonable estimates of the actual number of symbiont species associated with the host (using appropriate analytical techniques; cf. Gotelli & Colwell, 2001). Nevertheless, it has long been acknowledged that important characteristics of EMF communities vary with depth, but only in recent years have studies begun to clarify these details. For example, fungal hyphae show vertical niche differentiation (Dickie *et al.*, 2002), EMF community composition changes between soil horizons (Rosling *et al.*, 2003), ECM root tips and EMF extramatrical mycelium differ in their vertical structure (Genney *et al.*, 2006), and other depth-associated patterns continue to emerge (e.g. Egerton-Warburton *et al.*, 2003; Landeweert *et al.*, 2003; Baier *et al.*, 2006; Lindahl *et al.*, 2007; Courty *et al.*, 2008; Scattolin *et al.*, 2008; Beiler *et al.*, 2010; Clemmensen *et al.*, 2013; Taylor *et al.*, 2013). These observations, combined with our findings, substantiate earlier statements that current sampling methods provide a limited view of EMF assemblages (Taylor, 2002). Until more effort is spent sampling and characterising symbiont diversity and function at depth, we cannot know the true extent of these limitations.

Since deep roots are features of most ecosystems worldwide (Schenk & Jackson, 2005), the discoveries that could come with deeper sampling have the potential to profoundly change our outlook on patterns of EMF diversity and function. To illustrate, consider a recent and enlightening global-extent meta-analysis of local EMF diversity (Tedersoo *et al.*, 2012). Based on data from 55 published studies, total species richness (representing site-level species richness) was significantly associated with a number of climate-based predictor variables (e.g.

Pickles & Pither 2013

mean annual temperature, mean annual precipitation), and not surprisingly, number of samples and total sample volume. However, the meta-analysis included data gathered from a variety of host genera and ecosystem types, meaning that the rooting depths of hosts also varied substantially (see above). It would be interesting to determine if and how their findings would change if sampling was deeper, or was adjusted to account for site-specific rooting depths. Because different communities arise with increasing depth, we predict that deeper sampling will increase estimates of total richness and reveal significant changes in community composition. Perhaps every additional 50 cm of depth explored could provide as much richness again as that found in the organic horizon (as per Rosling *et al.*, 2003 & Landeweert *et al.*, 2003)? Based upon our findings, we speculate that the magnitude of this total increase will vary significantly with ecosystem type due to the differences in host rooting depth and density.

Variation in the rooting depth of a given host species is related to multiple factors including age, depth to bedrock, mean annual precipitation, mean annual potential evapotranspiration, and depth to the water table (Schenk 2005), all co-varying with ecosystem type. Thus, a Douglas-fir growing in seasonally dry evergreen forest is more likely to develop deep roots than one growing in a cool-temperate to sub-boreal region (cf. Schenk & Jackson, 2002). This has implications for sampling strategies (see below), and suggests that host species distributed across multiple ecosystem types, like Douglas-fir, may be associated with a much more diverse pool of EMF symbionts than current estimates indicate. This combination of varied rooting depths and soil environments provides a greater diversity of habitat to symbionts than do hosts whose distributions are predominantly restricted to a single ecosystem type (e.g. black spruce).

Another important finding concerns the thoroughness with which sampling methods are described in published articles. Of the 30 EMF studies published in the past 5 years in *New Phytologist*, 3 (10%) failed to report details about sampling depth. More generally, whereas some authors give detailed descriptions of the soil environment in relation to sampling strategy (e.g. Smith *et al.*, 2005; Ryberg *et al.*, 2009), depth information occasionally has to be derived

Pickles & Pither 2013

or is missing entirely. We suggest that where possible, details about sampling should be accompanied by estimates of average rooting depths at the site, for the host species of interest, even if these estimates are speculative. This would provide for better and more consistent estimates of realized sampling effort across studies.

Lastly, future research may not only require deeper sampling to minimise bias (depending upon the research objectives), but may also need to stratify sampling geographically according to potential rooting depth. Combining global estimates of soil depth (<http://www.fao.org/nr/land/soils/harmonized-world-soil-database/en/>) with global estimates of deep root distributions (Schenk & Jackson, 2005) and species' ranges (e.g. <http://esp.cr.usgs.gov/data/little/>) could help hone in on potential sampling regions, and ground penetrating radar technology (Sucre *et al.*, 2011) could be used to identify final sample locations. The logistical impediments associated with deep soil sampling (including cost; Harrison *et al.*, 2011) are daunting, but other research areas point to possible solutions, such as using drilling equipment to acquire ice or sediment cores (Nogu   *et al.*, 2013), or using excavation machinery such as a backhoe (Bornyasz *et al.*, 2005). These challenges are worth tackling given the potentially crucial roles that symbionts at depth may play in ecosystem function (e.g. Clemmensen *et al.* 2013; Kramer *et al.* 2013).

Acknowledgements

We thank Colin Scherer and Emma Walker for assistance with data mining, and Ian Dickie and two anonymous reviewers for their insightful comments on this manuscript. BJP acknowledges financial support from the Simard and Mohn labs at the University of British Columbia, Canada. JP acknowledges financial support from the Natural Sciences and Engineering Research Council of Canada (Discovery Grants program), the Canada Foundation for Innovation, and the I.K. Barber School of Arts and Sciences at the Okanagan campus of the University of British Columbia, Canada.

Pickles & Pither 2013

150

151 **References**

- 152 Baier R, Ingenhaag J, Blaschke H, Gottlein A, Agerer R. 2006. Vertical distribution of an
 153 ectomycorrhizal community in upper soil horizons of a young Norway spruce (*Picea abies* [L.]
 154 Karst.) stand of the Bavarian Limestone Alps. *Mycorrhiza* **16**: 197-206.
- 155 Beiler KJ, Durall DM, Simard SW, Maxwell SA, Kretzer AM. 2010. Architecture of the
 156 wood-wide web: *Rhizopogon* spp. genets link multiple Douglas-fir cohorts. *New Phytologist*
 157 **185**: 543-553.
- 158 Bornyasz MA, Graham RC, Allen MF. 2005. Ectomycorrhizae in a soil-weathered granitic
 159 bedrock regolith: Linking matrix resources to plants. *Geoderma* **126**: 141-160.
- 160 Buée M, Reich M, Murat C, Morin E, Nilsson RH, Uroz S, Martin F. 2009. 454
 161 pyrosequencing analyses of forest soils reveal an unexpectedly high fungal diversity. *New*
 162 *Phytologist* **184**: 449-456.
- 163 Canadell J, Jackson RB, Ehleringer JR, Mooney HA, Sala OE, Schulze E-D. 1996. Maximum
 164 rooting depth of vegetation types at the global scale. *Oecologia* **108**: 583-595.
- 165 Clemmensen KE, Bahr A, Ovaskainen O, Dahlberg A, Ekblad A, Wallander H, Stenlid J,
 166 Finlay RD, Wardle DA, Lindahl BD. 2013. Roots and associated fungi drive long-term carbon
 167 sequestration in Boreal forest. *Science* **339**: 1615-1618.
- 168 Coleman DC, Whitman WB. 2005. Linking species richness, biodiversity and ecosystem
 169 function in soil systems. *Pedobiologia* **49**: 479-497.
- 170 Courty P-E, Franc A, Pierrat J-C, Garbaye J. 2008. Temporal changes in the ectomycorrhizal
 171 community in two soil horizons of a temperate oak forest. *Applied and Environmental*
 172 *Microbiology* **79**: 5792-5801.
- 173 Dalpé Y, Diop TA, Plenchette C, Gueye M. 2000. Glomales species associated with surface
 174 and deep rhizosphere of *Faidherbia albida* in Senegal. *Mycorrhiza* **10**: 125-129.

Pickles & Pither 2013

- 175 Dickie IA, Moyerson B. 2008. Towards a global view of ectomycorrhizas. *New Phytologist*
 176 **180**: 263-265.
- 177 Dickie IA, Xu B, Koide RT. 2002. Vertical niche differentiation of ectomycorrhizal hyphae in
 178 soil as shown by T-RFLP analysis. *New Phytologist* **156**: 527-535.
- 179 Dickie IA, Martinez-Garcia LB, Koele N, Grelet G-A, Tylianakis JM, Peltzer DA, Richardson
 180 SJ. 2013. Mycorrhizas and mycorrhizal fungal communities throughout ecosystem
 181 development. *Plant & Soil* **367**: 11-39.
- 182 Egerton-Warburton LM, Graham RC, Hubbert KR. 2003. Spatial variability in mycorrhizal
 183 hyphae and nutrient and water availability in a soil-weathered bedrock profile. *Plant and Soil*
 184 **249**: 331-342.
- 185 Genney DR, Anderson IC, Alexander IA. 2006. Fine-scale distribution of pine
 186 ectomycorrhizas and their extramatrical mycelium. *New Phytologist* **170**: 381-390.
- 187 Gotelli NJ, Coldwell RK. 2001. Quantifying biodiversity: procedures and pitfalls in the
 188 measurement and comparison of species richness. *Ecology Letters* **4**: 379-391.
- 189 Harrison RB, Footen PW, Strahm BD. 2011a. Deep soil horizons: contribution and importance
 190 to soil carbon pools and in assessing whole-ecosystem response to management and global
 191 change. *Forest Science* **57**: 67-76.
- 192 Harrison RB, Richter D, Fox T. 2011b. Deep soils. *Forest Science* **57**: 1-2.
- 193 Horton TR, Bruns TD. 2001. The molecular revolution in ectomycorrhizal ecology: peeking
 194 into the black-box. *Molecular Ecology* **10**: 1855-1871.
- 195 Jackson RB, Canadell J, Ehleringer JR, Mooney HA, Sala OE, Schulze ED. 1996. A global
 196 analysis of root distributions for terrestrial biomes. *Oecologia* **108**: 389-411.
- 197 Jenkins MB, Virginia RA, Jarrell WM. 1988. Depth distribution and seasonal populations of
 198 Mesquite-nodulating *Rhizobia* in warm desert ecosystems. *Soil Science Society of America*
 199 *Journal* **52**: 1644-1650.

Pickles & Pither 2013

- 200 Kraft NJB, Comita LS, Chase JM, Sanders NJ, Swenson NG, Crist TO, Stegan JC, Vellend M,
 201 Boyle B, Anderson MJ *et al.* 2011. Disentangling the drivers of β diversity along latitudinal
 202 and elevational gradients. *Science* **333**: 1755-1758.
- 203 Kramer S, Marhan S, Haslwimmer H, Ruess L, Kandeler E. 2013. Temporal variation in
 204 surface and subsoil abundance and function of the soil microbial community in an arable soil.
 205 *Soil Biology and Biochemistry* **61**: 76-85.
- 206 Landeweert R, Leeftang P, Kuyper TW, Hoffland E, Rosling A, Wernars K, Smit E. 2003.
 207 Molecular identification of ectomycorrhizal mycelium in soil horizons. *Applied Environmental*
 208 *Microbiology* **69**: 327-333.
- 209 Lindahl BD, Ihrmark K, Boberg J, Trumbore SE, Högborg P, Stenlid J, Finlay RD. 2007.
 210 Spatial separation of litter decomposition and mycorrhizal nitrogen uptake in a boreal forest.
 211 *New Phytologist* **173**: 611-620.
- 212 Nogué S, de Nascimento L, Fernández-Palacios JM, Whittaker RJ, Willis KJ. 2013. The
 213 ancient forests of La Gomera, Canary Islands, and their sensitivity to environmental change.
 214 *Journal of Ecology* **101**: 368-377.
- 215 Perry DA, Borchers JG, Borchers SL, Amaranthus MP. 1990. Species migrations and
 216 ecosystem stability during climate change: the belowground connection. *Conservation Biology*
 217 **4**: 266-274.
- 218 Read DJ, Perez-Moreno J. 2003. Mycorrhizas and nutrient cycling in ecosystems – a journey
 219 towards relevance? *New Phytologist* **157**: 475-492.
- 220 Rosling A, Landeweert R, Lindahl BD, Larsson K-H, Kuyper TW, Taylor AFS Findlay RD.
 221 2003. Vertical distribution of ectomycorrhizal fungal taxa in a podzol soil profile. *New*
 222 *Phytologist* **159**: 775-783.
- 223 Ryberg M, Andreasen M, Björk RG. 2010. Weak habitat specificity in ectomycorrhizal
 224 communities associated with *Salix herbacea* and *Salix polaris* in alpine tundra. *Mycorrhiza* **21**:
 225 289-296.

Pickles & Pither 2013

- 226 Scattolin L, Montecchio L, Mosca E, Agerer R. 2008. Vertical distribution of the
 227 ectomycorrhizal community in the top soil of Norway spruce stands. *European Journal of*
 228 *Forest Research* **127**: 347-357.
- 229 Schenk HJ. 2005. Vertical vegetation structure below ground: Scaling from root to globe.
 230 *Progress in Botany* **66**: 341-373.
- 231 Schenk HJ, Jackson RB. 2002. The global biogeography of roots. *Ecological Monographs*
 232 **72**: 311-328.
- 233 Schenk HJ, Jackson RB. 2005 Mapping the global distribution of deep roots in relation to
 234 climate and soil characteristics. *Geoderma* **126**: 129-140.
- 235 Smith JE, McKay D, Brenner G, McIver J, Spatafora JW. 2005. Early impacts of forest
 236 restoration treatments on the ectomycorrhizal fungal community and fine root biomass in a
 237 mixed conifer forest. *Journal of Applied Ecology* **42**: 526-535.
- 238 Stone EL, Kalisz PJ. 1991. On the maximum extent of tree roots. *Forest Ecology &*
 239 *Management* **46**: 59 –102.
- 240 Sucre EB, Tuttle JW, Fox TR. 2011. The use of ground-penetrating radar to accurately
 241 estimate soil depth in rocky forest soils. *Forest Science* **57**: 59-66.
- 242 Taylor AFS. 2002. Fungal diversity in ectomycorrhizal communities: sampling effort and
 243 species detection. *Plant & Soil* **244**: 19-28.
- 244 Taylor DL, Hollingsworth TN, McFarland, Lennon NJ, Nusbaum C, Ruess RW. 2013. A first
 245 comprehensive census of fungi in soil reveals both hyperdiversity and fine-scale niche
 246 partitioning. *Ecology*, in press: <http://dx.doi.org/10.1890/12-1693.1>
- 247 Trumbore SE, Czimczik CI. 2008. An uncertain future for soil carbon. *Science* **321**: 1455-1456.
- 248 Vellinga EC, Wolfe BE, Pringle A. 2009. Global patterns of ectomycorrhizal introductions.
 249 *New Phytologist* **181**: 960-973.
- 250 Virginia RA, Jenkins MB, Jarell WM. 1986. Depth of root symbiont occurrence in soil. *Biology*
 251 *and Fertility of Soils* **2**: 127-130.

Pickles & Pither 2013



252 **Figure legends**

253 **Figure 1.** a. Proportion of maximum recorded rooting depth examined (\pm s.e.m.) across genera
254 using mean sampling depth derived from values reported in the literature (Table S1). Note
255 maximum y-axis value is a proportion of 0.15. b. Maximum rooting depths of selected host
256 species, with multiple bars (records) per species. Dashed red line represents average sampling
257 depth of EMF studies. In both panels, green bars indicate genera or species in the Pinaceae.
258 **Figure 2.** a. Sampled proportion of the mean depth at which 95% of ecosystem roots are
259 located, calculated using mean of sampling values reported in literature (Table S1). b.
260 Estimated mean depth (\pm s.e.m.) at which 95% of ecosystem roots are located using the
261 interpolated values of Schenk & Jackson (2002).

262

263 **Supporting Information**

264 **Table S1.** Citation, sample depth and host species for all ectomycorrhizal articles from the last
265 5 years of *New Phytologist* in which sampling depth was provided.



Proportion of mean rooting depth sampled



Mean ecosystem rooting depth (m)

