

Galloylated proanthocyanidins from shea (Vitellaria paradoxa) meal have potent anthelmintic activity against Ascaris suum

Article

Accepted Version

Ramsay, A., Williams, A., Thamsborg, S. and Mueller-Harvey, I. (2016) Galloylated proanthocyanidins from shea (*Vitellaria paradoxa*) meal have potent anthelmintic activity against *Ascaris suum*. *Phytochemistry*, 122. pp. 146-153. ISSN 0031-9422 doi: <https://doi.org/10.1016/j.phytochem.2015.12.005> Available at <https://centaur.reading.ac.uk/49230/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

To link to this article DOI: <http://dx.doi.org/10.1016/j.phytochem.2015.12.005>

Publisher: Elsevier

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online

Equations for quantifying extractable and unextractable PA:

PA calculation of extractable PA (ePA) in whole meal:

$$ePA \text{ in g/100g of DW of whole shea meal} = \left(\frac{\text{extract yield g}}{100 \text{ g whole meal}} \right) \times \left(\frac{PA \text{ g}}{100 \text{ g extract}} \right)$$

PA calculation of unextractable PA (uPA) in whole meal:

$$uPA \text{ in g/100g of DW of whole shea meal} = \left(\frac{\text{residue yield g}}{100 \text{ g whole meal}} \right) \times \left(\frac{PA \text{ g}}{100 \text{ g extract}} \right)$$

Equations for characterising PA compositions:

Thiolysis degrades PA and yields a 'global average' flavanol composition in the PA mixtures (Figure S1; Gea et al 2011):

The mean degree of polymerisation (mDP) is calculated as follows:

$$mDP = \frac{\text{Amount of extension and terminal flavanol units [mole]}}{\text{Amount of terminal flavanol units [mole]}}$$

The percentage of procyanidins (PC) within PA is calculated as follows:

$$\% \text{ PC} = \frac{C + EC + Cg + ECg \text{ units [mole]}}{\text{Total flavanols in PA [mole]}} \times 100$$

The percentage of prodelphinidins (PD) within PA is calculated as follows:

$$\% \text{ PD} = 100 \% - \% \text{ PC}$$

The percentage of *cis* flavan-3-ols within PA is calculated as follows:

$$\% \text{ cis} = \frac{EC + ECg + EGC + EGCg \text{ units [moles]}}{\text{Total flavanols in PA [mole]}} \times 100$$

The percentage of *trans* flavan-3-ols within PA is calculated as follows:

$$\% \text{ trans} = 100 \% - \% \text{ cis}$$

C = catechin, EC = epicatechin, EGC = epigallocatechin, Xg = flavan-3-ol gallate

Figure S1: Thiolytic degradation of galloylated and non-galloylated proanthocyanidins




Table S1: Flavan-3-ol composition of proanthocyanidins in shea meal as terminal and extension units (relative molar percentages; SD in parentheses; n = 3).). Data are presented so that terminal units add up to 100% and extension units add up to 100%.

Proanthocyanidins	Terminal units (%)						Extension units (%)							
	GC	EGC	GCg	EGCg	C	ECg	GC	EGC	GCg	EGCg	C	EC	Cg	ECg
Total PA	5.3	21.1	10.2	41.4	6.1	15.8	31.2	16.3	5.9	14.5	5.0	20.1	0.7	6.3
	(0.5)	(3.6)	(0.3)	(7.1)	(1.9)	(1.9)	(0.7)	(1.0)	(0.2)	(0.6)	(0.1)	(0.5)	(0.1)	(0.3)
Extractable PA	26.1	22.2	7.7	30.9	5.8	7.4	26.5	15.7	7.5	19.2	4.6	18.0	0.7	7.7
	(0.7)	(1.9)	(0.2)	(0.8)	(0.3)	(0.1)	(0.5)	(0.4)	(0.1)	(0.3)	(0.1)	(0.3)	(0.1)	(0.1)
Unextractable PA	5.9	13.3	9.0	49.6	5.5	16.7	29.1	15.5	5.3	15.1	5.2	22.1	0.6	7.2
	(0.2)	(0.9)	(0.2)	(0.7)	(0.6)	(0.6)	(0.8)	(0.4)	(0.1)	(0.4)	(0.1)	(0.6)	(0.1)	(0.2)

Table S2: Composition of flavan-3-ol terminal and extension units in three proanthocyanidin fractions (relative molar percentages; SD in parentheses; n = 3). Data are presented so that terminal units add up to 100% and extension units add up to 100%.

	Terminal units (%)						Extension units (%)							
	GC	EGC	GCg	EGCg	C	ECg	GC	EGC	GCg	EGCg	C	EC	Cg	ECg
Shea meal														
F1-fraction	36.1	24.2	3.4	23.9	7.6	4.8	20.3	27.1	7.2	12.8	6.8	20.5	0.7	4.6
	(0.2)	(0.1)	(0.1)	(0.2)	(0.1)	(0.1)	(2.3)	(4.1)	(1.2)	(0.2)	(0.1)	(0.4)	(0.1)	(0.1)
F2-fraction	17.0	10.4	15.6	40.4	4.5	12.1	27.3	12.3	8.1	21.2	4.6	16.6	1.1	8.7
	(0.3)	(0.1)	(0.1)	(0.2)	(0.1)	(0.1)	(0.2)	(0.4)	(0.1)	(0.2)	(0.1)	(0.1)	(0.1)	(0.1)
F3-fraction	18.8	12.2	15.1	40.6	5.7	7.7	13.7	33.3	3.4	20.8	4.4	14.9	0.7	8.7
	(1.2)	(0.5)	(0.8)	(2.1)	(0.3)	(4.7)	(5.6)	(2.6)	(0.2)	(1.4)	(0.2)	(0.9)	(0.1)	(0.6)

Table S3: 2D NMR (HSQC) chemical shifts (δ ppm) of the proanthocyanidin F2-fraction (DMSO- d_6 /pyridine- d_5 , 500 MHz).

Assignments*	^1H	^{13}C
2	5.25; 5.53	75.57; 67.83
3	4.02	71.19
4	4.83	35.91
6/8	6.00	95.98
2' PD/6' PD	6.57	106.01
2' PC	7.06	114.87
5' PC	6.75	114.87
6' PC	6.77	117.59
Gallic acid (2H)	7.03	108.97

* See Figure 1 for hydrogen and carbon numbers.