

# *Proanthocyanidins from Averrhoa bilimbi fruits and leaves*

Article

Accepted Version

Creative Commons: Attribution-Noncommercial-No Derivative Works 4.0

Ramsay, A. and Mueller-Harvey, I. (2016) Proanthocyanidins from Averrhoa bilimbi fruits and leaves. Journal of Food Composition and Analysis, 47. pp. 16-20. ISSN 0889-1575 doi: 10.1016/j.jfca.2015.12.004 Available at <https://centaur.reading.ac.uk/49827/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

To link to this article DOI: <http://dx.doi.org/10.1016/j.jfca.2015.12.004>

Publisher: Elsevier

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

[www.reading.ac.uk/centaur](http://www.reading.ac.uk/centaur)

**CentAUR**

Central Archive at the University of Reading

Reading's research outputs online

**Manuscript title: Procyanidins from *Averrhoa bilimbi* fruits and leaves**

A. Ramsay<sup>a\*</sup>, I. Mueller-Harvey<sup>a</sup>

<sup>a</sup>*Chemistry and Biochemistry Laboratory, School of Agriculture, Policy and Development,  
University of Reading, 1 Earley Gate, P O Box 236, Reading RG6 6AT, U.K*

\*Corresponding author (Tel: +44 (0)118 378 6619; Fax: +44 (0)118 935 2421; E-mail:

[aina.ramsay@hotmail.fr](mailto:aina.ramsay@hotmail.fr); [aina.ramsay@reading.ac.uk](mailto:aina.ramsay@reading.ac.uk))

**Procyanidins from *Averrhoa bilimbi* fruits and leaves**

A. Ramsay<sup>a\*</sup>, I. Mueller-Harvey<sup>a</sup>

<sup>a</sup>*Chemistry and Biochemistry Laboratory, School of Agriculture, Policy and Development,  
University of Reading, 1 Earley Gate, P O Box 236, Reading RG6 6AT, U.K*

\*Corresponding author (Tel: +44 (0)118 378 6619; Fax: +44 (0)118 935 2421; E-mail:  
[aina.ramsay@hotmail.fr](mailto:aina.ramsay@hotmail.fr); [aina.ramsay@reading.ac.uk](mailto:aina.ramsay@reading.ac.uk))

## Abstract

Proanthocyanidins from *Averrhoa bilimbi* fruits and leaves were analysed by thiolysis with benzyl mercaptan and high performance liquid chromatography - mass spectrometry and consisted of pure B-type procyanidins. These tannins consisted of almost pure homopolymers, with epicatechin accounting for most of the monomeric subunits in fruits (97%) and leaves (99%). Leaves contained more procyanidins (4.5 vs 2.2 g/100 g dry weight) with a higher mean degree of polymerisation (9 vs 6) than fruits. This study thus contributes information on the proanthocyanidins of a traditional food that can make an important contribution to the intake of compounds with antioxidant and health benefits. The fruits are prized for culinary purposes and the leaves are used in traditional medicine.

**Keywords:** *Averrhoa bilimbi*, cucumber tree, food analysis, proanthocyanidins, condensed tannins, gel-NMR, thiolysis, benzyl mercaptan

## 1. Introduction

Investigation into the phytochemical profiles of underutilized and/or wild foods is becoming increasingly important in the context of food security and tree foods are of particular interest, as trees are generally more resilient to periodic droughts and unseasonal weather events than crops. Underutilised foods can be especially valuable when staple foods are in short supply. Information on the contents of non-nutrients is needed to explore their bioactivities and dietary health benefits (Rush, 2001). Therefore, knowledge of the phytochemical composition of wild foods will allow local populations to better exploit local resources and their benefits (Scoones et al. 1992).

*Averrhoa bilimbi* (L.), commonly called the cucumber tree (Figure 1), belongs to the family of Oxalidaceae and grows in tropical regions (Central America, Asia and Caribbean Islands). The fruits are consumed locally in culinary preparations (fresh in salad or pickled) or as juice. The juice can also be used as a remedy to treat dental disorders, sore throats and stomach problems (Ariharan et al. 2012). *Averrhoa bilimbi* fruits have shown anti-obesity properties or anti-cholesterolemic activity (Ambili et al. 2009) and also antibacterial and antioxidant activities (Ashok Kumar et al. 2013). However, their high acidity (pH = 4) and high oxalate concentration (Morton et al. 1987) has led to renal failure after prolonged consumption of the juice in humans (Bakul et al. 2013). In terms of phytochemical compounds, the fruits are a good source of vitamin C (Ariharan et al. 2012) and various flavonoids (myricetin, luteolin, quercetin and apigenin) have been quantified (Koo Hui & Suhaila, 2001). Although the presence of tannins has been mentioned in the fruits (Ashkok Kumar et al. 2013, Hasanuzzaman et al. 2013), to our knowledge, proanthocyanidins have not previously been detected or characterised in *A. bilimbi* fruits or leaves. The leaves are traditionally used as a paste made with water for dermatological issues (skin rashes, itches, shingles, eczema, pimples) and against rheumatism (Ariharan et al. 2012). This information will be useful for

probing the health benefits of *A. bilimbi* fruits and leaves, for expanding food databases on proanthocyanidins (websites 1 and 2) and for enabling intake calculations, especially for populations consuming wild tropical and underutilised fruits and vegetables.

## **2. Materials and methods**

### *2.1. General*

Acetone (analytical reagent grade), acetonitrile (HPLC grade), dichloromethane (HPLC grade) and hydrochloric acid (37%, analytical reagent grade), were purchased from ThermoFisher Scientific Ltd (Loughborough, U.K.); ( $\pm$ )-taxifolin (98%); benzyl mercaptan (99%), epicatechin (EC) and catechin (C) ( $\geq 99\%$  HPLC) were purchased from Sigma-Aldrich (Poole, U.K.). Deionised water was obtained from a Milli-Q System (Millipore, Watford, U.K.).

### *2.2. Samples*

*Averrhoa bilimbi* leaves and fruits were harvested in December 2013 in a private botanical garden in Trois-Rivières, Guadeloupe, France. Any excess humidity was removed with kitchen paper, air-dried for a few hours, protected from direct light and immediately packed in an air-tight glass container and sent to Reading, U.K. by airplane (1-3 days). Upon arrival, leaves and fruits were freeze-dried and finely ground in an impeller SM1 cutting mill (Retsch, Haan, Germany) to pass a 1 mm sieve. The ground plant material was stored in the dark at room temperature.

### *2.3. Extraction and purification*

#### *2.3.1. Extractable proanthocyanidins*

Finely ground fruits (5.3 g) and leaves (5.5 g) were extracted using magnetic stirring for 1 h with acetone/water (125 mL; 7:3, v/v) and the solution was separated from the residue after filtration through a Büchner funnel. Acetone was removed under vacuum at 30 °C; the remaining aqueous solution was centrifuged for 3 min at 2045 x g and freeze-dried to give the extract (fruits = 1.6 g, yield = 31%; leaves = 0.8 g, yield = 14%). Acetone was allowed to evaporate from the plant residue in the fume cupboard overnight and protected from direct light before freeze-drying; these residues were used for the analysis of unextractable proanthocyanidins.

#### 2.4. Proanthocyanidin analysis

##### 2.4.1. Thiolytic of extractable proanthocyanidins

Acetone-water extracts (8 mg) were weighed in triplicates into screw cap glass tubes with a stirring magnet. Methanol (1.5 mL) was added followed by methanol acidified with concentrated HCl (3.3%; 500 µL) and benzyl mercaptan (50 µL). Tubes were capped and placed into a water bath at 40 °C for 1 h under vigorous stirring. The reaction was stopped by placing the tube in an ice bath for 5 min. Distilled water (2.5 mL) and the internal standard, taxifolin in methanol (500 µL; 0.1 mg/mL), were added and thoroughly mixed. The mixture was transferred into a 800 µL vial, closed with a crimp top and analyzed by HPLC-MS within 12 h (Ramsay et al. 2015).

##### 2.4.2. Thiolytic of in situ and unextractable proanthocyanidins

Whole freeze-dried fruits and leaves or the plant residues (200 mg), which remained after the aqueous acetone extraction, were reacted with the thiolytic reagent (2 mL methanol, 1 mL of 3.3% HCl in methanol, and 100 µL benzyl mercaptan) in triplicates as above. After the reaction, methanol (1 mL) was added to the mixture. The sample was mixed and centrifuged



at 2727 x g for 3 min and supernatant (1 mL) was transferred into another screw cap glass tube. Distilled water (9 mL) and internal standard, taxifolin in methanol (500 µL; 0.1 mg/mL), were added and thoroughly mixed. The mixture was transferred into a vial, closed with a crimp top and analysed by HPLC-MS as soon as possible or within the next 12 h.

## 2.5. Liquid chromatography-mass spectrometry (HPLC-MS) analysis

LC-MS was used to check for the presence of free flavan-3-ols in the plant materials and extract and to confirm the identity of terminal and extension units using an Agilent 1100 Series HPLC system and an API-ES instrument Hewlett Packard 1100 MSD detector (Agilent Technologies, Waldbronn, Germany). Samples (20 µl) were injected into the HPLC connected to an ACE C<sub>18</sub> column (3 µm; 250 x 4.6 mm; Hichrom Ltd, Theale, U.K.), which was fitted with a corresponding ACE guard column, at room temperature. The HPLC system consisted of a G1379A degasser, G1312A binary pump, G1313A ALS autoinjector, and G1314A VWD UV detector. Data were acquired with ChemStation software (version A 10.01 Rev. B.01.03). The flow rate was 0.75 ml/min using 1% acetic acid in water (solvent A) and HPLC-grade acetonitrile (solvent B). The following gradient programme was employed: 0-35 min, 36% B; 35-40 min, 36-50% B; 40-45 min, 50-100% B; 45-55 min, 100-0% B; 55-60 min, 0% B. Eluting compounds were recorded at 280 nm. Mass spectra were recorded in the negative ionisation scan mode between *m/z* 100 and 1000 using the following conditions: capillary voltage, -3000 V; nebuliser gas pressure, 35 psi; drying gas, 12 ml/min; and dry heater temperature, 350 °C (Ramsay & Mueller-Harvey, 2015). Flavan-3-ols and their benzyl mercaptan adducts were identified by their retention times and characteristic UV-VIS spectra between 220 and 595 nm. Peak areas of flavan-3-ols at 280 nm were integrated and quantified using molar response factors relative to taxifolin: 0.30 for catechin and epicatechin; 0.26 for their benzyl mercaptan adducts (Gea et al. 2011). This provided information on the

proanthocyanidin composition in terms of % terminal and % extension flavan-3-ol units (i.e. molar percentages). It also allowed calculation of the mean degree of polymerisation (mDP), % procyanidins (PC) and % *cis*- and *trans*-flavan-3-ols (molar percentages) (Gea et al. 2011).

## 2.6. Gel-NMR analysis

Samples were prepared as previously described (Grabber et al. 2013). Briefly, finely milled plant material (50 mg) was mixed in DMSO-*d*<sub>6</sub> (400 µL) and pyridine-*d*<sub>5</sub> (100 µL) and transferred to a 5 mm NMR tube. <sup>1</sup>H-<sup>13</sup>C correlation 2D NMR (HSQC) spectra were recorded at 27 °C on a Bruker Avance III 500 instrument equipped with TopSpin 2.4 software and a 5-mm BBI <sup>1</sup>H/<sup>13</sup>C gradient probe (Bruker, Coventry, U.K.). Spectral resonances were referenced to the residual signals of DMSO-*d*<sub>6</sub> (2.49 ppm for <sup>1</sup>H and 39.5 ppm for <sup>13</sup>C spectra) using 128 scans.

## 3. Results and Discussion

*Averrhoa bilimbi* fruits and leaves were analysed by thiolytic degradation with benzyl mercaptan for proanthocyanidin content and composition directly using the ground plant materials (i.e. *in situ* analysis) and also the aqueous acetone extracts and plant residues that remained from the solvent extractions. The thiolysis reaction released proanthocyanidin terminal units as flavan-3-ols and extension units as benzyl mercaptan derivatives, which were analysed by reverse-phase HPLC-MS (Ramsay & Mueller-Harvey, 2015). The proanthocyanidin contents and compositions are described in Table 1 for both fruits and leaves. Figures 3 and 4 illustrate the HPLC chromatograms of fruit and leaf proanthocyanidins after thiolysis. The total proanthocyanidin content in fruits is lower than in leaves (2.2 vs 4.5 g/100 g of dry weight). The average proanthocyanidin polymer size in fruits was also lower (mDP of 6 vs 9) than in leaves [*Note: no free flavan-3-ols could be detected in*

176 *the plant materials or extract before thiolysis*]. The key finding is that *A. bilimbi* fruits and  
177 leaves contained only pure procyanidins (PC) (Figure 2).  
178 Epicatechin accounted for 97% of the flavan-3-ol units in fruit proanthocyanidins and for  
179 99% of the leaf proanthocyanidins, with catechin accounting for the rest. Catechin and  
180 epicatechin occurred as terminal units in fruits and leaves, but epicatechin was the only  
181 extension unit. Catechin and epicatechin were assigned to peaks 1 and 2, respectively, at  
182 retention times of 23 min and 27 min (Figures 3 and 4), with ion fragments at  $m/z$  289.3 [ $M -$   
183  $H$ ] $^-$ . The epicatechin-benzyl mercaptan adduct was assigned to peak 3 at a retention time of  
184 43 min and generated ion fragments at  $m/z$  411.3 [ $M - H$ ] $^-$  and, after loss of the benzyl  
185 mercaptan molecule ( $-124$  amu) at  $m/z$  287.2.  
186 Unextractable proanthocyanidins were also investigated as they are often overlooked (Gea et  
187 al. 2011), yet their proportion can exceed extractable proanthocyanidins in foods and may  
188 thus represent a substantial amount of the dietary polyphenol intake (Pérez-Jiménez & Torres,  
189 2011). In fact, there were higher amounts of unextractable than extractable proanthocyanidins  
190 (fruits: 1.3 vs 0.8, leaves: 3.2 vs 1.3). The mDP values were also higher in the unextractable  
191 than the extractable proanthocyanidins (fruits: 6.7 vs 4.6, leaves: 13.7 vs 6.5) and agrees with  
192 our previous findings (Gea et al. 2011; Mechineni et al. 2014, Wang et al. 2015).  
193 A gel-NMR analysis ( $^1H - ^{13}C$  HSQC) was also applied directly to the milled leaves and  
194 fruits in order to verify the results from thiolysis. This analysis revealed distinct signals for  
195 procyanidins: signals at 6.7 and 120 ppm could be assigned to H/C-2'/5'/6' and signals at 6.0  
196 and 95 ppm were assigned to H/C-6 and H/C-8 (Figure 2). This confirmed that these  
197 proanthocyanidins were procyanidins and B-type linkages. The presence of A-type  
198 proanthocyanidins would have been indicated by signals at approximately H/C-3 (4.0/66  
199 ppm) and H/C-3 (4.5/27.9 ppm) but it was not detected (Appeldoorn et al. 2009).

Although proanthocyanidins have limited bioavailability and are relatively stable in the gastrointestinal tract (Serra et al. 2010), some evidence exists for their depolymerisation by intestinal microorganisms (Pérez-Maldonado & Norton et al. 1996; Touriño et al. 2009). Studies have also shown that procyanidins with lower mDP (< 4) are most likely absorbed in the colon after metabolisation by the gut microbiota and their metabolites could be detected in the plasma (Kerimi & Williamson, 2015). Proanthocyanidins and their metabolites can act as antioxidants *in vivo* (López-Andrés et al. 2013) and modulate key biological pathways *in vivo* (Nantz et al. 2013; Vertraetan et al. 2013).

#### **4. Conclusion**

This study revealed the presence of pure procyanidins in *A. bilimbi* fruits and leaves with a moderate average proanthocyanidin size. Epicatechin accounted for 94% to 97% of the flavan-3-ol subunits and these polymers had mean degrees of polymerisation that ranged from 5 to 14. Pure proanthocyanidins are not so common, especially in edible fruits. Therefore *A. bilimbi* fruits and leaves are potentially valuable sources for proanthocyanidins that could be used for future research into their nutritional and health benefits.

#### **Acknowledgements**

The authors would to thank Chris Drake, Honorata Ropiak and Christos Fryganas for their support in the laboratory and Mrs Anita Bazir for providing the plant samples.

#### **Funding sources**

This work was supported by the European Commission (Marie Curie Initial Training Network) under Grant PITN-GA-2011-289377, “LegumePlus”.

## References

- Ambili, S., Subramoniam, A., Nagarajan. N.S. (2009). Studies on the antihyperlipidemic properties of *Averrhoa bilimbi* fruit in rats. *Planta Medicinal*, 75, 55–58.
- Appeldoorn, M.M., Sanders, M., Vincken, J.P., Cheynier, V., Le Guernevé, C, Hollman, P.C.H., Gruppen, H. (2009). Efficient isolation of major procyanidin A-type dimers from peanut skins and B-type dimers from grape seeds. *Food Chemistry*, 117, 713–720.
- Ariharan, V.N., Kalirajan, K., Meena Devi, V.N., Nagendra Prasad, P. (2012). An exotic fruit which forms the new natural source for vitamin-C. *Rasayan Journal of Chemistry*, 5, 356–359.
- Ashok Kumar, K., Gousia, S.K., Anupama, M., Naveena Lavanya Latha J. (2013). A review on phytochemicals constituents and biological assays of *Averrhoa bilimbi*. *International Journal of Pharmacy and Pharmaceutical Science Research*, 3, 136–139.
- Bakul, G., Unni, V.N., Seethaleksmy, N.V., Mathew, A., Rajesh, R., Kurien, G., Rajesh, J., Jayaraj, P.M., Kishore, D.S., Jose, P.P. (2013). Acute oxalate nephropathy due to *Averrhoa bilimbi* fruit juice ingestion. *Indian Journal of Nephrology*, 23, 297–300.
- Gea, A., Stringano, E., Brown, R.H., Mueller-Harvey, I. (2011). *In situ* analysis and structural elucidation of sainfoin (*Onobrychis viciifolia*) tannins for high-throughput germplasm screening. *Journal of Agricultural and Food Chemistry*, 59, 495–503.

249 Grabber, J.H., Zeller, W.E., Mueller-Harvey, I. (2013). Acetone enhances the direct analysis  
 250 of procyanidin- and prodelphinidin-based condensed tannins in *Lotus* species by the  
 251 butanol–HCl–iron assay. *Journal of Agricultural and Food Chemistry* 61, 2669–2678.  
 252

253 Hasanuzzaman, M., Ramjan Ali, M., Marjan, H., Sourov, K., Mohammad Safiqul, I. (2013).  
 254 Evaluation of total phenolic content, free radical scavenging activity and phytochemical  
 255 screening of different extracts of *Averrhoa bilimbi* (fruits). *International Current*  
 256 *Pharmaceutical Journal* 2, 92–96.  
 257

258 Kerimi, A. & Williamson, G. (2015). The cardiovascular benefits of dark chocolate. *Vascular*  
 259 *Pharmacology*, 71, 11–15.  
 260

261 Koo Hui, M., Suhaila, M. (2001). Flavonoid (myricetin, quercetin, kaempferol, luteolin, and  
 262 apigenin) content of edible tropical plants. *Journal of Agricultural and Food Chemistry*, 49,  
 263 3106–3112.  
 264

265 López-Andrés P., Luciano G., Vasta V., Gibson T.M., Biondi L., Priolo A., Mueller-Harvey I.  
 266 (2013). Dietary quebracho tannins are not absorbed, but increase the antioxidant capacity of  
 267 liver and plasma in sheep. *British Journal of Nutrition*, 110, 632–639.  
 268

269 Mechineni, A., Kommuru, D.S., Gujja, S., Mosjidis, J. A., Miller, J. E., Burke, J. M., Ramsay,  
 270 A., Mueller-Harvey, I., Kannan G., Lee, J. H., Kouakou, B., Terrill, T. H. (2014). Effect of  
 271 fall-grazed sericea lespedeza (*Lespedeza cuneata*) on gastrointestinal nematode infections of  
 272 growing goats. *Veterinary Parasitology*, 204, 221–228.  
 273

274 Morton, J. (1987). Bilimbi. In Julia F. Morton, *Fruits of warm climates*. (p. 128–129). Miami,  
 275 FL, USA. Website (accessed 1 Oct 2015). <https://hort.purdue.edu/newcrop/morton/index.html>  
 276

277 Nantz, M., Rowe, C., Muller, C., Creasy, R., Colee, J., Khoo, C., Percival, S. (2013).  
 278 Consumption of cranberry polyphenols enhances human  $\gamma\delta$ -T cell proliferation and reduces  
 279 the number of symptoms associated with colds and influenza: a randomized, placebo-  
 280 controlled intervention study. *Nutrition Journal*, 12, 161– 170.

281

282 Pérez-Jiménez, J. & Torres, J.L. (2011). Analysis of nonextractable phenolic compounds in  
 283 foods: The current state of the art. *Journal of Agricultural and Food Chemistry*, 59, 12713–  
 284 12724.

285

286 Pérez-Maldonado, R. A., & Norton, B. W. (1996). The effects of condensed tannins from  
 287 *Desmodium intortum* and *Calliandra calothyrsus* on protein and carbohydrate digestion in  
 288 sheep and goats. *British Journal of Nutrition*, 76, 515-533.

289

290 Ramsay, A. & Mueller-Harvey, I. (2015). *Cassia alata* leaves are a good sources of  
 291 propelargonidins. *Natural Product Research*  
 292 (<http://dx.doi.org/10.1080/14786419.2015.1108976>).

293

294 Rush, D. (2001). Maternal nutrition and perinatal survival. *Journal of Health Population and*  
 295 *Nutrition*, 19, 220–264.

296

297 Scoones, I., Melnyk, M., Pretty, J.N. (1992). The hidden harvest: wild foods and agricultural  
 298 systems. A literature review and annotated bibliography. International Institute for

299 Environment and Development, London, UK.

300

301 Serra, A., Macia, A., Romero, M.-P., Valls, J., Blade, C., Arola, L. Motilva, M.-J. (2010).

302 Bioavailability of procyanidin dimers and trimers and matrix food effects in *in vitro* and *in*

303 *vivo* models. *British Journal of Nutrition*, 103, 944–952.

304

305 Touriño, S., Fuguet, E., Pilar Vinardell, M., Torres, J. L. (2009). Phenolic metabolites of

306 grape antioxidant dietary fiber in rat urine. *Journal of Agricultural and Food Chemistry*, 57,

307 11418–11426

308

309 Vertraeten, S. V., Jagers, G. K., Fraga, C. G., Oteiza, P. I. (2013). Procyanidins can interact

310 with Caco-2 cell membrane lipid rafts: Involvement of cholesterol. *Biochimica et Biophysica*

311 *Acta*, 1828, 2646–2653.

312

313 Wang, Y., McAllister, T. A., Acharya, S. (2015). Condensed tannins in sainfoin: composition,

314 concentration, and effects on nutritive and feeding value of sainfoin forage. *Crop Science*, 55,

315 13–22.

316

317 Website 1: <http://www.ars.usda.gov/SP2UserFiles/Place/80400525/Data/PA/PA.pdf> (accessed

318 30 Sep 2015).

319

320 Website 2: <http://phenol-explorer.eu/> (accessed 30 Sep 2015).

321

322



**Figure captions**

Fig. 1. *Averrhoa bilimbi* fruits.

**Fig. 2.** Structure of a procyanidin dimer (catechin–(4→8)–epicatechin).

**Fig. 3.** HPLC chromatogram at 280 nm after *in situ* thiolysis of proanthocyanidins from *Averrhoa bilimbi* fruits: **1**, catechin; **2**, epicatechin; **3**, epicatechin-benzyl mercaptan.

**Fig. 4.** HPLC chromatogram at 280 nm after *in situ* thiolysis of proanthocyanidins from *Averrhoa bilimbi* leaves: **1**, catechin; **2**, epicatechin; **3**, epicatechin-benzyl mercaptan.

346

347 **Table 1**348 Content and composition of *in situ*, extractable and unextractable proanthocyanidins in *Averrhoa bilimbi* fruits and leaves (n= 3).

Proanthocyanidins	Content	mDP	PC	<i>cis</i>	<i>trans</i>	Terminal units		Extension units
	(g/100 g DW)		(%)	(%)	(%)	(%)		(%)
						C	EC	EC
Fruits								
<i>In situ</i>	2.2 (0.1)	6.1 (0.1)	100	96.8 (0.2)	3.2 (0.2)	3.1 (0.2)	13.5 (0.5)	83.4 (0.2)
Extractable	0.8 (0.1)	4.6 (0.1)	100	94.4 (0.1)	5.6 (0.1)	5.7 (0.1)	15.9 (0.1)	78.4 (0.1)
Unextractable	1.3 (0.1)	6.7 (0.6)	100	96.9 (1.5)	3.1 (1.6)	3.2 (1.6)	11.8 (0.2)	85.0 (1.3)
Leaves								
<i>In situ</i>	4.5 (0.2)	9.2 (0.1)	100	99.5 (0.1)	0.5 (0.1)	0.5 (0.1)	10.4 (0.1)	89.1 (0.1)
Extractable	1.3 (0.1)	6.5 (0.3)	100	98.5 (0.3)	1.5 (0.3)	1.5 (0.3)	13.8 (0.5)	84.7 (0.8)
Unextractable	3.2 (0.1)	13.7 (0.1)	100	99.6 (0.1)	0.4 (0.1)	0.4 (0.1)	6.9 (0.2)	92.7 (0.1)

349

350 DW: dry weight; mDP: mean degree of polymerisation; PC: procyanidins; C: catechin (a 2,3-*trans* flavan-3-ol); EC: epicatechin (a 2,3-*cis*

351 flavan-3-ol); % represents relative molar percentages.

352