

A comprehensive meta-analysis of common genetic variants in autism spectrum conditions

Article

Published Version

Creative Commons: Attribution 4.0 (CC-BY)

Open Access

Warrier, V., Chee, V., Smith, P., Chakrabarti, B. ORCID: <https://orcid.org/0000-0002-6649-7895> and Baron-Cohen, S. (2015) A comprehensive meta-analysis of common genetic variants in autism spectrum conditions. *Molecular Autism*, 6 (1). 49. ISSN 2040-2392 doi: 10.1186/s13229-015-0041-0 Available at <https://centaur.reading.ac.uk/52481/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

Published version at: <http://dx.doi.org/10.1186/s13229-015-0041-0>

To link to this article DOI: <http://dx.doi.org/10.1186/s13229-015-0041-0>

Publisher: BioMed Central

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online

RESEARCH

Open Access



A comprehensive meta-analysis of common genetic variants in autism spectrum conditions

Varun Warrier^{1*†}, Vivienne Chee^{1†}, Paula Smith¹, Bhismadev Chakrabarti^{1,2*} and Simon Baron-Cohen^{1,3}

Abstract

Background: Autism spectrum conditions (ASC) are a group of neurodevelopmental conditions characterized by difficulties in social interaction and communication alongside repetitive and stereotyped behaviours. ASC are heritable, and common genetic variants contribute substantial phenotypic variability. More than 600 genes have been implicated in ASC to date. However, a comprehensive investigation of candidate gene association studies in ASC is lacking.

Methods: In this study, we systematically reviewed the literature for association studies for 552 genes associated with ASC. We identified 58 common genetic variants in 27 genes that have been investigated in three or more independent cohorts and conducted a meta-analysis for 55 of these variants. We investigated publication bias and sensitivity and performed stratified analyses for a subset of these variants.

Results: We identified 15 variants nominally significant for the mean effect size, 8 of which had *P* values below a threshold of significance of 0.01. Of these 15 variants, 11 were re-investigated for effect sizes and significance in the larger Psychiatric Genomics Consortium dataset, and none of them were significant. Effect direction for 8 of the 11 variants were concordant between both the datasets, although the correlation between the effect sizes from the two datasets was poor and non-significant.

Conclusions: This is the first study to comprehensively examine common variants in candidate genes for ASC through meta-analysis. While for majority of the variants, the total sample size was above 500 cases and 500 controls, the total sample size was not large enough to accurately identify common variants that contribute to the aetiology of ASC.

Keywords: Meta-analysis, Association, Genetic variants, Insertions, Autism spectrum conditions

Background

Autism spectrum conditions (ASC) are a group of neurodevelopmental conditions characterized by difficulties in social interaction and communication alongside unusually repetitive and stereotyped behaviour and unusually narrow interests [1]. ASC has an estimated heritability of around 50 % [2, 3], and common variants contribute to a significant proportion of the variability in the condition [3, 4]. ASC is polygenic and genetic variants, in addition to environmental, epigenetic and hormonal factors, contribute to ASC risk and phenotypic variability [5].

Sequencing and copy number variation analyses have identified a number of rare, highly penetrant, possibly causative variants. Strategies to identify common variants through genome-wide association studies have failed to produce consistent, replicable results across cohorts [5]. This may be attributed to many factors, including smaller than required sample size to adequately power these studies to identify variants with small effects. Over the last 15 years, a large number of studies have investigated common variants in candidate genes for ASC [6] typically investigating variants in a small number of genes using a relatively small sample size. These studies have provided some evidence of the association of a few genes with ASC, though they are not rigorous enough to definitively identify variants and results vary based on ethnicity, sample size, study methodology and clinical ascertainment [6].

* Correspondence: vw260@medschl.cam.ac.uk; b.chakrabarti@reading.ac.uk
Bhismadev Chakrabarti and Simon Baron-Cohen are equal senior co-authors.
†Equal contributors

¹Autism Research Centre, Department of Psychiatry, University of Cambridge, Douglas House, 18B Trumpington Road, Cambridge CB2 8AH, UK
Full list of author information is available at the end of the article

One method to investigate the underlying effect using summary level data is meta-analysis [7]. Though not without limitations, meta-analysis provides a fairly robust statistical framework to systematically analyse effect sizes [7]. Further, the combined power of a meta-analysis greatly exceeds the power of the individual studies in a meta-analysis [7].

In the field of psychiatric genetics, studies have comprehensively investigated existing candidate gene studies and used meta-analysis to investigate genetic associations [8–10]. In the field of autism genetics, such an overarching study is lacking and no study, to our knowledge, has provided a comprehensive overview of ASC genetics. To bridge this gap, we reviewed the existing literature for 552 genes implicated in ASC. Using a strict inclusion criteria, we identified common variants in 27 genes that were investigated in three or more independent cohorts. We performed meta-analyses, sensitivity analyses and subgroup analyses for these common variants and checked for publication bias in a subset of these common variants. This is the first comprehensive study of candidate gene associations in ASC.

Methods

Literature search and inclusion criteria

A preliminary literature search of genes associated with ASC was performed using SFARI gene (<https://gene.sfari.org/>) and HuGE Navigator (<http://hugenavigator.net/>). Since both of these databases do not completely document the available literature, we additionally searched PubMed, Scopus and Google Scholar. The search terms used were 'Gene name' or 'variant ID' and 'Autism' or 'Autistic Disorder' or 'Asperger Syndrome'.

Studies were included in the meta-analysis if: (1) they reported effect sizes or statistics to measure effect sizes and confidence intervals; (2) the studies were either a case-control association study or a transmission disequilibrium study of autism; (3) the variants did not deviate from Hardy-Weinberg Equilibrium (HWE) in the control group or if the sample size was too small to effectively calculate HWE due to sampling effect. Though we checked for HWE in family-based studies, this was not a requirement for including these studies as the study design overcomes the issue of population stratification; (4) cases had a diagnosis of an autism spectrum condition (Autism, PDDNOS, Asperger Syndrome) according to DSM-IV, DSM-5 or ICD-10 criteria; (5) the global minor allele frequency (MAF) of the variant investigated was greater than 0.01; (6) the studies were reported in English and (7) the common variants were investigated in independent cohorts. Authors of the articles were contacted if sufficient information was absent to use the data for meta-analysis. In addition to the published studies, we used unpublished genotype data from two cohorts from our research group at the Autism Research Centre, University

of Cambridge. These cohorts are labelled 'Chakrabarti [11]' and 'Warrier [12]' in the current study. The characteristics of the two cohorts are described elsewhere [11, 12]. Details of genotyping and statistical analysis are provided in Additional file 1. We did not include data from genome-wide association studies (GWAS) as there is an overlap between participants in the candidate gene association studies and the genome-wide association studies. Since we had access to only summary data, it was impossible to ascertain the degree of overlap and remove participants accordingly. Literature search and study inclusion was performed independently by two researchers (VC and VW) from March 2014 to September 2014.

Statistical analyses

Meta-analysis was performed only if variants were investigated in three or more independent cohorts. Family-based association tests (FBATs) studies were not included as effect sizes are not calculated in FBA. For variants investigated in five or more independent cohorts, we performed a complete meta-analysis. This included the calculation of effect size and publication bias, sensitivity analysis and subgroup analysis. For variants investigated in three to five independent cohorts, we performed a partial meta-analysis restricted to the calculation of mean effect size. We did not perform a meta-analysis for variants investigated in fewer than three cohorts as there was insufficient power to significantly investigate the underlying effect. For variants with P values <0.05 we calculated fail-safe N .

All analyses were performed using Comprehensive Meta-Analysis version 2.0 [13]. Meta-analysis was performed using the inverse-variance weighted method. Heterogeneity in the reported effects were examined using a fixed and a random effects model. Heterogeneity was measured using I^2 statistics in conjunction with Q-statistics. A fixed effect model was applied if the P value for Q-statistics was above 0.05 and I^2 was below 60. The random effects model was used if either the P value was below 0.05 or I^2 was above 60, as an I^2 above 60 indicates that 60 % of the total observed variation is due to true heterogeneity [7, 10].

Egger's regression in conjunction with a funnel plot was used to assess publication bias. Sensitivity analyses were performed by removing each study from the meta-analysis and calculating the mean effect size for the remaining studies. This analysis was used to assess the contribution of each study to the final weighted effect in the analysis. Additionally, for the variants with P values <0.05 , we computed both classic fail-safe N and Orwin's fail-safe N to check the number of studies required to make the P value non-significant and make the effect size trivial respectively. For Orwin's fail-safe N , the non-significant odds ratio (OR) was kept at 1.05 or 0.95 depending on the effect direction. While this is certainly not a trivial effect size, it

is difficult to identify variants with such small effects with precision given the sample sizes in the meta-analysis. Subgroup analysis was performed after stratifying based on ethnicity or study methodology to check if either of these variables affected the final effect size. We conducted the subgroup analysis only for variants investigated in five or more independent cohorts. Meta-analysis was performed only if there were at least three independent cohorts after stratification to account for power considerations.

OR and 95 % confidence intervals (CI) were used to calculate the mean effect size. For transmission disequilibrium tests (TDT), odds ratios were calculated according to methods laid out by Kazeem and Farall [14]. Where possible, OR and CI were calculated using allele numbers for case-controls (CC) and transmitted and non-transmitted numbers for TDT. Where information of OR and CI was provided for the complement allele of the allele investigated in the study, the log odds ratio (LOR) and standard error (SE) were calculated and used in the meta-analysis.

Age was not regarded a confounding variable as ASC is a neurodevelopmental condition, and genetic variations are largely invariant across lifespan. However, ASC has a male-female ratio of 5:1 [5], and sex is a potential confounding variable as gene expressions can vary based on sex. However, there was insufficient data to conduct a stratified analysis based on sex, so this is a limitation of the current study. Finally, due to the large number of studies carried out, we adopted a more conservative statistical significance threshold of 0.01. This is similar to what was used in a similar comprehensive meta-analysis of obsessive-compulsive disorder [10]. We did not carry out a Bonferroni correction as the sample for each variant investigated was very different, and as a result, multiple tests were not carried out on the same sample.

Analysis of the PGC dataset

While we did not choose to include data from available GWAS due to potential overlap of participants, we compared the results using the publicly available GWAS dataset from the Psychiatric Genomics Consortium (PGC). In the ASC cohort of the PGC dataset, 4788 trio cases and 4788 trio pseudocontrols as well as 161 cases and 526 controls have been genotyped. Details of the cohort, genotyping methods and statistical analysis are given elsewhere [15]. We searched for effect sizes and *P* values for variants with *P* values <0.05 in our meta-analysis. The autism PGC dataset is the largest available and accessible GWAS dataset for autism. The sample size of any of the variants investigated through meta-analysis in the study, except rs4141463 in *MACROD2*, is smaller than the sample size of the PGC autism dataset. Despite this, the PGC dataset is underpowered to detect variants with small effects. We were motivated to investigate the top variants in our study in the PGC dataset to ascertain if

the candidate variants were at least nominally significant (*P* < 0.05) and if the effect direction was concordant between the two samples.

Results

Literature review


We identified 463 genes that have been tested for genetic association using HuGE Navigator (as of August 2014). SFARI Gene reports 616 genes to be associated with autism (as of August 2014). Only 185 of these genes have been examined in ASC using genetic association studies. Of these, we identified 89 genes from the SFARI Gene list that were not included in the HuGE Navigator list, bringing the total list of potential genes to 552. We did not identify any additional genes from AutismKB database. Thus, we reviewed 552 genes in total for the meta-analysis.

Scopus, Google Scholar and PubMed were searched for publications relating to ASC and any of the 552 genes. We searched for common variations in these genes that have been investigated for ASC in at least three independent cohorts. Using the eligibility criteria outlined in the methods section, we identified 27 genes that could be taken forward for meta-analysis. In total, there were 58 common variants across these 27 genes that were investigated in our meta-analysis. Details of the studies included and excluded for the 27 genes are given in Additional file 1: Tables S1 and S2.

We next searched the literature for existing meta-analyses for the 58 variants and 27 genes in ASC, identifying existing meta-analyses for *OXTR* [16], *RELN* [17], *SLC6A4* [18], *HOXA1* [19], *HOXB1* [19] and *MTHFR* [20]. Detailed information of previous meta-analyses is provided in Additional file 1. As we had additional data and different inclusion criteria, we performed meta-analyses for all the variants in these six genes except rs723387731 in *HOXB1*, STin2 VNTR in *SLC6A4* and the GGC repeat in *RELN*. These three variants were excluded from the current meta-analyses as we could not identify additional data to add to the original meta-analyses. For the sake of comprehensiveness, we have included the data for these three variants in our table. Of the remaining 55 variants, we conducted a complete meta-analysis for 20 variants and a partial meta-analysis for 35 variants. A flow chart of the study protocol is given in Fig. 1.

Mean effect sizes

Effect sizes for 15 variants in 12 genes had *P* values below 0.05. Nine of these variants had a *P* value below 0.01. The most significant association was rs167771 in *DRD3* (OR = 1.822, *P* value = 9.08×10^{-6}). Seven other significant associations with *P* values <0.01 were in *CNTNAP2* (rs7794745, OR = 0.887, *P* value = 0.001), *RELN* (rs362691, OR = 0.832, *P* value = 3.93×10^{-5}), *OXTR* (rs2268491, OR = 1.31, *P* value = 0.004), *SLC25A12* (rs2292813, OR = 1.372, *P* value = 0.001 and rs2056202,



OR = 1.227, P value = 0.002), *EN2* (rs1861972, OR = 1.125, P value = 0.006) and *MTHFR* (rs1801133, OR = 1.370, P value = 0.010). As expected for common variants in ASC, the odds ratios for the alleles tested were small and lay between 0.781 (0.446–1.368) for *MAOA* uVNTR and 1.822 (1.398–2.375) for *DRD3* rs167771. Details of the variants analysed, model used and the P values are provided in Table 1. Forest plots for the nine most significant variants are in Additional file 1: Figures S1–S8.

Subgroup analyses

We performed subgroup analyses, stratifying by ethnicity and study methodology, for variants originally investigated in five or more independent cohorts. In the stratified analyses, six variants had P values below 0.05. Of these, the most significant three variants (rs2292813 and rs2056202-*SLC25A12*, rs362691-*RELN*) were also significant in the non-stratified analyses. Stratification did not increase the significance for these variants. A variant in *EN2* (rs1861973) was significant after stratifying based on both ethnicity (Caucasian only) and study methodology (TDT). Another variant in *EN2* (rs1861972) was significant after stratifying for study methodology (TDT). Finally, the *STIN2* variant in *SLC6A4* also exhibited a significant trend in the Caucasian-only subgroup. This result indicates that at least for a few variants implicated in ASC, ethnicity and study methodology can potentially influence the outcome. Results of the subgroup analyses are provided in Table 2. Forest plots for the significant and nominally significant subgroup analyses are provided in Additional file 1: Figures S9–S15.

Publication bias and sensitivity analyses

Publication bias was significant only for one variant, rs2254298 in *OXTR* (Egger's test (two-tailed) P value = 0.03). However, the mean effect size for the variant was not significant (P value = 0.425). Notably, sensitivity was significant for some variants. Of the nine variants with P values below 0.01, we performed sensitivity analyses on the six variants with data from more than five independent cohorts (rs7794745, rs362691, rs2292813, rs2056202, rs1861972, and rs1801133). For rs1801133, most studies contributed approximately equally, with the exception of two studies [21, 22]; both of these studies lowered the OR. A re-analysis of the data after removing either of the two studies decreased the P value of the OR (original P value = 0.010, P value after removing Park et al., 2014 [21] = 0.006; P value after removing Schmidt et al., 2011 [22] = 0.003). For rs2056202, the removal of data from one study [23] increased the P value from P value = 0.002 to P value = 0.088. Sensitivity was not an issue for the remaining four variants that were significant. However, of the nominally significant variants, sensitivity was an issue for rs4446909, rs736707 and rs1861972. Forest graphs of the sensitivity analyses for these five variants are provided in Additional file 1: Figures S16–S20.

Analysis of the PGC dataset

Of the 15 nominally significant variants in the current meta-analyses, 11 were genotyped in the PGC GWAS cohort, and none were found to be significant. Effect direction was concordant for 8 of the 11 variants between both the datasets. Effect sizes, as expected due to the larger sample size, were smaller in the PGC dataset for all the 11 variants, and the odds ratios were closer to 1. Total sample size was also not a significant predictor of concordance of effect direction between the two datasets. However, inspection of the datasets indicate that with the exception of rs2056202 in *SLC25A12*, the other three variants discordant for effect direction were analysed in small samples in the meta-analysis (see Table 2).

The lack of significance for 11 of the 15 variants in the PGC dataset forces us to re-evaluate the significance of the remaining four variants. For two variants, the classic fail-safe N is very small (three for rs4446909 in *ASMT*, and zero for rs4717806 in *STX1A*). The latter variant was analysed using a fixed effect model and becomes non-significant when analysed using a random effect model. For the remaining two variants (rs1861972 in *EN2* and rs362691 in *RELN*), the classic fail-safe N is above 10. The sample sizes, however, are modest. These analyses indicate that the first two variants are likely to be false positives. With rs1861972, the significance in P value is driven largely by the TDT-only subset in the original analysis (P value = 0.013, see Table 2). Both a case-control only subset and a Caucasian-only subset were not significant

Table 1 Summary of mean effect size analyses

S. No	Gene	Variants	Allele	Global MAF	Data sets	Mean OR (95% CI)	Z-Value	P-Value	Model (I2 value)	Total cases	Total controls	Trios	PGC P-value	Effect direction (odds ratio)	Classic fail-safe N	Orwin's fail safe N (OR = 1.05 or 0.95)
1	DRD3	rs167771	G vs A	G=0.4113	3	1.822 (1.398-2.375)	4.44	9.08E-06	Fixed effect (60)	580	754	0	0.6	discordant (0.980)	7	34
2	RELN	rs362691	C vs G	C=0.1210	8	0.832 (0.763-0.908)	-4.11	3.93E-05	Fixed effect (33.2)	765	765	303	NA	NA	12	21
3	SLC25A12	rs2292813	C vs T	T=0.2085	6	1.372 (1.161-1.621)	3.72	1.97E-04	Fixed effect (0)	465	450	1220	0.78	concordant (1.014)	5	25
4	CNTNAP2	rs7794745	A vs T	A=0.4946	4	0.887 (0.828-0.950)	-3.45	1.00E-03	Fixed effect (21.2)	322	524	2236	0.18	concordant (0.9594)	9	6
5	SLC25A12	rs2056202	T vs C	T=0.2420	8	1.227 (1.079 -1.396)	3.12	2.00E-03	Fixed effect (6.5)	756	1211	1220	0.99	discordant (0.9993)	6	26
6	OXTR	rs2268491	T vs C	T=0.2137	4	1.31 (1.092 -1.572)	2.91	4.00E-03	Fixed effect (0)	282	440	458	0.54	concordant (1.026)	3	19
7	EN2	rs1861972	A vs G	G=0.242	8	1.125 (1.035-1.224)	2.75	6.00E-03	Fixed effect (57.6)	669	1704	953	NA	NA	16	12
8	MTHFR	rs1801133	T vs C	A=0.2454	10	1.370 (1.079-1.739)	2.59	1.00E-02	Random effects (88.2)	2280	7235	0	0.57	concordant (1.018)	80	40
9	ASMT	rs4446909	G vs A	A=0.1741	5	1.195 (1.038-1.375)	2.48	1.30E-02	Fixed effect (0)	1066	1074	0	NA	NA	3	14
10	MET	rs38845	A vs G	A=0.3634	3	1.322 (1.013-1.724)	2.41	1.60E-02	Random effects (66.5)	405	594	419	0.2	concordant (1.04)	13	15
11	SLC6A4	rs2020936	T vs C	G=0.228	4	1.244 (1.036-1.492)	2.35	1.90E-02	Fixed effect (33.9)	0	0	1068	0.78	concordant (1.01)	3	14
12	STX1A	rs4717806	A vs T	A=0.2322	4	0.851 (0.741-0.978)	-2.28	2.30E-02	Fixed effect (35.3)	653	1007	375	NA	NA	0	9
13	RELN	rs736707	T vs C	G=0.3660	9	1.269 (1.030-1.563)	2.24	2.50E-02	Random effects (76.5)	975	1695	196	0.31	concordant (1.035)	126	48
14	PON1	rs662	A vs G	T=0.4571	3	0.794 (0.642-0.983)	-2.12	3.40E-02	Fixed effect (17.5)	334	641	0	0.07	discordant (1.058)	0	11
15	OXTR	rs237887	G vs A	G=0.3998	4	1.163 (1.002-1.349)	1.99	4.70E-02	Fixed effect (0)	282	440	458	0.94	concordant (1.002)	0	9
16	STX1A	rs6951030	G vs T	G=0.1771	4	1.383 (0.995-1.922)	1.93	5.40E-02	Random effects (76.7)	653	1007	375				
17	OXTR	rs2268493	C vs T	C=0.2049	3	0.845 (0.701-1.019)	-1.76	7.80E-02	Fixed effect (54.5)	574	1201	0				
18	ASMT	rs5989681	G vs C	NA	5	1.135 (0.984 - 1.308)	1.74	8.20E-02	Fixed effect (0)	1066	1074	0				
19	HOXB1					rs72338773*18	INS vs nINS	NA	8	1.36 (0.97- 1.33)	NA		1.18E- 01	Fixed Effect (NA)	362	448

Table 1 Summary of mean effect size analyses (Continued)

20	RELN	rs2073559	C vs T	C=0.4746	3	0.955 (0.900-1.014)	-1.5	1.35E-01	Fixed effect (64.5)	437	493	473
21	RELN	GGC repeat*16	NA	NA	7	1.11 (0.80-1.54)	NA	1.53E-01	Fixed effect (0)	878	1170	167
22	GLO1	rs2736654	A vs C	G=0.2873	4	1.307 (0.882 - 1.936)	1.34	1.82E-01	Random effects (68.7)	857	680	0
23	PON1	rs854560	A vs T	T=0.1827	3	1.140 (0.931 - 1.395)	1.27	2.05E-01	Fixed effect (0)	334	641	0
24	TPH2	rs11179000	T vs A	T=0.3988	3	1.130 (0.934-1.366)	1.26	2.08E-01	Fixed effect (0)	224	260	352
25	MET	rs1858830	G vs C	G=0.4575	8	0.905 (0.773-1.061)	-1.23	2.19E-01	Random effect (67.5)	1975	1589	798
26	OXTR	rs2268490	T vs C	T=0.2584	5	1.135 (0.920-1.400)	1.18	2.38E-01	Fixed effect (0)	292	761	458
27	OXTR	rs2301261	A vs G	T=0.1248	4	1.127 (0.889-1.430)	0.99	3.22E-01	Fixed effect (39.1)	650	1300	0
28	HOXA1	rs10951154	G vs A	C=0.2192	13	0.925 (0.791-1.081)	-0.98	3.28E-01	Fixed effect (35.7)	705	998	425
29	BDNF	rs6265	G vs A	T=0.2013	3	0.919 (0.763-1.107)	-0.89	3.72E-01	Fixed effect (0)	303	469	140
30	HTR2A	rs6311	A vs G	T=0.4435	6	0.871 (0.643-1.181)	-0.89	3.74E-01	Random effects (74.8)	179	313	396
31	ITGB3	rs5918	C vs T	C=0.0889	3	0.866 (0.630-1.191)	-0.88	3.77E-01	Fixed effect (37.1)	139	165	363
32	MAOA	uVNTR	short vs long	NA	3	0.781 (0.446 - 1.368)	-0.86	3.87E-01	Random effects (72)	436	469	0
33	MACROD2	rs4141463	T vs C	C=0.3818	7	0.913 (0.734-1.135)	-0.82	4.11E-01	Random effects (87.1)	1170	35307	1158
34	OXTR	rs2254298	A vs G	A=0.2071	5	0.813 (0.489-1.352)	-0.8	4.25E-01	Random effects (82.5)	650	1306	57
35	ASMT	rs6644635	C vs T	NA	4	1.056 (0.906 -1.230)	0.69	4.88E-01	Fixed effect (29.3)	788	819	0
36	SLC6A4	rs2020942	A vs G	T=0.2550	3	1.062 (0.881-1.281)	0.63	5.28E-01	Fixed effect (0)	0	0	678
37	OMG	rs11080149	A vs G	T=0.0409	4	0.847 (0.477 - 1.506)	-0.56	5.72E-01	Random effects (43.8)	65	131	431
38	ADA	rs7359837	G vs A	A=0.0282	3	1.375 (0.401 - 4.717)	0.51	6.13E-01	Random effects (89.1)	334	445	0
39	OXTR	rs237894	G vs C	C=0.1615	5	0.961 (0.818-1.129)	-0.48	6.26E-01	Fixed effect (4)	292	761	458
40	OXTR	rs53576	A vs G	A=0.3894	5	0.966 (0.839-1.113)	-0.48	6.31E-01	Fixed effect (44.9)	650	1300	57

Table 1 Summary of mean effect size analyses (*Continued*)

41	<i>OXTR</i>	rs2268494	A vs T	A=0.0683	4	1.076 (0.760 -1.510)	0.42	6.73E-01	Fixed effect (0)	76	99	458
42	<i>SLC6A4</i>	STin2 VNTR*17	12 vs 9/ 10	NA	8	1.129 (0.819–1.558)	NA	6.73E-01	Random effects (68.7)	0	0	814
43	<i>NF1</i>	GxAlu	9 vs non-9	NA	4	1.131 (0.633 - 2.022)	0.42	6.77E-01	Random effects (85.7)	262	312	0
44	<i>GRIK2</i>	rs2227281	T vs C	T=0.2738	4	0.929 (0.603-1.432)	-0.34	7.32E-01	Random effects (77.3)	0	0	508
45	<i>OXTR</i>	rs2268495	A vs G	A=0.2406	4	1.059 (0.763 - 1.468)	0.34	7.33E-01	Fixed effect (60.4)	282	446	458
46	<i>SHANK3</i>	rs9616915	C vs T	C=0.3433	3	0.974 (0.834 - 1.138)	-0.33	7.44E-01	Fixed effect (60.1)	340	863	308
47	<i>HTR2A</i>	rs6314	T vs G	A=0.0747	4	0.949 (0.691-1.304)	-0.32	7.47E-01	Fixed effect (18.3)	103	214	370
48	<i>CNTNAP2</i>	rs2710102	T vs C	A=0.4113	3	0.989 (0.924-1.059)	-0.31	7.60E-01	Fixed effect (17.3)	322	524	2051
49	<i>OXTR</i>	rs237885	G vs T	G=0.4884	6	0.981 (0.868 - 1.109)	-0.3	7.62E-01	Fixed effect (0)	574	1201	458
50	<i>COMT</i>	rs4680	Met vs Val (A vs G)	A=0.3692	5	0.982 (0.851-1.134)	-0.24	8.08E-01	Fixed effect (49)	814	741	35
51	<i>MTHFR</i>	rs1801131	C vs A	G=0.2494	6	0.979 (0.824-1.164)	-0.24	8.11E-01	Random effects (56.3)	1854	6819	0
52	<i>OXTR</i>	rs1042778	G vs A	T=0.4109	4	1.02 (0.849-1.225)	0.21	8.33E-01	Fixed effect (0)	282	440	458
53	<i>GRIK2</i>	rs2227283	A vs G	A=0.3275	4	0.967 (0.686-1.363)	-0.19	8.51E-01	Random effects (65.65)	0	0	508
54	<i>EN2</i>	rs3735653	T vs C	T=0.4097	4	1.007 (0.870-1.165)	0.09	9.28E-01	Fixed effect (0)	174	349	499
55	<i>NF1</i>	GxAlu	8 vs non-8	NA	4	0.982 (0.602 - 1.601)	-0.07	9.41E-01	Random effects (79.2)	262	312	0
56	<i>SLC6A4</i>	5-HTTLPR	short vs long	NA	17	0.994 (0.847-1.167)	-0.07	9.42E-01	Random effects (63.8)	0	0	2039
57	<i>HTR2A</i>	rs6313	T vs C	A=0.4413	3	1.007 (0.812-1.249)	0.07	9.47E-01	Fixed effect (0)	0	0	303
58	<i>EN2</i>	rs1861973	T vs C	T=0.2410	6	1.004 (0.775-1.300)	0.03	9.77E-01	Random effects (80.8)	669	1704	681

Rows highlighted in bold show variants with *P* values below 0.01

Table 2 Summary of subgroup analyses

S.No	Gene	Variant	Allele	Data sets	Subgroup	Mean OR (95% CI)	Z-Value	P-Value	Model
1	ASMT	rs4446909	G vs A	3	Caucasian	1.135 (0.886 - 1.454)	1	3.16E-01	Fixed
2	ASMT	rs5989681	G vs C	3	Caucasian	1.065 (0.841 - 1.349)	0.52	6.03E-01	Fixed
3	COMT	rs4680	A vs G	4	TDT	0.973 (0.840 - 1.128)	-0.36	7.17E-01	Fixed
4	EN2	rs1861973	T vs C	4	TDT	0.86 (0.791 - 0.954)	-2.94	3.00E-03	Fixed
5	EN2	rs1861973	T vs C	3	Caucasian	0.880 (0.801 - 0.969)	-2.26	9.00E-03	Fixed
6	EN2	rs1861972	A vs G	4	Case-control	1.186 (0.876 - 1.605)	1.11	2.69E-01	Random
7	EN2	rs1861972	A vs G	4	TDT	1.126 (1.025 - 1.238)	2.47	1.30E-02	Fixed
8	EN2	rs1861972	A vs G	4	Caucasian	1.118 (0.807 - 1.549)	1.32	1.86E-01	Fixed
9	HOXA1	rs10951154	A vs G	6	Case-control	0.876 (0.675 - 1.137)	-0.99	3.21E-01	Random
10	HOXA1	rs10951154	A vs G	6	Caucasian	0.887 (0.661 - 1.190)	-0.8	4.23E-01	Random
11	HOXA1	rs10951154	A vs G	7	TDT	0.963 (0.784 - 1.159)	-0.48	6.32E-01	Random
12	HTR2A	rs6311	A vs G	4	TDT	0.893 (0.602 - 1.325)	-0.56	5.73E-01	Random
13	HTR2A	rs6311	A vs G	3	Caucasian	0.929 (0.542 - 1.594)	-0.27	7.90E-01	Random
14	MACROD2	rs4141463	T vs C	5	Case-control	1.033 (0.944 - 1.131)	0.71	4.78E-01	Random
15	MET	rs1858830	G vs C	7	Case-control	0.889 (0.749 - 1.056)	-1.34	1.80E-01	Random
16	MET	rs1858830	G vs C	3	Italian	0.924 (0.592 - 1.444)	-0.35	7.29E-01	Random
17	MTHFR	rs1801133	T vs C	4	Caucasian	1.398 (1.249 - 1.565)	5.82	6.60E-02	Random
18	MTHFR	rs1801131	C vs A	3	Caucasian	0.904 (0.782 - 1.044)	-1.37	1.71E-01	Fixed
19	OXTR	rs237885	G vs T	3	Case-control	0.950 (0.817 - 1.106)	-0.65	5.11E-01	Fixed
20	OXTR	rs2268490	T vs C	3	TDT	1.281 (0.953 - 1.721)	1.64	1.01E-01	Fixed
21	OXTR	rs2254298	A vs G	4	Caucasian	0.664 (0.357 - 1.235)	-1.29	1.96E-01	Fixed
22	OXTR	rs2268490	T vs C	4	Caucasian	1.114 (0.882 - 1.409)	0.91	3.66E-01	Fixed
23	OXTR	rs237885	G vs T	4	Caucasian	1.039 (0.885 - 1.220)	0.47	6.40E-01	Fixed
24	OXTR	rs237885	G vs T	3	TDT	1.043 (0.846 - 1.285)	0.39	6.96E-01	Fixed
25	OXTR	rs2254298	A vs G	4	Case-control	1.034 (0.693 - 1.542)	0.16	8.69E-01	Fixed
26	RELN	rs362691	C vs G	6	Case-control	0.857 (0.783 - 0.939)	-3.32	1.00E-03	Fixed
27	RELN	rs736707	T vs C	8	Case-control	1.187 (0.953 - 1.479)	1.53	1.27E-01	Random
28	RELN	rs736707	T vs C	3	Caucasian	1.307 (0.843 - 2.025)	1.2	2.32E-01	Random
29	SLC25A12	rs2292813	C vs T	4	TDT	1.419 (1.158- 1.740)	3.52	7.33E-04	Fixed
30	SLC25A12	rs2056202	T vs C	5	TDT	1.275 (1.097 - 1.482)	3.17	2.00E-03	Fixed
31	SLC25A12	rs2056202	T vs C	3	Case-control	1.105 (0.862 - 1.416)	0.79	4.31E-01	Fixed
32	SLC25A12	rs2056202	T vs C	4	Caucasian	1.087 (0.873 - 1.355)	0.75	4.55E-01	Fixed
33	SLC6A4	5-HTTLPR	short vs long	5	Caucasian	0.960 (0.650 - 1.418)	-0.2	8.38E-01	Fixed
34	SLC6A4	STin2 VNTR	12 vs 9/10	4	Caucasian	1.492 (1.068 - 2.083)	2.34	1.90E-02	Fixed

Rows highlighted in bold show variants with *P* values below 0.05

(see Table 2). rs1861972 is in high LD with rs1861973 ($r^2 = 1$), and the two variants are separated by 152 base pairs. In this study, we used the random effects model to meta-analyse rs1861973 and it was not significant. Stratifying by both study methodology and ethnicity reduced the heterogeneity considerably, allowing us to use a fixed effect model. For rs1861973, both a Caucasian-only and a TDT-only subset were significant (see Table 2) but this variant was not significant in the larger Caucasian-only PGC

cohort. Additional research in a larger, well-powered sample is required to confirm the significance of the two variants.

Discussion

This is the first study to comprehensively investigate candidate gene association studies of common variants in ASC. Using two databases, we identified 552 genes that are reported to be implicated in ASC through genetic

association studies. We scanned the literature for these 552 genes and, using a strict inclusion criteria, we identified 27 genes that had sufficient data to perform a meta-analysis. Eight variants across seven genes were significant for combined effect sizes with P values below 0.01. Data for 11 variants was present in the PGC GWAS dataset. None of the 11 variants were significant in the PGC dataset though the majority of the variants were concordant for effect direction in both the datasets.

Effect sizes for most common variants are modest for ASC, and these results are consistent with this observation. However, there was no clear correlation between effect sizes in our dataset and the PGC dataset. Effect sizes were smaller in the PGC dataset. While most of the effects lay between 0.8 and 1.2, which is expected from GWAS data, for some variants, the effect was larger. Our most significant variant (rs167771) had data only from three studies and had a relatively high OR of 1.82 to 1.40–2.38. The small sample size for this variant inflated the OR making it significant. The effect direction was discordant for the variant in the PGC dataset, and it was not significant in this dataset.

While the sample sizes for most variants were competitive for candidate gene association studies (above 500 total cases and 500 total controls), these are not sufficient to accurately calculate effect sizes. Additionally, the different study methodologies and ethnicities contributed to heterogeneity in the sample which potentially confounded the analyses. It is clear from this study that significant heterogeneity exists for a large fraction of the variants tested. In fact, heterogeneity is significantly and positively correlated with the number of independent datasets included per variant in the analyses, indicating that the current study may not have uncovered all the heterogeneity. We were able to remove some of the heterogeneity after stratifying for ethnicity and study methodology, but heterogeneity influenced the results for some for the variants even after this. This indicates that other additional factors contribute to variance in the effect. One potential source of heterogeneity is finer population stratification. Fine-scale population stratification cannot be addressed in candidate gene association studies as these test only a few variants. Further, HWE which is used to check for population admixture among other issues is performed individually for each variant in these studies thereby failing to utilize multi-marker information to correct for population stratification. We were unable to stratify based on sex or clinical ascertainment two factors known to contribute to heterogeneity in ASC. It is unclear how clinical heterogeneity maps onto genetic heterogeneity in ASC. Existing genetic studies that stratify based on IQ or other clinical phenotype and subphenotypes have had limited success [24, 25]. The inability to completely identify sources of

heterogeneity forced us to choose between two models (fixed effect vs. random effects), when most variants are likely to have varying levels of heterogeneity. This is a significant concern for meta-analyses using candidate gene association studies. Even if sample sizes reach competitive levels, there are no techniques currently available that can accurately account for potential confounders such as ethnicity and study methodology. Both these issues can be satisfactorily addressed in GWAS.

Another cause for concern is the small number of genes with enough data to meta-analyse. Of 552 genes, we had data for only 27 of these, less than 5 %. None of the 27 genes analysed were ASC risk genes as predicted by DAWN [26]. Further, with the exception of *RELN* [27] and *SHANK3* [28], none of these genes have sufficient evidence to categorize them as risk genes using sequencing or copy number variation studies [27–31]. A few genes in the list of 552 genes but absent from the final list of 27 genes are predicted to be ASC risk genes. This includes *GABRB3*, *GRIN2B* and *SCN2A*. However, there was not enough evidence to evaluate the role of common variants in ASC for these genes through the current meta-analysis.

The majority of the studies analysed were of Caucasian ethnicity. We were able to stratify for a Caucasian ethnicity for some of the variants, but were not able to stratify for other ethnicities due to power considerations. It is also noteworthy that the PGC autism dataset used a Caucasian sample for analyses, and to our knowledge, there is no well-powered GWAS that investigates the role of common variants in autism in other ethnicities. Since the minor allele frequencies of the alleles tested and the variants tagged by these allele can vary depending on ethnicity, this makes it difficult to compare the results of the non-stratified meta-analyses with the PGC autism dataset. Replicating the top variants in well-powered samples from different ethnicities will help understand the ethnicity-specific risk for each variant.

The candidate gene association studies typically have small samples, which overestimate effect sizes. The lack of replication do not indicate that these loci do not contribute to the aetiology of ASC, but, rather, that there is insufficient evidence to implicate it in ASC. ASC is highly polygenic, and more than 49 % of its heritability can be attributed to common variants [3]. As effect size for each individual common variant are likely to be very modest and not likely to exceed an OR of 1.3, this indicates that there are several common variants that contribute to the condition. Disentangling this would require very large sample sizes, much larger than those in the current PGC autism GWAS. It is evident, from the current study, that candidate gene association studies in ASC have been underpowered to reliably detect causative variants with precision.

Conclusions

While recent studies [2, 3] have identified that common variants, *en masse*, contribute to a significant fraction of ASC, there have not been any sufficiently powered studies to date to identify important common variants. We attempted to address this issue using a meta-analysis of candidate gene association studies. Though this is the first comprehensive study of candidate gene association studies in ASC, it failed to identify causative variants—11 of 15 variants with *P* values <0.05 were not significant in a larger sample from the PGC. Data was unavailable for the remaining five variants in the PGC dataset. We discuss the potential issues with such an approach and underline the need for much larger sample sizes to accurately identify common variants that contribute to ASC.

Additional file

Additional file 1: Table S1. Studies included and study characteristics and **Table S2.** Studies Excluded; **Figures S1–S20.** Forest plots of significant variants from global and subgroup analysis, and sensitivity plots; details of data from our lab, details of previous meta-analyses, and references. (PDF 707 kb)

Abbreviations

ASC: autism spectrum conditions; CC: case-control; CI: confidence intervals; FBAT: family-based association test; GWAS: genome-wide association study; HWE: Hardy-Weinberg Equilibrium; LOR: log odds ratio; MAF: minor allele frequency; OR: odds ratio; PGC: Psychiatric Genomics Consortium; SE: standard error.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

VW, SBC and BC co-designed the study. VC, VW and PS analysed the data. SBC obtained the funding for the study. All authors wrote and edited the manuscript. All authors read and approved the manuscript.

Acknowledgements

We are grateful to Dr. Anitha Ayyappan Pillai, Prof. Elisabetta Trabetti and Dr. Wouter Staal for data-sharing. We thank Florina Uzefovsky for her critical comments and advice. This study was funded by grants from Target Autism Genome, the Autism Research Trust, Wellcome Trust Sanger Centre, and the Medical Research Council UK. VW is funded by the Nehru Trust for Cambridge University, St. John's College, and Cambridge Commonwealth Trusts. This study was submitted for the partial fulfilment of an MSc degree for VJC from Imperial College London, and a PhD degree for VW from the University of Cambridge.

Author details

¹Autism Research Centre, Department of Psychiatry, University of Cambridge, Douglas House, 18B Trumpington Road, Cambridge CB2 8AH, UK. ²Centre for Integrative Neuroscience and Neurodynamics, School of Psychology and Clinical Language Sciences, University of Reading, Reading, UK. ³CLASS Clinic, Cambridgeshire and Peterborough NHS Foundation Trust (CPFT), Cambridgeshire, UK.

Received: 23 February 2015 Accepted: 17 August 2015

Published online: 28 August 2015

References

1. American Psychiatric Association. Diagnostic and Statistical Manual of Mental Disorders: DSM-IV. 4th ed. Washington, DC: American Psychiatric Association; 1994.
2. Sandin S, Lichtenstein P, Kuja-Halkola R, Larsson H, Hultman CM, Reichenberg A. The familial risk of autism. *JAMA*. 2014;311:1770–7.
3. Gaugler T, Klei L, Sanders SJ, Bodea CA, Goldberg AP, Lee AB, et al. Most genetic risk for autism resides with common variation. *Nat Genet*. 2014;46:881–5.
4. Klei L, Sanders SJ, Murtha MT, Hus V, Lowe JK, Willsey AJ, et al. Common genetic variants, acting additively, are a major source of risk for autism. *Mol Autism*. 2012;3:9.
5. Lai MC, Lombardo MV, Baron-Cohen S. Autism. *Lancet*. 2014;383:896–910.
6. O'Riordan BJ, State MW. Autism genetics: strategies, challenges, and opportunities. *Autism Res*. 2008;1:4–17.
7. Borenstein M, Hedges LV, Higgins J, Rothstein H. Introduction to meta-analysis. Chichester: Wiley; 2009.
8. Badner JA, Gershon ES. Meta-analysis of whole-genome linkage scans of bipolar disorder and schizophrenia. *Mol Psychiatry*. 2002;7:405–11.
9. Munafò MR, Clark TG, Moore LR, Payne E, Walton R, Flint J. Genetic polymorphisms and personality in healthy adults: a systematic review and meta-analysis. *Mol Psychiatry*. 2003;8:471–84.
10. Taylor S. Molecular genetics of obsessive-compulsive disorder: a comprehensive meta-analysis of genetic association studies. *Mol Psychiatry*. 2013;18:799–805.
11. Chakrabarti B, Dudbridge F, Kent L, Wheelwright S, Hill-Cawthorne G, Allison C, et al. Genes related to sex steroids, neural growth, and social-emotional behavior are associated with autistic traits, empathy, and Asperger syndrome. *Autism Res*. 2009;2:157–77.
12. Warrier V, Baron-Cohen S, Chakrabarti B. Genetic variation in GABRB3 is associated with Asperger syndrome and multiple endophenotypes relevant to autism. *Mol Autism*. 2013;4:48.
13. Borenstein M, Hedges LV, Higgins J, Rothstein H. Comprehensive Meta-Analysis, 2.2050 edn. Englewood, NJ: Biostat; 2009.
14. Kazeem GR, Farrall M. Integrating case-control and TDT studies. *Ann Hum Genet*. 2005;69:329–35.
15. Cross Disorder Group of the Psychiatric Genomics Consortium. Identification of risk loci with shared effects on five major psychiatric disorders: a genome-wide analysis. *Lancet*. 2013;381:1371–9.
16. LoParo D, Waldman ID. The oxytocin receptor gene (OXTR) is associated with autism spectrum disorder: a meta-analysis. *Mol Psychiatry*. 2014 [Epub ahead of print].
17. Wang Z, Hong Y, Zou L, Zhong R, Zhu B, Shen N, et al. Reelin gene variants and risk of autism spectrum disorders: an integrated meta-analysis. *Am J Med Genet B Neuropsychiatr Genet*. 2014;165B:192–200.
18. Huang CH, Santangelo SL. Autism and serotonin transporter gene polymorphisms: a systematic review and meta-analysis. *Am J Med Genet B Neuropsychiatr Genet*. 2008;147B:903–13.
19. Song RR, Zou L, Zhong R, Zheng XW, Zhu BB, Chen W, et al. An integrated meta-analysis of two variants in HOXA1/HOXB1 and their effect on the risk of autism spectrum disorders. *PLoS ONE*. 2011;6:e25603.
20. Pu D, Shen Y, Wu J. Association between MTHFR gene polymorphisms and the risk of autism spectrum disorders: a meta-analysis. *Autism Res*. 2013;6:384–92.
21. Park J, Ro M, Pyun JA, Nam M, Bang HJ, Yang JW, et al. MTHFR 1298A>C is a risk factor for autism spectrum disorder in the Korean population. *Psychiatry Res*. 2014;215:258–9.
22. Schmidt RJ, Hansen RL, Hartiala J, Allayee H, Schmidt LC, Tancredi DJ, et al. Prenatal vitamins, one-carbon metabolism gene variants, and risk for autism. *Epidemiology*. 2011;22:476–85.
23. Ramoz N, Reichert JG, Smith CJ, Silverman JM, Bessalova IN, Davis KL, et al. Linkage and association of the mitochondrial aspartate/glutamate carrier SLC25A12 gene with autism. *Am J Psychiatry*. 2004;161:662–9.
24. Chaste P, Klei L, Sanders SJ, Hus V, Murtha MT, Lowe JK, et al. A genome-wide association study of autism using the Simons Simplex Collection: does reducing phenotypic heterogeneity in autism increase genetic homogeneity? *Biol Psychiatry*. 2015;77:775–84.
25. Liu XQ, Paterson AD, Szatmari P, Autism Genome Project Consortium. Genome-wide linkage analyses of quantitative and categorical autism subphenotypes. *Biol Psychiatry*. 2008;64:561–70.
26. Liu L, Lei J, Sanders SJ, Willsey AJ, Kou Y, Cicek AE, et al. DAWN: a framework to identify autism genes and subnetworks using gene expression and genetics. *Mol Autism*. 2014;5:22.

27. De Rubeis S, He X, Goldberg AP, Poultney CS, Samocha K, Cicek AE, et al. Synaptic, transcriptional and chromatin genes disrupted in autism. *Nature*. 2014;515:209–15.
28. Leblond CS, Nava C, Polge A, Gauthier J, Huguet G, Lumbroso S, et al. Meta-analysis of SHANK Mutations in Autism Spectrum Disorders: a gradient of severity in cognitive impairments. *PLoS Genet*. 2014;10:e1004580.
29. Betancur C. Etiological heterogeneity in autism spectrum disorders: more than 100 genetic and genomic disorders and still counting. *Brain Res*. 2011;1380:42–77.
30. Pinto D, Delaby E, Merico D, Barbosa M, Merikangas A, Klei L, et al. Convergence of genes and cellular pathways dysregulated in autism spectrum disorders. *Am J Hum Genet*. 2014;94:677–94.
31. Iossifov I, O’Roak BJ, Sanders SJ, Ronemus M, Krumm N, Levy D, et al. The contribution of de novo coding mutations to autism spectrum disorder. *Nature*. 2014;515:216–21.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

