

Simulating human and environmental exposure from hand-held knapsack pesticide application: Be-WetSpa-Pest, an integrative, spatially explicit modeling approach

Article

Accepted Version

Binder, C. R., Garcia-Santos, G., Andreoli, R., Diaz, J., Feola, G., Wittensoeldner, M. and Yang, J. (2016) Simulating human and environmental exposure from hand-held knapsack pesticide application: Be-WetSpa-Pest, an integrative, spatially explicit modeling approach. *Journal of Agricultural and Food Chemistry*, 64 (20). pp. 3999-4008. ISSN 0021-8561 doi: 10.1021/acs.jafc.5b05304 Available at <https://centaur.reading.ac.uk/53664/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

To link to this article DOI: <http://dx.doi.org/10.1021/acs.jafc.5b05304>

Publisher: American Chemical Society

including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online

Simulating human and environmental exposure from hand-held knapsack pesticide application: Be-WetSpa-Pest, an integrative, spatially explicit modeling approach

Claudia R. Binder, Glenda García-Santos, Romano Andreoli, Jaime Diaz, Giuseppe Feola, Moritz Wittensöldner, and Jing Yang

J. Agric. Food Chem., **Just Accepted Manuscript** • DOI: 10.1021/acs.jafc.5b05304 • Publication Date (Web): 30 Jan 2016

Downloaded from <http://pubs.acs.org> on February 1, 2016

Just Accepted

"Just Accepted" manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides "Just Accepted" as a free service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. "Just Accepted" manuscripts appear in full in PDF format accompanied by an HTML abstract. "Just Accepted" manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are accessible to all readers and citable by the Digital Object Identifier (DOI®). "Just Accepted" is an optional service offered to authors. Therefore, the "Just Accepted" Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the "Just Accepted" Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these "Just Accepted" manuscripts.



**Simulating human and environmental exposure from hand-held knapsack pesticide application:
Be-WetSpa-Pest, an integrative, spatially explicit modeling approach**

Claudia R. Binder^{1,2,3*}, Glenda García-Santos^{4*}, Romano Andreoli⁵, Jaime Diaz⁶, Giuseppe Feola⁷,
Moritz Wittensoeldner⁸, Jing Yang⁹

¹ Department of Geography, University of Zurich, Switzerland; ² Department of Geography, LMU-
University of Munich, Germany; ³ Institute for Environmental Engineering, ENAC, Ecole Polytechnique
Fédérale de Lausanne, Switzerland ⁴ Department of Geography, Alpen-Adria-University Klagenfurt,
Austria; ⁵ Agroscope, Switzerland; ⁶ Department of Sanitary and Environmental Engineering,
Universidad de Boyacá, Colombia; ⁷ Department of Geography and Environmental Science, University
of Reading, United Kingdom; ⁸ Brugg 3, Trogen Switzerland; ⁹ National Institute of Water and
Atmospheric Research, New Zealand

* Both authors contributed equally to the manuscript

Abstract

This paper presents an integrative and spatially explicit modeling approach for analyzing human and
environmental exposure from pesticide application of smallholders in the potato producing Andean
region in Colombia. The modeling approach fulfills the following criteria: (i) it includes environmental
and human compartments; (ii) it contains a behavioral decision-making model for estimating the effect
of policies on pesticide flows to humans and the environment; (iii) it is spatially explicit; and (iv) it is
modular and easily expandable to include additional modules, crops or technologies. The model was
calibrated and validated for the Vereda La Hoya and was used to explore the effect of different policy
measures in the region. The model has moderate data requirements and can be adapted relatively easy to
other regions in developing countries with similar conditions.

Keywords: integrative model, pesticide, exposure, environment, behavior, hydrology, developing countries

Introduction

World-wide pesticide use in crop production has constantly been increasing since the 1950s in total amount and in quantities per unit area [1,2]. Whereas in developed countries, mostly herbicides are applied and the environmental impacts, e.g., fish toxicity, soil damage, are of major concern [3-6] in developing countries, the bulk of pesticides used are insecticides and fungicides and human mortality and morbidity due to exposure to pesticides has become a concern as it significantly affects the livelihood of small farmers [7,8].

Several tools have been developed to assess and reduce the impact of pesticides on the environment and on human health [9]. Of special interest in our case is the development and use of models. A first string of research develops environmental pesticide fate models. They analyze the distribution of pesticides within the environment, including plant, soil, water and in some cases through air (drift) (PEARL [10], CESMOS, BASINS (HSPF) ,[11]SWAT [12], Wet_hydro [13], etc.). A particular group of models studies explicitly the contamination of water bodies. Models in this category are for example DRIPS, EXAMS, PIRANHA a/b/c (see [14] for a review). Other models, such as empirical curves, focus on pesticide airborne or drift deposition and were developed in the temperate region for terrestrial mechanized boom sprayers [15-19]. Some of these models perform optimally after calibration for the case of hand-held knapsack sprayers, mostly used in developing countries in the tropics, recently demonstrated by García-Santos et al. [20]. More sophisticated models focusing on pesticide exchange with the atmosphere after applied to soils and crops are dynamic and physically based [21]. The most advanced model within pesticide emission models is the PestLCI model [22,23] and was developed for use in agricultural life cycle assessment following mechanized spray application at local scale [24]. As

these models have all been developed for special needs and conditions of use, they differ in their spatial and temporal scales, time step, spatially explicitness, processes modeled, data requirements, complexity and output (see [14] for a review). Furthermore, most of these models have large data requirements and have never been applied in developing countries [25] and none of these models includes potential human exposure of the applicators as an output.

Within a second string of research, tools and models have been developed to assess human exposure to pesticide use. The tools range from qualitative assessment of human exposure, focusing mostly on dermal exposure (e.g., DERM [26]; EASE [27]; PHED [28]; COSHH [29]; DREAM [30]; RISKOFDERM [31]; STOFENMANAGER [32]; see [33,34] for a review), to sophisticated quantitative simulation models of pesticide emission in the air (Plume Model “Gaussian Plume” [35]; Gaussian Diffusion Model (GDM) [36]; Model for Risk assessment of pesticide drift damage [37]; One-Box Model [38]). These models focus specifically on human exposure and do not include environmental effects. The only tools in which both the environment and human health issues are included are empirical studies [39] and indicator based assessments (see [40] for a review). However, these assessment methods are usually neither dynamic nor spatially explicit e.g. accumulation issues, feedbacks and self-organization processes as part of the system dynamics are neglected. Furthermore, another disadvantage of empirical point-based static approaches is that the evaluation of probability is usually insufficiently considered [41] and key parameters might be not considered since they were developed in different contexts [42].

A third string of models combines agricultural production models (including pesticide use and to some extent environmental fate models) with economic models; an area where there has been significant progress in the last years (see [43] for a review; [44,45]). The integration of models ranges from linear programming models to spatially explicit multiple scales and multiple goal models [46,47]. A few

models analyze the effect of environmental degradation, e.g. erosion, water contamination on farmers' income and on the economic system at a regional level (e.g., ECECMOD [48]; and SAM [49]). The trade-off model [50] is the only integrative model that has been applied to pesticide management of potatoes in the Andes Region. It combines bio-physical models with econometric-process simulation models and provides an integrated analysis of tradeoffs between economic and environmental indicators. However, it has been found that farmers decision-making is often affected by parameters other than economic ones such as norms and traditions [51,52] and thus behavioral models are required which, on the one hand, simulate farmers' behavior and estimate the impacts of policies on pesticide use and, on the other hand, can be linked to spatially explicit pesticide models estimating the impact of behavioral change on human and environmental exposure. Finally, most models have high data requirements that cannot be met in developing countries and thus less data demanding models are required [40].

This study contributes to the development of integrated models in the area of pesticide management. The modeling approach, Be-WetSpa-Pest, (i) integrates a hydrological model with an extended pesticide emission model considering environmental and human compartments; (ii) is coupled to a behavioral model, for estimating the effect of different policies on farmers pest control behavior, affecting pesticide distribution in the environment and onto the applicator; and (iii) has already been applied to Vereda La Hoya, Colombia, a region with low data availability.

We combine a farmer behavioural model [53], with a spatially explicit hydrology model, WetSpa [54], and a pesticide emission model, PestLCI [22]. We assess the potential human and environmental fate of pesticides due to adoption of different policies in low mechanized cultivation of potato in the Colombian Andes (a tropical region).

Methods

Be-WetSpa-Pest¹: Model structure. Figure 1 presents the basic model structure composed of input data, core model processing and the output data. The strength of the Be-WetSpa-Pest is its modular structure, preferred over a fully integrated model approach as it facilitates the use, inclusion and adaptation of disciplinary models in the shape of modules, i.e. hydrological, fate and behavioral model, to different study areas (see also [50]), additional crops, pesticides and technologies.

<Figure 1>

Input data. To run the model four types of data are needed (Fig. 1 and 2 in Database box; Table 1 in 01_Supporting information): “global parameters” as parameters required for the hydrological WetSpa model and pesticide emission PestLCI model, “climate data”, i.e., representative meteorological data of the catchment, “spatial metadata”, i.e., spatially explicit land use, catchment, weather station and appropriation data, “pesticide data”, i.e., physical and chemical properties of applied pesticides, and “external socio-economic factors”, i.e., policies and crop rotation. The input data are stored in form of different GIS layers (Fig.2 File system box; Table 3 in 01_Supporting information). Ownership information was entered in GIS based on the latest Cadastral map [55]. Physical and chemical properties of applied pesticides within “pesticide data” were obtained from the PestLCI database and own empirical field experiments in similar soils [56,57].

<Figure 2>

Model processing. The core model, so-called Be-WetSpa-Pest model, is composed of a behavioral model and the WetSpa-Pest model. The “Behavioral model” predicts pesticide type (most prominent three fungicides and insecticides for the case study), amount applied, and application frequency per farmer and plot in the study area [53]: this information is used in the pesticide emission’s model as total pesticide applied (P_A , in equation 1).

¹Beta version available upon request.

113 A set of influencing factors contributes in determining the probability of a farmer to adopt a fungicide
114 and insecticide application type respectively (see section 3 of 02_supporting information). Concerning
115 fungicides, the influencing factors are training delivered by pesticide producing companies, the
116 proportion of household income coming from agriculture, the sense of compliance with the prescriptive
117 social norm (i.e. other farmers recommendations), the plot area, and being member of a farmer
118 cooperative. Concerning insecticides, the influencing factors are farmer's educational level, the presence
119 of significant sources of non-agricultural income for the household, training delivered by pesticide
120 producing companies, and the plot area ([53] and 02_supporting information). The default values of
121 these factors (i.e. those used in the baseline scenario) in WetSpa-Pest correspond to those observed in
122 Vereda La Hoya [52,53].

123 The model allows for modifying the value of selected factors for each farmer, thus simulating different
124 types of interventions on pesticide use in the area. Interventions that WetSpa-Pest allows to be simulated
125 are: i) intensification of training delivered by pesticide producing companies; ii) organization of farmers
126 in a cooperative; iii) modification of household non-agricultural income sources; iv) increase/decrease of
127 farmer sense of compliance with the prescriptive social norm. It has to be noted that the policies
128 intervene on those farmers who in the baseline scenario did not already participate in a training program
129 or who were not intervened. In the behavioral model, a farmer can adopt one among three different
130 fungicide and insecticide application types that correspond to decreasing input effectiveness levels [53].

131 The application types differ in terms of i) intensity of application (i.e. quantity of active ingredients per
132 unit of surface), ii) number of applications per agricultural cycle, and iii) class of active ingredients
133 applied, in particular with respect to the share of carbamates and pyrethroids over the total quantity of
134 fungicide and insecticide ([53] and Table 5 in 01_Supporting information). Furthermore there is the
135 option to use the option "crop rotation" based on [58,59].

136 The behavioural model is coupled to the WetSpa-Pest model and provides the interface for entering and
137 assessing policy recommendations. Application frequencies (i.e. average number of pesticide application
138 per phase of agricultural cycle) is also associated to the adoption of a fungicide and of an insecticide
139 application type, respectively. This information is used to determine application dates, which can be
140 distributed either randomly (using the `math.random` algorithm in which the values are chosen
141 pseudorandomly with approximately uniform distribution from the range [0.0 - 1.0]) or evenly within
142 each phase of the agricultural cycle excluding Sundays and days with rain (i.e. conditions by which
143 farmers from the study area do not apply pesticides). In addition, the first application date in a cycle
144 must be on a Tuesday (on even applications), and the application time must be between 06:00 and 17:00,
145 i.e. the time in which farmers usually work in the fields (see section 4.2 of 02_Supporting information).
146 The empirical data on pesticide use per farmer and plot proceed from the survey of 25% of the registered
147 farmers in the catchment area [52]. Therefore, an algorithm was developed to extrapolate the data from
148 these known data plots to all other agricultural plots in the study area. For details on the extrapolation
149 algorithm see section 5 of 02_Supporting information.

150 The second part of Be-WetSpa-Pest model is the WetSpa-Pest model. WetSpa-Pest is a fully distributed,
151 spatially explicit hydrology and emission pesticide model based on the WetSpa model by Liu and Smedt
152 [54] and the here modified PestLCI model [22], respectively. The model WetSpa is a GIS-based
153 distributed hydrological model for flood prediction and water balance simulation on a catchment scale
154 (for more details on the water flows on a cell basis see [54]). It was developed by the Free University of
155 Brussels and can be downloaded for free (<http://www.vub.ac.be/WetSpa/>). The pesticide emission model
156 PestLCI was developed in Denmark by Birkved and Hauschild [22] (updated version in [23]) to provide
157 information for the estimation of pesticide mass in the environment (air, surface water and groundwater)
158 outside of the sprayed field after aircraft, boom spray (pull tractor) or soil injection application

159 techniques to be included in life cycle assessment. Its modular structure allows adaptation to conditions
160 for different regions and agricultural practices and techniques. A human exposure component was added
161 and drift fraction was modified for hand-held spray pest application (more details below).

162 Thus, Wetspa-Pest simulates simultaneously the dynamics and balances of water, energy and pesticide
163 on a grid (cell) basis at catchment scale. It uses as input data the type, amount, and frequency of
164 pesticides use per plot estimated based with the behavioural model [53]. As in WetSpa, WetSpa-Pest
165 simulates simultaneously water processes i.e. snow processes (freezing and melting), canopy
166 interception and potential evapotranspiration, infiltration, percolation, surface runoff generation,
167 interflow, groundwater flow and river routing, pesticide flows distribution into the compartments air,
168 water, soil and canopy, based on the modified PestLCI model and farmer's exposure fraction based on
169 own field experiments [60].

170 In our study, each cell (30 x 30 m) is a unit process (technosphere in life cycle assessment studies),
171 equivalent to an agricultural field where only one crop grows, which is cultivated by one farmer with
172 spatially uniform pesticide application. The cell is vertically divided into different environmental
173 compartments i.e., air (100 m vertical), canopy, soil (soil surface, soil matrix and groundwater) (1 m soil
174 depth), as in PestLCI 2.0 [23] and additionally includes a human compartment i.e., applicator. When a
175 pesticide leaves the unit process (cell), it is considered an emission. The model takes into account
176 emissions to air, surface water and groundwater compartment like in PestLCI and additionally emissions
177 to soils outside of the cell as pesticide soil deposition from drift, as harvest (leaf uptake) and as
178 applicator (human exposure).

179 The primary pesticide distribution processes are those taking place during pesticide spray application as
180 described by the equation (1):

$$181 \quad P_A = (P_V + P_D + P_H + P_L + P_S) P_A / 100 \quad (1)$$

182 where P_A is the total pesticide (active ingredient) applied (kg active ingredient/ha), P_V is the pesticide
183 fraction which stays in the air of the plot (%), P_D is the pesticide drifted by wind out of the sprayed plot
184 (%), P_H is the pesticide that reaches the applicator (human exposure) (%), P_L is the pesticide fraction
185 deposited on the leaves (canopy) (%) and P_S is the pesticide directly reaching the soil (%).

186 Total pesticide applied (P_A): The quantity (kg/ha) and type of active ingredients applied in a plot, and
187 the frequency of application are determined through the behavioral model (see above) developed by
188 Feola and Binder [52,53].

189 Pesticide staying in the air (P_V): The value for the pesticide staying in the air depends on the type of
190 pesticide and meteorological conditions of the area. It can be entered into the model as a specific fixed
191 value. In the case of Vereda La Hoya the value of 1 % was used given the measurements in the field
192 [59].

193 Pesticide drifted by wind (P_D): Hand-held spray is not a considered technique in the PestLCI or PestLCI
194 2.0. For the case of hand-held spray in potato production, it is found that drift is higher as compared to
195 conventional boom spray with tractor [61] and therefore the here modified PestLCI includes two
196 possibilities a fix drift fraction of 3.1% of the applied dose as derived from García-Santos et al. [62] or
197 calculation of drift for the first 20 m outside the plot in function of distance through the optimized
198 IMAG drift calculator (v 1.1) by Holterman and Zande [16] (in PestLCI v2) after García-Santos et al.
199 [61] (optimized parameters: $a = 29$, $b = -6.8$, $c = 18.35$, $d = -0.44$). Other equations in function of wind
200 speed may reflect a more realistic scenario in cases where spray is conducted under wind conditions
201 above 2 m s^{-1} , available in García-Santos et al. [61]. This could be added into the model structure.

202 Pesticide reaching the human (P_H) (not in PestLCI 2.0): P_H is the fraction of the pesticide reaching the
203 farmer's clothes and is calculated by using a fixed fraction of 1% of the total pesticide derived from
204 empirical measurements [60]. The type of protection equipment used determines the final exposure of

the farmers [60]. In the case we applied the model, Vereda La Hoya, farmers use gummy boots, long trousers and a T-shirt covered by a “ruana” or “poncho”.

Pesticide fraction deposited on the leaves (canopy) P_L : P_L in Eq. 3 depends, besides the amount of pesticide drifted away and the amount landing on the applicator, on the leaf area covered and thus on the growth stage of the plant,

$$P_L = (P_A - P_H - P_D - P_V) * v_i \quad (3)$$

where v_i is the interception fraction and represents the growth stage of the plant [63].

Pesticide reaching directly the soil (P_S): P_S is calculated as the balance from the total amount of pesticide applied and the pesticide reaching the other compartments,

$$P_S = P_A - P_V - P_D - P_H - P_L \quad (4)$$

After the plot is treated, the applied pesticide is redistributed in the environment and degraded. The secondary modelled processes by WetSpa-Pest after the pesticide application are infiltration into soil, percolation into the groundwater (WestSpa model), and pesticide outflow of the watershed through surface runoff (fraction of pesticide in the top soil liquid phase) and groundwater flow (fraction of pesticide in the soil matrix liquid phase) (modified PestLCI) (see equations in 03_Supporting information and coefficients in 01_Supporting information Table 2). Pesticide loss through macropore flow and tillage (considered in PestLCI 2.0) is not modelled but could be incorporated into the model.

Output data. The output data is composed first of hydrological times series, including infiltration, percolation, and evaporation, and surface runoff. The data is used to calibrate the WetSpa part of the model. Second, spatially explicit pesticide concentration data onto the applicator, the crop (as harvest), soil surface, soil matrix, surface runoff and groundwater is generated. Furthermore, for a specific plot, the same information can be obtained as time series (Fig. 2).

Case study. Be-WetSpa-Pest was implemented in a typical Andean mountainous region, the department (“Departamento”) of Boyacá, which contributes to ca. 26% of the national potato production and to 45% at regional level despite its low productivity, and its land tenants are mainly smallholders (95% of the workforce) [20, 70]. The same area of study has been focus of recent studies on human and environmental exposure from hand-held knapsack pesticide applications [20,39,40,44,45, 52,53,60,61,62,64,69]. The study area Vereda La Hoya has 840 ha and is located in the district La Hoya of the community Tunja (Colombia) at a height of about 2800 to 3200 m a.s.l. It lies in the cold climate thermal floor zone, with a total mean annual rainfall of 620 ± 20 mm. The area has prevailing south-east winds with average wind speed of 1.8 ± 1.39 m s⁻¹ and a maximum of 7.6 m s⁻¹ (data from this study). Meteorological variables were registered every 15 min for 3 years, from October 2008 until October 2011, at 3 m above ground, using a low cost automatic meteorological station, Davis Vantage Pro-2, installed within the watershed because no representative weather information was found in the national net (IDEAM).

The moisture regime of the soil is ustic and soil texture according to US-Soil taxonomy is sandy loam as described by García-Santos and Keller-Forrer [64] and [65]. Average pH of the soil is 5.03 ± 0.31 . Total organic carbon is 9.51 ± 3.93 % (Walkley Black method) and bulk density is 0.84 ± 0.1 gr cm⁻³ [64,66];

The seasonality of water discharge is caused mainly by variations in rainfall events in May and October, ranging from less than 10 l s⁻¹ in pre-event situations to above 60 l s⁻¹ during spring and autumn. Discharge regularly intermits during summer. Water flow at the outlet was measured using an ultrasonic Doppler sensor (Unidata STARFLOW) calibrated with a propeller. Measured water velocity was multiplied by the cross section to obtain water flow (l s⁻¹).

Orographic characterization of the catchment was obtained through a digital elevation model. A geographic information system (GIS) was used to store digitized land use representing the watershed characteristics with a grid of 100 x 100 m and to convert the information to raster [55].

Main crop production in Vereda La Hoya is potato. The catchment lies within the second highest potato producing region in Colombia, Boyacá, after Nariño [67]. In the region, mostly *solanum tuberosum spp* is produced. The potato cycle last about 6-7 months and the average yield is low with about 7 ton/ ha and cycle [53]. Farmers cultivate an average of 3 ha, which are subdivided into small plots, being to a large extent distant from each other within the catchment and most located on terrains, which are not appropriate for mechanization.

Pests are controlled through the application of insecticides and fungicides (see [53] for details) during the entire cycle using hand-held sprayers, lever-operated knapsack sprayer.

Model calibration and validation of the hydrological module. The hydrological module was calibrated following the principles of the guidelines for WetSpa calibration by Liu and Smedt [54]. First, a rough calibration was made separately for the WetSpa model using the stream flow data from the study area. Calibration data were from 4.9.2010 - 17.10.2010 and 28.10.2010 - 28.11.2010). These periods included a precipitation event with a return period of 25 years. The gap is due to missing discharge and meteorological data. Second the parameters: correction factor for evapotranspiration; surface runoff exponent; threshold rainfall intensity; interflow scaling factor, and baseflow recession coefficient were calibrated using the 3 months with highest rain intensity (September to November) during 2010 (see also [54]). Third, the hydro-meteorological data from 29.11.2010 to 19.12.2010 was used for validation of the model. To evaluate the goodness of fit of the modeled discharge during the calibration and validation period, we used five statistical criteria (Table 1): the model bias [54] which is the relative mean difference between predicted and observed stream flows (0 represents a perfect fit);

the Nash-Sutcliffe efficiency [68] cited in [54], which is used to evaluate how good the model simulates the stream flow (1 indicates a perfect fit); the RMSE (root mean square error); the Pearson coefficient of determination (R^2); and the standard error.

<Table 1>

Model validation of the pesticide module in soil. To validate the pesticide module of the WetSpa-Pest, we modeled the concentration in soil of an active ingredient used (i) widely and (ii) in high dosage in the study area. Chlorpyrifos corresponded to these characteristics and is used to protect the crop from three typical pests threatening the potato crop in the study area: the soil-dwelling larvae of the Andean weevil (*Premnotrypes vorax*), the late blight fungus (*Phytophthora infestans*) and the Guatemalan potato moth (*Tecia solanivora*) [53]. The crop cycle period extended from the 20th September until the 28th December 2009. The day and time of pesticide application, day of planting and day of harvest represent real conditions as reported by the farmer. Predicted concentrations were compared to previously measured concentrations in the same area [69]. The calculated concentrations in soil showed a high agreement with the measured values along the different stages of growth of potatoes with an r^2 of 0.82 (Figure 3). The modeled values lied to a large extent within the error margin of the measured concentrations.

<Figure 3>

Simulation scenarios. To show some of the results the model can generate, we used Be-WetSpa-Pest to simulate three scenarios, i.e. a “baseline scenario”, ‘training by companies’ and ‘cooperative’ respectively. The latter two were developed to exemplify how the model can be used to assess the effect of policies on pesticide use, and environmental and human exposure. We show the results for the use of the active ingredient Mancozeb, applied six times with different time spans in between the applications, from 20.07.2009 to 20.12.2009 for a specific field in the study area.

294 All scenarios simulate fungicide and insecticide application over 4 agricultural cycles of potato
295 production (see Table 4 and Table 5 in 01_Supporting information). The first and second cycle
296 correspond to a baseline scenario, in which self-reported pesticide application rates are used [53]. The
297 initial values of parameters in the behavioral model (see section above) also correspond to observed
298 values in the study area [53]. The third and fourth agricultural cycle served to run the “training by
299 companies” and “cooperative” separately.

300 **Results and discussion: application of the Be-WetSpa-Pest model**

301 **The baseline scenario.** Figure 4 shows the simulation results for a specific plot of the active ingredient
302 Mancozeb fungicide ($C_8H_{12}MnN_4S_8Zn$), an ethylenebisdithiocarbamate (CAS Number:8018-01-7).
303 On the top X-Axis and the right Y-Axis, precipitation on the plot is depicted, on the lower X-Axis and
304 the left Y-Axis, the concentration of Mancozeb is shown. The first application (20.8.2009) took place at
305 the start of the growth phase. The amount of Mancozeb intersected by the canopy is low, a large share of
306 the Mancozeb applied (about 83 %) end in the topsoil and some infiltration occurs when there is
307 precipitation. In the larger growth stage (20.10.2009), the plant intercepts a higher percentage of the
308 pesticide leading to a lower amount reaching the topsoil (up to a share of about 48% for plant and
309 topsoil). The amount infiltrating through the topsoil to the soil is rather below <4% and is favored is by
310 precipitation due to the wash off from canopy to topsoil and from topsoil to soil. The total amount of
311 Mancozeb accumulated in the soil is low and zero at the end of the cycle. On contrary, the concentration
312 in the topsoil decreases slowly and one could potentially encounter residues even when the next cycle
313 starts. This is not only due to the application itself, but also to the withering and degradation of the
314 vegetation stubbles after harvesting the potatoes.

315 <Figure 4>

Figure 5 shows the spatial distribution of the pesticide concentration in the watershed for Mancozeb at the end of the four simulated cycles (the same case as above). Red (Hotspots) are the plots with high Mancozeb concentrations (between 4-8 kg/ha), in the green plots no pesticides were applied at all. Hotspots of topsoil concentrations higher than 1 kg/ha can only be found in a few places, however some of them being close to a stream. The light green plots show the pesticide concentrations due to drift. On these plots no pesticides were applied but we could simulate low concentrations of Mancozeb. The contaminated plots are spatially interconnected, which is due to similar cultivation practices on neighboring plots and the drift of pesticides to neighboring plots.

<Figure 5>

Scenario analysis. In the policy scenario ‘training by companies’ we simulate the effect of a training program held by pesticide producing companies. We considered only the farmers who had no intervention in the baseline scenario. For these farmers, the introduction of the policy modifies the probability of adopting one of three fungicide and insecticide application types respectively. This potentially determines a variation (i.e. increase) of pesticide released into the environment. The results show a marginal increase in productivity, but a significant increase in fungicide and insecticide use ([53]). In the policy scenario ‘cooperative’ we simulate the effect of the participation of farmers in producer cooperatives. The results show that this scenario significantly improves the efficacy of insecticide use, that is, it reduces insecticide use rates [53].

Figure 6 shows the total amount of active ingredient of two fungicides and two insecticides applied in the total area during one cycle (140 days) for the baseline and the two scenarios mentioned above. Regarding the insecticides, Carbofuran and Permethrin, almost no differences can be found between the scenarios, implying that these policy measures will not be effective in reducing exposure to pesticides in the case studied. For the fungicides Mancozeb and Cymoxanil the training by pesticide producing

companies significantly increases the amount of pesticides applied, whereas joining a cooperative reduces the amount of fungicides applied by almost 90%. This suggests that fostering cooperatives and providing training and production support through the cooperatives might be a good way forward to reduce the amount of fungicides applied. Even though this variable had a strong estimated effect on fungicide use it was not significant for insecticides use and was therefore not included in the equation. That is, insecticide and fungicide use are determined by different sets of factors, and even when there is one factor that influences both pesticide and insecticide use (e.g. the proportion of household income coming from agriculture), the coefficient is not the same in both equations for the two behaviors (as estimated in the statistical models [53]). This indicates the necessity for a disaggregated behavioral analysis for pesticide use rather than an overall analysis looking at the amount of money spent on pesticides.

<Figure 6>

When looking at the spatial distribution of Mancozeb (Figures 7 a,b), the amount of hotspots (>1 kg/ha) increases in the case of the training scenario, whereas the areas, where pesticide amounts higher than 1kg/ha can be found in the topsoil is markedly reduced in the cooperative scenario. In particular, the concentration of Mancozeb in the areas close to the river is reduced. This result reflects the necessity of a spatially explicit analysis (see also [50]) to provide decision support for reducing surface water contamination by decreasing the pesticide concentration on plots close to surface water areas.

<Figure 7>

Model use recommendations and further model developments.

This paper presented Be-WetSpa-Pest, a simulation model that (i) integrates a hydrological model with an extended pesticide emission model considering environmental and human compartments; and (ii) is coupled to a behavioral model, for estimating the effect of different policies on farmers' pest control

362 behavior, affecting pesticide distribution in the environment and onto the applicator. The model was
363 calibrated and validated for the case of Vereda La Hoya in Colombia and provided valuable results on
364 the effect of policies on humans and environmental exposure to pesticides. We suggest the model to be
365 applicable for similar cases in developing countries, where there is low data availability. However, we
366 recommend, for the environmental part, to measure the key input variables for climate data like daily
367 precipitation and temperature from a nearby station, and to obtain a DEM of the area to be studied for
368 the spatial data.

369 It was shown that changes in farmers' behavior play a significant role for environmental and human
370 exposure and that policies affect fungicide and insecticide use in a different way. To apply Be-WetSpa-
371 Pest to other regions, thus, we consider that (i) the behavioral model should be validated. In a similar
372 cultural background as the case study, we expect that the influencing factors might be the same, even
373 though their impact (i.e. the estimated coefficient in the deterministic equation) might differ. In other
374 cultural contexts, it is likely that not only the influencing factors' estimated effects, but also the type of
375 influencing factors (e.g. social, economic, technical) may be different from the one validated for this
376 case study. Therefore, it is recommended that a behavioral study is carried out allowing for validating
377 the behavioral model, estimating the effect of the influencing factors, and measure the initial values of
378 quantity and type of pesticide used.

379 Furthermore, the model has some potential for expansion and further development:

380 a) *Adapting to other pesticide application techniques and human behavior*: The model can be easily
381 adapted to new application techniques. Thereby, the transfer-coefficients developed and applied for
382 other pesticide models can be used as input values for the coefficients to the environmental
383 compartments. For estimating the flow to the human compartment, however, experiments should be
384 performed to estimating the amount of pesticides ending on human body. Similarly, the behavioral

model will have to be adapted as the behavioral factors that were used in this case study may not explain pesticide application decisions in cases in which the application technique is not the knapsack sprayer. The theoretical framework used to inform the model in this case study [52] may be used to inform the study of pesticide application in different contexts, which will result in different behavioral equations, thus helping to adapt Be-Wetspa-Pest to other pesticide application techniques.

b) *Including pesticide uptake by plants through the root zone:* As a further development of the model a module on pesticide uptake into the crop through the root zone should be considered. This would allow to model a further channel of exposure to human health, namely through food consumption. Be-WetSpa-Pest models concentration of pesticide in the soil and therefore this can be used as input variable into a pesticide crop uptake model [69].

c) *Including different irrigation systems:* Some agricultural areas might have water reservoirs for agricultural purposes in addition to rain. The movement of the pesticides in soil might be affected in different ways depending on the irrigation techniques.

d) *Including second order degradation of the active ingredient:* One caveat of the model is that we considered for keeping the model easy manageable only the first order degradation of the active ingredient. We consider that given the high data uncertainties in developing countries, this is the right decision to take. If the model were adapted to regions with better data quality, it might be adequate to evaluate to which extent the inclusion of second order degradation of pesticides might make sense.

e) *Include horizontal redistribution in surface water:* In all compartments, except groundwater, the redistribution of pesticides, which is linked to water is vertically. This implies that horizontal

distribution from plot to plot is not considered (and might be also very marginal). Future models could consider and complement the vertical redistribution with that horizontal redistribution.

f) *Modeling farmers' behavior dynamically*: In Be-WetSpa-Pest, farmer decision making is modeled through a deterministic equation. This is partly due to the fact that it was not possible to quantify the influence of farmers' perception of e.g. pesticide infestation levels, or environmental quality on pesticide application decisions [53]. In fact, little evidence exist that farmers in Vereda La Hoya changed their pesticide application decisions because of the perception of environmental impact of pesticide use. However, in Be-WetSpa-Pest there is further scope for internalizing farmers' behaviour, i.e. to model not only the impact of pesticide application on the environment, but also how farmers respond to pesticide distribution and concentration in the environment should this be relevant for the study area to which the model is applied [53].

g) *Crop rotation*: There is an option in the model to implement crop rotation. This allows for accounting for different amounts of pesticide applied depending on the crops produced in the region.

h) *Modeling the effect of climate scenarios*: Be-WetSpa-pest presents no limitation to input meteorological data (rainfall, temperature, wind speed and direction) from downscaled climate scenarios. This application might be of relevance for the risk assessment analysis of climate impacts.

Acknowledgements. The early part of this work was carried out while the authors were in the Unit for Social and Industrial Ecology, Department of Geography, University of Zurich, Switzerland. This project was funded by the Swiss National Science Foundation (SNSF) through the project number 110807. The authors thank Dr. Ronnie Jursake for valuable comments, Lisa Ketzer and Sophia Lohmayer for formatting and for editing, the students from the University of Boyacá for their support in the field work.

Supporting information description.

Supporting information includes i) input variables and coefficients in WetSpa-Pest model, GIS layers derived from the DEM (Digital Elevation Model), simulated scenarios and fungicide application types (01 – Supporting information: Table 1, Table 2, Table 3, Table 4, Table 5), ii) detailed description of how pesticide application data and integration of farmer decision models into the WetSpa-Pest model are generated (02 - Supporting information: Generation pesticide application) and iii) detailed description and equations on the secondary distribution of pesticides in the modified PestLCI (03 - Supporting information: Redistribution and degradation processes of the pesticides).

References

1. Carvalho, F.P. Agriculture, pesticides, food security and food safety. *Environmental Science & Policy* **2006**, *9*, 685–692.
2. World Health Organization, W.H.O. *Public health impact of pesticides use in agriculture*. W.H.O.: Geneva, 1990.
3. Ecobichon, D.J. Pesticide use in developing countries. *Toxicology* **2001**, *160*, 27–33.
4. Younes, M.; Galal-Gorchev, H. Pesticides in drinking water — a case study. *Food and Chemical Toxicology* **2000**, *38*, 87–90.
5. Carlomagno, M.; Mathó, C.; Cantou, G.; Sanborn, J.R.; Last, J.A.; Hammock, B.D.; Roel, A.; González, D.; González-Sapienza, G. A clomazone immunoassay to study the environmental fate of the herbicide in rice (*oryza sativa*) agriculture. *Journal of agricultural and food chemistry* **2010**, *58*, 4367-4371.

- 452 6. Zhang, R.; Cui, J.; Zhu, H.M.; Yang, H. Effect of dissolved organic matters on napropamide
453 availability and ecotoxicity in rapeseed (*brassica napus*). *Journal of agricultural and food*
454 *chemistry* **2010**, *58*, 3232-3240.
- 455 7. Wilson, C.; Tisdell, C. Why farmers continue to use pesticides despite environmental, health and
456 sustainability costs. *Ecological Economics* **2001**, *39*, 449–462.
- 457 8. *International code of conduct on the distribution and use of pesticides: guidance on pest and*
458 *pesticide management policy development*. Food and Agriculture Organization of the United
459 Nations: Rome, 2010.
- 460 9. Rice, P.J.; Rice, P.J.; Arthur, E.L.; Barefoot, A.C. Advances in pesticide environmental fate and
461 exposure assessments. *Journal of agricultural and food chemistry* **2007**, *55*, 5367-5376.
- 462 10. Leistra, M.; Van der Linden, A.; Boesten, J.; Tiktak, A.; Van den Berg, F. *Pearl model for*
463 *pesticide behaviour and emissions in soil-plant systems: Description of the processes in focus*
464 *pearl v 1.1. 1*. Alterra, Green World Research Wageningen: 2001.
- 465 11. Bicknell, B.R.; Imhoff, J.C.; Kittle Jr, J.L.; Jobes, T.; Donigian Jr, A.; Johanson, R. Hydrological
466 simulation program-fortran: Hspf version 12 user's manual (computer program manual). *AQUA*
467 *TERRA Consultants, Mountain View, California* **2001**.
- 468 12. Neitsch, S.; Arnold, J.; Kiniry, J.; Williams, J. Soil and water assessment tool (swat) theoretical
469 documentation. Grassland soil and water research laboratory. *Agricultural Research Service,*
470 *Blackland Research Center, Texas Agricultural Experiment Station* **2005**.
- 471 13. Metta, J.W.P.; Bolte, J.; Vaché, K.; Jepson, P.; Jenkins, J.; Mc Donnel, J. Wethydro: A
472 watershed-scale, fully distributed model of pesticide fate/transport. *Geological Society of*
473 *America Abstract with Programs* **2006**, *38*, 40.

- 474 14. Quilbé, R.; Rousseau, A.N.; Villeneuve, J.P. Gibsi: an integrated modelling software for
475 watershed management. *Geophysical Research Abstracts* **2006**.
- 476 15. Ganzelmeier, H.; Rautmann, D.; Spangenberg, R.; Streloke, M. Studies on the spray drift of plant
477 protection products: Mitteilungen aus der biologischen bundesanstalt für land- und
478 forstwirtschaft. **1995**.
- 479 16. Holterman, H.; van De Zande, J. Imag draft report. Imag drift calculator v 1.1, user manual.
480 2003.
- 481 17. Holterman, H.J.; van de Zande, J.C.; Porskamp, H.A.J.; Huijsmans, J.F.M. Modelling spray drift
482 from boom sprayers. *Computers and Electronics in Agriculture* **1997**, *19*, 1–22.
- 483 18. Nuyttens, D.; de Schamphelre, M.; Baetens, K.; Sonck, B. The influence of operator-controlled
484 variables on spray drift from field crop sprayers. *Transactions of the ASABE* **2007**, *50*, 1129–
485 1140.
- 486 19. Baetens, K.; Ho, Q.T.; Nuyttens, D.; de Schamphelre, M.; Endalew, A.M.; Hertog, M.; Nicolai,
487 B.; Ramon, H.; Verboven, P. A validated 2-d diffusion–advection model for prediction of drift
488 from ground boom sprayers. *Atmospheric Environment* **2009**, *43*, 1674–1682.
- 489 20. García-Santos, G.; Feola, G.; Nuyttens, D.; Diaz, J. Drift from the use of hand-held knapsack
490 pesticide sprayers in boyacá (colombian andes). *Journal of agricultural and food chemistry*
491 **2015**, *x*, xx-xx. DOI: 10.1021/acs.jafc.5b03772
- 492 21. Scholtz, M.; Voldner, E.; McMillan, A.; Van Heyst, B. A pesticide emission model (pem) part i:
493 Model development. *Atmospheric Environment* **2002**, *36*, 5005–5013.
- 494 22. Birkved, M.; Hauschild, M.Z. Pestlci—a model for estimating field emissions of pesticides in
495 agricultural lca. *Ecological Modelling* **2006**, *198*, 433–451.

- 496 23. Dijkman, T.J.; Birkved, M.; Hauschild, M.Z. Pestlci 2.0: A second generation model for
497 estimating emissions of pesticides from arable land in lca. *The International Journal of Life*
498 *Cycle Assessment* **2012**, *17*, 973–986.
- 499 24. van Zelm, R.; Larrey-Lassalle, P.; Roux, P. Bridging the gap between life cycle inventory and
500 impact assessment for toxicological assessments of pesticides used in crop production.
501 *Chemosphere* **2014**, *100*, 175-181.
- 502 25. Andreoli, R. *Analysing the pesticide fate model wetspa-pest for its application in "vereda la*
503 *hoya, colombia": In context of the project: "Reducing human health and environmental risks*
504 *from pesticide use: Integrating decisionmaking and spatial risk assessment models: The case of*
505 *vereda la hoya, colombia*. Geographisches Institut der Universität Zürich: 2010.
- 506 26. Blanco, L.E.; Aragón, A.; Lundberg, I.; Wesseling, C.; Nise, G. The determinants of dermal
507 exposure ranking method (derm): A pesticide exposure assessment approach for developing
508 countries. *The Annals of Occupational Hygiene* **2008**, *52*, 535–544.
- 509 27. Cherrie, J.W.; Tickner, J.; Friar, J. *Evaluation of further development of the ease model 2.0*. HSE
510 Books: 2003; p viii, 127.
- 511 28. Dosemeci, M.; Alavanja, M.C.R.; Rowland, A.S.; Mage, D.; Hoar Zahm, S.; Rothman, N.;
512 Lubin, J.H.; Hoppin, J.A.; Sandler, D.P.; Blair, A. A quantitative approach for estimating
513 exposure to pesticides in the agricultural health study. *The Annals of Occupational Hygiene*
514 **2002**, *46*, 245–260.
- 515 29. Garrod, A.N.I.; Rajan-Sithamparamanadarajah, R. Developing coshh essentials: Dermal exposure,
516 personal protective equipment and first aid. *Annals of Occupational Hygiene* **2003**, *47*, 577–588.

- 517 30. Van-Wendel-de-Joode, B.; Brouwer, D.H.; Vermeulen, R.; van Hemmen, J.J.; Heederik, D.;
518 Kromhout, H. Dream: A method for semi-quantitative dermal exposure assessment. *The Annals*
519 *of Occupational Hygiene* **2003**, *47*, 71–87.
- 520 31. van Hemmen, J.J.; Auffarth, J.; Evans, P.G.; Rajan-Sithamparanadarajah, B.; Marquart, H.;
521 Oppl, R. Riskofderm: Risk assessment of occupational dermal exposure to chemicals. An
522 introduction to a series of papers on the development of a toolkit. *The Annals of Occupational*
523 *Hygiene* **2003**, *47*, 595–598.
- 524 32. Marquart, H.; Heussen, H.; Le Feber, M.; Noy, D.; Tielemans, E.; Schinkel, J.; West, J.; van der
525 Schaaf, D. 'Stoffenmanager', a web-based control banding tool using an exposure process model.
526 *Annals of Occupational Hygiene* **2008**, *52*, 429–441.
- 527 33. Lesmes-Fabian, C.; Teubl, S.; Binder, C.R. Evaluation of models for dermal exposure
528 assessment in farming systems in developing countries. *journal of Environmental Engineering*
529 *and Ecological Science* **2014**, *3*, 1.
- 530 34. Lesmes-Fabian, C. Correction: Dermal exposure assessment to pesticides in farming systems in
531 developing countries: Comparison of models. *Int. J. Environ. Res. Public health* **2015**, *12*, 4670–
532 4696. *International journal of environmental research and public health* **2015**, *12*, 9264.
- 533 35. Gil, Y.; Sinfort, C. Emission of pesticides to the air during sprayer application: A bibliographic
534 review. *Atmospheric Environment* **2005**, *39*, 5183–5193.
- 535 36. Craig, I.P. The gds model - a rapid computational technique for the calculation of aircraft spray
536 drift buffer distances. *Computers and Electronics in Agriculture* **2004**, *43*, 235–250.
- 537 37. de Schampheleire, M.; P., S.; Brusselman, E.; Sonck, S. Risk assessment of pesticide spray drift
538 damage in belgium. *Crop Protection* **2007**, *26*, 602–611.

- 539 38. Keil, C.B. A tiered approach to deterministic models for indoor air exposures. *Applied*
540 *Occupational and Environmental Hygiene* **2000**, *15*, 145–151.
- 541 39. García-Santos, G.; Scheiben, D.; Binder, C.R. The weight method: A new screening method for
542 estimating pesticide deposition from knapsack sprayers in developing countries. *Chemosphere*
543 **2011**, *82*, 1571–1577.
- 544 40. Feola, G.; Rahn, E.; Binder, C.R. Suitability of pesticide risk indicators for less developed
545 countries: A comparison. *Agriculture, Ecosystems & Environment* **2011**, *142*, 238–245.
- 546 41. Balderacchi, M.; Trevisan, M. Comments on pesticide risk assessment by the revision of
547 directive eu 91/414. *Environmental Science and Pollution Research* **2010**, *17*, 523–528.
- 548 42. Meli, S.M.; Renda, A.; Nicelli, M.; Capri, E. Studies on pesticide spray drift in a mediterranean
549 citrus area. *Agronomie* **2003**, *23*, 667–672.
- 550 43. Rossing, W.A.H.; Zander, P.; Josien, E.; Groot, J.C.J.; Meyer, B.C.; Knierim, A. Integrative
551 modelling approaches for analysis of impact of multifunctional agriculture: A review for france,
552 germany and the netherlands. *Agriculture, Ecosystems & Environment* **2007**, *120*, 41–57.
- 553 44. Feola, G.; Sattler, C.; Saysel, A.K. Simulation models in farming systems research: Potential and
554 challenges. In *Farming systems research into the 21st century: The new dynamic*, Darnhofer, I.;
555 Gibbon, D.; Dedieu, B., Eds. Springer Dordrecht, Netherlands, 2012; pp 281–306.
- 556 45. Binder, C.R.; Feola, G.; Steinberger, J.K. Considering the normative, systemic and procedural
557 dimensions in indicator-based sustainability assessments in agriculture. *Environmental Impact*
558 *Assessment Review* **2010**, *30*, 71–81.
- 559 46. Rossing, W.A.H.; Opdam, P.; van der Knaap, W.; Grashof-Bokdam, C. Landscape prototypes for
560 multifunctional farming - seeking synergy between functional biodiversity and other green
561 services at field, farm and landscape scales. *IOBC-WPRS Bull* **2003**, *26*, 129–134.

- 562 47. Groot, J.C.J.; Rossing, W.A.H.; Jellema, A.; Stobbelaar, D.J.; Renting, H.; van Ittersum, M.K.
563 Exploring multi-scale trade-offs between nature conservation, agricultural profits and landscape
564 quality — a methodology to support discussions on land-use perspectives. *Agriculture,*
565 *Ecosystems & Environment* **2007**, *120*, 59-70.
- 566 48. Vatn, A.; Bakken, L.; Botterweg, P.; Romstad, E. Ececmmod: An interdisciplinary modelling
567 system for analyzing nutrient and soil losses from agriculture. *Ecological Economics* **1999**, *30*,
568 189–206.
- 569 49. Belcher, K.W.; Boehm, M.M.; Fulton, M.E. Agroecosystem sustainability: A system simulation
570 model approach. *Agricultural Systems* **2004**, *79*, 225–241.
- 571 50. Stoorvogel, J.; Antle, J.M.; Crissman, C.; Bowen, W. The tradeoff analysis model: Integrated
572 bio-physical and economic modeling of agricultural production systems. *Agricultural Systems*
573 **2004**, *80*, 43-66.
- 574 51. Schoell, R.; Binder, C.R. System perspectives of experts and farmers regarding the role of
575 livelihood assets in risk perception: Results from the structured mental model approach. *Risk*
576 *analysis : an official publication of the Society for Risk Analysis* **2009**, *29*, 205–222.
- 577 52. Feola, G.; Binder, C.R. Towards an improved understanding of farmers' behaviour: The
578 integrative agent-centred (iac) framework. *Ecological Economics* **2010**, *69*, 2323–2333.
- 579 53. Feola, G.; Binder, C.R. Identifying and investigating pesticide application types to promote a
580 more sustainable pesticide use. The case of smallholders in boyacá, colombia. *Crop Protection*
581 **2010**, *29*, 612–622.
- 582 54. Liu, Y.B.; Smedt, F.d. *Wetspa extension, a gis-based hydrologic model for flooded prediction and*
583 *watershed management. Documentation and user manual.* 2004.

- 584 55. Instituto Geográfico Agustín Codazzi, I.G.A.C. Urban and rural cadastral actualization for the
585 municipality of tunja. Departamento Nacional de Estadística (DANE): 2005a.
- 586 56. Mosquera-Vivas, C.S.; García-Santos, G.; Celis-Ossa, R.E.; Hellweg, S.H.; Juraske, R.;
587 González-Murillo, C.A.; Guerrero-Dallos, J.A. Adsorption, desorption and degradation
588 behaviour of the pesticides pyrimethanil, dimethomorph and chlorpyrifos through a tropical
589 colombian agricultural soil profile. *Unpublished a*.
- 590 57. Mosquera-Vivas, C.S.; Martinez-Cordon, M.J.; García-Santos, G.; Guerrero-Dallos, J.A.
591 Adsorption-desorption studies of tebuconazole in tropical colombian agricultural soils.
592 *Unpublished b*.
- 593 58. Tuchschnid, M. *Field survey: Roundtable with the farmers. Vereda la hoya, colombia*. 2004.
- 594 59. Tuchschnid, M. Gis-basierende räumliche risikoabschätzung vom pestizideinsatz in der
595 landwirtschaft fallstudie: Vereda la hoya, kolumbien. Diplomarbeit, ETH (Eidgenössische
596 Technische Hochschule Zürich), Zurich, 2004.
- 597 60. Lesmes-Fabian, C.; García-Santos, G.; Leuenberger, F.; Nuyttens, D.; Binder, C.R. Dermal
598 exposure assessment of pesticide use: The case of sprayers in potato farms in the colombian
599 highlands. *The Science of the Total Environment* **2012**, *430*, 202–208.
- 600 61. García-Santos, G.; Feola, G.; Nuyttens, D.; Diaz, J. Drift from the use of handled knapsack
601 pesticide sprayers in boyacá (colombian andes). *This issue*.
- 602 62. García-Santos, G.; Scheiben, D.; Binder, C.R. The weight method: A new screening method for
603 estimating pesticide deposition from knapsack sprayers in developing countries. *Chemosphere*
604 **2011**, *82*, 1571-1577.

- 605 63. Linders, J.; Mensink, H.; Stephenson, G.; Wauchope, D.; Racke, K. Foliar interception and
606 retention values after pesticide application. A proposal for standardized values for environmental
607 risk assessment (technical report). *Pure and applied chemistry* **2000**, *72*, 2199–2218.
- 608 64. García-Santos, G.; Keller-Forrer, K. Avoidance behaviour of *eisenia fetida* to carbofuran,
609 chlorpyrifos, mancozeb and metamidophos in natural soils from the highlands of colombia.
610 *Chemosphere* **2011**, *84*, 651–656.
- 611 65. Binder, C.; Patzel, N. Preserving tropical soil organic matter at watershed level. A possible
612 contribution of urban organic wastes. *Nutrient Cycling in Agroecosystems* **2001**, *61*, 171-181.
- 613 66. Vargas, E.M. *Evaluación de la lixiviación de pesticidas hacia las aguas subterráneas de la*
614 *vereda de la hoya del municipio de tunja, por medio del modelo pearl 2.2.2. Trabajo de grado*
615 *por universidad de boyacá, ingeniería sanitaria y ambiental*. Colombia, 2005.
- 616 67. Ministerio de Agricultura y Desarrollo Rural, M. *La cadena de la papa en colombia. Una mirada*
617 *global de su estructura y dinámica 1991-2005*. Ministerio de Agricultura y Desarrollo Rural:
618 Columbia, 2006.
- 619 68. Nash, J.; Sutcliffe, J.V. River flow forecasting through conceptual models part i—a discussion of
620 principles. *Journal of hydrology* **1970**, *10*, 282-290.
- 621 69. Juraske, R.; Mosquera Vivas, C.S.; Erazo-Velásquez, A.; García-Santos, G.; Berdugo-Moreno,
622 M.B.; Diaz-Gomez, J.; Binder, C.R.; Hellweg, S.; Guerrero-Dallos, J.A. Pesticide uptake in
623 potatoes: Model and field experiments. *Environmental Science & Technology* **2011**, *45*, 651–
624 657.
- 625 70. Gobernación de Boyacá Direccion de Desarrollo Agropecuario. Acuerdo de competitividad de la
626 cadena agroalimentaria de la papa, para el Departamento de Boyacá. 2004, Tunja, Colombia.

627 **Tables captions**

Table 1: Goodness of fit coefficients for the calibration and validation periods for water discharge.

Figures captions

Figure 1: Be-Westspa-Pest model structure including input, core model and output data.

Figure 2: Representation of the detailed software design of the integrative approach Be-WetSpa-Pest.

Figure 3: Comparison of measured (blue) and modeled (green) concentrations of Chlorpyrifos in soil (mg/kg soil) along the different growth stages of a potato field within the study region (La Hoya, Boyaca). Vertical bars denote standard deviation.

Figure 4: Simulation results for pesticide application of Mancozeb (six applications) on a sample field, for a total cycle of 140 days (method even). The triangles show the application dates. The dashed line and the continuous line show the soil respectively the topsoil concentration. The dotted line shows the concentration on the canopy.

Figure 5: Spatial distribution of Mancozeb concentration in the watershed. Four consecutives cycles were simulated (baseline scenario, method even). In the green marked areas no pesticides were applied during this simulation run.

Figure 6: Total amount of pesticide applied in one cycle (140 days) and the whole area.
i: insecticide; f: fungicide

Figure 7: Spatial distribution of Mancozeb concentration in the watershed. Four consecutives cycles were simulated (8a: training scenario; 8b: cooperative scenario, method even). In the green marked areas no pesticides were applied during this simulation run.

Tables

Table 1:


Coefficient	Calibration	Validation
Model bias	-0.162	0.151
Nash-Sutcliffe efficiency	0.994	0.9966
RMSE	0.233	0.2375
Pearson coefficient	0.848	0.956
Standard error	0.012	0.008


651

652

Figures graphics

Figure 1:



667 **Figure 2:**

668

669

Figure 3:





Figure 4:






680 **Figure 5:**

Figure 6:




707 **Figure 7:**

719 TOC graphic



Training by companies

Cooperative



720