

*Remember, remember the fifth of
November: was that thunder I heard or
not?*

Article

Accepted Version

Owens, M. J. ORCID: <https://orcid.org/0000-0003-2061-2453>
(2016) Remember, remember the fifth of November: was that
thunder I heard or not? *Weather*, 71 (6). pp. 134-137. ISSN
0043-1656 doi: <https://doi.org/10.1002/wea.2725> Available at
<https://centaur.reading.ac.uk/63950/>

It is advisable to refer to the publisher's version if you intend to cite from the
work. See [Guidance on citing](#).

To link to this article DOI: <http://dx.doi.org/10.1002/wea.2725>

Publisher: Wiley

All outputs in CentAUR are protected by Intellectual Property Rights law,
including copyright law. Copyright and IPR is retained by the creators or other
copyright holders. Terms and conditions for use of this material are defined in
the [End User Agreement](#).

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online

Remember, remember the fifth of November: Was that thunder I heard or not?

M.J. Owens¹

¹Space and Atmospheric Electricity Group, Department of Meteorology, University of Reading, Earley Gate, PO Box 243, Reading RG6 6BB, UK

Abstract

“Thunder days” are simple records of thunderstorm activity, logging whether a human observer heard thunder on a particular day or not. Despite their low dynamic range and inherent subjectivity, thunder days are invaluable as the only long-term observations of thunderstorm occurrence, with some records stretching back into the 19th century. Thunder days, however, are potentially susceptible to false positives, particularly from explosions. Thus one might expect UK thunder days to show anomalously high counts on New Year’s Eve and days around 5 November, Bonfire Night, both of which are celebrated with large firework displays across the country. It is demonstrated that Met Office records of thunder days between 1980 and 2010 do not show any significant increase in thunder reporting around 5 November or 31 December. In fact, the days around 5 November exhibit the largest reduction in the amount of reported thunder relative to annual climatology. While meteorological variability cannot be completely ruled out, this result is suggestive of observer bias; It is speculated that human observers, armed with a priori knowledge of the likelihood of false positives, “second guess” themselves to a greater degree around 5 November than the rest of the year. In fact, the data suggest they should trust in their ability to correctly discriminate between thunder and fireworks.

Introduction

The only long-term observations of thunderstorm activity, extending back more than 100 years, are “thunder days,” wherein an observer records 1 or a 0 depending on whether or not (they think) they have heard thunder that day [e.g., *Brooks*, 1925; *Changnon*, 1985; *Kitagawa*, 1989]. Despite the low dynamic range (e.g., a storms with 1 or 1000 lightning strokes will both simply register a 1 in thunder days) and the inherent subjectivity of such measurements, they're invaluable for long-term studies of thunderstorm occurrence, which could vary as a result of global warming [*Romps et al.*, 2014] or even changes in the space environment [*Stringfellow*, 1974; *Owens et al.*, 2014].


Thunder day observations are potentially susceptible to false positives, such as vehicle noise or natural/anthropogenic explosions being wrongly attributed to thunder [e.g., *Rampino*, 1989]. Thus increasing urbanisation and industrialisation may result in a long-term change in the noise level and hence bias in the data. On shorter timescales, “Bonfire” or “Guy Fawkes” night, as well as New Year’s Eve, in the UK (5 November) are obvious candidates for false-positive thunder identification.

Data and analysis

Thunder day records produced by professional observers at UK Met Office meteorological stations are available from “Met Office Integrated Data Archiving System (MIDAS) land and marine surface stations (1853–current),” made available by the NCAS British Atmospheric Data Centre (<http://badc.nerc.ac.uk>). The measurement is a simple one: Met Office observers simply record a thunder day on any day they hear thunder [*Lewis*, 1991]. There is no formal training in discriminating between thunder and false positives, and there is no stipulation to use any additional instrumentation for verification [J. Wilkinson, personal communication]. In practice, however, on the suspected identification of thunder observers may, on occasion, also consult radar data or radio lightning observations (or “spherics”) if/when available [P. Inness, personal communication]. Conversely, it is not possible for a thunder day to be recorded without the observer having heard thunder, thus

44 distance bright fireworks confused with lightning flashes should not result in false positives in the
45 thunder day record.

46 Prior to 1950, thunder days were recorded in the climatological returns of manned UK stations, with
47 monthly totals available in the Monthly Weather Report. In the MIDAS daily dataset, thunder day
48 records routinely began in 1950, with the number of manned UK stations making such observations
49 was limited to fewer than 10 until 1957 and fewer than 100 until 1971. After this date, there were
50 approximately 400 stations making thunder day observations, though it unfortunately slowly tapered
51 off after 2000, to fewer than 200 again at the end of 2010 [see also *Perry and Hollis, 2005* and Figure
52 1a]. Due to the difficulty in discriminating between an observation of no thunder and no thunder
53 observations, any station making a single observation of thunder in a given year is assumed to be
54 actively observing and included in the analysis. T , the fraction of UK observing stations which recorded
55 thunder on a given day, is constructed from individual station data.




56

57 Figure 1: (a) The number of stations making thunder day observations. (b) The daily (black and left-hand axis) and annual
58 (red and right-hand axis) values of T , the fraction of UK stations reporting thunder on a given day.

59

60 Multi-shell rockets and slow-burn composite fireworks are most likely to be mistaken for thunder
61 (Pyrosociety.org.uk, personal communication). It is difficult to obtain data on the widespread use of
62 such fireworks in the UK, but anecdotally at least, they appear to have been in wide-spread public use
63 for UK Bonfire night celebrations since at least the early 1980s. Thus this study primarily considers the
64 period 1980–2010. Changing the start year for this interval by ± 10 years does not qualitatively change
65 the results reported in this study, as discussed later. Figure 1b shows daily (black) and annual (red) T .
66 As expected, there is a very strong seasonal variation in T . There is some evidence in a change in the
67 annual mean T associated with the increase in number of stations from 1950 through to the early
68 1970s. After this time, there does not appear to be any immediate correspondence between changes
69 in the number of stations and annual T , suggesting that the spatial distribution of reporting stations is
70 not changing dramatically through the period of interest (1980–2010).



71

72 **Figure 2: A composite of the annual mean variation in T for the period 1980–2010. Black lines show the daily values, red**
73 **lines show 50-day running means. The blue dashed line shows 5 November (for non-leap years).**

74 In order to compare the reporting of thunder on 5 November with the rest of the year, it is necessary
75 to subtract the strong annual variation which dominates thunderstorm occurrence. Mean T for each
76 day of year is computed over the whole 1980–2010 period (the black lines in Figure 2). In order to
77 make an annual climatology, a 50-day running mean of the annual composite is computed (the red
78 line in Figure 2), denoted T_{50} . The size of this smoothing window does not qualitatively affect the
79 results presented here, as long as it is greater than ~ 10 days and shorter than ~ 6 months. The
80 climatological deviation is then taken to be the difference between the smoothed and unsmoothed
81 value (i.e., $T - T_{50}$). As the magnitude of the climatological deviation will be much greater during times
82 of higher average T (summer) than lower average T (winter), the fractional climatological deviations,
83 ΔT , are computed (i.e., $\Delta T = [T - T_{50}] / T_{50}$).

84 Organised firework displays and private firework use may occur a number of days around 5 November,
85 particularly in years when it falls during the middle of the week and the celebrations are shifted either
86 forward or backward to the weekend. If it assumed that fireworks use is focussed on the Friday or
87 Saturday closest to 5 November, then on average this will mean fireworks use is within ± 1.3 days on


88 fireworks night. To capture this, a 3-day running mean is applied to ΔT , yielding $\Delta T3$. This may miss
89 some activity in years when fireworks celebrations are split between the weekends before/after 5
90 November, but this is preferable to over-smoothing the whole data set and removing the possible
91 signal.

92 An annual composite of $\Delta T3$, centred on 5 November so as to allow for leap days, is shown for the
93 1980–2010 period in Figure 3b. In this annual composite, there is clearly no evidence for an increase
94 in the false positive reporting of thunder due to fireworks. In fact, 5 November is the lowest mean $\Delta T3$
95 value.

96 The most basic estimate of the probability of the lowest $\Delta T3$ value falling on 5 November purely by
97 chance is that it is $1/N$, in this case approximately 1 in 365 (or closer to 1 in 100 if the 3-day smoothing
98 window is considered). But it is more important to quantify the probability that such a value ($\langle \Delta T3 \rangle =$
99 -0.67) falls within the natural meteorological variability present in the data. The following “Monte
100 Carlo” approach is taken. Instead of using 5 November in each year as the $t=0$ time for the composite
101 to produce $\langle \Delta T3 \rangle$, random times throughout each year are selected and a new $\langle \Delta T3 \rangle$ computed. This
102 is done 10,000 times. The grey-shaded panels on Figure 3 show the bands which contain 90, 95 and
103 99% of the $\langle \Delta T3 \rangle$ values about these randomly selected times. It can be seen that 5 November is
104 below the 99% band, also suggesting approximately a 1 in 100 probability of the low value occurring
105 purely by chance. (Note, however, that this test implicitly assumes the annual variation has been
106 adequately removed by the climatology.) Figure 4b shows the probability density function of $\langle \Delta T3 \rangle$
107 for the 1980–2010 period, using a kernel density estimate. It can be seen that while 5 November is
108 the lowest $\langle \Delta T3 \rangle$ value, it sits at the edge of the overall variability, rather than being an extreme
109 outlier. Finally, as a simple test of the robustness of this signal, individual decades of data within this
110 period are considered. For the years 1980–1989, 1991–2000 and 2001–2010, $\langle \Delta T3 \rangle = -0.64$, -0.51
111 and -0.86 , and the rank within the distribution is 15, 53 and 2 of 365, respectively. So in all cases, 5
112 November is well below climatology (though the climatology itself will be poorly described for such
113 short intervals).


114 Note that Figure 3b also shows a large positive peak in $\langle \Delta T3 \rangle$ at $t = 55$ days, i.e., 30 December. It is
115 only significant relevant to the meteorological variation at the 95% level and only present in the 1990–
116 1999 data, not in the 1980s or 2000s. Its possible relation to fireworks displays associated with New
117 Year’s Eve celebrations is discussed later.

118



119

120 **Figure 3:** Mean fractional climatological deviation of thunder days, $\Delta T3$, about 5 November. Panel (a) shows the period
 121 1957–1979. Panel (b) shows the period 1980–2010. The grey-shaded areas shows the intervals containing 90, 95 and 99%
 122 of the variations from a Monte Carlo sampling of $\Delta T3$ in the given period. Note that in panel (b) there is no evidence for
 123 an enhanced false positive rate around 5 November. In fact, $\Delta T3$ on 5 November is the lowest observed, suggesting a
 124 systematic under-reporting of thunder.



125

126 **Figure 4:** Probability density functions, computed from kernel density, of $\Delta T3$ for the (a) 1957–1979 and (b) 1980–2010
 127 periods. The grey-shaded areas shows the intervals containing 90, 95 and 99% of the variations from a Monte Carlo
 128 sampling of ΔT in the given period. The red lines show the 5 November.

129 As a comparison to the 1980–2010 period, wherein UK fireworks use around 5 November is expected
 130 to be widespread, the earlier period, 1957-1979, is considered here. 1957 is taken as the start date as
 131 the number of stations rises to approximately 100 (see Figure 1a). Unfortunately, there is a large jump
 132 in the number of stations in the early 1970s, but limiting the period to 1973–1979 reduces the
 133 meaningfulness of the climatology required to compute $\Delta T3$, which is even less desirable. With this

134 limitation in mind, Figures 3a and 4a show the analysis for 1957–1979. From the asymmetric
135 probability distribution function, it is clear the change in number of stations is having an effect.
136 Nevertheless, it can be seen that 5 November for this periods lies somewhere within the middle of the
137 $\langle \Delta T3 \rangle$ variability. Also of note is that there is one point below the 99% lower bound ($\langle \Delta T3 \rangle = -0.62$ at
138 $+162$ days from 5 November, i.e., 25 or 26 April). This is to be expected as part of the normal
139 meteorological variability (i.e., by definition, 1 point in 100 should be outside the 99% band on
140 average).

141 Discussion and conclusions

142 Audible thunder records, or “thunder days,” are expected to be susceptible to false positives,
143 particularly from explosions. Thus in the UK, one might reasonably expect an over-reporting of
144 thunder on and around 5 November. Fireworks experts were spilt on the issue, with some expressing
145 doubt that shells and rockets would be mistaken for thunder, while others argued that fireworks are,
146 on occasion, mistaken for thunder, particularly multi-shell fireworks and slow-burning compositions
147 [Pyrosociety.org.uk, personal communication]. In fact, no increase in thunder, relative to climatology,
148 is found for the 1980–2010 period when the signal is expected to be most pronounced. Similarly, there
149 is not strong evidence for an enhancement in thunder reporting on New Year’s Eve, as the peak is on
150 30 December and celebrations are not traditionally spread around that time. Furthermore, the peak
151 is only present in the 1990–1999 interval and not 1980–1989 or 2000–2009. Thus there is no direct
152 evidence for false positives in the thunder day data as a result of fireworks.

153 In fact, around 5 November for 1980–2010, there has been a dearth of thunder reported (relative to
154 the climatological variation). A similar drop in reported thunder is not found for 31 December, but
155 New Year’s Eve firework displays are a more recent phenomenon and personal firework use remains
156 more limited. The apparent lack of thunder for 5 November is at the edge of the observed variability
157 in the data, so chance or simple meteorological variation cannot be completely ruled out. But an
158 alternative explanation of observer bias seems more plausible. Audible thunder records are not
159 compiled by an unthinking listening device or computer algorithm. They are put together by human
160 observers, which come bundled with “a priori” knowledge that around 5 November there are a lot of
161 loud noises which, they believe, can be mistaken for thunder. Thus they will be more likely to “second
162 guess” [Pliske et al., 2004] any potential thunder observation as a false positive resulting from loud
163 fireworks. The results presented here suggest observers are actually playing it too safe. The actual
164 probability of an observer mistaking fireworks for thunder is much lower than the observer assumes.
165 Thus observers should trust more in their ability to discriminate between thunder and fireworks.

166 Acknowledgements

167 MO is part-funded by Science and Technology Facilities Council (STFC) grant number ST/M000885/1
168 and acknowledges support from the Leverhulme Trust through a Philip Leverhulme Prize.

169 References

- 170 Brooks, C. E. P. (1925), *The distribution of thunderstorms over the globe*, HM Stationery Office.
171 Changnon, S. A. (1985), Secular variations in thunder-day frequencies in the twentieth century, *J.*
172 *Geophys. Res.*, 90(D4), 6181-6194.
173 Kitagawa, N. (1989), Long-term variations in thunder-day frequencies in Japan, *J. Geophys. Res.*,
174 94(D11), 13183-13189.
175 Lewis, R. P. W. (1991), *Meteorological glossary*, 6th Edition ed., Her Majesty's Stationary Office,
176 London.
177 Owens, M., C. Scott, M. Lockwood, L. Barnard, R. Harrison, K. Nicoll, C. Watt, and A. Bennett (2014),
178 Modulation of UK lightning by heliospheric magnetic field polarity, *Env. Res. Lett.*, 9(11), 115009.

179 Perry, M., and D. Hollis (2005), The generation of monthly gridded datasets for a range of climatic
180 variables over the UK, *International Journal of Climatology*, 25(8), 1041-1054.
181 Pliske, R. M., B. Crandall, and G. Klein (2004), Competence in Weather Forecasting, in *Psychological*
182 *investigations of competence in decision making*, edited by J. Shanteau and P. Johnson, Cambridge
183 University Press, Cambridge.
184 Rampino, M. R. (1989), Distant effects of the Tambora Eruption of April 1815: An eyewitness account,
185 *Eos, Transactions American Geophysical Union*, 70(51), 1559-1559.
186 Roms, D. M., J. T. Seeley, D. Vollaro, and J. Molinari (2014), Projected increase in lightning strikes in
187 the United States due to global warming, *Science*, 346(6211), 851-854.
188 Stringfellow, M. F. (1974), Lightning incidence in Britain and the solar cycle, *Nature*, 249, 332-333,
189 doi:10.1038/249332a0.

190