

The use of asparaginase to reduce acrylamide levels in cooked food

Article

Accepted Version

Creative Commons: Attribution-Noncommercial-No Derivative Works 4.0

Xu, F., Oruna-Concha, M.-J. and Elmore, J. S. (2016) The use of asparaginase to reduce acrylamide levels in cooked food. *Food Chemistry*, 210. pp. 163-171. ISSN 0308-8146 doi: <https://doi.org/10.1016/j.foodchem.2016.04.105> Available at <https://centaur.reading.ac.uk/64518/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

To link to this article DOI: <http://dx.doi.org/10.1016/j.foodchem.2016.04.105>

Publisher: Elsevier

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online

The use of asparaginase to reduce acrylamide levels in cooked food

Fei Xu, Maria-Jose Oruna-Concha, J. Stephen Elmore*

Department of Food and Nutritional Sciences, University of Reading, Whiteknights,
Reading RG6 6AP, UK

E-mail addresses:

Mr Fei Xu: f.xu@pgr.reading.ac.uk

Dr Maria-Jose Oruna-Concha; M.J.Oruna-Concha@reading.ac.uk

Dr Stephen Elmore j.s.elmore@reading.ac.uk (*corresponding author; tel.: +118-3787455)

RESEARCH HIGHLIGHTS

Strategies to reduce acrylamide in cooked food often rely on a reduction in the Maillard reaction.

Mitigation procedures that modify the Maillard reaction may negatively affect flavour and colour.

Asparaginase may reduce acrylamide formation, while maintaining sensory quality.

This review collates research on the use of enzymes to mitigate acrylamide formation.

25

26 ABSTRACT

27 Strategies proposed for reducing the formation of the suspected carcinogen acrylamide in
28 cooked foods often rely on a reduction in the extent of the Maillard reaction, in which
29 acrylamide is formed from the reaction between asparagine and reducing sugars. However,
30 the Maillard reaction also provides desirable sensory attributes of cooked foods. Mitigation
31 procedures that modify the Maillard reaction may negatively affect flavour and colour. The
32 use of asparaginase to convert asparagine to aspartic acid may provide a means to reduce
33 acrylamide formation, while maintaining sensory quality. This review collates research on the
34 use of enzymes, asparaginase in particular, to mitigate acrylamide formation. Asparaginase is
35 a powerful tool for the food industry and it is likely that its use will increase. However, the
36 potential adverse effects of asparaginase treatment on sensory properties of cooked foods and
37 the need to achieve sufficient enzyme–substrate contact remain areas for future research.

38

39 **Key words:** Acrylamide, asparaginase, enzymes, asparagine, reducing sugars, Maillard
40 reaction

41

42

43 **1. Introduction**

44 It is now over ten years since the Swedish Food Authority and the University of
45 Stockholm confirmed the existence of the suspected carcinogen acrylamide in a variety of
46 heated foods (Tareke, Rydberg, Karlsson, Eriksson, & Tornqvist, 2002). Several months after
47 the announcement, researchers showed that acrylamide is formed from asparagine and
48 reducing sugars during the Maillard reaction (Mottram, Wedzicha, & Dodson, 2002; Stadler
49 et al., 2002). As shown in Figure 1, asparagine and reducing sugars take part in a conjugation
50 reaction resulting in the formation of *N*-glycosylasparagine, which as a result of high
51 temperature treatment will form a decarboxylated Schiff base. The decarboxylated Schiff
52 base may decompose directly to form acrylamide or may hydrolyse to form 3-
53 aminopropionamide (Hedegaard, Frandsen, & Skibsted, 2008). 3-Aminopropionamide is also
54 believed to be an important precursor of acrylamide (Granvogl & Schieberle, 2006).

55 Since 2002, the food industry worldwide has collaborated with scientists, in order to
56 reduce the levels of acrylamide in cooked foods. Mitigation techniques can be separated into
57 three different types. Firstly, starting materials low in acrylamide precursors can be used to
58 reduce the acrylamide in the final product. Secondly, process conditions may be modified, in
59 order to decrease the amount of acrylamide formation. Thirdly, post-process intervention
60 could be used to reduce acrylamide (Pedreschi, Mariotti, & Granby, 2014). While the third
61 approach is not widely considered, an example is the use of supercritical CO₂ extraction to
62 reduce acrylamide levels in coffee. Almost 80% of the acrylamide was removed using this
63 technique (Banchero, Pellegrino, & Manna, 2013), although further sensory tests are needed
64 to validate the effect on food quality.

65 This review will describe the main mitigation strategies used for acrylamide but will focus
66 on the use of enzymes, in particular asparaginase, to reduce levels of acrylamide precursors.
67 A more general review on acrylamide mitigation has been recently published (Friedman,

68 2015), while another recent review has covered the latest studies on the sources, purification,
69 and characterisation of L-asparaginase and its application in both the pharmaceutical and
70 food industries (Zuo, Zhang, Jiang, & Mu, 2015a).

71

72 2. Acrylamide mitigation strategies

73 2.1 Raw materials

74 Decreasing the amounts of acrylamide precursors will have a huge impact on final
75 acrylamide production (Zyzak et al., 2003). However, the effect will be dependent on the
76 relative levels of precursors. If total reducing sugars are present at higher levels than
77 asparagine in a food, reduction in asparagine will have the greater effect on acrylamide
78 formation, and *vice versa*. Numerous papers have demonstrated that acrylamide formation is
79 proportional to reducing sugar concentrations in potato (Elmore, Briddon, Dodson,
80 Muttucumar, Halford, & Mottram, 2015; Ohara-Takada et al., 2005; Vinci, Mestdagh, & De
81 Meulenaer, 2012), while in cereals, such as rye and wheat, acrylamide formation is
82 proportional to asparagine content (Curtis et al., 2010; Halford, Curtis, Muttucumar, Postles,
83 Elmore, & Mottram, 2012). Hence potato varieties low in reducing sugars and cereal varieties
84 low in asparagine are sought. Storage may increase levels of reducing sugars in stored
85 potatoes, particularly under low-temperature conditions (Rak, Navarro, & Palta, 2013), while
86 levels of fertilisation, for example nitrogen and sulfur, may have effects on reducing sugars
87 and asparagine levels in both potatoes and cereals (Elmore, Mottram, Muttucumar, Dodson,
88 Parry, & Halford, 2007; Muttucumar et al., 2006; Muttucumar, Powers, Elmore, Mottram,
89 & Halford, 2013). It is clear, however, that little or no acrylamide will form in the absence of
90 asparagine, while other components of the food matrix, such as lipid-derived aldehydes
91 (Zamora & Hidalgo, 2008), amino acids such as serine and threonine (Shu, 1999), and other

92 carbonyl-containing molecules (Hamzalioglu & Gokmen, 2012; Zamora, Delgado, &
93 Hidalgo, 2011), can react with asparagine to form acrylamide.

94 *2.2 Process-based mitigation*

95 Initial mitigation methods involved the control of processing conditions; for instance,
96 lowering pH, reducing cooking temperature and shortening the processing time (Palazoğlu &
97 Gokmen, 2008). Although these methods achieved an effective reduction of acrylamide,
98 sensory properties of the food were compromised. As the Maillard reaction begins when food
99 is heated, the first option in this type of mitigation method is to lower the temperature and
100 time of heating. However, as the Maillard reaction is also responsible for generating desirable
101 taste and smell in cooked food, sensory properties become unacceptable when cooking
102 temperature is substantially reduced (Masi, Dinnella, Barnaba, Navarini, & Monteleone,
103 2013).

104 Besides temperature, another important parameter, pH, has also been studied. In a model
105 system, the acrylamide content will reach a maximum amount when the pH is around 8,
106 which is near to the pK_a value of asparagine and leads to an enhancement in the initial steps
107 of acrylamide formation (Rydberg, Eriksson, Tareke, Karlsson, Ehrenberg, & Tornqvist,
108 2003). Several authors have reduced acrylamide formation by reducing the pH, using
109 compounds such as citric acid (Gama-Baumgartner, Grob, & Biedermann, 2004), although
110 product quality has suffered generally using this approach (Vinci, Mestdagh, & De
111 Meulenaer, 2012).

112 *2.3 Use of additives*

113 Adding other chemicals prior to or after heating could also decrease the final acrylamide
114 amount in a cooked product. For example, adding glycine before heating will compete with
115 asparagine to lower the final acrylamide amount (Bråthen & Knutsen, 2005). It could also be

116 added after the Maillard reaction, to react with acrylamide directly, thus lowering its amount
117 in the final product (Liu, Man, Zhu, Hu, & Chen, 2013). However, this method also has
118 negative effects on the sensory properties of the product, as added glycine will react with
119 reducing sugars to increase levels of odour-active alkylpyrazines (Low, Parker, & Mottram,
120 2007)

121 The addition of divalent cations has also been shown to be an effective means of reducing
122 acrylamide. Nixtamalisation, the traditional cooking of corn grains in calcium hydroxide
123 solution prior to milling, is traditionally used in the preparation of tortillas (Salazar,
124 Arambula-Villa, Luna-Barcenas, Figueroa-Cardenas, Azuara, & Vazquez-Landaverde, 2014).
125 A calcium chloride dip for potatoes reduced acrylamide in French fries by 95%, with no
126 adverse effects on product quality reported. The effect is due to the divalent cations inhibiting
127 formation of the Schiff base (Gökmen & Şenyuva, 2007). Monovalent cations at low
128 concentrations may also exhibit a mitigation effect. Addition of 1–2% sodium chloride to a
129 bread mix led to a substantial reduction in acrylamide in baked rolls. Higher salt
130 concentrations inhibited yeast growth, resulting in increased acrylamide formation (Claus,
131 Mongili, Weisz, Schieber, & Carle, 2008). A 2% sodium chloride dip pre-treatment (60 min,
132 room temperature) resulted in a 78% reduction in acrylamide in fried potato discs (Sansano,
133 Juan-Borras, Escriche, Andres, & Heredia 2015).

134 Antioxidants could also be added to inhibit the formation of acrylamide. However, effects
135 were variable in the several studies carried out during the past decade, due to the various
136 types of antioxidants used (Jin, Wu, & Zhang, 2013). For instance, rosemary added to corn or
137 olive oil could effectively lower the amount of acrylamide in fried potato slices (Becalski,
138 Lau, Lewis, & Seaman, 2002), while some commonly used antioxidants, such as BHT,
139 sesamol and Vitamin E, had an enhanced effect on acrylamide formation in cooked meat
140 (Tareke, 2003). The reducing or promoting effects could be attributed to differences in

141 reaction conditions, antioxidant dosage and different reaction pathways.

142 *2.4 Enzymatic approaches for acrylamide reduction*

143 Fermentation methods use specific microorganisms to consume the asparagine or reducing
144 sugar before the food processing step (Sadd, Hamlet, & Liang, 2008). For instance, a starter
145 medium containing lactic acid bacteria was used in the preparation of wholemeal rye bread
146 and substantially reduced acrylamide levels in the final product (Bartkiene, Jakobsonė,
147 Juodeikiene, Vidmantienė, Pugajeva, & Bartkevics, 2013b). As well as lowering the pH, the
148 lactic acid bacteria reduced the levels of reducing sugars in the dough.

149 However, there are several points that need to be considered in a fermentation approach.
150 To begin with, the temperature and pH need to be controlled in order to maximise the activity
151 of the microorganism. Even if the reducing sugar consumed was added back after processing,
152 the sensory quality of the final product may still be affected by the fermentation step
153 (Bartkiene, Jakobsonė, Juodeikiene, Vidmantienė, Pugajeva, & Bartkevics, 2013a). Secondly,
154 fermentation predominantly works in bakery products, with limited application application in
155 potato-based products and coffee (Kamkar, Qajarbeygi, Jannat, Babaei, Misaghi, & Aghaei,
156 2015).

157 The use of an enzymatic approach to modify reaction pathways was first proposed by
158 Amrein et al. (2004), who used asparaginase to hydrolyse asparagine to aspartic acid and
159 ammonia (Ciesarová, Kiss, & Boegl, 2006). This approach is considered to be effective
160 because asparagine is not considered a major contributor to the overall flavour and colour of
161 cooked foods (Parker, Balagiannis, Higley, Smith, Wedzicha, & Mottram, 2012), so desirable
162 sensory properties are maintained.

163 3. Asparaginase

164 Asparaginase (L-asparagine amidohydrolases EC 3.5.1.1) is an enzyme widely distributed
165 in animals, plants and living organisms (Wriston, 1985). It has been shown that asparaginase
166 catalyses the hydrolysis of asparagine into aspartic acid and ammonia by hydrolysing the
167 amide group in the side chain of asparagine (Hendriksen, Kornbrust, Ostergaard, & Stringer,
168 2009). Aspartic acid will then enter the citric acid cycle, playing a vital role in amino acid
169 metabolism. Asparagine is responsible for nitrogen storage in most plants; therefore,
170 asparaginase plays an important role in energy utilisation (Sieciechowicz, Joy, & Ireland,
171 1988).

172 L-Asparaginase has been used as a therapeutic treatment for certain kinds of cancer, such
173 as leukaemia (Bushman, Palmieri, Whinna, & Church, 2000). In leukaemia sufferers
174 malignant cells depend more on exogenous asparagine and glutamine to survive than normal
175 cells. Asparaginase injected into the bloodstream hydrolyses free asparagine into aspartic acid
176 and ammonia, and glutamine to glutamic acid and ammonia (Friedman, 2003). In this way,
177 the growth of malignant cells is inhibited.

178 L-Asparaginase is an intracellular enzyme which is obtained from a variety of
179 microorganisms: *Escherichia coli*, *Erwinia carotovora*, *Bacillus* sp., *Enterobacter aerogenes*,
180 *Corynebacterium glutamicum*, *Pseudomonas stutzeri* and *Candida utilis* (Qin & Zhao, 2003).
181 For pharmaceutical uses, L-asparaginase is typically obtained from *Escherichia coli* (Pritsa &
182 Kyriakidis, 2001). However, the production of the enzyme is complex with low yield. Until
183 now, there is no medium that has been specifically established for the optimum production of
184 L-asparaginase from different microorganisms. To maximise enzyme production, each
185 organism has its own optimum conditions. Research on *E. coli* using response surface
186 methodology achieved a 10-fold enhancement in asparaginase production (Kenari,
187 Alemzadeh, & Maghsodi, 2011).

188 Most asparaginases are quite specific for asparagine. Optimal activity is usually achieved
189 at pH 5–7 and 37 °C. However, as glutamine has similar structure to asparagine, some
190 enzymes also have a low activity towards glutamine (Krasotkina, Borisova, Gervaziev, &
191 Sokolov, 2004). A small group of enzymes, called glutaminase-asparaginases, have activities
192 for both asparagine and glutamine but prefer glutamine as a substrate (Roberts, Holcenberg,
193 & Dolowy, 1972). Crystallographic study has shown that both types of asparaginase,
194 common asparaginase and glutaminase-asparaginase, have the same basic structure and
195 catalytic mechanism but differ in working conditions (pH and temperature) (Yao, Yasutake,
196 Morita, & Tanaka, 2005). Researchers believe that glutaminase activity caused by
197 glutaminase-asparaginase will exert serious adverse effects on human health, such as liver
198 dysfunction, pancreatitis and leucopenia (Mahajan, Saran, Kameswaran, Kumar, & Saxena,
199 2012). Therefore, this specific type of asparaginase should be strictly avoided in the food
200 industry.

201 Commercially, there are two asparaginase products currently available for acrylamide
202 mitigation in the food industry. These are PreventASETM from DSM (Heerlen, The
203 Netherlands) and Acrylaway[®] from Novozymes A/S (Bagsvaerd, Denmark). PreventASETM
204 was the first, launched in 2007. It was obtained after analysing the gene sequence of
205 *Aspergillus niger* and produced recombinant in the *Aspergillus niger* host. It has an acidic
206 profile (optimum pH 4–5, temperature 50 °C). Acrylaway[®] on the other hand, is obtained
207 from *Aspergillus oryzae* and has an almost neutral profile (optimum pH 7, temperature 37
208 °C).

209 Regarding safety, these enzymes are produced by specific fungal strains of *A. oryzae* and
210 *A. niger*, fungi that have been widely used in commercial products for several decades and
211 have been proved to be safe by JECFA (JECFA, 2007). Acrylaway[®] and PreventASETM have
212 shown high specificity and therefore minimum activity towards glutamine and other amino

213 acids. Ultimately, these enzymes will be deactivated during the heating process, ensuring
214 their safe application in foodstuffs (Hendriksen et al., 2009).

215 Asparaginase has received “generally recognized as safe” status from the US government.
216 It has also been given a favourable evaluation as a food additive by the Joint FAO/WHO
217 Expert committee (JEFCA, 2007) and it is currently used in several countries, including
218 United States, Australia, New Zealand, China, Russia, Mexico and several European
219 countries. As different dosages of asparaginase will be used in different types of food, there is
220 no unified standard for the maximum dosage.

221 3.1. Use of asparaginase in acrylamide mitigation

222 As shown in Table 1, over the last decade there have been numerous studies monitoring
223 the reduction of acrylamide formation by means of asparaginase treatment. The first study
224 was carried out by Zyzak et al. in 2003, immediately after the formation mechanism was
225 revealed. However, Zyzak’s research was focused on the formation mechanism rather than
226 the mitigation efficiency. He used commercial asparaginase from Aldrich (A2925 from
227 *Erwinia chrysanthemi*), 50 U added to 60 g of mashed potato slurry (15 g potato, 45 g water),
228 to hydrolyse the asparagine, in order to verify that asparagine is indeed the precursor of
229 acrylamide. The asparaginase achieved an 88% asparagine reduction that led to 99%
230 acrylamide reduction in a microwaved mashed potato snack, heated at full power until brown
231 (Zyzak et al., 2003).

232 The following year, the first paper on the use of asparaginase as an acrylamide mitigation
233 method was published. Asparaginase (from *E. coli*, 4 U/kg) added to gingerbread hydrolysed
234 approximately 75% of the free asparagine, leading to a 55% acrylamide reduction in the final
235 product. The acrylamide-reduced product was identical to a control product in both colour
236 and taste (Amrein, Schönbächler, Escher, & Amadò, 2004). Though this enzyme application

237 formed only a small part of the research, it stressed the advantage of the enzymatic method
238 on mitigating acrylamide while maintaining the organoleptic properties of the product.

239 Ciesarová et al. (2006) set up a model system to examine the importance of all the related
240 factors, such as temperature, dosage and application time. However, there were insufficient
241 time and temperature points studied to determine optimum activity. Applying asparaginase to
242 dried potato powder led to a 90% acrylamide reduction in cooked product. However, instead
243 of considering the effect of the cut and shape of the potato products, this research focused
244 more on potato varieties. Although this research showed great success in acrylamide
245 reduction, the agronomic factors were discussed more than the enzyme itself.

246 Pedreschi, Kaack, and Granby (2008) were the first to publish results using a commercial
247 asparaginase (Acrylaway®). They established that the optimum temperature and pH for this
248 enzyme were 60 °C and 7.0, respectively. A reduction of 67% in acrylamide was achieved in
249 French fries under these conditions. In this study, the importance of blanching and
250 temperature control of asparaginase treatment was highlighted. It is known that blanching
251 will change the microstructure of the potato strips and increase the contact probability of
252 asparaginase and asparagine (Lisińska, Tajner-Czopek, & Kalum, 2007), so blanching is
253 highly recommended for increasing the performance of the enzyme.

254 Another study by the same group focused on the combination of asparaginase
255 (Acrylaway®) and conventional blanching, alongside their individual usage. Blanching using
256 hot water at 85 °C to treat the potato tuber samples for 3.5 min was compared with enzymatic
257 mitigation using an asparaginase solution (10000 ASNU/L) at 50 °C for 20 min. One ASNU
258 is defined as the amount of asparaginase that produces one micromole of ammonia per
259 minute under standard conditions (pH 7; 37 °C). Experimental results showed that blanching
260 and enzyme treatments have a similar effect on acrylamide reduction (17%). By combining
261 the two methods, almost 90% of acrylamide was mitigated. The authors assumed that the

262 microstructure of the potato tissues was changed in the blanching process, causing the
263 asparagine in the cell to have a more effective interaction with the enzyme outside the cell
264 (Pedreschi, Mariotti, Granby, & Risum, 2011). Although acrylamide in this research was
265 significantly reduced, no sensory analysis of the product was performed.

266 In 2009, another study involving Acrylaway® was carried out on a much wider range of
267 foods, including gingerbread, crispbread, semi-sweet biscuits, French fries and crisps
268 (Hendriksen et al., 2009). Again the optimum conditions of temperature and pH were 60 °C
269 and 7.0. In this study, other factors were also taken into consideration, depending on the food
270 matrix. In semi-sweet biscuits, the dosage was the variable and the temperature was set at 40
271 °C. Asparaginase treatment took place at the dough resting time before the biscuits were
272 baked at 260 °C for 5.5 min (Fig. 2a). For the crispbread trial, the temperature was held at 10,
273 15, or 20 °C for 30 or 60 min and the dosage was set at 2100 ASNU/kg of flour. Then the
274 crispbread was baked at 250 °C for 11 min (Fig. 2b). By changing the dosage of enzyme and
275 time, the influences of each factor are revealed. Besides dosage and temperature, water
276 content is another important factor in a cereal-based product like gingerbread, as higher water
277 activity will provide sufficient contact for the enzyme with the substrate. Therefore, in order
278 not to compromise final product sensory quality, a higher water content is recommended,
279 although a further drying step may be needed subsequently. For potato-based products, a
280 reduction was achieved in both French fries and crisps. Potatoes made into French fries were
281 treated with 10500 ASNU/L and then fried for 3 min at 175 °C. Sliced potatoes used to make
282 potato crisps were treated with various concentrations of enzyme for 15 min at 40 °C. Then
283 frying was conducted for 2.5 min at 180 °C. In the French fries experiment, the authors
284 prepared a sample set with a one-minute dip and 20 minutes soaking in the enzyme bath.
285 Although the acrylamide reduction in the one-minute dip (59% maximum) was less than for
286 the samples in the 20-min soak (85% maximum), the results were still meaningful for the

287 practical continuous process. Results indicated a broad range of enzyme applications. The
288 authors suggested that by combining modified processing conditions with an enzymatic
289 approach, acrylamide could be mitigated at a fairly low cost. The key point in this research is
290 that the authors tried to assess all related variables and generate specific solutions for each
291 type of product from an enormous data-set.

292 Also in 2009, Kukurová, Morales, Bednáriková, and Ciesarová (2009) used two levels of
293 Acrylaway® (100 U/kg and 500 U/kg flour) in the preparation of fried bread rolls. The
294 asparaginase treatment (15 min, 37 °C) removed at least 96% of the asparagine from the
295 dough; acrylamide could not be quantified in the fried rolls with Acrylaway® added, while
296 levels of 215 µg per kg were present in control rolls fried at 200 °C for 8 min. The same
297 group also studied acrylamide formation in cookies treated with Acrylaway® (500 U/kg
298 flour) and different raising agents (Kukurová, Ciesarová, Mogol, Açar, & Gökmen, 2013).
299 The raising agents increased the pH of the dough, reducing the effectiveness of the
300 asparaginase in reducing asparagine. When applied for less than 30 minutes, the asparagine
301 had no effect on the sensory properties of the cookies.

302 Hendriksen, Budolfson, and Baumann (2013) used Acrylaway® to study potato and cereal
303 products and also carried out the first experiments on the effect of asparaginase on
304 acrylamide in coffee. For cereal products, the mitigation efficiency reached 95% in lebkuchen
305 and 90% in tortilla chips, whereas in coffee a 70–80% reduction was achieved. In potato
306 products, experiments on potato tuber snack pellets and French fries were carried out. In
307 potato tuber snack pellets, the enzyme was added directly to a dough based on 29% potato
308 starch, 27.6% potato granules, 15% potato flakes, 1.4% salt and 27% water, while in the
309 French fries test, Bintje or Maris Piper potatoes were manually peeled and cut into 8 mm × 8
310 mm strips. Two innovative points of this study stands out. First, asparaginase is added at the
311 disodium acid pyrophosphate (SAPP) dipping stage rather than as a separate step. SAPP is

312 commonly used in potato processing to prevent after-cooking darkening. Integrating the two
313 treatments could reduce time and cost. Secondly, an industrial scale trial was carried out to
314 test the mitigation efficiency of asparaginase in continuous processing. Such trials will push
315 the industrial application of asparaginase forward. Reductions in potato product were
316 comparatively low, due to the insufficient contact of asparagine and enzyme. However, the
317 industrial scale experiment (8 tonne/h) still achieved satisfactory results; a 43% reduction in
318 10 mm × 10 mm and 53% reduction in 7 mm × 7 mm potato pieces was achieved. A dye-
319 based experiment indicated that asparaginase could only penetrate 1 mm into the potato,
320 again highlighting the importance of incorporating a blanching step when treating potatoes
321 (Hendriksen et al., 2013).

322 For coffee, increased acrylamide mitigation could be achieved by incubating the wetted
323 green beans. Typically, green coffee beans are steamed to decrease the caffeine content. The
324 decaffeination process is usually carried out by a water or solvent partition system. Firstly,
325 green coffee beans are steamed to make the caffeine available. Then, a solvent is used to
326 extract the caffeine. Finally, the green beans are steamed again to remove any residual
327 solvent (Spiller, 1997). Hence, asparaginase could be infused during these steps with minor
328 changes to the processing conditions. A laboratory-scale experiment indicated that a low
329 dosage (2000–6000 ASNU) of asparaginase could achieve 55–74% acrylamide reduction in
330 coffee beans (Hendriksen et al., 2013), while work in our laboratory showed that both the
331 steaming step and the asparaginase treatment caused a reduction in free asparagine when the
332 coffee was roasted, which was reflected in acrylamide losses from 69% to 86% using dosages
333 of 2600 to 20000 ASNU, respectively (Xu, Khalid, Oruna-Concha, & Elmore, 2015).

334 The effect of asparaginase on acrylamide mitigation in biscuits has also been examined by
335 Anese, Quarta, and Frias (2011). The authors used asparaginase levels from 100 to 900
336 ASNU (Acrylaway®) with 20–54 °C incubation temperature and 10–30 min incubation time,

337 15 treatments in all. By analysing the results from each treatment, the influence of each factor
338 was considered and optimum conditions could be obtained. This study not only demonstrated
339 a method that could assess the effect of the enzyme but compared the effect obtained with the
340 cost for each treatment. This paper also contained valuable advice on the practical application
341 of asparaginase. For instance, acrylamide development was at a minimum at intermediate
342 asparaginase concentrations and increased asparaginase addition did not significantly affect
343 the colour of the final product.

344 We are only aware of one publication where PreventASe™ was used as the asparaginase
345 source. A solution of 500 ASNU in 10 mL of water was spread onto the surface of a
346 wheat/oat bread loaf prior to baking. The enzyme was effective during a proofing step of 15
347 min at 32 °C. This treatment led to a 46% reduction in the acrylamide content of the baked
348 bread crust (Ciesarová et al., 2014).

349 3.2. *New sources of asparaginase*

350 Recent papers have identified new sources of asparaginase for acrylamide reduction.
351 Tuncel, Yılmaz, and Şener (2010) used asparaginase from a vegetable source. They
352 germinated pea flour to enhance asparaginase activity and remove beany flavour. The pea
353 flour was finely ground and then added to wheat flour in white wheat bread, wheat bran bread
354 and wholegrain white bread at three different levels (1%, 3% and 5%). The bread was baked
355 at 220 °C for 22–25 min. In white wheat bread acrylamide reduction was less than 10% in all
356 cases, while addition of 5% pea flour to bran bread and grain bread reduced acrylamide levels
357 in crust by 57% and 68%, respectively. Although the sensory panel showed detectable
358 differences in the final products, there was no significant negative impact on sensory
359 properties. The extraction of asparaginase from fungus is relatively costly; therefore this
360 approach provides an effective alternative means to produce asparaginase.

361 Asparaginase with low glutaminase activity was successfully extracted from *Bacillus*
362 *licheniformis* and was used to reduce acrylamide in fried potato strips by up to 80%
363 (Mahajan, Saran, Kameswaran, Kumar, & Saxena, 2012).

364 Asparaginase produced from *Cladosporium* sp. was used in the crumb and crust of sweet
365 bread (Kumar, Shimray, Indrani, & Manonmani, 2014). The dosage used varied from 50–300
366 U. However, the units were not defined. Reductions of 97% and 73% were achieved at 300 U
367 in bread crust and crumb, respectively. The authors also measured the formation of 5-
368 (hydroxymethyl)furfural, which is another potential toxicant formed in the Maillard reaction,
369 and it was also decreased. This research also showed a new possible source for asparaginase,
370 although yield data were not disclosed.

371 Asparaginase has also been extracted from *Rhizomucor miehei* (Huang, Liu, Sun, Yan, &
372 Jiang, 2014). The extracted asparaginase, designated as RmAsnase, was optimally active at
373 pH 7.0 and 45 °C and was stable at this temperature for 30 min. RmAsnase was cloned and
374 expressed in *Escherichia coli* and proved highly specific towards asparagine. The researchers
375 demonstrated that a low concentration of asparaginase (0.5 U/g flour) had much better
376 mitigation efficiency in bread (40%) than biscuits (15%). However, an 80% reduction was
377 achieved in both products at 10 U/g flour. Overall, this new enzyme showed remarkable
378 potential both as an acrylamide mitigator and also in leukaemia therapy.

379 Asparaginase from food-grade *Bacillus subtilis* was applied to potato chips (Onishi,
380 Prihanto, Yano, Takagi, Umekawa, & Wakayama, 2015). Compared to a control sample, 40
381 U of asparaginase led to 80% reduction in acrylamide. One unit of the enzyme was defined as
382 the amount that catalysed the formation of 1 μ mol ammonia per min; additional details were
383 not provided by the authors. They suggested that BAsnase, as the enzyme was christened,
384 could be used to spray potatoes prior to cooking at home. Similar reductions in acrylamide in
385 French fries were also obtained using an asparaginase from *Thermococcus zilligii*. The

386 purified enzyme displayed a maximum activity at pH 8.5 and 90 °C (Zuo, Zhang, Jiang, &
387 Mu, 2015b) and retained 70% of its original activity after 2 hours incubation at 85 °C.
388 Another asparaginase with high temperature stability was recently isolated from *Pyrococcus*
389 *furiosus* (Kundu, Bansal, & Mishra, 2013). In addition, at the end of 2013 Novozymes
390 launched Acrylaway® HighT, which is specifically designed for higher temperature
391 processing ([http://www.novozymes.com/en/news/news-archive/Pages/New-Novozymes?-
392 solution-enables-acrylamide-mitigation-in-even-more-product-categories--.aspx](http://www.novozymes.com/en/news/news-archive/Pages/New-Novozymes?-solution-enables-acrylamide-mitigation-in-even-more-product-categories--.aspx).) Enzymes
393 stable at such temperatures can be incorporated into the blanching step of a commercial
394 process, which would increase their applicability.

395 Immobilised asparaginase may not convert as much asparagine as free asparaginase but its
396 stability is improved, meaning that blanching water containing immobilised enzyme can be
397 re-used several times without loss of asparaginase activity. The asparaginase is immobilised
398 by crosslinking with glutaraldehyde on an inert silica-based carrier (Hendriksen, Puder, &
399 Olsen, 2014).

400 3.3. *Asparaginase activity under different conditions*

401 To quantify the effect of asparaginase in food applications, its activity needs to be
402 determined. For the two commercially available enzymes (Acrylaway® by Novozymes and
403 PreventASe™ by DSM), two different methods to determine the activity of the enzyme have
404 been used. Both methods are based on measuring the ammonia that is generated from the
405 asparagine hydrolysis. However, in the method used to measure the activity of Acrylaway,
406 ammonia subsequently reacts with α -ketoglutarate to form L-glutamic acid. The reaction is
407 catalysed by glutamate dehydrogenase in the presence of NADH, which is oxidised to NAD⁺
408 with the concomitant loss of absorbance measured at 340 nm. The asparaginase activity is
409 measured as the rate of NADH consumption under standard conditions (pH = 7; 37 °C). The

410 activity of asparaginase is expressed in ASNU activity units. One ASNU is defined as the
411 amount of asparaginase that produces one micromole of ammonia per minute under standard
412 conditions (Hendriksen et al., 2009). The activity of PreventASe™ is measured by a different
413 method. The liberated ammonia subsequently reacts with phenol nitroprusside and alkaline
414 hypochlorite resulting in a blue colour. This is known as the Berthelot reaction (Rhine, Sims,
415 Mulvaney, & Pratt, 1998). The activity of asparaginase is determined by measuring the
416 absorbance of the reaction mixture at 600 nm. Asparaginase activity is expressed in ASPU
417 activity units. One ASPU is defined as the amount of asparaginase that liberates one
418 micromole of ammonia from L-asparagine per minute under standard conditions (pH = 5.0;
419 37 °C).

420 By setting up a standard activity determination method, the activity of the enzyme under
421 different conditions of pH and temperature can be measured. From the work done by
422 Hendriksen's group in Novozymes, it was shown that Acrylaway® has almost two times the
423 activity at 60 °C compared to its activity at 37 °C. Also, the activity of the enzyme at pH 7 is
424 almost two times its activity at pH 5. These differences in activity should be considered prior
425 to the application of the enzyme to a substrate.

426 3.4. *Practicalities of asparaginase application*

427 The practicalities of asparaginase application can be assessed from three different aspects:
428 raw material composition, processing and commerce.

429 The amount of asparagine in a foodstuff should be considered when deciding the dosage
430 of asparaginase. For example, concentrations of free asparagine in potato may vary by a
431 factor of 10 or more, and can be affected by variety and growing conditions, (Halford et al.,
432 2012b).

433 When using asparaginase in food manufacture, factors like temperature, time and substrate

434 ratio are of importance. For example, the optimal temperature for Acrylaway® is around 60
435 °C and its activity will decrease significantly above this temperature (Hendriksen et al.,
436 2009). Hence asparaginase will be denatured and inactivated during food processing. The
437 dwell time for enzyme and foodstuff before heat treatment should be optimised before
438 application. Enzyme–substrate ratio is also an important factor and dosage of the enzyme
439 should be determined, so that maximum mitigation is achieved with minimum enzyme
440 concentration. Extra water may be used to ensure the delivery of the enzyme (Whitehurst &
441 Van Oort, 2009).

442 Even though asparaginase has advantages over other mitigation methods, its use by
443 manufacturers may not be commercially viable at present. There are also issues with the
444 industrial application of asparaginase in a continuous process, which can achieve good results
445 in a relatively short time. Pilot-scale experiments have been carried out using a continuous
446 process for French fries production (Hendriksen et al., 2009). High levels of acrylamide
447 reduction were achieved (60–85% in French Fries and 60% in potato chips). However, more
448 research is needed to better incorporate asparaginase usage into industrial-scale food
449 production. To maximise the overall effect of the enzymatic method, pre- and post-treatment
450 procedures may need to be adapted; for instance, reduction of starting material dimensions
451 and blanching before enzyme treatment and modification of the process conditions after the
452 enzyme treatment by, for example, changing cooking temperature and pH.

453

454 **4. Conclusions**

455 Asparaginase has become a powerful tool for acrylamide mitigation in the food industry.
456 With the success of commercial products, it is likely that asparaginase will be used more and
457 more. The first commercially available “acrylamide-free” product, biscuits treated with

458 Preventase™, was announced to launch shortly in Germany for Christmas, 2008. This
459 information was released by DSM Food Specialities, although the manufacturer's name was
460 not disclosed (Foodingredients1st, 2008). However, there was no more news subsequently.

461 The potential adverse effects of asparaginase treatment on sensory properties of cooked
462 foods and the need to achieve sufficient enzyme–substrate contact are areas for future
463 research. However, if the application of asparaginase becomes commercially attractive, its
464 use alongside raw materials low in asparagine may provide the solution to the acrylamide
465 problem.

466

467 **5. References**

- 468 Amrein, T. M., Schönbacher, B., Escher, F., & Amadó, R. (2004). Acrylamide in
469 gingerbread: Critical factors for formation and possible ways for reduction. *Journal of*
470 *Agricultural and Food Chemistry*, 52(13), 4282-4288.
- 471 Anese, M., Quarta, B., & Frias, J. (2011a). Modelling the effect of asparaginase in reducing
472 acrylamide formation in biscuits. *Food Chemistry*, 126(2), 435-440.
- 473 Anese, M., Quarta, B., Peloux, L., & Calligaris, S. (2011b). Effect of formulation on the
474 capacity of l-asparaginase to minimize acrylamide formation in short dough biscuits.
475 *Food Research International*, 44(9), 2837-2842.
- 476 Banhero, M., Pellegrino, G., & Manna, L. (2013). Supercritical fluid extraction as a
477 potential mitigation strategy for the reduction of acrylamide level in coffee. *Journal of*
478 *Food Engineering*, 115(3), 292-297.
- 479 Bartkiene, E., Jakobsone, I., Juodeikiene, G., Vidmantiene, D., Pugajeva, I., & Bartkevics, V.
480 (2013a). Effect of lactic acid fermentation of lupine wholemeal on acrylamide content
481 and quality characteristics of wheat-lupine bread. *International Journal of Food*
482 *Sciences and Nutrition*, 64(7), 890-896.
- 483 Bartkiene, E., Jakobsone, I., Juodeikiene, G., Vidmantiene, D., Pugajeva, I., & Bartkevics, V.
484 (2013b). Study on the reduction of acrylamide in mixed rye bread by fermentation
485 with bacteriocin-like inhibitory substances producing lactic acid bacteria in
486 combination with *Aspergillus niger* glucoamylase. *Food Control*, 30(1), 35-40.
- 487 Becalski, A., Lau, B. P. Y., Lewis, D., & Seaman, S. W. (2002). Acrylamide in Foods:
488 Occurrence, Sources, and Modeling. *Journal of Agricultural and Food Chemistry*,
489 51(3), 802-808.
- 490 Bråthen, E., & Knutsen, S. H. (2005). Effect of temperature and time on the formation of
491 acrylamide in starch-based and cereal model systems, flat breads and bread. *Food*

492 *Chemistry*, 92(4), 693-700.

493 Bushman, J. E., Palmieri, D., Whinna, H. C., & Church, F. C. (2000). Insight into the
494 mechanism of asparaginase-induced depletion of antithrombin III in treatment of
495 childhood acute lymphoblastic leukemia. *Leukemia Research*, 24(7), 559-565.

496 Ciesarová, Z., Kiss, E., & Boegl, P. (2006). Impact of L-asparaginase on acrylamide content
497 in potato products. *Journal of Food and Nutrition Research*, 45(4), 141-146.

498 Ciesarová, Z., Kukurová, K., Mikušová, L., Basil, E., Polakovičová, P., Duchoňová, L.,
499 Vlček, M., & Šturdík, E. (2014). Nutritionally enhanced wheat-oat bread with reduced
500 acrylamide level. *Quality Assurance and Safety of Crops & Foods*, 6(3), 327-334.

501 Claus, A., Mongili, M., Weisz, G., Schieber, A., & Carle, R. (2008). Impact of formulation
502 and technological factors on the acrylamide content of wheat bread and bread rolls.
503 *Journal of Cereal Science*, 47(3), 546-554.

504 Curtis, T. Y., Powers, S. J., Balagiannis, D., Elmore, J. S., Mottram, D. S., Parry, M. A. J.,
505 Rakszegi, M., Bedö, Z., Shewry, P. R., & Halford, N. G. (2010). Free amino acids and
506 sugars in rye grain: Implications for acrylamide formation. *Journal of Agricultural
507 and Food Chemistry*, 58, 1959-1969.

508 Elmore, J. S., Briddon, A., Dodson, A. T., Muttucumaru, N., Halford, N. G., & Mottram, D.
509 S. (2015). Acrylamide in potato crisps prepared from 20 UK-grown varieties: Effects
510 of variety and tuber storage time. *Food Chemistry*, 182, 1-8.

511 Elmore, J. S., Mottram, D. S., Muttucumaru, N., Dodson, A. T., Parry, M. A. J., & Halford, N.
512 G. (2007). Changes in free amino acids and sugars in potatoes due to sulfate
513 fertilization and the effect on acrylamide formation. *Journal of Agricultural and Food
514 Chemistry*, 55(13), 5363-5366.

515 Foodingredients1st. (2008). First "Acrylamide-Free" Biscuits Will be Launched in Germany.
516 In).<http://www.foodingredientsfirst.com/search-results/First-Acrylamide-Free->

517 Biscuits-Will-Be-Launched-in-Germany.html

518 Friedman, M. (2003). Chemistry, biochemistry, and safety of acrylamide. A review. *Journal*
519 *of Agricultural and Food Chemistry*, 51(16), 4504-4526.

520 Friedman, M. (2015). Acrylamide: inhibition of formation in processed food and mitigation
521 of toxicity in cells, animals, and humans. *Food & Function*, 6(6), 1752-1772.

522 Gama-Baumgartner, F., Grob, K., & Biedermann, M. (2004). Citric acid to reduce acrylamide
523 formation in French fries and roasted potatoes. *Mitteilungen aus*
524 *Lebensmitteluntersuchung und Hygiene*, 95(1), 110-117.

525 Gökmen, V., & Şenyuva, H. Z. (2007). Acrylamide formation is prevented by divalent cations
526 during the Maillard reaction. *Food Chemistry*, 103(1), 196-203.

527 Granvogl, M., & Schieberle, P. (2006). Thermally generated 3-aminopropionamide as a
528 transient intermediate in the formation of acrylamide. *Journal of Agricultural and*
529 *Food Chemistry*, 54(16), 5933-5938.

530 Halford, N. G., Curtis, T. Y., Muttucumaru, N., Postles, J., Elmore, J. S., & Mottram, D. S.
531 (2012a). The acrylamide problem: a plant and agronomic science issue. *Journal of*
532 *Experimental Botany*, 63(8), 2841-2851.

533 Halford, N. G., Muttucumaru, N., Powers, S. J., Gillatt, P. N., Hartley, L., Elmore, J. S., &
534 Mottram, D. S. (2012b). Concentrations of free amino acids and sugars in nine potato
535 varieties: effects of storage and relationship with acrylamide formation. *Journal of*
536 *Agricultural and Food Chemistry*, 60(48), 12044-12055.

537 Hamzalioglu, A., & Gökmen, V. (2012). Role of bioactive carbonyl compounds on the
538 conversion of asparagine into acrylamide during heating. *European Food Research*
539 *and Technology*, 235(6), 1093-1099.

540 Hedegaard, R. V., Frandsen, H., & Skibsted, L. H. (2008). Kinetics of formation of
541 acrylamide and Schiff base intermediates from asparagine and glucose. *Food*

542 *Chemistry*, 108(3), 917-925.

543 Hendriksen, H. V., Kornbrust, B. A., Ostergaard, P. R., & Stringer, M. A. (2009). Evaluating
544 the potential for enzymatic acrylamide mitigation in a range of food products using an
545 asparaginase from *Aspergillus oryzae*. *Journal of Agricultural and Food Chemistry*,
546 57(10), 4168-4176.

547 Hendriksen, H. V., Budolfson, G., & Baumann, M. J. (2013). Asparaginase for acrylamide
548 mitigation in food. *Aspects of Applied Biology*, 116, 41–50.

549 Hendriksen, H. V.; Puder, K.; Olsen, A. G. Method for reducing the level of asparagine in a
550 food material. International Patent No. WO/2014/161935, October 9th, 2014.

551 [http://www.novozymes.com/en/news/news-archive/Pages/New-Novozymes'-solution-
552 enables-acrylamide-mitigation-in-even-more-product-categories--.aspx](http://www.novozymes.com/en/news/news-archive/Pages/New-Novozymes'-solution-
552 enables-acrylamide-mitigation-in-even-more-product-categories--.aspx). Accessed 6th
553 April, 2016.

554 Huang, L., Liu, Y., Sun, Y., Yan, Q., & Jiang, Z. (2014). Biochemical characterization of a
555 novel L-asparaginase with low glutaminase activity from *Rhizomucor miehei* and Its
556 application in food safety and leukemia treatment. *Applied and Environmental*
557 *Microbiology*, 80(5), 1561-1569.

558 JECFA (2007). Compendium of Food Additive Specifications. Monograph No. 4. Rome:
559 FAO.

560 Jin, C., Wu, X., & Zhang, Y. (2013). Relationship between antioxidants and acrylamide
561 formation: A review. *Food Research International*, 51(2), 611-620.

562 Kamkar, A., Qajarbeygi, P., Jannat, B., Babaei, A. H. H., Misaghi, A., & Aghae, E. M.
563 (2015). The inhibitory role of autolysed yeast of *Saccharomyces cerevisiae*, vitamins
564 B-3 and B-6 on acrylamide formation in potato chips. *Toxin Reviews*, 34(1), 1-5.

565 Kenari, S. L. D., Alemzadeh, I., & Maghsodi, V. (2011). Production of L-asparaginase from
566 *Escherichia coli* ATCC 11303: Optimization by response surface methodology. *Food*

567 *and Bioproducts Processing*, 89(4), 315-321.

568 Krasotkina, J., Borisova, A. A., Gervaziev, Y. V., & Sokolov, N. N. (2004). One-step
569 purification and kinetic properties of the recombinant L-asparaginase from *Erwinia*
570 *carotovora*. *Biotechnology and Applied Biochemistry*, 39, 215-221.

571 Kukurová, K., Morales, F. J., Bednáriková, A., & Ciesarová, Z. (2009). Effect of L-
572 asparaginase on acrylamide mitigation in a fried-dough pastry model. *Molecular*
573 *Nutrition & Food Research*, 53(12), 1532-1539.

574 Kukurová, K., Ciesarová, Z., Mogol, B. A., Açar, O. C., & Gökmen, V. (2013). Raising
575 agents strongly influence acrylamide and HMF formation in cookies and conditions
576 for asparaginase activity in dough. *European Food Research and Technology*, 237(1),
577 1-8.

578 Kumar, N. S. M., Shimray, C. A., Indrani, D., & Manonmani, H. K. (2014). Reduction of
579 acrylamide formation in sweet bread with L-asparaginase treatment. *Food and*
580 *Bioprocess Technology*, 7(3), 741-748.

581 Kundu, B.; Bansal, S.; Mishra, P. Mutants of L-asparaginase. US Patent No.20130330316,
582 12th December, 2013.

583 Lisińska, G., Tajner-Czopek, A., & Kalum, L. (2007). The effects of enzymes on fat content
584 and texture of French fries. *Food Chemistry*, 102(4), 1055-1060.

585 Liu, J., Man, Y., Zhu, Y., Hu, X., & Chen, F. (2013). Simultaneous analysis of acrylamide and
586 its key precursors, intermediates, and products in model systems by liquid
587 chromatography-triple quadrupole mass spectrometry. *Analytical Chemistry*, 85(19),
588 9262-9271.

589 Low, M. Y., Parker, J. K., & Mottram, D. S. (2007). Mechanisms of alkylpyrazine formation
590 in a potato model system containing added glycine. *Journal of Agricultural and Food*
591 *Chemistry*, 55(10), 4087-4094.

592 Mahajan, R. V., Saran, S., Kameswaran, K., Kumar, V., & Saxena, R. K. (2012). Efficient
593 production of L-asparaginase from *Bacillus licheniformis* with low-glutaminase
594 activity: optimization, scale up and acrylamide degradation studies. *Bioresource*
595 *Technology*, 125, 11-16.

596 Masi, C., Dinnella, C., Barnaba, M., Navarini, L., & Monteleone, E. (2013). Sensory
597 properties of under-roasted coffee beverages. *Journal of Food Science*, 78(8), S1290-
598 1300.

599 Mottram, D. S., Wedzicha, B. L., & Dodson, A. T. (2002). Acrylamide is formed in the
600 Maillard reaction. *Nature*, 419(6906), 448-449.

601 Muttucumar, N., Halford, N. G., Elmore, J. S., Dodson, A. T., Parry, M., Shewry, P. R., &
602 Mottram, D. S. (2006). Formation of high levels of acrylamide during the processing
603 of flour derived from sulfate-deprived wheat. *Journal of Agricultural and Food*
604 *Chemistry*, 54(23), 8951-8955.

605 Muttucumar, N., Powers, S. J., Elmore, J. S., Mottram, D. S., & Halford, N. G. (2013).
606 Effects of nitrogen and sulfur fertilization on free amino acids, sugars, and
607 acrylamide-forming potential in potato. *Journal of Agricultural and Food Chemistry*,
608 61(27), 6734-6742.

609 Ohara-Takada, A., Matsuura-Endo, C., Chuda, Y., Ono, H., Yada, H., Yoshida, M.,
610 Kobayashi, A., Tsuda, S., Takigawa, S., Noda, T., Yamauchi, H., & Mori, M. (2005).
611 Change in content of sugars and free amino acids in potato tubers under short-term
612 storage at low temperature and the effect on acrylamide level after frying. *Bioscience*,
613 *Biotechnology and Biochemistry*, 69(7), 1232-1238.

614 Onishi, Y., Prihanto, A. A., Yano, S., Takagi, K., Umekawa, M., & Wakayama, M. (2015).
615 Effective treatment for suppression of acrylamide formation in fried potato chips
616 using L-asparaginase from *Bacillus subtilis*. *3 Biotech*, 5(5), 783-789.

617 Palazoğlu, T. K., & Gökmen, V. (2008). Reduction of acrylamide level in French fries by
618 employing a temperature program during frying. *Journal of Agricultural and Food*
619 *Chemistry*, 56(15), 6162-6166.

620 Parker, J. K., Balagiannis, D. P., Higley, J., Smith, G., Wedzicha, B. L., & Mottram, D. S.
621 (2012). Kinetic model for the formation of acrylamide during the finish-frying of
622 commercial French fries. *Journal of Agricultural and Food Chemistry*, 60(36), 9321-
623 9331.

624 Pedreschi, F., Kaack, K., & Granby, K. (2008). The effect of asparaginase on acrylamide
625 formation in French fries. *Food Chemistry*, 109(2), 386-392.

626 Pedreschi, F., Mariotti, M. S., & Granby, K. (2014). Current issues in dietary acrylamide:
627 formation, mitigation and risk assessment. *Journal of the Science of Food and*
628 *Agriculture*, 94(1), 9-20.

629 Pedreschi, F., Mariotti, S., Granby, K., & Risum, J. (2011). Acrylamide reduction in potato
630 chips by using commercial asparaginase in combination with conventional blanching.
631 *LWT-Food Science and Technology*, 44(6), 1473-1476.

632 Pritsa, A. A., & Kyriakidis, D. A. (2001). L-Asparaginase of *Thermus thermophilus*:
633 Purification, properties and identification of essential amino acids for its catalytic
634 activity. *Molecular and Cellular Biochemistry*, 216(1-2), 93-101.

635 Qin, M., & Zhao, F. S. (2003). L-asparaginase release from *Escherichia coli* cells with
636 aqueous two-phase micellar systems. *Applied Biochemistry and Biotechnology*,
637 110(1), 11-21.

638 Rak, K., Navarro, F. M., & Palta, J. P. (2013). Genotype x storage environment interaction
639 and stability of potato chip color: Implications in breeding for cold storage chip
640 quality. *Crop Science*, 53(5), 1944-1952.

641 Rhine, E. D., Sims, G. K., Mulvaney, R. L., & Pratt, E. J. (1998). Improving the Berthelot

642 reaction for determining ammonium in soil extracts and water. *Soil Science Society of*
643 *America Journal*, 62(2), 473-480.

644 Roberts, J., Holcenberg, J. S., & Dolowy, W. C. (1972). Isolation, crystallization, and
645 properties of *Achromobacteraceae* glutaminase-asparaginase with antitumor activity.
646 *Journal of Biological Chemistry*, 247(1), 84-90.

647 Rydberg, P., Eriksson, S., Tareke, E., Karlsson, P., Ehrenberg, L., & Tornqvist, M. (2003).
648 Investigations of factors that influence the acrylamide content of heated foodstuffs.
649 *Journal of Agricultural and Food Chemistry*, 51(24), 7012-7018.

650 Sadd, P. A., Hamlet, C. G., & Liang, L. (2008). Effectiveness of methods for reducing
651 acrylamide in bakery products. *Journal of Agricultural and Food Chemistry*, 56(15),
652 6154-6161.

653 Salazar, R., Arambula-Villa, G., Luna-Barcenas, G., Figueroa-Cardenas, J. D., Azuara, E., &
654 Vazquez-Landaverde, P. A. (2014). Effect of added calcium hydroxide during corn
655 nixtamalization on acrylamide content in tortilla chips. *LWT-Food Science and*
656 *Technology*, 56(1), 87-92.

657 Sansano, M., Juan-Borras, M., Escriche, I., Andres, A., & Heredia, A. (2015). Effect of
658 pretreatments and air-frying, a novel technology, on acrylamide generation in fried
659 potatoes. *Journal of Food Science*, 80(5), T1120-T1128.

660 Shu, C.-K. (1999). Pyrazine formation from serine and threonine. *Journal of Agricultural and*
661 *Food Chemistry*, 47, 4332-4335.

662 Sieciechowicz, K. A., Joy, K. W., & Ireland, R. J. (1988). The metabolism of asparagine in
663 plants. *Phytochemistry*, 27(3), 663-671.

664 Spiller, G. A. (1997). *Caffeine*: London: Taylor & Francis.

665 Stadler, R. H., Blank, I., Varga, N., Robert, F., Hau, J., Guy, P. A., Robert, M.-C., & Riediker,
666 S. (2002). Acrylamide from Maillard reaction products. *Nature*, 419, 449-450.

667 Tareke, E. (2003). *Identification and Origin of Potential Background Carcinogens:*
668 *Endogenous Isoprene and Oxiranes, Dietary Acrylamide*. Ph.D. thesis, Department of
669 Environmental Chemistry, University of Stockholm.

670 Tareke, E., Rydberg, P., Karlsson, P., Eriksson, S., & Tornqvist, M. (2002). Analysis of
671 acrylamide, a carcinogen formed in heated foodstuffs. *Journal of Agricultural and*
672 *Food Chemistry*, 50(17), 4998-5006.

673 Tuncel, N. B., Yilmaz, N., & Şener , E. (2010). The effect of pea (*Pisum sativum* L.)-
674 originated asparaginase on acrylamide formation in certain bread types. *International*
675 *Journal of Food Science and Technology*, 45(12), 2470-2476.

676 Vinci, R. M., Mestdagh, F., & De Meulenaer, B. (2012). Acrylamide formation in fried potato
677 products – present and future, a critical review on mitigation strategies. *Food*
678 *Chemistry*, 133(4), 1138-1154.

679 Whitehurst, R. J., & Van Oort, M. (2009). *Enzymes in Food Technology*: Oxford, UK: Wiley-
680 Blackwell

681 Wriston, J. C. (1985). Asparaginase. *Methods in Enzymology*, 113, 608-618.

682 Xu, F.; Khalid, P.; Oruna-Concha, M.-J.; Elmore, J. S. (2015). Effect of asparaginase on
683 flavour formation in roasted coffee. In: A.J. Taylor, & D. S. Mottram (Eds.), *Flavour*
684 *Science* (563–566). Packington, United Kingdom: Context Products Ltd.

685 Yao, M., Yasutake, Y., Morita, H., & Tanaka, I. (2005). Structure of the type I L-asparaginase
686 from the hyperthermophilic archaeon *Pyrococcus horikoshii* at 2.16 angstrom
687 resolution. *Acta Crystallographica Section D-Biological Crystallography*, 61, 294-
688 301.

689 Zamora, R., Delgado, R. M., & Hidalgo, F. J. (2011). Strecker aldehydes and alpha-keto
690 acids, produced by carbonyl-amine reactions, contribute to the formation of
691 acrylamide. *Food Chemistry*, 128(2), 465-470.

692 Zamora, R., & Hidalgo, F. J. (2008). Contribution of lipid oxidation products to acrylamide
693 formation in model systems. *Journal of Agricultural and Food Chemistry*, 56(15),
694 6075-6080.

695 Zuo, S., Zhang, T., Jiang, B., & Mu, W. (2015a). Recent research progress on microbial L-
696 asparaginases. *Applied Microbiology and Biotechnology*, 99, 1069-1079.

697 Zuo, S., Zhang, T., Jiang, B., & Mu, W. (2015b). Reduction of acrylamide level through
698 blanching with treatment by an extremely thermostable L-asparaginase during French
699 fries processing. *Extremophiles*, 19(4), 841-851.

700 Zyzak, D. V., Sanders, R. A., Stojanovic, M., Tallmadge, D. H., Eberhart, B. L., Ewald, D. K.,
701 Gruber, D. C., Morsch, T. R., Strothers, M. A., Rizzi, G., & Villagran, M. D. (2003).
702 Acrylamide formation mechanisms in heated foods. *Journal of Agricultural and Food*
703 *Chemistry*, 51(4782-4787).


704 Zyzak, D. V., Sanders, R. A., Stojanovic, M., Tallmadge, D. H., Eberhart, B. L., Ewald, D. K.,
705 Gruber, D. C., Morsch, T. R., Strothers, M. A., Rizzi, G. P., & Villagran, M. D.
706 (2003). Acrylamide formation mechanism in heated foods. *Journal of Agricultural*
707 *and Food Chemistry*, 51(16), 4782-4787.

708

709 Figure 1 Mechanism of acrylamide formation (adapted from Parker, Balagiannis, Higley,
 710 Smith, Wedzicha, and Mottram (2012)).

711

712





713

714

715

716 Figure 2 (a) Effect of asparaginase (Acrylaway®) dose at pH 7 and 40 °C on acrylamide formation in semi-sweet biscuits; biscuits were baked at
717 260 °C for 5.5 min. (b) Effect of asparaginase (Acrylaway®; 2100 ASNU/kg of flour) incubation conditions on acrylamide formation in
718 crispbread; crispbread was baked at 250 °C for 11 min (adapted from Hendriksen, Kornbrust, Ostergaard, and Stringer (2009)).

719



720

721

722

Table 1 Results published on enzymatic mitigation of acrylamide using asparaginase

foodstuff	enzyme source	enzyme dosage	processing conditions	acrylamide reduction	reference
potato	<i>Escherichia coli</i>	not stated	not stated	99%	Zyzak et al. (2003)
gingerbread	<i>E. coli</i>	4 U/kg	various time/temperature combinations	55%	Amrein et al.(2004)
potato	<i>E. coli</i>	0.2–1 U/g	180 °C, 20 min	50–90%	Ciesarová et al. (2006)
French fries	<i>A. oryzae</i>	10000 ASNU/L*	175 °C, 3 min	67%	Pedreschi et al. (2008)
semi-sweet biscuits, ginger biscuits, crispbread, French fries, potato crisps	<i>A. oryzae</i>	various dosages	various time/temperature combinations	semi-sweet biscuits: 65–84% ginger biscuits: 34–90% crispbread: 84–92% French fries: 59% potato crisps: 60%	Hendriksen et al. (2009)
fried dough model system	<i>A. oryzae</i>	100, 500, 1000 U/kg	180 or 200 °C; 4, 6 or 8 min	90%	Kukurová et al. (2009)
bread	enzymes from <i>Pisum sativum</i> L.	not stated	220 °C, 22–25 min	wheat bran bread: 57% whole-grain bread: 68%	Tuncel et al. (2010)
potato chips	<i>A. oryzae</i>	10000 ASNU/L	170 °C, 5 min	90%	Pedreschi et al. (2011)
biscuits	<i>A. oryzae</i>	100–900 U/kg	200 °C, final moisture content 2%	7–88%	Anese et al. (2011a)
biscuits	<i>A. oryzae</i>	900 U/kg	200 °C, final moisture content 2%	69%	Anese et al. (2011b)
potato	<i>Bacillus licheniformis</i>	30 IU/mL	175 °C, 15 min	80%	Mahajan et al. (2012)
lebkuchen, tortilla chips, potato snack, French fries, coffee	<i>A. oryzae</i>	various dosages	lebkuchen: 200 °C, 14 min tortilla chips: 190 °C, 60 s	lebkuchen: 95% tortilla chips: 90%	Hendriksen et al. (2013).

			French fries: 175 °C, 3 min others not specified	potato snack: 40% French fries: 57% coffee: 55–74%	
cookies	<i>A. oryzae</i>	500 U/kg	205 °C, 11 or 15 min	23–75%	Kukurová et al. (2013)
wheat-oat bread	<i>Aspergillus niger</i>	500 U	220, 230 and 250 °C; 10, 30 and 40 min	90%	Ciesarová et al. (2014)
sweet bread	<i>Cladosporium sp.</i>	50–300 U	220 °C; 25 min	sweet bread crust: 97% sweet bread crumb: 73%	Kumar et al. (2014)
biscuits, bread	<i>Rhizomucor miehei</i>	0.5–10 U	200 °C; 15 min	biscuits: 81.6% bread: 94.2%	Huang et al. (2014)
potato crisps	<i>Bacillus subtilis</i>	0–40 U	170 °C; 90 s	80%	Onishi et al. (2015)
French fries	<i>Thermococcus zilligii</i>	0–20 U	175 °C; 5 min	80%	Zuo et al. (2015b)

* ASNU is defined as the amount of asparaginase that produces 1 µmol of ammonia per min under the conditions of the assay (pH = 7 ± 0.005; 37 ± 0.5 °C) using

Acrylaway®