

*Similarities and differences in the
historical records of lava dome-building
volcanoes: implications for understanding
magmatic processes and eruption
forecasting*

Article

Accepted Version

Creative Commons: Attribution-Noncommercial-No Derivative Works 4.0

Sheldrake, T. E., Sparks, R. S. J., Cashman, K. V., Wadge, G. and Aspinall, W. P. (2016) Similarities and differences in the historical records of lava dome-building volcanoes: implications for understanding magmatic processes and eruption forecasting. *Earth Science Reviews*, 160. pp. 240-263. ISSN 0012-8252 doi: 10.1016/j.earscirev.2016.07.013 Available at <https://centaur.reading.ac.uk/66290/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

To link to this article DOI: <http://dx.doi.org/10.1016/j.earscirev.2016.07.013>

Publisher: Elsevier

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other

copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online

Title: Similarities and differences in the historical records of lava dome-building volcanoes: implications for understanding magmatic processes and eruption forecasting.

Authors:

Sheldrake, T. E. ^{a,* 1}

Sparks, R.S.J. ^a

Cashman, K.V. ^a

Wadge, G. ^b

Aspinall, W.P. ^a

^a School of Earth Sciences, University of Bristol, Wills Memorial Building, Queen's Road, Bristol, BS8 1RJ, UK

^b Department of Meteorology, University of Reading, Reading, RG6 6AL, UK

* Thomas.sheldrake@unige.ch (corresponding author)

¹ Current address: Section of Earth and Environmental Sciences, University of Geneva, rue des Maraîchers 13, Geneva CH-1205, Switzerland

Abstract:

A key question for volcanic hazard assessment is the extent to which information can be exchanged between volcanoes. This question is particularly pertinent to hazard forecasting for dome-building volcanoes, where effusive activity may persist for years to decades, and may be punctuated by periods of repose, and sudden explosive activity. Here we review historical eruptive activity of fifteen lava dome-building volcanoes over the past two centuries, with the goal of creating a hierarchy of exchangeable (i.e., similar) behaviours. Eruptive behaviour is classified using empirical observations that include patterns of SO₂ flux, eruption style, and magma composition. We identify two eruptive regimes: (i) an *episodic* regime where eruptions are much shorter than intervening periods of repose, and degassing is temporally correlated with lava effusion; and (ii) a *persistent* regime where eruptions are comparable in length to periods of repose and gas emissions do not correlate with eruption rates. A corollary to these two eruptive regimes is that there are also two different types of repose: (i) inter-eruptive repose separates episodic eruptions, and is characterised by negligible gas emissions and (ii) intra-eruptive repose is observed in persistently active volcanoes, and is characterised by

39 continuous gas emissions. We suggest that these different patterns of can be used to
40 infer vertical connectivity within mush-dominated magmatic systems. We also note
41 that our recognition of two different types of repose raises questions about
42 traditional definitions of historical volcanism as a point process. This is important,
43 because the ontology of eruptive activity (that is, the definition of volcanic activity in
44 time) influences both analysis of volcanic data and, by extension, interpretations of
45 magmatic processes. Our analysis suggests that one identifying exchangeable traits
46 or behaviours provides a starting point for developing robust ontologies of volcanic
47 activity. Moreover, by linking eruptive regimes to conceptual models of magmatic
48 processes, we illustrate a path toward developing a conceptual framework not only
49 for comparing data between different volcanoes but also for improving forecasts of
50 eruptive activity.

51

52 Keywords: Lava-dome volcanoes; Exchangeable behaviours; Persistent; Episodic;
53 Magmatic processes; Forecasting.

54
55

56 1. Introduction

57

58 Volcanic activity can be manifested in many different ways. From a volcanic risk
59 perspective one important variety of eruptive activity is extrusion of lava domes at
60 intermediate and silicic volcanoes. Recurrent hazards associated with dome-
61 building activity include: pyroclastic flows and volcanic blasts associated with the
62 collapse of lava domes and edifice instability; fountain-fed pyroclastic flows
63 associated with Vulcanian to sub-Plinian explosions; and copious tephra fall .
64 Worldwide, such volcanic activity has been responsible for over two thirds of
65 volcanic fatalities since 1600 C.E. (Auker et al., 2013).

66

67 Within the Smithsonian Global Volcanism Program (GVP) database there are 205
68 recorded dome-building volcanoes that have been active in the Holocene (Siebert et
69 al., 2010). Of these, 117 have erupted in the last millennium and 89 have erupted
70 since 1900 C.E. (Ogburn et al., 2015). Historical eruptions have lasted many months,
71 years or even decades (Newhall and Melson, 1983; Sparks, 1997; Ogburn et al.,
72 2015). Over historical timescales volcanic activity can be regarded as continuous,
73 albeit fluctuating, but may also include complex episodic and sometimes cyclic
74 fluctuations in intensity, duration, frequency and eruptive style.

75

76 Lava dome formation requires particular conditions, which suggests that magmatic
77 processes at dome-building volcanoes have shared characteristics. Specifically, the
78 lavas of dome-building volcanoes have low average eruption rates ($\sim 10^{-1}$ to 10^{-2}

79 km³ yr⁻¹) and high viscosities (10⁶ to 10¹¹ Pa s; [Yokayama, 2005](#)) that are commonly
80 associated with high groundmass crystallinity ([Cashman, 1992](#)) and, consequently,
81 substantial yield strength ([Calder et al., 2015](#)). Nevertheless, dome-building
82 volcanoes can exhibit markedly different eruptive histories, including both the
83 duration of individual eruptive episodes and the potential for explosive activity. This
84 variability reflects the general conceptual tensions in volcanology where: (1) there
85 is a belief that individual volcanoes are unique, as exhibited by the complex nature
86 of their eruptive records, and (2) the concept that eruptive activity is driven by
87 common magmatic processes that produce certain eruptive styles and volcano
88 morphologies ([Cashman & Biggs, 2014](#)).

89

90 In this review we identify characteristics of fifteen lava-dome building volcanoes
91 that are similar (exchangeable) or unique (not exchangeable), as well as those that
92 are common only to a sub-group of volcanic records. In volcanology, for example,
93 the concept of exchangeable characteristics can be used to define the common traits
94 for all volcanoes, and to infer the conceptual system that this definition represents.
95 Using this idea, the basic exchangeable characteristics of a volcanic system - implied
96 by the definition of a volcano by [Borgia et al. \(2010\)](#) - are simply magma, eruption,
97 and edifice. We ally to this the idea that the volcanic system (and thus the
98 conceptual construct of volcanism) should be hierarchically organized, such that
99 identifying and characterizing different hierarchies allows individual volcanoes to
100 be distinguished in space and time ([Szakács, 2010](#)). For this reason, we develop a
101 hierarchy of different eruptive behaviours using observations from the historical

records of fifteen well-characterised dome-building volcanoes. By characterising exchangeable behaviours we can assess inaccessible elements (e.g., the magmatic system) from observed elements (e.g., surface phenomena). A similar approach is employed in medical sciences, where individuals (i.e. humans) are unique, but different groups of humans are known to have similar health traits (Spiegelhalter, 1986; Best et al., 2013).

Using a hierarchical construct for eruptive behaviours at dome-building volcanoes we consider the conceptual system that can explain the different sets of shared traits and characteristics. Specifically, we ask whether the diversity in behaviours can be explained by subsystems of the magmatic system (e.g., shallow crustal reservoirs) or whether it requires a more holistic view of crustal magmatic processes (i.e., a transcrustal reservoir system that extends from the surface through the crust and into the mantle). This approach allows us to evaluate emerging paradigms for eruptive activity based on the destabilisation and reorganisation of igneous mush systems (e.g., Cashman and Giordano, 2014; Christopher et al., 2015), and to interpret the role of connectivity within a magmatic system on the pattern and style of eruptive activity at dome-building volcanoes.

An additional application of our study relates to the implications of a hierarchical construct on the analysis of volcanic datasets. An important issue relates to the concept of volcanic activity as a point-process of discrete events as this influences how magmatic processes are interpreted and how probabilistic forecasts are made.

We also examine the implications of different patterns of eruptive behaviour on forecasting the activity of one volcano using observations from other (perhaps better characterized) volcanoes of the same type. We discuss the issues when selecting evidence to make eruptive forecasts and contextualize this in regards of forecasting the onset of eruptive activity.

2. Data

The fifteen dome-building volcanoes selected for this review are listed in Figure 1. Our selection is governed by the quality of available data and relevant observations, and guided by the principle that our dataset should contain volcanoes that are well-characterised, have long records of activity and have been recently active. All fifteen volcanoes sit in arc environments, and erupt magmas that are hydrous and intermediate in composition. As volcanic gas emissions are an important aspect of dome-building volcanism, we also include one volcano characterised by persistent gas emissions but no recent eruptive activity.

To enable comparison of similar dome-building behaviour, we restricted the selection to volcanoes of intermediate composition, thus omitting domes formed by the eruption of crystal-poor rhyolites (e.g., Chaiten 2008; [Pallister et al., 2013](#)). To ensure that the eruptive records are complete and not affected by recording biases ([Coles and Sparks, 2006](#); [Deligne et al., 2010](#)), we review patterns of eruptive activity only back to 1800 C.E. (Fig. 2), as prior to this date each of the individual

eruptive records is assumed to be incomplete. However, recent advances in the ability to monitor and observe eruptive activity (Cashman and Sparks, 2013) mean that much of the data derive from eruptive activity in the late 20th and early 21st centuries. Data sources include eruption databases (Siebert et al., 2010; Ogburn, 2013), peer-reviewed publications (e.g., journal articles, professional publications), and observatory data and databases of volcanic unrest (e.g., WOVOdat; <http://www.wovodat.org/>). Detailed profiles for the volcanoes can be found in the supplementary material.

Data are collated for two purposes: (i) as empirical evidence of long-term behaviours at dome-building volcanoes, and (ii) as a semi-quantitative measure of their behaviour. Empirical evidence includes observations of phenomenological behaviour, magmatic degassing, and the bulk rock characteristics of erupted products. In contrast to focussed studies at the individual volcanoes, we do not use the observations as direct evidence of specific magmatic processes or characteristics of the respective magmatic systems. Instead, we use them only to subdivide individual volcanoes into groups that reflect their long-term eruptive behaviour. We then examine geophysical (seismicity and deformation) and petrological observations within groups to compare the behaviour of the magmatic systems within and between volcano groups.

2.1. Phenomenological behaviour

Dome-building volcanoes exhibit a range of effusive and explosive behaviours (Newhall and Melson, 1983; Sparks, 1997; Ogburn et al., 2015). By definition, however, the main eruptive activity involves protracted lava dome extrusion, with extrusive phases that may last from months to many years; our reference volcanoes have also experienced periods of quiescence of months to decades. During times of activity, lava discharge rates can be estimated from ground-based and satellite-based techniques (e.g., Sparks, 1997; van Manen et al., 2010) and used to characterise the intensity of dome growth phases. The effusion rate, together with the magma viscosity, determines whether lava moves away from the vent as a lava flow or builds either an ever-larger dome and talus apron or a near-solid lava spine (Watts et al., 2002; Cashman et al., 2008). Where lava accumulates over the vent, the increase in magma-static head creates a backpressure that can resist extrusion and influence the longer-term dynamics of the magmatic system (Stasiuk et al. 1993; Scandone et al., 2007).

Phases of extrusive activity can be interspersed with more explosive activity, including Strombolian, Vulcanian and sub-Plinian eruption styles. The intensity and explosivity of eruptive activity can be characterised using phenomenological observations such as ash column height, pyroclastic run-out, tephra fall deposit volumes, some of which can serve as proxies for magnitude, intensity and explosion style (Newhall and Self, 1982). Infrequently, dome-building volcanoes also have large-magnitude explosions, including Plinian eruptions and lateral blasts (Ogburn et al., 2015).

193

194 2.2. Magmatic degassing

195 As magma ascends through the crust, volatiles exsolve and rise to the surface
196 (Wallace, 2003; 2005; Oppenheimer et al., 2011). The most abundant volatile
197 species are H₂O and CO₂. SO₂, however, is the most commonly monitored volatile
198 because it is a trace gas in the atmosphere and thus its concentration can be readily
199 measured using remote sensing techniques (Rose et al., 2000; Edmonds et al., 2003;
200 Galle et al., 2003; Carn et al., 2013). SO₂ fluxes are quantified using ultraviolet
201 absorption spectra and measured in tonnes per day (t/d); some data for the last few
202 decades are sporadically available for most of the study volcanoes.

203

204 Prior to the development of ultraviolet spectroscopic techniques, gas fluxes were
205 estimated by sampling fumarolic gases (Giggenbach, 1996). Introduction of the
206 correlation spectrometer (COSPEC) in the 1970s allowed SO₂ flux measurements,
207 although early measurements were prone to large errors (Oppenheimer et al.,
208 2011). More recently, fluxes have been estimated from differential optical
209 absorption spectroscopy (DOAS; Platt and Stutz, 2008). A major advantage of this
210 method is that spatially distributed multi-beam instruments can provide precise
211 estimates for plume velocity, which significantly reduces measurement errors of
212 flux (Oppenheimer et al., 2011, and references within). Importantly, however,
213 measurements are restricted to sunlight hours only and the quality of gas data from
214 all remote sensing techniques depends on meteorological conditions (e.g., low
215 humidity and no clouds).

216

217 Terrestrial-based spectroscopic measurements are not feasible for measuring
218 volatile emissions in major explosive events because abundant ash masks the
219 signals. Consequently, the mass of gas released during large eruption events is
220 measured using satellite-based techniques and then converted to fluxes (Carn and
221 Prata, 2010; Carn et al., 2013).

222

223 *2.3. Bulk rock observations*

224 In our data set, the products of dome building volcanoes range in composition from
225 basaltic andesite (~52-57 wt.% SiO₂) to dacite (~64-69 wt.% SiO₂; Table 1). Whilst
226 it is impossible to observe the long-term dynamics of magmatic systems directly,
227 macroscopic observations and bulk rock analysis can be used to interpret the
228 compositional, and potentially the physical, structure of magmatic systems (e.g.,
229 Barclay et al., 2010; Larsen et al., 2010; Coombs et al., 2013; Scott et al., 2013;
230 Turner et al., 2013).

231

232 Magma rheology is a major determinant of physical behaviour, particularly at
233 shallow depths where flow at the surface may be inhibited by high yield strength.
234 Magma is a multiphase system and consequently its rheology is complex (Mader et
235 al., 2013). Rheology is strongly controlled by the crystallinity of the magmas, which
236 is typically high in intermediate arc magmas. Crystallization is further increased by
237 syn-ascent decompression and degassing and is thus modulated by eruption rate,
238 the pressure of shallow storage prior to eruption, the bulk composition of the

magma and kinetic factors associated with bubble dynamics (Jaupart and Vergnolle, 1989; Geschwind and Rutherford, 1995; Nakada and Motomura, 1999; Hammer et al., 2000; Cashman and McConnell, 2005; Divoux et al., 2009; Wright et al., 2012). Exsolved gas can also lead to marked rheological variations as functions of bubble size distribution and bubble content (Manga et al., 1998; Mader et al., 2013). The interplay between magma ascent, decompression, gas exsolution, crystallization and rheology can lead to complex episodic behaviours (e.g., Jaupart and Allegre, 1991; Melnik and Sparks, 1999; Michaut et al. 2013).

2.4. Geophysical observations

For each volcano we report common geophysical observations; for consistency, we omit specialised observations (e.g., strain meters, broadband seismicity) made at only one or two volcanoes. Geophysical monitoring observations are susceptible to spatial and temporal biases associated with network capacities and technological constraints at volcano observatories (Sparks et al., 2012). Therefore, it is important to understand these biases and thus the robustness and validity of comparing records. Spatial biases arise from variations in monitoring capacities due to both resource availability and accessibility. Temporal biases are associated with advances in technology that improve observation thresholds and the precision of measurements. These are discussed in more detail with reference to the particular observables.

2.4.1. Seismicity

Volcanic seismicity can be categorised either by its physical cause, if occurring at the surface (e.g., rockfalls, lahars, pyroclastic flows, etc.), or its waveform and frequency content if originating from within the crust (e.g., high or low-frequency signals; Chouet, 1996; Neuberg, 2000; McNutt, 2005; Chouet and Matoza, 2013). High-frequency (volcano-tectonic) events have recognisable P and S wave first arrivals and are attributed to brittle fracturing related to opening of new pathways for either magma or magmatic fluids (Kilburn, 2003). Low-frequency (long-period and hybrid) events are associated with movement of magma and magmatic fluids (McNutt, 2005). Seismicity is most commonly associated with eruptive activity but is also observed during periods of quiescence, that is, when a volcano is in a non-eruptive state, and can be diagnostic of incipient unrest (Phillipson et al., 2013) or post-eruptive tectonic stress recovery (e.g., Barker and Malone, 1991).

Although seismicity can be characterised using a range of metrics, we focus on the number of events (daily counts) as this is the most commonly recorded observation across the volcanoes in the dataset. We do not compare absolute numbers of seismic events or cumulative seismic moment between volcanoes due to recording biases associated with variations in network capacities and sensitivities (e.g., number of and type of instruments). Instead we compare patterns of total seismicity and the relative frequency of different types of events, primarily long-period and volcano-tectonic earthquakes.

2.4.2. Deformation

The episodic and sometimes repetitive nature of eruptive activity at many dome-building volcanoes commonly manifests as time-varying deformation of the crust that can be monitored at the surface using geodetic techniques. Great variability in instrumentation and network design in the near-field monitoring of ground deformation, however, makes direct comparisons difficult. For this reason, we focus only on far-field deformation (> 5 km from the vent). These data also provide useful constraints on deeper magmatic processes. Far-field deformation can be measured by geodetic networks (using GPS), although these measurements require ground-based support and are restricted to only a few of the volcanoes in our dataset. On the other hand, Interferometric Synthetic Aperture Radar (InSAR) techniques using satellite-based instruments provide a global approach for observing far-field deformation (Biggs et al., 2014). By combining observations from these two methods we compare patterns of deformation (i.e. whether the volcano is inflating, deflating or neither) between different volcanoes and relate deformation behaviour to eruptive and non-eruptive phases of activity.

2.5. Petrology

Petrologic data provide information on the homogeneity of the magmatic system, temporal changes of magma composition and the extent to which eruptive activity is influenced by the ascent of discrete magma batches. Of particular interest is evidence for the interaction of different magmas, which can occur at a range of scales. Macroscopic evidence for magma mingling includes enclaves or compositional banding in erupted products. Microscopic details of geochemical

interactions provide information on the nature and timing of mingling events. Analysis of individual crystals and their melt inclusions provides information on both intrinsic and extrinsic properties (e.g., temperature, pressure and volatile inventories) of magma storage regions (e.g., Nakamura, 1995; Zellmer et al., 2003a; Dirksen et al., 2006; Humphreys et al., 2006; Costa et al., 2013). Finally, petrological analyses and U-series geochemistry can constrain the timescales of magmatic processes (e.g., Volpe and Hammond, 1991; Zellmer et al., 2003b; Cooper and Reid, 2008; Dosseto et al., 2008; Claiborne et al., 2010) that control and sustain eruptive activity at dome-building volcanoes. Quantification of groundmass characteristics (crystallinity, crystal size and shape) can further constrain rates of magma ascent to the surface (e.g., Hammer et al., 2000; Toramaru et al. 2008; Wright et al., 2012).

3. Patterns of eruptive activity at dome-building volcanoes

We identify in our dataset two types of long-term behaviour defined by the relative time a volcano remains in a state of eruption or repose (i.e. non-eruption): (1) activity is *episodic* when time scale of eruption is much less than the time scale of repose; and (2) activity is *persistent*, when the time scale of eruption is comparable to that of repose (Fig. 3a). Identification of episodic and persistent regimes represents the first sub-level in our hierarchical construct of historical dome-building volcanism (Fig. 4).

Episodic and persistent behaviour can be manifested over different timescales (Fig. 4) and, over time, individual volcanoes can show both types of behaviour (Fig. 3c). Over the examined time period of the past 200 years, for example, many dome-building volcanoes are characterised by episodic behaviour; however, within that broad description, some have remained in a persistent regime for multiple decades. We characterise these volcanoes as belonging to a mixed regime. Over very long time periods, all the volcanoes in our sample can be viewed as mixed.

Patterns of SO₂ degassing also provide additional insight into long-term patterns of volcanic activity. The largest volumes of SO₂ emissions are always associated with major explosive events (e.g., [Carn and Prata, 2010](#); [Werner et al., 2013](#)). Two patterns of less energetic degassing can be defined as: (1) SO₂ flux that is closely correlated with eruptions (Fig. 3b) and (2) degassing that is not correlated with eruptive activity (Fig. 3a). Correlated degassing is common at volcanoes in an episodic regime; here both gas and magma fluxes decrease with time after an initial (often explosive) maximum (Fig. 5a). Poor correlation between degassing and eruptive activity, in contrast, is typical of persistent activity (Fig. 5b). The correlation of degassing patterns with eruptive behaviour suggests that magmatic degassing constitutes an important distinction between persistent and episodic regimes (e.g., [Whelley et al., 2015](#)).

Finally, we use differences in degassing behaviour to distinguish two states of repose: (1) inter-eruptive repose separates episodic eruptions and is characterised

by negligible degassing (Fig 3a;5a); and (2) intra-eruptive repose occurs in the persistent regime and is characterised by sustained degassing (Fig 3b;5b). We also identify a non-eruptive degassing regime to describe dome-building volcanoes that remain in a state of long-term repose (~decades) characterised by low levels of persistent degassing.

3.1. Episodic regime

Volcanoes in an episodic regime are characterised by periods of eruptive activity separated by much longer periods of repose. The onset of eruptive episodes is explosive, with high magma discharge rates. Both magma discharge rates and SO₂ fluxes decrease with time during eruptive periods (e.g., Fig. 6). During eruptive periods, the later stages of activity are typically characterised by low extrusion rates and associated extensive syn-eruptive crystallisation that combine to produce lava spines (e.g., [Watts et al., 2002](#); [Cashman et al., 2008](#)). We distinguish two different timescales for episodic activity in historical records (Fig 4): (1) volcanoes where eruptive episodes last several years, and (2) volcanoes where eruptive episodes last a few months at most. These two subgroups can be further distinguished by the homogeneity or heterogeneity of erupted magma compositions.

3.1.1. Eruptive episodes lasting years

Two volcanoes in this review have experienced episodic activity lasting several years (Fig. 2; UNZ, PEL). In both cases, lava compositions are broadly homogeneous. The duration of inter-eruptive periods of repose is multiple decades or longer.

(a) *Mount Unzen*, Japan (UNZ), is a complex dacitic volcano that last erupted near-continuously from 1991-1995 (Fig. 2). No previous historic activity is known although a major sector collapse event of an older dome occurred in 1792 (Ui et al., 2000). Between 1991 and 1995, the composition of eruptive products was ~65 wt.% SiO₂ (Nakada and Motomura, 1999) and the average lava effusion rate was ~1 m³s⁻¹, with higher rates (~4-6 m³s⁻¹) during the eruption onset. Extrusion rates generally diminished with time, although a secondary peak was observed in 1993 (Nakada et al., 1999; Fig. 6). SO₂ fluxes averaged 137 t/d, were correlated with extrusion rate and diminished soon after eruptive activity ceased (Hirabayashi et al., 1995).

(b) *Mont Pelée*, Martinique (Fig. 1), is an andesitic volcano that has erupted infrequently (Fig. 2). The best-recorded eruptive activity occurred in the early part of the 20th century, between 1902-05 and 1929-32 (Lacroix, 1904; Perret, 1937; Tanguy, 1994). During both periods, lava fluxes decreased from >10 m³s⁻¹ to ~1 m³s⁻¹ (Tanguy, 2004), with later stages characterised by spine extrusion (Lacroix, 1904; Perret, 1937). The composition of eruptive products from Mont Pelée is quite homogeneous at 62 wt.% (Fichaut et al., 1989b; Gourgaud et al., 1989; Smith and Roobol, 1990).

3.1.2. Eruptive episodes lasting months

Two volcanoes in this review have experienced eruptive episodes lasting a few months (Fig. 2; RED, AUG). In contrast to the volcanoes in the previous group, the duration of inter-eruptive periods of repose is several years to a few decades. Additionally, lavas of different composition are erupted contemporaneously.

(c) *Mount Redoubt*, USA (RED), is an andesitic volcano that has erupted intermittently on four separate occasions since 1902 (Fig. 2). The most recent eruptive episodes have been in 1989-90 and 2009, each lasting for several months (Miller and Chouet, 1994; Bull and Buurman, 2013). During each eruptive episode eruptive products ranged from 57 to 63 wt.% SiO₂, with the later stages involving the more silicic lava (Nye et al., 1994; Coombs et al., 2013). SO₂ degassing is highly correlated with periods of eruptive activity. In 1989 -1990, extrusion rates varied from 2.1 to 26 m³s⁻¹, with average dome growth occurring at ~5.8 m³s⁻¹ (Miller, 1994). Similar extrusion rates were observed in 2009 (2.2 -35 m³s⁻¹) although the average rate was slightly higher at ~9.5 m³s⁻¹ (Diefenbach et al., 2013). In both cases the initial activity was the most explosive and the extrusion rate declined during eruptive activity (Miller, 1994; Diefenbach et al., 2013). Initial explosive activity in 2009 was associated with the largest SO₂ fluxes (~3000 to ~17000 t/d). Subsequent activity involved more continuous extrusion with SO₂ fluxes ≤3000 t/d (Hobbs et al., 1991; Casadevall et al., 1994; Werner et al., 2013). In both 1990 and 2009, it took several years for SO₂ fluxes to

return to undetectable levels after eruptive activity ceased (Doukas, 1995; Werner et al., 2013).

(d) *Mount Augustine*, USA (AUG), is an andesitic volcano that has had nine known eruptive episodes since 1812, with the most recent in 1976, 1986 and 2006 (Fig 2), each lasting for several months (Swanson and Kienle, 1988; Power et al., 2006; Power and Lalla, 2010). The composition of the erupted magma has ranged from 56 to 64 wt.% SiO₂, with more silicic magma preferentially erupted later in each eruptive episode (Harris, 1994; Roman et al., 2006; Larsen et al., 2010). During the 2006 eruptive activity, magma fluxes varied from 2 to 22 m³s⁻¹ (Coombs et al., 2010). Notably, in contrast to other volcanoes in episodic regimes, the final stages of eruptive activity at Augustine in 2006 were characterised by elevated discharge rates and the formation of lava flows, although discharge rates were still lower than at the onset of eruptive activity (Coombs et al., 2010). Magmatic degassing is correlated with eruptive activity, with the largest fluxes commonly associated with explosive activity (Stith et al., 1978, Rose et al., 1988; McGee et al., 2010). In 2006, however, the highest SO₂ fluxes (~9000 t/d) were associated with a brief hiatus in eruptive activity, although SO₂ fluxes were high (~3000 t/d) throughout the eruptive episode (McGee et al., 2010), and it took 1-2 years after the end of eruptive episodes in 1986 and 2006 for SO₂ fluxes to return to undetectable levels (Symonds et al., 1990; Doukas, 1995; McGee et al., 2010).

443

444 3.2. *Persistent regime*

445 We identify eight volcanoes in this review that have remained in a persistent regime
446 for decades or longer. Volcanoes in a persistent regime exhibit broadly consistent
447 behaviour associated with stable long-term lava fluxes. For example, although rates
448 of lava effusion at Bezymianny, Kamchatka, have varied over the short term, they
449 have been approximately constant over the past several decades (Fig. 5). The
450 eruptive activity of an individual volcano can also show ‘typical’ (repeatable)
451 patterns of behaviour, as illustrated by Santiaguito, Guatemala, where typical
452 behaviour comprises “small to moderate explosions of steam and ash, small
453 pyroclastic flows... and effusion of blocky lava domes and flows” (Scott et al., 2012).
454 Typical intermittent behaviour at Merapi, Indonesia, in contrast, is characterised by
455 eruptive activity that is “low in explosivity with VEI-3 or less ... [that] involve the
456 formation of a lava dome” (Ratdomopurbo et al., 2013).

457

458 We distinguish two different variants of long-term persistent behaviour (Fig. 4).
459 Firstly, there are volcanoes that have remained in a persistent regime at least the
460 19th century. These volcanoes produce lavas with an approximately constant bulk
461 composition. Secondly, there are volcanoes that have entered a persistent regime
462 following a long period of in a state of repose. Volcanoes in this group typically have
463 bulk compositions that show a decrease in SiO₂ content with time.

464

465 3.2.1. *Long-term persistent regimes*

Four of the dome-building volcanoes in this study have been in a persistent regime throughout the 19th, 20th and 21st centuries; these volcanoes are characterised by frequent, intermittent phases of dome-growth (Fig. 2; MER, COL, LAS, SHI). The style of eruptive activity is generally consistent through time and characterised by definable ‘typical’ behaviour, except for rare large-magnitude explosions (Fig. 2). Interestingly, these explosive events commonly involve magma that is more mafic than erupted during the effusive phases. Activity at each volcano is described in detail below.

(a) *Merapi*, Indonesia (MER), is a basaltic andesite volcano that has been in an eruptive state every few years since at least the 18th century. Eruptive activity is characterised by minor explosions associated with the extrusion of viscous lava domes and coulées that can collapse to form block-and-ash pyroclastic flows (Voight et al., 2000). Lava extrusion rates are approximately constant over historical records at $\sim 0.5 \text{ m}^3\text{s}^{-1}$ (Siswowardoyo et al., 1995). Persistent effusive activity has been punctuated by at least two major explosions that have produced high-energy pyroclastic density currents (Surono et al., 2012). The bulk rock lava composition ranges from 52 to 56 wt.% SiO₂ (Andreastuti et al., 2000; Gertisser and Keller, 2003) and shows no temporal trend, although explosive events appear to involve deeply sourced, volatile-rich magmas (Costa et al., 2013), which may be more mafic (Gertisser and Keller, 2003). SO₂ degassing is continuous with fluxes between 50 and 250 t/d (Humaida, 2008), although instantaneous fluxes can be much

larger ($\sim 10,000$'s t/d) during major explosive events (Surono et al., 2012).
Importantly, SO₂ fluxes and eruptive activity appear decoupled, with SO₂ flux
peaks observed during inter-eruptive periods, and sometimes associated
with ash venting (Ratdomopurbo et al., 2013).

(b) *Colima*, Mexico (COL), is an andesite volcano that has been erupting
intermittently since the 18th century. Periods of intra-eruptive repose
normally last on the order of years, although longer periods without
apparent eruptive activity have followed major explosive events in 1818 and
1913. These longer periods of repose probably involved endogenous growth
below the crater rim (Robin et al., 1991; González et al., 2002), so we infer
that Colima remained in a persistent regime during post-explosion periods.
Eruptive activity is characterised by lava dome extrusion, Vulcanian
explosions and occasional block-and-ash flows (Zobin et al., 2002). Short-
term lava effusion rates vary from <1 to >5 m³s⁻¹ (Varley et al., 2010), but
long-term averages are poorly constrained. The lava composition ranges
from 59 to 62 wt.% SiO₂ with no clear temporal trend (Luhr and Carmichael,
1980; 1990; Savov et al., 2008), except that products of major explosive
events are more mafic (SiO₂ = 55-58 wt.%; Luhr and Carmichael, 1990; Reubi
and Blundy, 2009; Crummy et al., 2014). SO₂ degassing is continuous, with
fluxes typically between 50 and 1000 t/d (Casadevall et al., 1984; Engberg,
2009), although sometimes as high as 5000 t/d (Taran et al., 2002; Varley
and Taran, 2003). Magmatic degassing appears decoupled from eruptive

activity (Zobin et al., 2008), but the largest SO₂ fluxes are associated with more explosive events (Taran et al., 2002).

(c) *Lascar*, Chile (LAS), is an andesitic volcano that has been erupting intermittently at yearly to decadal timescales throughout much of its history. Lava dome growth has been confined within a large summit crater. Four periods of near-continuous dome growth occurred between 1984 and 1993; each culminated in lava dome subsidence and explosive events, including a Plinian explosion in April 1993 (Matthews et al., 1997). Long-term lava extrusion rates are poorly constrained but are likely to be < 0.1 m³s⁻¹ (Matthews et al., 1997). Since 1993, activity has comprised episodic Vulcanian explosions that have decreased in both intensity and frequency; the last explosion occurred in 2007. Juvenile pyroclasts from 1993 can be separated by composition into two groups: 57.6-58.7 or 60.4-61.4 wt.% SiO₂ (Matthews et al., 1999); similarities to previously erupted lavas (Deruelle, 1985) suggest that the magma composition has remained constant throughout its history. *Lascar* has exhibited continuous fumarolic activity (Casertano, 1963; Gardeweg & Medina, 1994) with recent SO₂ fluxes sustained between 150 and 940 t/d (Henney et al., 2012, Menard et al., 2014). During more explosive activity, fluxes have reached 2300 t/d (Andres et al., 1991; Mather et al., 2004). SO₂ fluxes have shown an irregular pattern of degassing during periods of intra-eruptive repose and therefore appear decoupled from magma flux (Menard et al., 2014).

(d) *Shiveluch*, Russia (SHI) is an andesitic volcano that has been erupting intermittently since a major explosive event in 1854. Even prior to 1854, sparse observations suggest that periods of repose lasted no more than a few decades. Recent phases of eruptive activity have varied in duration from months to several years, and Shiveluch has been in a near-continuous eruptive state since 2000 (Belousov, 1995; Zharinov and Demyanchuk, 2008). Between 1980 and 2007 the average lava discharge rate was $\sim 0.4 \text{ m}^3\text{s}^{-1}$, although fluxes fluctuated considerably (Zharinov and Demyanchuk, 2008). Explosive activity has been of variable magnitude, with major Plinian events in 1854 and 1964 (Belousov, 1995). The eruptive products contain 56-62 wt.% SiO_2 and show no temporal trends (Dirksen et al., 2006; Humphreys et al., 2006; Gorbach and Portnyagin, 2011). Fumarolic activity has been sustained throughout both eruptive activity and intra-eruptive repose (Belousov, 1995; Gorelchik et al., 1997; Zharinov and Demyanchuk, 2008), but SO_2 fluxes have not been documented.

3.2.2. Long-duration repose preceding a long-term persistent regime

Two volcanoes in this study have initiated persistent behaviour after explosive eruptions that followed a long period in a state of repose (\sim millennia; Fig. 2; SAN, BEZ). The onset of a persistent regime at these volcanoes is characterised by Plinian and lateral blast explosions. In contrast to the previous group, the most evolved

pyroclasts in this group are associated with major explosive events; the SiO₂ content of subsequent lavas decreases systematically through time.

(e) *Santiaguito (Santa Maria)*, Guatemala (SAN), is a dome complex that has been active since 1922; effusive activity followed the Plinian eruption of its parent volcano, Santa Maria, in 1902 (Rose, 1972). Effusive activity has been nearly continuous at long-term rates of $\sim 0.46 \text{ m}^3\text{s}^{-1}$, with marked fluctuations that have been classified into eight distinct phases (Rose, 1973; Harris et al., 2003; Scott et al., 2013). Each phase has initiated with high rates ($0.5\text{--}2.1 \text{ m}^3\text{s}^{-1}$) and has been followed by low, sustained extrusion rates of $< 0.2 \text{ m}^3\text{s}^{-1}$ (Harris et al., 2003; Ebmeier et al., 2012). The lavas are dacitic to silicic andesite in composition, with SiO₂ contents that have decreased systematically from ~ 66 to ~ 62 wt.% since 1922. SO₂ degassing is continuous with average fluxes between 80 and 120 t/d (Andres et al., 1993; Rodríguez et al., 2004).

(f) *Bezmyanny*, Russia (BEZ), is an andesite volcano that has been erupting near-continuously to intermittently since a lateral blast and associated sector collapse in 1956 (Belousov et al., 2007). Between 1956 and 1977, eruptive activity was limited to periods of endogenous lava dome growth associated with sustained fumarolic activity (Gorshkov, 1959; Bogoyavlenskaya et al., 1985; Belousov, 1996). After 1977, dome growth occurred exogenously and included occasional explosions (van Manen et al., 2010). More recently,

eruptive phases have decreased in duration and have become increasingly explosive (West, 2013). The long-term average extrusion rate was $0.6 \text{ m}^3\text{s}^{-1}$ between 1956 and 1976 (Belousov et al., 2002) and 1993 to 2008 (van Manen et al., 2010). Since 1956 the eruptive products have become steadily less evolved with time, varying from 60.4 to 56.8 wt.% SiO_2 (Bogoyavlenskaya et al., 1985; Turner et al., 2013). SO_2 degassing has been sustained. Fluxes have been measured at 140 to 280 t/d during three campaigns conducted during periods of low eruptive activity (Lopez et al., 2013). These measurements are not sufficient to assess relations between degassing and magma discharge.

3.3. Mixed eruptive regime

Persistent and episodic regimes can manifest over different timescales at individual volcanoes. Consequently, the historical records of some dome-building volcanoes exhibit patterns of eruptive activity that are characteristic of both regimes: they exhibit persistent behaviour over several decades but are also characterised by long periods of inter-eruptive repose. We identify four volcanoes that fit this category and define them as ‘mixed’ regime volcanoes (Fig. 2; MSH, SHV, TUN, POP).

The eruptive behaviour at these volcanoes varies markedly, with persistent activity over short timescales but episodic activity over timescales of decades to centuries and persistent activity over shorter timescales. Mixed activity is sufficiently varied, however, that it cannot be considered exchangeable. For example, Mount St Helens

showed persistent activity throughout most of the 1980's with degassing that was well correlated temporally with lava extrusion. Tungurahua, in contrast, has remained in a persistent regime since 1999, with degassing that has been poorly correlated with lava extrusion. A common observation at all of these volcanoes, however, is intermittent ash venting.

(a) *Mount St. Helens*, USA (MSH), is a dacitic volcano that has experienced two eruptive episodes in recent times: 1980 to 1986, and 2004 to 2008 (Swanson and Holcomb, 1990; Scott et al., 2008), following an inter-eruptive period of repose lasting 136 years (Fig. 2). Eruptive activity in 1980 initiated with endogenous growth of the edifice (Lipman and Mullineaux, 1981) that caused a major flank collapse accompanied by sub-Plinian explosive activity (Voight et al., 1983; Glicken, 1998). This was followed by sub-Plinian to Vulcanian explosions in the summer of 1980 that steadily decreased in magnitude and duration (Scandone and Malone, 1985). Subsequent effusive activity transitioned between discrete and continuous eruptions of variably crystalline lavas (Cashman, 1992). Between 1980 and 1986, extrusion rates varied from 1.4 to 40 m³s⁻¹, with a long-term average of ~ 0.4 m³s⁻¹ (Anderson and Fink, 1990; Swanson and Holcomb, 1990). Renewed continuous effusion in 2004 occurred at rates that decreased steadily until 2008, with a maximum of < 5.9 m³s⁻¹ and a long-term average of 0.1 m³s⁻¹ (Schilling et al., 2008; Major et al., 2009). Between 1980 and 1986 magma compositions were broadly homogeneous at 62-64 wt.% SiO₂ (Cashman,

1992; Pallister et al., 1992; Blundy et al., 2008; Pallister et al., 2008). Lavas erupted between 2004 and 2008 were similarly homogenous at 63-65 wt.% SiO₂ (Blundy et al., 2008; Pallister et al., 2008). During both eruptive periods, degassing was continuous and largely coupled with magma extrusion. The largest SO₂ fluxes were associated with explosive activity in the early 1980's, when they frequently exceeded 1000 t/d (Gerlach and McGee, 1994). The lowest SO₂ fluxes (~70 t/d) were associated with the dome-building activity in 1982-86 and 2004-2008 (Gerlach and McGee, 1994; Gerlach et al., 2008). Following the cessation of each eruptive episode, SO₂ fluxes decreased rapidly to negligible levels. In the 1990's, however, detectable gas emissions (Gerlach et al., 2008) were observed concurrently with elevated shallow VT seismicity and explosive emissions of non-juvenile tephra (Mastin, 1994).

(b) *Soufrière Hills Volcano*, Montserrat (SHV), is an andesitic volcano that erupted in 1995 following several centuries of no eruptive activity. Since 1995 it has exhibited intermittent activity with five phases of eruptive activity lasting several months to years (Young et al., 1998; Sparks and Young, 2002; Wadge et al., 2010; 2014), with the last phase ending in 2010. The eruptive activity has included lava dome extrusion, block-and-ash flows and Vulcanian explosions; periods of repose have been characterised by ash venting and continuous degassing (Wadge et al., 2014). The time-averaged lava extrusion has been 3 m³s⁻¹, although rates exceeding 10 m³s⁻¹ have characterised some phases of dome extrusion (Wadge et al., 2010; Wadge et

649 al., 2014). The SiO₂ content of historically erupted products has varied from
650 58 to 62 wt.% (Murphy et al., 2000; Zellmer et al., 2003b; Barclay et al., 2010;
651 Christopher et al., 2014). The average SO₂ emission rate from 1995 to 2010
652 was ~530 t/d (Christopher et al., 2010) and largely decoupled from eruptive
653 activity (Christopher et al., 2010; Edmonds et al., 2010; Christopher et al.,
654 2015). Soufrière Hills Volcano continues to degas at ~ 430 t/d (Christopher
655 et al., 2015). During periods of intra-eruptive repose, peaks in degassing of
656 several thousand t/d have been associated with bursts in seismicity (VTs)
657 and are sometimes accompanied by ash venting (Cole et al., 2014).

658

659 (c) *Tungurahua*, Ecuador (TUN), erupted in 1999 following 81 years of no
660 eruptive activity. Slow lava extrusion and frequent explosive activity during
661 phases of eruptive activity have limited lava dome growth. Between 1999
662 and 2006 *Tungurahua* alternated between explosive (Strombolian to
663 Vulcanian) eruptions and relatively quiet periods dominated by ash venting
664 and fumarolic activity. The most explosive activity occurred during July and
665 August 2006 (Arellano et al., 2008), after which activity returned to frequent
666 low-intensity Strombolian explosions (Steffke et al., 2010). Whilst the magma
667 supply rate has varied over timescales of months (Wright et al., 2012), the
668 long-term emission rate of ash has been approximately constant at >0.2
669 m³s⁻¹, and possibly >0.4 m³s⁻¹ (Le Pennec et al., 2012). The eruptive products
670 have compositions of 56-59 wt.% SiO₂ and show no systematic variation with
671 time or eruptive style (Samaniego et al., 2011), except that major explosive

672 events in 1866 and 2006 have included a minor dacitic component
673 (Samaniego et al., 2011). Between 1999 and 2006, SO₂ fluxes varied from
674 several hundred to thousands of t/d; degassing has been largely decoupled
675 from eruptive activity (Arellano et al., 2008), although since 2006 daily SO₂
676 fluxes have decreased and appear to be better correlated with eruptive
677 activity.

678

679 (d) *Popocatepetl*, Mexico (POP), has experienced several periods of eruptive
680 activity in the 20th century. Most recently, eruptive activity was renewed in
681 1994 and has involved repeated periods of dome growth that have
682 culminated in explosive eruptions and dome collapse. Extrusion rates have
683 ranged from 0.5 to 4.1 m³s⁻¹ during dome-growth in 1996 and 1997; the
684 long-term average has been 0.24 m³s⁻¹ (Delgado-Granados et al., 2001). Prior
685 to 1995, Popocatepetl last erupted between 1920 and 1927 (Delgado-
686 Granados et al., 2001) followed by several decades of minor degassing and
687 ash venting (Brennan, 2007). Pyroclasts erupted between 1996 and 1998
688 ranged in bulk composition from ~ 59 to 64 wt.% SiO₂ (Athanasopoulos,
689 1997; Straub and Martin-Del Pozzo, 2001), with all compositions erupted
690 contemporaneously (Witter et al., 2005). In 1994, average SO₂ fluxes were
691 several thousand t/d. Similarly high SO₂ fluxes (30,000-50,000 t/d) marked
692 explosive activity between 1996 and 1998 (Goff et al., 1998; Delgado-
693 Granados et al., 2001). DOAS measurements of the plume in 2006 provide an

694 average flux of 2450 t/d, with large daily variations not always associated
695 with eruptive activity (Grutter et al., 2008).

696
697 *3.4. Non-eruptive degassing regime*

698 At volcanoes that have remained in a persistent regime throughout the 20th and 21st
699 centuries (section 3.1.1), fumarolic activity may be sustained during periods of
700 repose lasting years or even decades (e.g., Lascar; Gardeweg & Medina, 1994). One
701 volcano in our database has not erupted during the 20th and 21st centuries but has
702 exhibited sustained and persistent degassing of SO₂.

703
704 (a) *Kudryavy (Moyorodake/ Medvezhia)*, Russia (KUD), is a basaltic andesite
705 volcano that has been in a persistent state of high temperature fumarolic
706 degassing and phreatic activity since its last magmatic eruption in 1883
707 (Fischer et al., 1998; Korzhinsky et al., 2002). The only measurements come
708 from a single campaign in 1995, which measured SO₂ fluxes of 73 ±15 t/d
709 (Fischer et al., 1998).

710
711 **4. Magmatic behaviour in persistent and episodic regimes**

712
713 Geochemical analysis of erupted products and geophysical observations can provide
714 semi-empirical evidence for different magmatic processes. We summarise these
715 data for the fifteen dome-building volcanoes, with a particular focus on systematic
716 variations in the behaviour of volcanoes in the different regimes.

717

4.1. Interaction of magmas

Evidence of mixing and mingling between different batches of magma are observed in all 14 volcanoes in our database that have erupted in the 20th century (Table 2 and references therein). Different magma batches typically vary in composition, although interactions are also observed between magmas or melts that are similar in composition but differ in temperature and crystallinity (Cashman and Blundy, 2013; Costa et al., 2013; Troll et al., 2013). Evidence for magma interaction over short timescales (days to years) is ubiquitous and includes: (1) disequilibrium mineral assemblages; (2) disequilibria between mineral assemblages and matrix glass; and (3) phenocryst zoning (Table 2). Zoning patterns, in particular, provide evidence that magma mixing is sustained over a range of times. Discrete magma mixing events may be associated with single explosive events (Pallister et al., 2008; Samaniego et al., 2011; Scott et al., 2013) or individual phases of effusive activity lasting months (Dirksen et al., 2006). Frequent and near-continuous magma mixing may accompany sustained lava effusion (Nakamura, 1995; Barclay et al., 2010; Turner et al., 2013).

The degree of mixing ranges from contemporaneous eruption of different magma compositions to the eruption of lavas that are homogeneous in bulk composition but heterogeneous on a thin section scale. Evidence for incomplete mixing includes banded lava or pumice, or mafic enclaves in more silicic host lavas. Where incomplete mixing is observed, historical activity tends to be episodic with moderate to long periods of inter-eruptive repose. Persistent activity, in contrast,

tends to produce homogeneous lavas; here evidence for magma mixing is preserved only at the micro-scale, in melt inclusions, disequilibrium mineral assemblages, polymodal mineral compositions, and phenocryst zonation (Table 2).

4.2. Geophysical observations

4.2.1. Seismicity

Similar patterns of seismicity are observed across all the volcanoes in this review, with no apparent correlation with eruptive regime. Most volcanic earthquakes occur prior to and during eruptive activity. Renewed eruptive activity is generally preceded by elevated VT seismicity, with elevated LP seismicity immediately prior to eruption initiation. Levels of LP seismicity are highest at volcanoes in persistent regimes where degassing rates are high (e.g., Lascar, Popocatépetl; Asch et al., 1996). Hybrid events (LP seismicity with clear P & S wave arrivals) are commonly associated with dome-growth (e.g., Miller et al., 1998; Umakoshi et al., 2008).

Once a volcano has remained in a state of repose for more than a few months, the level of seismicity decreases, although episodic increases in VT seismicity are common and are often associated with elevated degassing and ash venting (Mastin, 1994; Ratdomopurbo et al., 2013; Budi-Santoso et al., 2013; Sernageomin, 2013; Cole et al., 2014). Seismic crises can occur during inter-eruptive repose; these may last for several months to several years with multiple felt earthquakes and no eruption of magma (Japan Meteorological Agency, 1996, Young et al., 1998).

4.2.2. Deformation

Geodetic measurements of far-field deformation are more common at volcanoes in an episodic regime than at those in a persistent regime (Table 3), although this apparent correlation could be coincidental, since many of the volcanoes in our dataset that exhibit episodic behaviour are located in developed countries, which tend to have well-established monitoring and research capabilities (e.g., USA and Japan). Alternatively, volcanoes in a persistent regime may lack far-field observations because only near-field observations are required for short-term forecasting. At episodic volcanoes, periods of repose may show inflation, whereas deflation is primarily associated with phases of dome growth (Table 3). The timescales of inflation vary from years (e.g., Augustine, Redoubt; Cervelli et al., 2010; Grapenthin et al., 2013a) to decades (e.g., Augustine, Unzen; Kohno et al., 2008; Lee et al., 2010). Soufrière Hills Volcano, which has remained in a persistent regime since 1995, also exhibits cycles of far-field inflation and deflation coincident with eruptive and non-eruptive cycles of months to years (Odbert et al., 2014a). Where persistent behaviour includes short phases of lava effusion and explosive eruption (e.g., Bezymianny, Merapi, Colima), InSAR measurements suggest negligible far-field deformation (Chaussard et al., 2013; Grapenthin et al., 2013b).

5. Conceptual magmatic models for dome-building volcanism

The interpretation of magmatic processes and their relation to volcanism requires a conceptual model for volcanic activity. From this perspective, understanding the

787 geometry of pre-eruptive magma storage is critical. A widespread, but not universal,
788 observation about dome-building volcanoes is that magma is supplied from storage
789 regions in the shallow crust (Table 5 and references therein), which has stimulated
790 models of eruptive activity modulated by shallow magma chambers (Gourgaud et
791 al., 1989; Murphy et al., 2000; Mora et al., 2002; Humphreys et al., 2008; Roberge et
792 al., 2009; Larsen et al., 2010; Samaniego et al., 2011; Shcherbakov et al., 2011;
793 Coombs et al., 2013; Turner et al., 2013). There is also evidence, however, for deeper
794 levels of magma storage, including mid- to lower crustal earthquakes associated
795 with volcanism (McNutt, 2005; Power et al., 2013), deep sources of deformation
796 (Pritchard and Simons, 2002; Elsworth et al., 2008), and deep sources of gas (Troll
797 et al., 2013; Hautmann et al., 2014; Christopher et al. 2015). Petrological and
798 geochemical data help to quantify the importance of deep igneous processes
799 (Hildreth, 2004; Troll et al., 2013; Edmonds et al., 2014), including mineral
800 assemblages that record multiple crystallisation depths (Matthews et al., 1994;
801 Martel et al., 1998; Scott et al., 2012; Cashman and Blundy, 2013; Turner et al.,
802 2013) and geochronology evidence for long crustal residence times (Volpe and
803 Hammond, 1991; Zellmer et al., 2003b; Cooper and Reid, 2008; Dosseto et al., 2008;
804 Claiborne et al., 2010). Finally, tomographic images of arc volcanoes suggest magma
805 storage occurs at different depths throughout the crust (e.g., Koulakov et al., 2013).

806
807 Here we place geochemical and geophysical evidence for transcrustal magmatic
808 systems in the context of our categorisation of temporal variations in the historical
809 records of lava dome-building volcanoes. Specifically, we address the question of the

810 extent to which observed regimes are consistent with non-linear processes
811 associated with a shallow magma chamber, or whether they require involvement of
812 vertically extensive crustal processes. Importantly, our aim is not to attribute the
813 behaviour of an individual volcano or eruptive event to either paradigm, but instead
814 to investigate the extent to which different eruptive regimes may reflect
815 fundamentally different subsurface conditions, at least with regard to the extent and
816 connectivity of individual magma lenses. We conclude that whilst storage of magma
817 in the upper crust exerts an important control on when and what eruptive activity
818 occurs, over historical timescales different patterns of volcanism can be better
819 ascribed to a conceptual model based on complex behaviours of vertically extensive
820 magma storage regions.

821 822 *5.1. Shallow chamber paradigm*

823 A common model for eruptive activity at dome-building volcanoes is a shallow melt-
824 dominated magma chamber that is replenished from depth and periodically
825 discharges magma (Fig. 7). In this paradigm, intrusion of mafic magma from depth is
826 assumed to trigger the eruption of shallow magma bodies (Gourgaud et al., 1989;
827 Murphy et al., 2000; Mora et al., 2002; Humphreys et al., 2008; Roberge et al., 2009;
828 Larsen et al., 2010; Samaniego et al., 2011; Shcherbakov et al., 2011; Coombs et al.,
829 2013; Turner et al., 2013). The concept of mafic triggers derives primarily from
830 near-ubiquitous evidence for magma mixing (Table 2). Intruding mafic magma also
831 provides an explanation for observations of excess SO₂ (that is, emission of SO₂ in
832 excess of amounts dissolved in the erupted magma; Andres et al., 1991; Wallace,

833 2003; Shinohara, 2008; Christopher et al., 2010; Wallace and Edmonds, 2011), as
834 SO₂ is much more soluble in mafic magmas than in silicic magmas (Wallace, 2005).
835 Petrologic evidence for shallow magma storage comes from saturation pressures
836 recorded in melt inclusions, as well as phase assemblages consistent with storage
837 pressures \leq 200 MPa (e.g., Moore and Carmichael, 1998; Blundy and Cashman,
838 2001; Couch et al., 2001).

839

840 The modulating effect of shallow magmatic systems on eruptive processes is
841 supported by geophysical data. Deflation during eruptive periods can be related to
842 magma discharge from upper- or mid-crustal magma chambers (Nishi et al., 1999;
843 Elsworth et al., 2008; Cervelli et al., 2010; Mattioli et al., 2010; Grapenthin et al.,
844 2013a). Furthermore, most seismicity associated with unrest and eruptive activity is
845 restricted to depths of <10 kilometres (Ratdomopurbo and Poupinet, 2000; Moran
846 et al., 2008; Power and Lalla, 2010; Thelan et al., 2010; Petrosino et al., 2011).

847 Seismicity is commonly inferred to record the stress effects of the formation of
848 magma transport pathways (Kilburn, 2003; Scandone et al., 2007) and rise of
849 magmatic fluids from shallow magma chambers (Neuberg, 2000; McNutt, 2005;
850 Chouet and Matoza, 2013). Shallow seismicity is also associated with shallow
851 magma intrusion (Moran et al., 2011), pressurisation and pre-eruptive inflation.

852

853 Patterns of recharge have been used to explain pulsatory and cyclic behaviour
854 (Melnik and Sparks, 1999; Barmin et al., 2002). Indeed it is likely that volcanism is
855 modulated, jointly, by different parts of the volcanic system, including shallow

magma chambers. However, because the mechanism for replenishment in the shallow chamber paradigm is poorly understood, it cannot completely explain the hierarchy of common behaviours and similar patterns and styles of eruptive activity.

5.2. Transcrustal destabilisation

A shallow magma chamber can be envisaged as the upper manifestation of a much larger transcrustal system (Marsh, 2000; Cañón-Tapia and Walker, 2004), which may extend throughout the crust and even into the mantle (Fig. 8). Such a conceptual model implies that mechanisms for unrest and eruption may involve more complex processes than discrete intrusions. Specifically, magmatic systems can be viewed as comprising extensive bodies of crystal-rich magma (mush) with interspersed lenses of melt and magmatic fluids that are formed by repeated intrusion of mafic melts from the mantle (Solano et al., 2012; Connolly and Podladchikov, 2013; Christopher et al. 2015). From this perspective, melt and fluid layers are susceptible to destabilisation, and reorganisation of these layers may provide a trigger for eruptive activity in mafic (Tarasewicz et al., 2012; Neave et al., 2013) and large caldera systems (Cashman and Giordano, 2014). Similarly, transcrustal processes can explain apparently anomalous activity in some dome-building volcanoes (Christopher et al., 2015), whilst also providing a source of deep magma and magmatic fluids. Key is the concept of the meta-stability of transcrustal magmatic systems and destabilisation events that involve either all or part of the melt-bearing region (Fig. 8a,b), with or without contemporaneous eruptive activity (Fig. 8c).

879

880 Temporal and spatial variations in the susceptibility of vertically extensive
881 magmatic systems to destabilisation can also explain long-term patterns of eruptive
882 activity at dome-building volcanoes. First we return to the question of mafic
883 eruption triggers, particularly as evidenced by varying intensities of
884 magma mixing in the eruptive products. Mixing has long been used to describe the
885 homogenisation of two melts, as manifested in linear two-element geochemical
886 diagrams. Mixing, however, is increasingly viewed as involving complex interactions
887 between melts and crystal mushes (Blundy et al., 2008; Humphreys et al., 2009;
888 Cashman and Blundy, 2013). From this perspective, the role of mixing as a primary
889 mechanism of eruption triggering is less clear. In fact, mixing may be an effect, as
890 much as a cause, of eruptive activity, particularly if triggered initially by
891 destabilisation of the magmatic system. Destabilisation could occur from the bottom
892 up, with deep level disturbances propagating into the upper crust (e.g., Christopher
893 et al., 2015). Alternatively destabilisation could propagate downward, driven by a
894 downward propagating decompression wave caused by early eruptive activity (e.g.,
895 Tarasewicz et al., 2012). In either case, destabilisation of a complex magmatic
896 system can force interaction among melt lenses and intervening crystal mush zones
897 (e.g., Cashman and Giordano, 2014).

898

899 Another important aspect of dome-building volcanoes in hydrous arc system relates
900 to the evolution and migration of volatiles. Fractionation of deeply sourced arc
901 basalts (Annen et al., 2006) can cause sulphur saturation of more evolved felsic

melts in the middle and lower crust (Wallace, 2005). This occurs because, although sulphur is highly soluble in basaltic melts, it is much less soluble in felsic melts (Lesne et al., 2011). As a consequence, SO₂ degassing can start deep within the crust, well below levels of shallow magma storage. The same is true of CO₂, where strong pressure-dependence may promote CO₂ exsolution throughout the crust (e.g., Blundy et al., 2010). Different volatile elements can therefore be fractionated and stored independently at multiple crustal levels during inter-eruptive periods of repose. Separation of volatiles from their parental magmas during these periods of repose can explain both the excess SO₂ degassing and decoupling of gas and magma fluxes observed in dome-building volcanoes in the persistent regime. Ascent of magmatic fluids from depth can also explain decoupling of shallow seismicity from eruptive activity (Moran, 1994; Roman et al., 2004; Girona et al., 2014; Hautmann et al., 2014; Christopher et al., 2015). Similarly, deep (20 to 40 km), long period earthquakes in arcs can be explained by exsolution and migration of insoluble gases like CO₂ (McNutt, 2005; Nichols et al., 2011). Finally, independent rise of magmatic fluids may cause the surface deformation observed at passively degassing volcanoes (Girona et al., 2014), and can help to explain varying timescales of far-field inflation at dome-building volcanoes.

5.3. Persistent dome-building behaviour

The persistent regime combines pulsatory phases of effusive eruption and homogeneous magma compositions with sustained, and decoupled, degassing (section 3.1.1), and is typical of ‘open’ system behaviour (e.g., Chaussard et al.,

925 2013). These observations appear to require a dynamically connected, through-
926 going magmatic system to sustain a persistent regime, especially over long
927 timescales. Large explosive eruptions in these systems involve magma that is more
928 mafic (deeper, more volatile-rich) than that produced during effusive activity.
929 Transitions between persistent shallow-seated effusive behaviour and intermittent
930 deep-seated explosions thus suggest that magmatic systems at these volcanoes are
931 vertically extensive and (transiently) dynamically connected, at least to mid-crustal
932 levels (Fig. 8a). More generally, rapid transport of deep, mafic and volatile-rich
933 magmas is commonly invoked for paroxysmal events at open-system basaltic
934 volcanoes (e.g., Métrich et al., 2010; Sides et al., 2014).

935

936 Eruptive activity at a second group of volcanoes in the persistent regime (section
937 3.1.2) reactivated with major explosive events that followed long periods of inter-
938 eruptive repose. In these volcanoes, the explosively erupted magma is more evolved
939 than subsequent extrusive lavas, which show gradual decreases in SiO₂ with time.
940 Progressive variation in the composition of erupted products can be explained by a
941 vertically extensive and connected magmatic system, although a more traditional
942 zoned magma chamber model (e.g., Scott et al., 2013) cannot be excluded on the
943 basis of these characteristics alone. Most important from a volcanic hazards
944 perspective, however, are the compositional homogeneity and paucity of mafic
945 enclaves (Scott et al., 2013; Turner et al., 2013) that characterise activity. This
946 suggests that these persistently active volcanoes have relatively stable magmatic

947 systems that are less susceptible to large-scale destabilisation than during inter-
948 eruptive periods of repose.

949

950 The observation that explosive eruptions may be either more or less evolved than
951 magma erupted effusively from the same system provides insight into explosive
952 eruption triggers. ‘Top-down’ destabilisation is observed in cases of edifice collapse
953 following either a long duration in a state of inter-eruptive repose (Bezymianny,
954 Santiaguito, Mount St. Helens) or sustained effusive activity and dome growth
955 (Lascar). Top-down triggering taps evolved magma from high in the crust. ‘Bottom-
956 up’ destabilisation, in contrast, explains explosive events that appear to be triggered
957 by the rapid rise of deep-derived magmas (Merapi, Colima, Shiveluch).

958

959 Persistent eruptive regimes require that the magmatic system is ‘open’, or vertically
960 connected. Under these conditions, eruptive activity may be neither strictly ‘top
961 down’ nor ‘bottom up’ but instead reflect the intrinsic instability of complex
962 magmatic systems. One mechanism of instability relates to the behaviour of crystal-
963 melt suspensions, which segregate to form separate layers of melt and/or volatiles.
964 We suggest that these (unstable) layers can reorganise rapidly to trigger abrupt
965 changes in eruption patterns. Layer destabilisation may occur because of external
966 triggers, such as regional tectonics or eruptions of neighbouring volcanoes (e.g.,
967 [Walter et al., 2007](#); [De la Cruz-Reyna et al., 2010](#); [Biggs et al., 2016](#)). Alternatively,
968 passive volatile release during a state of repose may cause the pressure distribution
969 sufficiently to cause replenishment of magma from depth ([Girona et al., 2015](#)). Such

mechanisms are not restricted to dome-building volcanoes, and have been observed at basaltic arc systems that are vertically well-connected and exhibit complex feedback mechanisms for magma discharge (e.g., Stromboli; Ripepe et al., 2015).

5.4. Episodic dome-building behaviour

Dome-building volcanoes that show episodic behaviour are characterised by diminishing eruption rates with time and correlations between lava extrusion and volatile emission. Both characteristics are indicative of closed system behaviour, which likely reflects the formation and ascent of discrete magma batches. In many of these volcanoes, however, there is evidence for the interaction of different melts (Table 3), which argues against discrete melt batches. In fact, volcanoes in an episodic regime that erupt frequently (e.g., Augustine, Redoubt) erupt a wide range of compositions during any individual eruption. This suggests that small melt batches evolve independently and interact only during eruptions (e.g., Roman et al., 2006). More homogeneous magma compositions produced by volcanoes that erupt less frequently (e.g., Mont Pelée, Unzen), in contrast, suggests that magma mixing may occur prior to, as well as during, eruptive episodes (Browne et al., 2006).

A magmatic model based on the shallow chamber paradigm suggests that if magmas are generated at a constant rate at depth, then the duration a volcano remains in a state of repose will control the volume of magma components (volatiles, melt, and crystal mush) that can accumulate; this time-dependent volume may, in turn, influence the duration a volcano remains in an eruptive state. In contrast, under the

transcrustal paradigm, variations in frequency and duration of eruptive episodes could reflect patterns of destabilisation within the deeper system. Stability may be controlled by physical properties, such as the size of magmatic systems, or fundamental parameters such as the flux of magma at depth (Caricchi et al., 2014).

5.5. Large-magnitude explosive eruptions

The dynamic nature of eruptive activity at dome-building volcanoes suggests that past behaviour is likely to influence stability of the magmatic system, and future patterns of eruptive activity. For example, edifice collapse associated with large magnitude explosions is known to reduce storage pressures (Pinel & Albino, 2013) and enable the eruption of denser, more mafic magmas, which would otherwise stall at shallow depths (Pinel & Jaupart, 2000; 2005). Indeed, volcanoes in our dataset where the onset of eruptive activity involved edifice collapse may well have shown different long-term patterns of eruptive activity if the onset of eruptive activity had been effusive. Conversely, where edifice collapse occurred after a long duration in a state of repose (~millenia), persistent activity appears to last for many decades (e.g. Bezymianny, Santiaguito; Fig. 2). Removal of the edifice during these large magnitude events thus appears to destabilise the system (Pinel & Albino, 2013).

A different situation occurred at Mount St. Helens in 1980, where the initial explosive eruption was related to edifice collapse, but the prior repose interval was only slightly more than a century. In this case, persistent behaviour continued for only six years. It is noteworthy that the volcano reactivated between 2004-2008

(Fig. 2) after two intervening episodes of inferred recharge from deeper in the system (Moran, 1994; Musumeci et al., 2002). The limited persistent activity of Mount St. Helens compared to Bezymianny and Santiaguito may be simply a result of shorter inter-eruptive repose, which could limit the accumulation of eruptible magma. Alternatively, it may be related to the dacitic composition of magma at Mount St. Helens, compared to the andesitic magmas of Bezymianny and Santiaguito.

6. Conceptualising volcanism in time

Records of eruptive activity inform our understanding of magmatic processes and are commonly the basis for forecasts of eruptive activity. Traditionally, volcanism is conceptualised as a series of discrete eruptions (Siebert et al., 2010) that are characterised by measureable properties such as magnitude, duration, intensity and eruptive style (Mercalli, 1907; Newhall and Self, 1982; Pyle, 2000). The intervals between eruptions are usually referred to as repose periods and at these times the volcano is commonly interpreted to be in a dormant state. This ontology of volcanic activity as a point process stems from geological records that comprise a punctuated series of distinct deposits, and historical records that are biased towards occasional memorable, and generally explosive, individual events (Szakács and Cañón-Tapia, 2010).

A different perspective emerges from our analysis of long-term eruptive behaviours at fifteen well-studied dome-building volcanoes. Instead of identifying discrete eruptions, we suggest that periods of eruptive activity be classified in the context of the eruptive history. For example, at two different volcanoes, periods of dome extrusion may have similar lava volumes, rates of extrusion, and duration, but can occur in very different situations (e.g., as period of episodic activity or a phase of lava extrusion in a persistent regime). Including time as a key parameter highlights the shortcomings of viewing volcanoes as in only either an “eruptive” or “non-eruptive” state. Critically, this ontology of volcanic activity should influence interpretation of both volcanic data and inferred magmatic processes.

The evidence for different states of repose provided by our case studies suggests that lava dome-building volcanoes can be characterised by three, rather than two, states: (i) a state of dormancy without abnormal geochemical or geophysical signals (inter-eruptive); (ii) an active state in which magma is erupted; and (iii) a state of unrest where perturbations in the system at depth cause marked and measurable departures from a background (dormant) state (intra-eruptive). Historical records allow volcano classification by one, two or all three of these states. Over geological timescales, we assume all volcanoes experience periods of dormancy or inter-eruptive repose periods. Intra-eruptive repose periods can be more difficult to identify, and present the greatest challenges for volcanic hazard assessment.

Inter-eruptive repose occurs at volcanoes that show episodic behaviour, meaning that they conform more closely to the traditional interpretation of volcanism as a sequence of discrete eruptions. The duration of inter-eruptive repose can vary from many years (e.g., Augustine, Redoubt) to centuries (e.g., Mount Unzen), but in all cases the volcano is deemed to be in a dormant state between eruptive periods. Volcanoes classified as dormant can move into the unrest state with increases in geophysical (e.g., seismicity, and deformation) and fumarolic activity. For example, prior to 1992, Soufrière Hills Volcano had been in a dormant state for over 350 years, but had moved into a state of unrest in 1896-97, 1933-37 and 1966-67, as evidenced by elevated fumarolic activity and intermittent seismic crises (Shepherd et al., 1971; Odbert et al., 2014b). Similar seismic crises were also observed throughout the 20th century at Mt Unzen prior to eruption onset in 1991 (Japan Meteorological Agency, 1996).

Intra-eruptive repose is observed at volcanoes in a persistent regime where intervals between pulses of eruptive activity can last for months to years or even decades, especially following major explosive events (e.g., Bezymianny, Colima, Lascar, Santiaguito). At these volcanoes, however, periods of repose are characterised by sustained degassing, intermittent seismicity and ash venting, all of which indicate magmatic unrest that is not consistent with dormancy. Importantly, unrest under these conditions does not imply imminent eruptive activity, as observed in the example of Kudryavy where a persistent state of high temperature

fumarolic degassing and phreatic activity is inferred since its last magmatic eruption in 1883 (Fischer et al., 1998; Korzhinsky et al., 2002).

By characterising exchangeable traits of volcanic behaviour, we demonstrate that the case histories in this review challenge the depiction of volcanism as a point process in time, and raise questions about what it means to say that a volcano is dormant and how to view periods of non-eruptive volcanic unrest. Importantly, several of our case study volcanoes show unrest signals that are greatly elevated after eruptive activity, in comparison to unrest signals when a volcano is in a period of longer dormancy (e.g., Merapi, Lascar, Bezymianny). For this reason, we suggest that the state of unrest be used to classify volcanic activity, with the caveat that it is important to recognise when the distinction between unrest and dormancy is determined by a change in detection thresholds and not by true changes in the state of a magmatic system.

The conceptualisation of eruptions as discrete events has been, and still is, fundamental to volcano classification, volcano databases, data selection in probabilistic forecasts and the interpretation of magmatic processes. The GVP database (Siebert et al., 2010) is the only comprehensive global compilation of active volcanoes, and is widely used to characterise volcanism, inform interpretations of volcanic processes and provide evidence for eruptive forecasts. The catalogue is predicated, however, on viewing volcanism as an alternation of two different events, repose period and eruption. The GVP further defines repose as any

cessation in eruptive activity that exceeds 3 months. This definition works well for some of our case studies (e.g., Augustine, Redoubt), but is problematic for volcanoes showing prolonged intermittent activity (e.g., Bezymianny, Mount St. Helens, Merapi, Soufrière Hills Volcano). More critically, the GVP database structure does not record information that is useful for both characterising and interpreting states of eruption and unrest.

7. Information exchangeability in forecasting volcanic activity

In recent decades probabilistic methods have become established as the principal approach to forecasting volcanic activity. Importantly, they can capture both aleatory and epistemic uncertainties and include multiple strands of evidence and different kinds of data (e.g., Newhall and Hoblitt, 2002; Aspinall et al., 2003; Marzocchi et al., 2004; Sparks and Aspinall, 2004; Neri et al., 2008; Sobradelo et al., 2013; Aspinall and Woo, 2014; Hincks et al., 2014; Sobradelo and Martí, 2015). Probabilistic approaches, however, have highlighted specific challenges associated with eruptive forecasts at dome-building volcanoes. The most acute problem relates to a lack of data, especially at volcanoes with infrequent eruptive activity in episodic regimes. The issue of sparse data, however, can also manifest at volcanoes in a persistent regime, when forecasting a long period of dormancy. Consequently, an important question in volcanology is whether observations from a number of well-studied volcanoes can be used to reduce uncertainty associated with a lack of data at an individual volcano. This is especially pertinent with the development of global

databases (e.g., Smithsonian GVP; La MEVE; WovoDAT) and global approaches to data collection (e.g., Biggs et al., 2014; Carn et al., 2016).

Importantly, the principle of using observations from multiple volcanoes requires an assumption of information or data exchangeability (e.g., Bebbington, 2014; Sheldrake, 2014). From a Bayesian perspective, exchangeability requires a (subjective) level of similarity, but importantly, does not require the behaviours of the objects to be identical (Bernado, 1996; Gelman et al., 2013). Hence, similar behaviours and traits based on phenomenological observations identified in this review could be a basis for assumptions of exchangeability.

7.1. Approaches to assuming exchangeability

One approach to the problem of limited data is through expert judgement (Aspinall and Cooke, 2013), where experienced scientists assess key parameters and likelihoods of future events based upon their own knowledge, experience and judgements. In principle, the experts should also estimate the uncertainty of their likelihood assessment (Aspinall, 2010). Issues of exchangeable data arise when comparisons with other volcanoes enter into these discussions, at least informally. In many volcano emergencies, for example, such assessments are *ad hoc* and executed largely through unstructured discussion within a volcano observatory team. These efforts can be improved by formalised methods for pooling expert judgements, as illustrated by hazard assessments for Soufrière Hills Volcano (Wadge and Aspinall, 2014). Importantly, the experience of an expert in previous volcanic

crises will likely influence their views. This illustrates a major disadvantage in the informal approach, where the basis for assessment may be anecdotal and biased towards previously witnessed discrete events. Moreover, even the most experienced volcanologist is unlikely to have witnessed more than a handful of eruptive events, so these comparisons warrant a more rigorous approach to identifying appropriate analogue volcanoes and to what extent comparisons are justified.

Broad classifications for volcano ‘type’ based on characteristics such as morphology (Rittmann, 1962; Siebert et al., 2010) or eruptive style (e.g., Hawaiian, Strombolian, Peléean, Vulcanian and Plinian; Bullard, 1962) provide a natural framework for assumptions of exchangeability. However, as the analysis in this review has outlined, the historical records of dome-building volcanoes are only partially exchangeable. Thus, whilst exchangeability may be assumed based on volcano ‘type’ (e.g., lava-dome building), the limitations and sources of aleatory uncertainty of probabilistic forecasts that arise from this assumption must be addressed by identifying both the underlying conceptual model and the common process that together form the basis for exchangeability. It is equally important to recognise key differences when applying exchangeability. This is evident in a cladistics analysis of Japanese arc volcanoes (Hone et al., 2007) that identified three broad volcano types grouped by composition, eruptive products and morphological characteristics. Differences are also identified in a study of magnitude-frequency relations that treats separately closed- and open-vent stratovolcanoes (Whelley et al., 2015).

1174 7.2. Volcanic unrest

1175 The concept of exchangeability can be used to interpret volcanic unrest, which is an
1176 almost a ubiquitous precursor to volcanic activity. Signs of unrest are typically
1177 monitored using geodetic, geophysical and geochemical surveys (e.g., Swanson et al.,
1178 1983; Sparks, 2003; Sandri et al., 2004; Jaquet et al., 2006; Chouet and Matoza,
1179 2013). Critically, these monitoring data are used to infer magmatic processes (e.g.,
1180 Voight, 1988; Kilburn, 2003; Smith et al., 2007; Lavallée et al., 2008), an approach
1181 that requires implicit, if not explicit, comparisons with unrest from previous activity.

1182

1183 The simplest approach to comparing volcanic unrest among volcanoes is to consider
1184 all signals of unrest as weakly exchangeable, with variations in the duration, pattern
1185 and occurrence the result of aleatory uncertainty, reflecting the natural variability of
1186 volcanic systems. A stronger assumption of exchangeability compares signs of
1187 unrest between volcanoes of a specific type (e.g., Phillipson et al., 2013), with the
1188 underlying assumption that different types of volcanoes should behave in similar
1189 ways. Our work shows, however, that even particular volcano ‘types’ can vary
1190 greatly in behaviour. In particular, we have shown that intra-repose unrest of a
1191 volcano in a persistent regime may reflect a very different state of activity than
1192 inter-repose unrest in the episodic regime, which may herald the onset of explosive
1193 activity. In this way, our categorization of eruptive activity at dome-building
1194 volcanoes as episodic (closed-system) or persistent (open-system) could help to
1195 further refine classifications of unrest, particularly with regard to the problem of
1196 distinguishing between non-eruptive unrest and unrest related to reawakening of a

volcano in repose (e.g., [Phillipson et al., 2013](#)). Furthermore, by attempting to understand differences in episodic and persistent behaviour in terms of magmatic processes, this provides an opportunity to interpret patterns of volcanic unrest in terms of these magmatic processes, rather than purely the outcome of eruptive activity (e.g., [Hincks et al., 2014](#)).

8. Conclusions

We have shown that dome-building volcanoes show two fundamentally different patterns of eruptive behaviours that we term episodic and persistent. Episodic behaviour is characterised by discrete episodes comprising an explosive onset followed by effusion and dome formation. In this regime, explosively erupted magma may have more evolved compositions than later-erupted lava. Excess gas emissions may be observed during explosive activity, but SO₂ fluxes are correlated with the eruption of lava and diminish to negligible levels following the end of each eruptive episode. Persistent behaviour, in contrast, is characterised by frequent (~yearly) phases of eruptive activity and sustained gas fluxes during periods of intra-eruptive repose. Erupted material is often compositionally homogeneous, except during explosive (paroxysmal) eruptions, which often involve deep, more primitive, magma compositions. Alternatively, at volcanoes that have not erupted for a long time (~millenia), large explosive Plinian eruptions can be followed by persistent behaviour where lava compositions become less evolved with time. Importantly, all volcanic activity is episodic if viewed over sufficiently long times.

1220

1221 We explain the variety of episodic and persistent behaviour through the lens of
1222 vertically extensive magmatic systems, where the extent of connectivity within the
1223 system dictates episodic or persistent behaviour (e.g., [Christopher et al., 2015](#)).
1224 Importantly, open-system behaviour involves transient, dynamically triggered
1225 magma transfer from depth but continuous gas transfer through the system.
1226 Episodic behaviour, in contrast, records eruption and gas loss from a magma batch
1227 that is quickly isolated from deeper (mid-crustal) reservoir. An interesting question
1228 relates to the importance of volatiles and volatile-rich melts in determining the
1229 stability of a magmatic system, particularly transitions between episodic and
1230 persistent regimes, and eruption triggering in episodic regimes (e.g., [Borisova et al.,](#)
1231 [2014](#); [Christopher et al., 2015](#); [Girona et al., 2015](#)).

1232

1233 From a hazard forecasting perspective, our 15 case studies show that dome-building
1234 volcanic activity cannot be characterised by a point process. This observation
1235 highlights a key ontological issue for volcanology. Discrete eruptive events can
1236 appear similar in nature in both an episodic and persistent regime, but are
1237 associated with different states of repose and long-term behaviour. Therefore, when
1238 analysing volcanic data, and interpreting magmatic processes, it is important to
1239 characterise eruptive activity in the context of the longer-term behaviour of a
1240 volcanic system. We have shown that gas data, in particular, may help to
1241 discriminate between inter- and intra-eruptive repose. Also important are patterns

of seismicity, which provide information on the depth and volume of magma storage (e.g., [White and McCausland, 2016](#)).

Also important for hazard forecasting is developing a method to determine how monitoring data from well-observed volcanoes can be used to inform interpretations of monitoring data from periods of unrest at less-studied volcanoes. Such an approach is feasible, but requires an understanding of the extent to which the monitoring data can be considered exchangeable. We suggest that exchangeability can be formalised by assessing temporal patterns in volcanic phenomena (especially relative patterns of eruption, degassing and repose), even if the datasets have different spatial and temporal data. From a theoretical standpoint, linking assumptions of exchangeability (e.g., episodic vs. persistent) to conceptual models of volcanic systems (e.g., closed vs. open) provides a mechanism to interpret monitoring data using a framework of magmatic processes.

Importantly, the approach employed in this review cannot be used to identify unique magmatic processes at individual volcanoes, and in that sense cannot replace ‘in-depth’ studies of individual volcanic systems. However, it provides a conceptual framework for interpreting common processes at dome-building volcanoes. From a broader perspective, our work demonstrates the value of constructing a hierarchical framework for volcanic activity based on exchangeable behaviours. We suggest that this approach could be extended to volcanoes with other types of characteristic activity, and thus provides a holistic approach to analysing global volcanic records.

1265

1266 *Acknowledgements*

1267 Many thanks to Prof. Jonty Rougier in the School of Mathematics, University of
1268 Bristol, who provided advice on the definition and application of exchangeability,
1269 both in a general context and more specifically at volcanoes.

1270

1271 We also thank two anonymous reviewers whose revisions and suggestions helped
1272 us more clearly explain the methodology that has been used, and clarify specific
1273 aspects of the discussion.

1274

1275 TES and RSJS were supported by a European Research Grant, Voldies. WPA was
1276 supported in part by the Natural Environment Research Council through the
1277 Consortium on Risk in the Environment: Diagnostics, Integration, Benchmarking,
1278 Learning and Elicitation (CREDIBLE; NE/J017450/1). KVC was supported by the
1279 AXA Research Fund and a Royal Society Wolfson Merit Award.

1280

1281

1282 **References:**

1283

1284 Almeev, R. R., Holtz, F., Ariskin, A. A., Kimura, J-I., 2013. Storage conditions of
1285 Bezymianny Volcano parental magmas: results of phase equilibria experiments at
1286 100 and 700 MPa. *Contributions to Mineralogy and Petrology* 166, 1389-1414.

1287 Anderson, S., Fink, J., 1990. The Development and Distribution of Surface Textures at
1288 the Mount St. Helens Dome. In: Fink, J. H. (Ed.), *Lava Flows and Domes*. Vol. 2 of
1289 *IAVCEI Proceedings in Volcanology*. Springer Berlin Heidelberg, pp. 25-46.

1290 Andreastuti, S., Alloway, B., Smith, I., 2000. A detailed tephrostratigraphic
1291 framework at Merapi Volcano, Central Java, Indonesia: implications for eruption
1292 predictions and hazard assessment. *Journal of Volcanology and Geothermal*
1293 *Research* 100, 51-67.

1294 Andres, R., Rose, W., Stoiber, R., Williams, S., Matías, O., Morales, R., 1993. A
1295 summary of sulfur dioxide emission rate measurements from Guatemalan
1296 volcanoes. *Bulletin of Volcanology* 55, 379-388.

1297 Andres, R., Rose, W., Kyle, P., DeSilva, S., Francis, P., Gardeweg, M., Roa, H. M., 1991.
1298 Excessive sulfur dioxide emissions from Chilean volcanoes. *Journal of Volcanology*
1299 *and Geothermal Research* 46, 323-329.

1300 Annen, C., Blundy, J. D., Sparks, R. S. J., 2006. The Genesis of Intermediate and Silicic
1301 Magmas in Deep Crustal Hot Zones. *Journal of Petrology* 47, 505-539.

1302 Arellano, S., Hall, M., Samaniego, P., Le Pennec, J-L., Ruiz, A., Molina, I., Yepes, H.,
1303 2008. Degassing patterns of Tungurahua volcano (Ecuador) during the 1999-2006

1304 eruptive period, inferred from remote spectroscopic measurements of SO₂
 1305 emissions. *Journal of Volcanology and Geothermal Research* 176, 151-162.

1306 Asch, G., Wylegalla, K., Hellweg, M., Seidl, D., Rademacher, H., 1996. Observations of
 1307 rapid-fire event tremor at Lascar volcano, Chile. *Annals of Geophysics* 39.

1308 Aspinall, W., 2010. A route to more tractable expert advice. *Nature* 463, 294-295.

1309 Aspinall, W., Woo, G., 2014. Santorini unrest 2011-2012: an immediate Bayesian
 1310 belief network analysis of eruption scenario probabilities for urgent decision
 1311 support under uncertainty. *Journal of Applied Volcanology* 3.

1312 Aspinall, W. P., Cooke, R. M., 2013. Expert elicitation and judgement. In: *Risk and*
 1313 *Uncertainty Assessment for Natural Hazards*. Cambridge University Press, Ch. 4, pp.
 1314 64-99.

1315 Aspinall, W., Woo, G., Voight, B., Baxter, P., 2003. Evidence-based volcanology:
 1316 application to eruption crises. *Journal of Volcanology and Geothermal Research* 128,
 1317 273-285.

1318 Athanasopoulos, P., 1997. The origin and ascent history of the 1996 dacitic dome,
 1319 Volcán Popocatepetl, Mexico. B.Sc. thesis, University of Manitoba, Winnipeg.

1320 Atlas, Z. D., Dixon, J. E., Sen, G., Finny, M., Martin-Del Pozzo, A. L., 2006. Melt
 1321 inclusions from Volcán Popocatepetl and Volcán de Colima, Mexico: Melt evolution
 1322 due to vapor-saturated crystallization during ascent. *Journal of Volcanology and*
 1323 *Geothermal Research* 153, 221-240.

1324 Auker, M., Sparks, R. S., Siebert, L., Crosweller, H. S., Ewert, J., 2013. A statistical
 1325 analysis of the global historical volcanic fatalities record. *Journal of Applied*
 1326 *Volcanology* 2.

1327 Barclay, J., Herd, R. A., Edwards, B. R., Christopher, T., Kiddle, E. J., Plail, M., Donovan,
 1328 A., 2010. Caught in the act: Implications for the increasing abundance of mafic
 1329 enclaves during the recent eruptive episodes of the Soufrière Hills Volcano,
 1330 Montserrat. *Geophysical Research Letters* 37.

1331 Barclay, J., Rutherford, M. J., Carroll, M. R., Murphy, M. D., Devine, J. D., Gardner, J.,
 1332 Sparks, R. S. J., 1998. Experimental phase equilibria constraints on pre-eruptive
 1333 storage conditions of the Soufrière Hills magma. *Geophysical Research Letters* 25,
 1334 3437-3440.

1335 Barker, S. E., Malone, S. D., 1991. Magmatic system geometry at Mount St. Helens
 1336 modeled from the stress field associated with post-eruptive earthquakes. *Journal of*
 1337 *Geophysical Research: Solid Earth* 96, 11883-11894.

1338 Barmin, A., Melnik, O., Sparks, R., 2002. Periodic behavior in lava dome eruptions.
 1339 *Earth and Planetary Science Letters* 199, 173-184.

1340 Beauducel, F., Cornet, F. H., 1999. Collection and three-dimensional modeling of GPS
 1341 and tilt data at Merapi volcano, Java. *Journal of Geophysical Research: Solid Earth*
 1342 104, 725-736.

- 1343 Bebbington, M., 2014. Long-term forecasting of volcanic explosivity. *Geophysical*
1344 *Journal International* 197, 1500-1515.
- 1345 Belousov, A., 1996. Deposits of the 30 March 1956 directed blast at Bezymianny
1346 volcano, Kamchatka, Russia. *Bulletin of Volcanology* 57, 649-662.
- 1347 Belousov, A. B., 1995. The Shiveluch volcanic eruption of 12 November 1964-
1348 explosive eruption provoked by failure of the edifice. *Journal of Volcanology and*
1349 *Geothermal Research* 66, 357-365.
- 1350 Belousov, A., Voight, B., Belousova, M., 2007. Directed blasts and blast-generated
1351 pyroclastic density currents: a comparison of the Bezymianny 1956, Mount St
1352 Helens 1980, and Soufrière Hills, Montserrat 1997 eruptions and deposits. *Bulletin*
1353 *of Volcanology* 69, 701- 740.
- 1354 Belousov, A., Voight, B., Belousova, M., Petukhin, A., 2002. Pyroclastic surges and
1355 flows from the 8-10 May 1997 explosive eruption of Bezymianny volcano,
1356 Kamchatka, Russia. *Bulletin of Volcanology* 64, 455-471.
- 1357 Bernardo, J. M. (1996). The concept of exchangeability and its applications. *Far East*
1358 *Journal of Mathematical Sciences* 4, 111-121.
- 1359 Best, N., Ashby, D., Dunstan, F., Foreman, D., McIntosh, N., 2013. A Bayesian approach
1360 to complex clinical diagnoses: a case-study in child abuse. *Journal of the Royal*
1361 *Statistical Society: Series A (Statistics in Society)* 176, 53-96.
- 1362 Biggs, J., Robertson, E., Cashman, K., 2016, The lateral extent of volcanic interactions
1363 during unrest and eruption. *Nature Geoscience* 9, 308-311.
- 1364 Biggs, J., Ebmeier, S. K., Aspinall, W. P., Lu, Z., Pritchard, M. E., Sparks, R. S. J., Mather,
1365 T. A., 2014. Global link between deformation and volcanic eruption quantified by
1366 satellite imagery. *Nature Communications* 5.
- 1367 Blundy, J., Cashman, K., Rust, A., Witham, F., 2010. A case for CO₂-rich arc magmas.
1368 *Earth and Planetary Science Letters* 290, 289-301.
- 1369 Blundy, J., Cashman, K. V., Berlo, K., 2008. Evolving Magma Storage Conditions
1370 Beneath Mount St. Helens Inferred from Chemical Variations in Melt Inclusions from
1371 the 1980-1986 and Current (2004-2006) Eruptions. In: Sherrod, D. R., Scott, W. E.,
1372 Stauffer, P. H. (Eds.), *A Volcano Rekindled: The Renewed Eruption of Mount St.*
1373 *Helens, 2004-2006. U.S. Geological Survey Professional Paper* 1750, Ch. 33, pp. 755-
1374 790.
- 1375 Blundy, J., Cashman, K., 2001. Ascent-driven crystallisation of dacite magmas at
1376 Mount St Helens, 1980-1986. *Contributions to Mineralogy and Petrology* 140, 631-
1377 650.
- 1378 Bogoyavlenskaya, G., Braitseva, O., Melekestsev, I., Kiriyanov, V., Miller, C. D., 1985.
1379 Catastrophic eruptions of the directed-blast type at Mount St. Helens, Bezymianny
1380 and Shiveluch volcanoes. *Journal of Geodynamics* 3, 189-218.
- 1381 Borgia, A., Aubert, M., Merle, O., van Wyk de Vries, B., 2010. What is a volcano?
1382 *Geological Society of America Special Papers* 470, 1-9.

- 1383 Borisova, A Y., Toutain, J-P, Dubessy, J., Pallister, J., Zwick, A., Salvi, S., 2014. H₂O-CO₂-
1384 S fluid triggering the 1991 Mount Pinatubo climactic eruption (Philippines). *Bulletin*
1385 *of Volcanology* 76.
- 1386 Bullard, F. M., 1962. *Volcanoes in History, in Theory, in Eruption*. Austin, University
1387 of Texas Press.
- 1388 Brennan, C., 2007. *The far side of the sky*. Dankat Publishing.
- 1389 Browne, B. L., Eichelberger, J. C., Patino, L. C., Vogel, T. A., Dehn, J., Uto, K., Hoshizumi,
1390 H., 2006. Generation of Porphyritic and Equigranular Mafic Enclaves During Magma
1391 Recharge Events at Unzen Volcano, Japan. *Journal of Petrology* 47, 301-328.
- 1392 Budi-Santoso, A., Lesage, P., Dwiyono, S., Sumarti, S., Subandriyo, Surono, Jousset, P.,
1393 Metaxian, J-P., 2013. Analysis of the seismic activity associated with the 2010
1394 eruption of Merapi Volcano, Java. *Journal of Volcanology and Geothermal Research*
1395 261, 153-170.
- 1396 Bull, K. F., Buurman, H., 2013. An overview of the 2009 eruption of Redoubt Volcano,
1397 Alaska. *Journal of Volcanology and Geothermal Research* 259, 2-15.
- 1398 Cabral-Cano, E., Correa-Mora, F., Meertens, C., 2008. Deformation of Popocatepetl
1399 volcano using GPS: Regional geodynamic context and constraints on its magma
1400 chamber. *Journal of Volcanology and Geothermal Research* 170, 24-34.
- 1401 Calder, E. S., Lavallée, Y., Kendrick, J. E., Bernstein, M., 2015. Chapter 18-Lava Dome
1402 Eruptions . In: Sigurdsson, H. et al., (Eds.), *The Encyclopedia of Volcanoes*, 2nd
1403 Edition. Academic Press, pp. 343-362.
- 1404 Cañón-Tapia, E., Walker, G. P. L., 2004. Global aspects of volcanism: the perspectives
1405 of “plate tectonics” and “volcanic systems”. *Earth-Science Reviews* 66, 163-182.
- 1406 Caricchi, L., Annen, C., Blundy, J., Simpson, G., Pinel, V., 2014. Frequency and
1407 magnitude of volcanic eruptions controlled by magma injection and buoyancy.
1408 *Nature Geoscience* 7, 126- 130.
- 1409 Carn, S. A., Clarisse, L., Prata, A. J., 2016. Multi-decadal satellite measurements of
1410 global volcanic degassing. *Journal of Volcanology and Geothermal Research* 311, 99-
1411 134.
- 1412 Carn, S. A., Krotkov, N. A., Yang, K., Krueger, A. J., 2013. Measuring global volcanic
1413 degassing with the Ozone Monitoring Instrument (OMI). Geological Society, London,
1414 Special Publications 380.
- 1415 Carn, S. A., Prata, F. J., 2010. Satellite-based constraints on explosive SO₂ release
1416 from Soufrière Hills Volcano, Montserrat. *Geophysical Research Letters* 37 (19).
- 1417 Casadevall, T., Doukas, M., Neal, C., McGimsey, R., Gardner, C., 1994. Emission rates of
1418 sulfur dioxide and carbon dioxide from Redoubt Volcano, Alaska, during the 1989 -
1419 1990 eruptions. *Journal of Volcanology and Geothermal Research* 62, 519-530.
- 1420 Casadevall, T. J., Rose, W. I., Fuller, W. H., Hunt, W. H., Hart, M. A., Moyers, J. L.,
1421 Woods, D. C., Chuan, R. L., Friend, J. P., 1984. Sulfur dioxide and particles in quiescent

1422 volcanic plumes from Poás, Arenal, and Colima Volcanos, Costa Rica and Mexico.
1423 *Journal of Geophysical Research: Atmospheres* 89, 9633-9641.

1424 Casertano, L., 1963. Catalogue of the active volcanoes of the world; Part XV, Chilean
1425 continent. IAVCEI 55pp.

1426 Cashman, K., Biggs, J., 2014. Common processes at unique volcanoes - a
1427 volcanological conundrum. *Frontiers in Earth Science* 2.

1428 Cashman, K. V., Giordano, G., 2014. Calderas and magma reservoirs. *Journal of*
1429 *Volcanology and Geothermal Research* 288, 28-45.

1430 Cashman, K., Blundy, J., 2013. Petrological cannibalism: the chemical and textural
1431 consequences of incremental magma body growth. *Contributions to Mineralogy and*
1432 *Petrology* 166, 703-729.

1433 Cashman, K. V., Sparks, R. S. J., 2013. How volcanoes work: A 25 year perspective.
1434 *Geological Society of America Bulletin*.

1435 Cashman, K. V., Thornber, C. R., Pallister, J. S., 2008. From Dome to Dust: Shallow
1436 Crystallization and Fragmentation of Conduit Magma During the 2004-2006 Dome
1437 Extrusion of Mount St. Helens, Washington. In: Sherrod, D. R., Scott, W. E., Stauffer, P.
1438 H. (Eds.), *A Volcano Rekindled: The Renewed Eruption of Mount St. Helens, 2004-*
1439 *2006. U.S. Geological Survey Professional Paper 1750, Ch. 19, pp. 387-414.*

1440 Cashman, K., McConnell, S., 2005. Multiple levels of magma storage during the 1980
1441 summer eruptions of Mount St. Helens, WA. *Bulletin of Volcanology* 68, 57-75.

1442 Cashman, K. V., 1992. Groundmass crystallization of Mount St. Helens dacite, 1980-
1443 1986: a tool for interpreting shallow magmatic processes. *Contributions to*
1444 *Mineralogy and Petrology* 109, 431-449.

1445 Cervelli, P. F., Fournier, T. J., Freymueller, J. T., Power, J. A., Lisowski, M., Pauk, B. A.,
1446 2010. Geodetic Constraints on Magma Movement and Withdrawal During the 2006
1447 Eruption of Augustine Volcano. In: Power, J. A., Coombs, M. L., Freymueller, J. T.
1448 (Eds.), *The 2006 Eruption of Augustine Volcano, Alaska. U.S. Geological Survey*
1449 *Professional Paper 1769, Ch. 17, pp. 427-452.*

1450 Chaussard, E., Amelung, F., Aoki, Y., 2013. Characterization of open and closed
1451 volcanic systems in Indonesia and Mexico using InSAR time series. *Journal of*
1452 *Geophysical Research: Solid Earth* 118, 3957-3969.

1453 Chouet, B. A., 1996. Long-period volcano seismicity: its source and use in eruption
1454 forecasting. *Nature* 380, 309-316.

1455 Chouet, B. A., Matoza, R. S., 2013. A multi-decadal view of seismic methods for
1456 detecting precursors of magma movement and eruption. *Journal of Volcanology and*
1457 *Geothermal Research* 252, 108-175.

1458 Christopher, T., Blundy, J., Cashman, K., Cole, P., Edmonds, M., Smith, P., R.S.J, S.,
1459 Stinton, A., 2015. Crustal-scale degassing due to magma system destabilisation and
1460 magma-gas decoupling at Soufrière Hills Volcano, Montserrat. *Geochemistry,*
1461 *Geophysics, Geosystems* 16, 2797-2811.

1462 Christopher, T. E., Humphreys, M. C. S., Barclay, J., Genareau, K., De Angelis, S. M. H.,
1463 Plail, M., Donovan, A., 2014. Petrological and geochemical variation during the
1464 Soufrière Hills eruption, 1995 to 2010. Geological Society, London, Memoirs 39, 317-
1465 342.

1466 Christopher, T., Edmonds, M., Humphreys, M., Herd, R., 2010. Volcanic gas emissions
1467 from Soufrière Hills Volcano, Montserrat 1995-2009, with implications for mafic
1468 magma supply and degassing. Geophysical Research Letters 37.

1469 Claiborne, L. L., Miller, C. F., Flanagan, D. M., Clynne, M. A., Wooden, J. L., 2010. Zircon
1470 reveals protracted magma storage and recycling beneath Mount St. Helens. *Geology*
1471 38, 1011- 1014.

1472 Cole, P. D., Smith, P., Komorowski, J-C., Alfano, F., Bonadonna, C., Stinton, A.,
1473 Christopher, T., Odbert, H. M., Loughlin, S., 2014. Ash venting occurring both prior to
1474 and during lava extrusion at Soufrière Hills Volcano, Montserrat, from 2005 to 2010.
1475 Geological Society, London, Memoirs 39, 71-92.

1476 Coles, S. G., Sparks, R. S. J., 2006. Extreme value methods for modelling historical
1477 series of large volcanic magnitudes. In: Mader, H. M., Coles, S. G., Connor, C. B.,
1478 Connor, L. J. (Eds.), *Statistics in Volcanology*. Vol. 1 of Special Publications of IAVCEI.
1479 Geological Society, London, pp. 47-56.

1480 Connolly, J., Podladchikov, Y., 2013. A Hydromechanical Model for Lower Crustal
1481 Fluid Flow. In: *Metasomatism and the Chemical Transformation of Rock*. Lecture
1482 Notes in Earth System Sciences. Springer Berlin Heidelberg, pp. 599-658.

1483 Coombs, M. L., Sisson, T. W., Bleick, H. A., Henton, S. M., Nye, C. J., Payne, A. L.,
1484 Cameron, C. E., Larsen, J. F., Wallace, K. L., Bull, K. F., 2013. Andesites of the 2009
1485 eruption of Redoubt Volcano, Alaska. *Journal of Volcanology and Geothermal*
1486 *Research* 259, 349-372.

1487 Coombs, M. L., Bull, K. F., Vallance, J. W., Schneider, D. J., Thoms, E. E., Wessels, R. L.,
1488 McGimsey, R. G., 2010. Timing, Distribution, and Volume of Proximal Products of the
1489 2006 Eruption of Augustine Volcano. In: Power, J. A., Coombs, M. L., Freymueller, J. T.
1490 (Eds.), *The 2006 Eruption of Augustine Volcano, Alaska*. U.S. Geological Survey
1491 Professional Paper 1769, Ch. 8, pp. 145-186.

1492 Cooper, K. M., Reid, M. R., 2008. Uranium-series Crystal Ages. *Reviews in Mineralogy*
1493 *and Geochemistry* 69, 479-544.

1494 Costa, F., Andreastuti, S., de Maisonrouve, C. B., Pallister, J. S., 2013. Petrological
1495 insights into the storage conditions and magmatic processes that yielded the
1496 centennial 2010 Merapi explosive eruption. *Journal of Volcanology and Geothermal*
1497 *Research* 261, 209-235.

1498 Couch, S., Sparks, R., Carroll, M., 2001. Mineral disequilibrium in lavas explained by
1499 convective self-mixing in open magma chambers. *Nature* 411, 1037-1039.

1500 Crummy, J. M., Savov, I. P., Navarro-Ochoa, C., Morgan, D. J., Wilson, M., 2014. High-K
1501 Mafic Plinian Eruptions of Volcán de Colima, Mexico. *Journal of Petrology* 55, 2155-
1502 2192.

1503 De la Cruz-Reyna, S., Tárrega, M., Ortiz, R., Martínez-Bringas, A., 2010. Tectonic
 1504 earthquakes triggering volcanic seismicity and eruptions. Case studies at
 1505 Tungurahua and Popocatépetl volcanoes. *Journal of Volcanology and Geothermal*
 1506 *Research* 193, 37-48.

1507 Delgado-Granados, H., González, L. C., Sánchez, N. P., 2001. Sulfur dioxide emissions
 1508 from Popocatépetl volcano (Mexico): case study of a high-emission rate, passively
 1509 degassing erupting volcano. *Journal of Volcanology and Geothermal Research* 108,
 1510 107-120.

1511 Deligne, N. I., Coles, S. G., Sparks, R. S. J., 2010. Recurrence rates of large explosive
 1512 volcanic eruptions. *Journal of Geophysical Research: Solid Earth* 115.

1513 Deruelle, B., 1985. Le Volcan Lascar: Geologie et Petrologie. IV Congreso Geologico
 1514 Chileno, Agosto 1985.

1515 DeShon, H. R., Thurber, C. H., Rowe, C., 2007. High-precision earthquake location and
 1516 threedimensional P wave velocity determination at Redoubt Volcano, Alaska.
 1517 *Journal of Geophysical Research: Solid Earth* 112.

1518 Diefenbach, A. K., Bull, K. F., Wessels, R. L., McGimsey, R. G., 2013. Photogrammetric
 1519 monitoring of lava dome growth during the 2009 eruption of Redoubt Volcano.
 1520 *Journal of Volcanology and Geothermal Research* 259, 308-316.

1521 Dirksen, O., Humphreys, M., Pletchov, P., Melnik, O., Demyanchuk, Y., Sparks, R.,
 1522 Mahony, S., 2006. The 2001-2004 dome-forming eruption of Shiveluch volcano,
 1523 Kamchatka: Observation, petrological investigation and numerical modelling.
 1524 *Journal of Volcanology and Geothermal Research* 155, 201-226.

1525 Divoux, T., Bertin, E., Vidal, V., Géménar, J.C., 2009. Intermittent outgassing through a
 1526 non-Newtonian fluid. *Physical Review E* 79.

1527 Dosseto, A., Turner, S. P., Sandiford, M., Davidson, J., 2008. Uranium-series isotope
 1528 and thermal constraints on the rate and depth of silicic magma genesis. *Geological*
 1529 *Society, London, Special Publications* 304, 169-181.

1530 Doukas, M., 1995. A compilation of sulfur dioxide and carbon dioxide emission-rate
 1531 data from Cook Inlet volcanoes (Redoubt, Spurr, Iliamna, and Augustine), Alaska
 1532 during the period from 1990 to 1994. Tech. rep., U.S. Geological Survey Open-File
 1533 Report OF 95-0055, 15 pp.

1534 Ebmeier, S., Biggs, J., Mather, T., Elliott, J., Wadge, G., Amelung, F., 2012. Measuring
 1535 large topographic change with InSAR: Lava thicknesses, extrusion rate and
 1536 subsidence rate at Santiaguito volcano, Guatemala. *Earth and Planetary Science*
 1537 *Letters* 335-336, 216-225.

1538 Edmonds, M., Humphreys, M. C. S., Hauri, E. H., Herd, R. A., Wadge, G., Rawson, H.,
 1539 Ledden, R., Plail, M., Barclay, J., Aiuppa, A., Christopher, T. E., Giudice, G., Guida, R.,
 1540 2014. Chapter 16: Pre-eruptive vapour and its role in controlling eruption style and
 1541 longevity at Soufrière Hills Volcano. *Geological Society, London, Memoirs* 39, 291-
 1542 315.

1543 Edmonds, M., Aiuppa, A., Humphreys, M., Moretti, R., Giudice, G., Martin, R. S., Herd,
1544 R. A., Christopher, T., 2010. Excess volatiles supplied by mingling of mafic magma at
1545 an andesite arc volcano. *Geochemistry, Geophysics, Geosystems* 11.

1546 Edmonds, M., Herd, R., Galle, B., Oppenheimer, C., 2003. Automated, high time-
1547 resolution measurements of SO₂ flux at Soufrière Hills Volcano, Montserrat. *Bulletin*
1548 *of Volcanology* 65, 578-586.

1549 Elsworth, D., Mattioli, G., Taron, J., Voight, B., Herd, R., 2008. Implications of Magma
1550 Transfer Between Multiple Reservoirs on Eruption Cycling. *Science* 322, 246-248.

1551 Engberg, E., 2009. SO₂ Emissions at Volcan de Colima, 2003-2007. Master's Thesis,
1552 Michigan Technological University.

1553 Fichaut, M., Marcelot, G., Clocchiatti, R., 1989a. Magmatology of Mt. Pelée
1554 (Martinique, F.W.I.). II: petrology of gabbroic and dioritic cumulates. *Journal of*
1555 *Volcanology and Geothermal Research* 38, 171-187.

1556 Fichaut, M., Maury, R., Traineau, H., Westercamp, D., Joron, J., Gourgaud, A., Coulon,
1557 C., 1989b. Magmatology of Mt. Pelée (Martinique, F.W.I.). III: Fractional
1558 crystallization versus magma mixing. *Journal of Volcanology and Geothermal*
1559 *Research* 38, 189-213.

1560 Fischer, T. P., Giggenbach, W. F., Sano, Y., Williams, S. N., 1998. Fluxes and sources of
1561 volatiles discharged from Kudryavy, a subduction zone volcano, Kurile Islands. *Earth*
1562 *and Planetary Science Letters* 160, 81-96.

1563 Galle, B., Oppenheimer, C., Geyer, A., McGonigle, A. J., Edmonds, M., Horrocks, L.,
1564 2003. A miniaturised ultraviolet spectrometer for remote sensing of SO₂ fluxes: a
1565 new tool for volcano surveillance. *Journal of Volcanology and Geothermal Research*
1566 119, 241-254.

1567 Gardeweg, M. C., Medina, E., 1994. La erupcion subpliniana del 19-20 de Abril de
1568 1993 del Volcan Lascar, N de Chile. *Actas 7th Congreso Geologico Chileno*, Santiago,
1569 1:299-304.

1570 Gelman, A., Carlin, J. B., Stern, H. S., Rubin, D. B., 2004. *Bayesian Data Analysis*. 2nd
1571 Edition. Chapman & Hall/CRC, Boca Raton.

1572 Gerlach, T. M., McGee, K. A., Doukas, M. P., 2008. Use of Digital Aerophotogrammetry
1573 to Determine Rates of Lava Dome Growth, Mount St. Helens, Washington, 2004-
1574 2005. In: Sherrod, D. R., Scott, W. E., Stauffer, P. H. (Eds.), *A Volcano Rekindled: The*
1575 *Renewed Eruption of Mount St. Helens, 2004-2006*. U.S. Geological Survey
1576 Professional Paper 1750, Ch. 26, pp. 543-572.

1577 Gerlach, T. M., McGee, K. A., 1994. Total sulfur dioxide emissions and pre-eruption
1578 vaporsaturated magma at Mount St. Helens, 1980-88. *Geophysical Research Letters*
1579 21, 2833-2836.

1580 Gertisser, R., Keller, J., 2003. Temporal variations in magma composition at Merapi
1581 Volcano (Central Java, Indonesia): magmatic cycles during the past 2000 years of
1582 explosive activity. *Journal of Volcanology and Geothermal Research* 123, 1-23.

1583 Geschwind, C-H., Rutherford, M., 1995. Crystallization of microlites during magma
 1584 ascent: the fluid mechanics of 1980-1986 eruptions at Mount St Helens. *Bulletin of*
 1585 *Volcanology* 57, 356-370.

1586 Giggenbach, W., 1996. Chemical Composition of Volcanic Gases. In: *Monitoring and*
 1587 *Mitigation of Volcano Hazards*. Springer Berlin Heidelberg, pp. 221-256.

1588 Girona, T., Costa, F., Schubert, G., 2015. Degassing during quiescence as a trigger of
 1589 magma ascent and volcanic eruptions. *Scientific Reports* 5.

1590 Girona, T., Costa, F., Newhall, C., Taisne, B., 2014. On depressurization of volcanic
 1591 magma reservoirs by passive degassing. *Journal of Geophysical Research: Solid*
 1592 *Earth* 119, 8667- 8687.

1593 Glicken, H., 1998. Rockslide-debris avalanche of May 18, 1980, Mount St. Helens
 1594 volcano. Washington: *Bulletin of the Geological Survey of Japan* 49, 55-106.

1595 Goff, F., Janik, C. J., Delgado, H., Werner, C., Counce, D., Stimac, J. A., Siebe, C., Love, S.
 1596 P., Williams, S. N., Fischer, T., Johnson, L., 1998. Geochemical surveillance of
 1597 magmatic volatiles at Popocatepetl volcano, Mexico. *Geological Society of America*
 1598 *Bulletin* 110, 695-710.

1599 González, M. B., Ramírez, J. J., Navarro, C., 2002. Summary of the historical eruptive
 1600 activity of Volcán De Colima, Mexico 1519-2000. *Journal of Volcanology and*
 1601 *Geothermal Research* 117, 21-46.

1602 Gorbach, N., Portnyagin, M., 2011. Geology and petrology of the lava complex of
 1603 Young Shiveluch Volcano, Kamchatka. *Petrology* 19, 134-166.

1604 Gorelchik, V., Shirokov, V., Firstov, P., Chubarova, O., 1997. Shiveluch volcano:
 1605 seismicity, deep structure and forecasting eruptions (Kamchatka). *Journal of*
 1606 *Volcanology and Geothermal Research* 78, 121-137.

1607 Gorshkov, G., 1959. Gigantic eruption of the volcano Bezymianny. *Bulletin*
 1608 *Volcanologique* 20, 77-109.

1609 Gottsmann, J., Odbert, H., 2014. The effects of thermomechanical heterogeneities in
 1610 island arc crust on time-dependent preeruptive stresses and the failure of an
 1611 andesitic reservoir. *Journal of Geophysical Research: Solid Earth* 119, 4626-4639.

1612 Gourgaud, A., Fichaut, M., Joron, J-L., 1989. Magmatology of Mt. Pelée (Martinique,
 1613 F.W.I.). I: Magma mixing and triggering of the 1902 and 1929 Pelean nuées ardentes.
 1614 *Journal of Volcanology and Geothermal Research* 38, 143-169.

1615 Grapenthin, R., Freymueller, J. T., Kaufman, A. M., 2013a. Geodetic observations
 1616 during the 2009 eruption of Redoubt Volcano, Alaska. *Journal of Volcanology and*
 1617 *Geothermal Research* 259, 115-132.

1618 Grapenthin, R., Freymueller, J. T., Serovetnikov, S. S., 2013b. Surface deformation of
 1619 Bezymianny Volcano, Kamchatka, recorded by GPS: the eruptions from 2005 to
 1620 2010 and long-term, long-wavelength subsidence. *Journal of Volcanology and*
 1621 *Geothermal Research* 263, 58-74.

1622 Grutter, M., Basaldud, R., Rivera, C., Harig, R., Junkerman, W., Caetano, E., Delgado-
 1623 Granados, H., 2008. SO₂ emissions from Popocatepetl volcano: emission rates and
 1624 plume imaging using optical remote sensing techniques. *Atmospheric Chemistry and*
 1625 *Physics* 8, 6655- 6663.

1626 Hammer, J., Cashman, K., Voight, B., 2000. Magmatic processes revealed by textural
 1627 and compositional trends in Merapi dome lavas. *Journal of Volcanology and*
 1628 *Geothermal Research* 100, 165-192.

1629 Harris, A. J., Rose, W. I., Flynn, L. P., 2003. Temporal trends in lava dome extrusion at
 1630 Santiaguito 1922-2000. *Bulletin of Volcanology* 65, 77-89.

1631 Harris, G. W., 1994. The petrology and petrography of lava from the 1986 eruption
 1632 of Augustine volcano. Master's Thesis, Fairbanks, Alaska, 131 p.

1633 Hautmann, S., Witham, F., Christopher, T., Cole, P., Linde, A. T., Sacks, I. S., Sparks, R.
 1634 S. J., 2014. Strain field analysis on Montserrat (W.I.) as tool for assessing permeable
 1635 flow paths in the magmatic system of Soufrière Hills Volcano. *Geochemistry,*
 1636 *Geophysics, Geosystems* 15, 676-690.

1637 Henney, L., Rodríguez, L., Watson, I., 2012. A comparison of SO₂ retrieval techniques
 1638 using mini-UV spectrometers and ASTER imagery at Lascar volcano, Chile. *Bulletin*
 1639 *of Volcanology* 74, 589-594.

1640 Hildreth, W., 2004. Volcanological perspectives on Long Valley, Mammoth Mountain,
 1641 and Mono Craters: several contiguous but discrete systems. *Journal of Volcanology*
 1642 *and Geothermal Research* 136, 169-198.

1643 Hincks, T., Komorowski, J.-C., Sparks, S., Aspinall, W., 2014. Retrospective analysis of
 1644 uncertain eruption precursors at La Soufrière volcano, Guadeloupe, 1975-77:
 1645 volcanic hazard assessment using a Bayesian Belief Network approach. *Journal of*
 1646 *Applied Volcanology* 3, 3.

1647 Hirabayashi, J., Ohba, T., Nogami, K., Yoshida, M., 1995. Discharge rate of SO₂ from
 1648 Unzen volcano, Kyushu, Japan. *Geophysical Research Letters* 22, 1709-1712.

1649 Hobbs, P. V., Radke, L. F., Lyons, J. H., Ferek, R. J., Coffman, D. J., Casadevall, T. J., 1991.
 1650 Airborne measurements of particle and gas emissions from the 1990 volcanic
 1651 eruptions of Mount Redoubt. *Journal of Geophysical Research: Atmospheres* 96,
 1652 18735-18752.

1653 Hone, D., Mahony, S., Sparks, R., Martin, K., 2007. Cladistic analysis applied to the
 1654 classification of volcanoes. *Bulletin of Volcanology* 70, 203-220.

1655 Humaida, H., 2008. SO₂ Emission Measurement By Doas (Differential Optical
 1656 Absorption Spectroscopy) and Cospec (Correlation Spectroscopy) At Merapi
 1657 Volcano (Indoensia). *Indonesian Journal of Chemistry* 8, 151-157.

1658 Humphreys, M., Christopher, T., Hards, V., 2009. Microlite transfer by disaggregation
 1659 of mafic inclusions following magma mixing at Soufrière Hills volcano, Montserrat.
 1660 *Contributions to Mineralogy and Petrology* 157, 609-624.

- 1661 Humphreys, M., Blundy, J., Sparks, R., 2008. Shallow-level decompression
1662 crystallisation and deep magma supply at Shiveluch Volcano. *Contributions to*
1663 *Mineralogy and Petrology* 155, 45-61.
- 1664 Humphreys, M. C. S., Blundy, J. D., Sparks, R. S. J., 2006. Magma Evolution and Open-
1665 System Processes at Shiveluch Volcano: Insights from Phenocryst Zoning. *Journal of*
1666 *Petrology* 47, 2303-2334.
- 1667 Japan Meteorological Agency, 1996. Unzendake, National Catalogue of the Active
1668 Volcanoes in Japan, second Edition. Japan Meteorological Agency, in Japanese.
- 1669 Jaquet, O., Carniel, R., Sparks, S., Thompson, G., Namar, R., Cecca, M. D., 2006. DEVIN:
1670 A forecasting approach using stochastic methods applied to the Soufrière Hills
1671 Volcano. *Journal of Volcanology and Geothermal Research* 153, 97-111.
- 1672 Jaupart, C., Allègre, C. J., 1991. Gas content, eruption rate and instabilities of eruption
1673 regime in silicic volcanoes. *Earth and Planetary Science Letters* 102, 413-429.
- 1674 Jaupart, C., Vergnolle, S., 1989. The generation and collapse of a foam layer at the
1675 roof of a basaltic magma chamber. *Journal of Fluid Mechanics*, 203, 347-380.
- 1676 Kienle, J., Lalla, D., Pearson, C., Barrett, S., 1979. Search for shallow magma
1677 accumulations at Augustine volcano. Tech. rep., Geophysical Institute, University of
1678 Alaska Fairbanks, final report to U.S. Department of Energy, 157 pp.
- 1679 Kilburn, C. R., 2003. Multiscale fracturing as a key to forecasting volcanic eruptions.
1680 *Journal of Volcanology and Geothermal Research* 125, 271-289.
- 1681 Kohno, Y., Matsushima, T., Shimizu, H., 2008. Pressure sources beneath Unzen
1682 Volcano inferred from leveling and GPS data. *Journal of Volcanology and Geothermal*
1683 *Research* 175, 100-109.
- 1684 Korzhinsky, M. A., Botcharnikov, R. E., Tkachenko, S. I., Steinberg, G. S., 2002. Decade-
1685 long study of degassing at Kudriavyy volcano, Iturup, Kurile Islands (1990-1999): Gas
1686 temperature and composition variations, and occurrence of 1999 phreatic eruption.
1687 *Earth, Planets and Space* 54, 337-347.
- 1688 Koulakov, I., Gordeev, E. I., Dobretsov, N. L., Vernikovskiy, V. A., Senyukov, S.,
1689 Jakovlev, A., Jaxybulatov, K., 2013. Rapid changes in magma storage beneath the
1690 Klyuchevskoy group of volcanoes inferred from time-dependent seismic
1691 tomography. *Journal of Volcanology and Geothermal Research* 263, 75-91, Magma
1692 System Response to Edifice Collapse.
- 1693 Lacroix, A., 1904. La Montagne Pelée et ses éruptions. Masson et Cie, Paris.
- 1694 Larsen, J. F., Nye, C. J., Coombs, M. L., Tilman, M., Izbekov, P., Cameron, C., 2010.
1695 Petrology and Geochemistry of the 2006 Eruption of Augustine Volcano. In: Power, J.
1696 A., Coombs, M. L., Freymueller, J. T. (Eds.), *The 2006 Eruption of Augustine Volcano,*
1697 *Alaska. U.S. Geological Survey Professional Paper* 1769, Ch. 15, pp. 335-382.
- 1698 Lavallée, Y., Meredith, P. G., Dingwell, D. B., Hess, K-U., Wassermann, J., Cordonnier,
1699 B., Gerik, A., Kruhl, J. H., 2008. Seismogenic lavas and explosive eruption forecasting.
1700 *Nature* 453, 507-510.

1701 Le Pennec, J-L., Ruiz, G. A., Ramón, P., Palacios, E., Mothes, P., Yepes, H., 2012. Impact
 1702 of tephra falls on Andean communities: The influences of eruption size and weather
 1703 conditions during the 1999-2001 activity of Tungurahua volcano, Ecuador. *Journal*
 1704 *of Volcanology and Geothermal Research* 217-218, 91-103.

1705 Lee, C-W., Lu, Z., Jung, H-S., Won, J-S., Dzurisin, D., 2010. Surface Deformation of
 1706 Augustine Volcano, 1992-2005, from Multiple-Interferogram Processing Using a
 1707 Refined Small Baseline Subset (SBAS) Interferometric Synthetic Aperture Radar
 1708 (InSAR) Approach. In: Power, J. A., Coombs, M. L., Freymueller, J. T. (Eds.), *The 2006*
 1709 *Eruption of Augustine Volcano, Alaska*. U.S. Geological Survey Professional Paper
 1710 1769, Ch. 18, pp. 453-467.

1711 Lesne, P., Kohn, S. C., Blundy, J., Witham, F., Botcharnikov, R. E., Behrens, H., 2011.
 1712 Experimental Simulation of Closed-System Degassing in the System Basalt-H₂O-CO₂-
 1713 S-Cl. *Journal of Petrology* 52, 1737-1762.

1714 Lipman, P. W., Mullineaux, D. R., 1981. *The 1980 Eruptions of Mount St. Helens,*
 1715 *Washington*. U.S. 1250. Geological Survey Professional Paper.

1716 Lisowski, M., Dzurisin, D., Denlinger, R. P., Iwatsubo, E. Y., 2008. Analysis of GPS-
 1717 Measured Deformation Associated with the 2004-2006 Dome-Building Eruption of
 1718 Mount St. Helens, Washington. In: Sherrod, D. R., Scott, W. E., Stauffer, P. H. (Eds.), *A*
 1719 *Volcano Rekindled: The Renewed Eruption of Mount St. Helens, 2004-2006*. U.S.
 1720 Geological Survey Professional Paper 1750, Ch. 15, pp. 301-334.

1721 Lopez, T., Ushakov, S., Izbekov, P., Tassi, F., Cahill, C., Neill, O., Werner, C., 2013.
 1722 Constraints on magma processes, subsurface conditions, and total volatile flux at
 1723 Bezymianny Volcano in 2007-2010 from direct and remote volcanic gas
 1724 measurements. *Journal of Volcanology and Geothermal Research* 263, 92-107.

1725 Luhr, J. F., 2002. Petrology and geochemistry of the 1991 and 1998-1999 lava flows
 1726 from Volcán de Colima, México: implications for the end of the current eruptive
 1727 cycle. *Journal of Volcanology and Geothermal Research* 117, 169-194.

1728 Luhr, J. F., Carmichael, I. S., 1990. Petrological monitoring of cyclical eruptive activity
 1729 at Volcán Colima, Mexico. *Journal of Volcanology and Geothermal Research* 42, 235-
 1730 260.

1731 Luhr, J. F., Carmichael, I. S., 1980. The Colima Volcanic Complex, Mexico. I. Post-
 1732 Caldera Andesites From Volcán Colima. *Contributions to Mineralogy and Petrology*
 1733 71, 343-372.

1734 Mader, H., Llewellyn, E., Mueller, S., 2013. The rheology of two-phase magmas: A
 1735 review and analysis . *Journal of Volcanology and Geothermal Research* 257, 135-
 1736 158.

1737 Major, J., Dzurisin, D., Schilling, S., Poland, M., 2009. Monitoring lava-dome growth
 1738 during the 2004-2008 Mount St. Helens, Washington, eruption using oblique
 1739 terrestrial photography. *Earth and Planetary Science Letters* 286, 243-254.

1740 Manga, M., Castro, J., Cashman, K. V., Loewenberg, M., 1998. Rheology of bubble-
 1741 bearing magmas. *Journal of Volcanology and Geothermal Research* 87, 15-28.

- 1742 Marsh, B. D., 2015. Chapter 8-Magma Chambers . In: Sigurdsson, H. et al., (Eds.), The
1743 Encyclopedia of Volcanoes, 2nd Edition. Academic Press, pp. 343-362.
- 1744 Martel, C., Ali, A. R., Poussineau, S., Gourgaud, A., Pichavant, M., 2006. Basalt-
1745 inherited microlites in silicic magmas: Evidence from Mount Pelée (Martinique,
1746 French West Indies). *Geology* 34, 905-908.
- 1747 Martel, C., Pichavant, M., Bourdier, J-L., Traineau, H., Holtz, F., Scaillet, B., 1998.
1748 Magma storage conditions and control of eruption regime in silicic volcanoes:
1749 experimental evidence from Mt. Pelée. *Earth and Planetary Science Letters* 156, 89-
1750 99.
- 1751 Marzocchi, W., Sandri, L., Gasparini, P., Newhall, C., Boschi, E., 2004. Quantifying
1752 probabilities of volcanic events: the example of volcanic hazard at Vesuvius. *Journal*
1753 *of Geophysical Research* 109, 3-20.
- 1754 Mastin, L. G., 1994. Explosive tephra emissions at Mount St. Helens, 1989-1991: The
1755 violent escape of magmatic gas following storms? *Geological Society of America*
1756 *Bulletin* 106, 175- 185.
- 1757 Mather, T. A., Tsanev, V. I., Pyle, D. M., McGonigle, A. J. S., Oppenheimer, C., Allen, A.
1758 G., 2004. Characterization and evolution of tropospheric plumes from Lascar and
1759 Villarrica volcanoes, Chile. *Journal of Geophysical Research: Atmospheres* 109.
- 1760 Matthews, S. J., Sparks, R. S. J., Gardeweg, M. C., 1999. The Piedras Grandes-Soncor
1761 Eruptions, Lascar Volcano, Chile; Evolution of a Zoned Magma Chamber in the
1762 Central Andean Upper Crust. *Journal of Petrology* 40, 1891-1919.
- 1763 Matthews, S. J., Gardeweg, M. C., Sparks, R. S. J., 1997. The 1984 to 1996 cyclic
1764 activity of Lascar Volcano, northern Chile: cycles of dome growth, dome subsidence,
1765 degassing and explosive eruptions. *Bulletin of Volcanology* 59, 72-82.
- 1766 Matthews, S. J., Jones, A. P., Gardeweg, M. C., 1994. Lascar Volcano, Northern Chile;
1767 Evidence for Steady-State Disequilibrium. *Journal of Petrology* 35, 401-432.
- 1768 Mattioli, G. S., Herd, R. A., Strutt, M. H., Ryan, G., Widiwijayanti, C., Voight, B., 2010.
1769 Long term surface deformation of Soufrière Hills Volcano, Montserrat from GPS
1770 geodesy: Inferences from simple elastic inverse models. *Geophysical Research*
1771 *Letters* 37.
- 1772 McGee, K. A., Doukas, M. P., McGimsey, R. G., Neal, C. A., Wessels, R. L., 2010. Seismic
1773 Precursors to Volcanic Explosions During the 2006 Eruption of Augustine Volcano.
1774 In: Power, J. A., Coombs, M. L., Freymueller, J. T. (Eds.), *The 2006 Eruption of*
1775 *Augustine Volcano, Alaska. U.S. Geological Survey Professional Paper* 1769, Ch. 26,
1776 pp. 609-627.
- 1777 McNutt, S. R., 2005. Volcanic Seismology. *Annual Review of Earth and Planetary*
1778 *Sciences* 33, 461-491.
- 1779 Melnik, O., Sparks, R. S. J., 1999. Nonlinear dynamics of lava dome extrusion. *Nature*,
1780 37-41.

- 1781 Menard, G., Moune, S., Vlastélic, I., Aguilera, F., Valade, S., Bontemps, M., González, R.,
1782 2014. Gas and aerosol emissions from Lascar volcano (Northern Chile): Insights into
1783 the origin of gases and their links with the volcanic activity. *Journal of Volcanology*
1784 and *Geothermal Research* 287, 51-67.
- 1785 Mercalli, G., 1907. *Vulcani Attivi Della Terra*. Milano: Ulrico Hoepli.
- 1786 Métrich, N., Bertagnini, A., Di Muro, A., 2010. Conditions of Magma Storage,
1787 Degassing and Ascent at Stromboli: New Insights into the Volcano Plumbing System
1788 with Inferences on the Eruptive Dynamics. *Journal of Petrology* 51, 603-626.
- 1789 Michaut, C., Ricard, Y., Bercovici, D., Sparks, R. S. J., 2013. Eruption cyclicity at silicic
1790 volcanoes potentially caused by magmatic gas waves. *Nature Geoscience* 6, 856-860.
- 1791 Miller, A. D., Stewart, R. C., White, R. A., Luckett, R., Baptie, B. J., Aspinall, W. P.,
1792 Latchman, J. L., Lynch, L. L., Voight, B., 1998. Seismicity associated with dome growth
1793 and collapse at the Soufriere Hills Volcano, Montserrat. *Geophysical Research*
1794 *Letters* 25, 3401-3404.
- 1795 Miller, T., Chouet, B., 1994. The 1989-1990 eruptions of Redoubt Volcano: an
1796 introduction. *Journal of Volcanology and Geothermal Research* 62, 1-10.
- 1797 Miller, T. P., 1994. Dome growth and destruction during the 1989-1990 eruption of
1798 Redoubt volcano. *Journal of Volcanology and Geothermal Research* 62, 197-212.
- 1799 Molina, I., Kumagai, H., Le Pennec, J. L., Hall, M., 2005. Three-dimensional P-wave
1800 velocity structure of Tungurahua Volcano, Ecuador. *Journal of Volcanology and*
1801 *Geothermal Research* 147, 144-156.
- 1802 Moore, G., Carmichael, I. S. E., 1998. The hydrous phase equilibria (to 3 kbar) of an
1803 andesite and basaltic andesite from western Mexico: constraints on water content
1804 and conditions of phenocryst growth. *Contributions to Mineralogy and Petrology*
1805 130, 304-319.
- 1806 Mora, J., Macías, J., Saucedo, R., Orlando, A., Manetti, P., Vaselli, O., 2002. Petrology of
1807 the 1998-2000 products of Volcán de Colima, México. *Journal of Volcanology and*
1808 *Geothermal Research* 117, 195-212.
- 1809 Moran, S. C., Newhall, C., Roman, D. C., 2011. Failed magmatic eruptions: late-stage
1810 cessation of magma ascent. *Bulletin of Volcanology* 73, 115-122.
- 1811 Moran, S. C., Malone, S. D., Qamar, A. I., Thelen, W. A., Wright, A. K., Caplan-Auerbach,
1812 J., 2008. Seismicity Associated with Renewed Dome Building at Mount St. Helens,
1813 2004-2005. In: Sherrod, D. R., Scott, W. E., Stauffer, P. H. (Eds.), *A Volcano Rekindled:*
1814 *The Renewed Eruption of Mount St. Helens, 2004-2006*. U.S. Geological Survey
1815 Professional Paper 1750, Ch. 2, pp. 27-60.
- 1816 Moran, S. C., 1994. Seismicity at Mount St. Helens, 1987-1992: Evidence for
1817 repressurization of an active magmatic system. *Journal of Geophysical Research:*
1818 *Solid Earth* 99, 4341- 4354.

1819 Musumeci, C., Gresta, S., Malone, S. D., 2002. Magma system recharge of Mount St.
1820 Helens from precise relative hypocenter location of microearthquakes. *Journal of*
1821 *Geophysical Research*, 107.

1822 Murphy, M. D., Sparks, R. S. J., Barclay, J., Carroll, M. R., Brewer, T. S., 2000.
1823 Remobilization of Andesite Magma by Intrusion of Mafic Magma at the Soufrière
1824 Hills Volcano, Montserrat, West Indies. *Journal of Petrology* 41, 21-42.

1825 Nakada, S., Motomura, Y., 1999. Petrology of the 1991-1995 eruption at Unzen:
1826 effusion pulsation and groundmass crystallization. *Journal of Volcanology and*
1827 *Geothermal Research* 89, 173-196.

1828 Nakada, S., Shimizu, H., Ohta, K., 1999. Overview of the 1990-1995 eruption at Unzen
1829 Volcano. *Journal of Volcanology and Geothermal Research* 89, 1-22.

1830 Nakamura, M., 1995. Continuous mixing of crystal mush and replenished magma in
1831 the ongoing Unzen eruption. *Geology* 23, 807-810.

1832 Neave, D. A., Passmore, E., MacLennan, J., Fitton, G., Thordarson, T., 2013. Crystal-
1833 Melt Relationships and the Record of Deep Mixing and Crystallization in the AD
1834 1783 Laki Eruption, Iceland. *Journal of Petrology*.

1835 Neri, A., Aspinall, W., Cioni, R., Bertagnini, A., Baxter, P., Zuccaro, G., Andronico, D.,
1836 Barsotti, S., Cole, P., Esposti Ongaro, T., Hincks, T., Macedonio, G., Papale, P., Rosi, M.,
1837 Santacroce, R., Woo, G., 2008. Developing an Event Tree for probabilistic hazard and
1838 risk assessment at Vesuvius. *Journal of Volcanology and Geothermal Research* 178,
1839 397-415.

1840 Neuberg, J., 2000. Characteristics and causes of shallow seismicity in andesite
1841 volcanoes. *Philosophical Transactions of the Royal Society of London A:*
1842 *Mathematical, Physical and Engineering Sciences* 358, 1533-1546.

1843 Newhall, C. G., Hoblitt, R. P., 2002. Constructing event trees for volcanic crises.
1844 *Bulletin of Volcanology* 64, 3-20.

1845 Newhall, C., Melson, W., 1983. Explosive activity associated with the growth of
1846 volcanic domes. *Journal of Volcanology and Geothermal Research* 17, 111-131.

1847 Newhall, C. G., Self, S., 1982. The volcanic explosivity index (VEI) an estimate of
1848 explosive magnitude for historical volcanism. *Journal of Geophysical Research:*
1849 *Oceans* 87, 1231- 1238.

1850 Nichols, M., Malone, S., Moran, S., Thelen, W., Vidale, J., 2011. Deep long-period
1851 earthquakes beneath Washington and Oregon volcanoes. *Journal of Volcanology and*
1852 *Geothermal Research* 200, 116-128.

1853 Nishi, K., Ono, H., Mori, H., 1999. Global positioning system measurements of ground
1854 deformation caused by magma intrusion and lava discharge: the 1990-1995
1855 eruption at Unzendake volcano, Kyushu, Japan. *Journal of Volcanology and*
1856 *Geothermal Research* 89, 23-34.

- 1857 Núñez-Cornú, F., Nava, F., De la Cruz-Reyna, S., Jiménez, Z., Valencia, C., García-
1858 Arthur, R., 1994. Seismic activity related to the 1991 eruption of Colima Volcano,
1859 Mexico. *Bulletin of Volcanology* 56, 228-237.
- 1860 Nye, C. J., Swanson, S. E., Avery, V. F., Miller, T. P., 1994. Geochemistry of the 1989-
1861 1990 eruption of Redoubt volcano: Part I. Whole-rock major and trace-element
1862 chemistry. *Journal of Volcanology and Geothermal Research* 62, 429-452.
- 1863 Odbert, H. M., Ryan, G. A., Mattioli, G. S., Hautmann, S., Gottsmann, J., Fournier, N.,
1864 Herd, R. A., 2014a. Volcano geodesy at the Soufrière Hills Volcano, Montserrat: a
1865 review. *Geological Society, London, Memoirs* 39, 195-217.
- 1866 Odbert, H. M., Stewart, R. C., Wadge, G., 2014b. Cyclic phenomena at the Soufrière
1867 Hills Volcano, Montserrat. *Geological Society, London, Memoirs* 39, 41-60.
- 1868 Ogburn, S., Loughlin, S., Calder, E., 2015. The association of lava dome growth with
1869 major explosive activity (VEI 4): DomeHaz, a global dataset. *Bulletin of Volcanology*
1870 77.
- 1871 Ogburn, S. E., 2013. DomeHaz: Dome-forming eruptions database. VHub.
1872 <https://vhub.org/groups/domedatabase>.
- 1873 Oppenheimer, C., Scaillet, B., Martin, R. S., 2011. Sulfur Degassing From Volcanoes:
1874 Source Conditions, Surveillance, Plume Chemistry and Earth System Impacts.
1875 *Reviews in Mineralogy and Geochemistry* 73, 363-421.
- 1876 Ozerov, A., Ariskin, A., Kyle, P., Bogoyavlenskaya, G., Karpenko, S., 1997.
1877 Petrologicalgeochemical model for genetic relationships between basaltic and
1878 andesitic magmatism of Klyuchevskoy and Bezymianny volcanoes, Kamchatka.
1879 *Petrology* 5, 550-569.
- 1880 Pallister, J. S., Diefenbach, A. K., Burton, W. C., Muñoz, J., Griswold, J. P., Lara, L. E.,
1881 Lowenstern, J. B., Valenzuela, C. E., 2013. The Chaitén rhyolite lava dome: Eruption
1882 sequence, lava dome volumes, rapid effusion rates and source of the rhyolite
1883 magma. *Andean Geology* 40, 277-294.
- 1884 Pallister, J. S., Thornber, C. R., Cashman, K. V., Clynne, M. A., Lowers, H. A., Mandeville,
1885 C. W., Brownfield, I. K., Meeker, G. P., 2008. Petrology of the 2004-2006 Mount St.
1886 Helens Lava Dome-Implications for Magmatic Plumbing and Eruption Triggering. In:
1887 Sherrod, D. R., Scott, W. E., Stauffer, P. H. (Eds.), *A Volcano Rekindled: The Renewed*
1888 *Eruption of Mount St. Helens, 2004-2006*. U.S. Geological Survey Professional Paper
1889 1750, Ch. 30, pp. 647-702.
- 1890 Pallister, J. S., Hoblitt, R. P., Crandell, D. R., Mullineaux, D. R., 1992. Mount St. Helens a
1891 decade after the 1980 eruptions: magmatic models, chemical cycles, and a revised
1892 hazard assessment. *Bulletin of Volcanology* 54, 126-146.
- 1893 Perret, F., 1937. The eruption of Mt. Pelée, 1929-1932. Carnegie Institute of
1894 Washington Publication, v. 458, 126 pp.

- 1895 Petrosino, S., Cusano, P., La Rocca, M., Galluzzo, D., Orozco-Rojas, J., Bretón, M.,
1896 Ibáñez, J., Del Pezzo, E., 2011. Source location of long period seismicity at Volcán de
1897 Colima, México. *Bulletin of Volcanology* 73, 887-898.
- 1898 Phillipson, G., Sobradelo, R., Gottsmann, J., 2013. Global volcanic unrest in the 21st
1899 century: An analysis of the first decade. *Journal of Volcanology and Geothermal*
1900 *Research* 264, 183- 196.
- 1901 Pinel, V., Albino, F., 2013. Consequences of volcano sector collapse on magmatic
1902 storage zones: Insights from numerical modeling . *Journal of Volcanology and*
1903 *Geothermal Research* 252, 29-37.
- 1904 Pinel, V., Jaupart, C., 2005. Some consequences of volcanic edifice destruction for
1905 eruption conditions . *Journal of Volcanology and Geothermal Research* 145, 68-80.
- 1906 Pinel, V., Jaupart, C., 2000. The effect of edifice load on magma ascent beneath a
1907 volcano. *Philosophical Transactions of the Royal Society of London A: Mathematical,*
1908 *Physical and Engineering Sciences* 358, 1515-1532.
- 1909 Plail, M., Barclay, J., Humphreys, M. C. S., Edmonds, M., Herd, R. A., Christopher, T. E.,
1910 2014. Chapter 18 Characterization of mafic enclaves in the erupted products of
1911 Soufrière Hills Volcano, Montserrat, 2009 to 2010. *Geological Society, London,*
1912 *Memoirs* 39, 343-360.
- 1913 Platt, U., Stutz, J., 2008. Differential Absorption Spectroscopy. In: *Differential Optical*
1914 *Absorption Spectroscopy. Physics of Earth and Space Environments.* Springer Berlin
1915 Heidelberg, pp. 135-174.
- 1916 Power, J. A., Lalla, D. J., 2010. Seismic Observations of Augustine Volcano, 1970-
1917 2007. In: Power, J. A., Coombs, M. L., Freymueller, J. T. (Eds.), *The 2006 Eruption of*
1918 *Augustine Volcano, Alaska.* U.S. Geological Survey Professional Paper 1769, Ch. 1, pp.
1919 3-40.
- 1920 Power, J. A., Stihler, S. D., Chouet, B. A., Haney, M. M., Ketner, D. M., 2013. Seismic
1921 observations of Redoubt Volcano, Alaska-1989-2010 and a conceptual model of the
1922 Redoubt magmatic system. *Journal of Volcanology and Geothermal Research* 259,
1923 31-44.
- 1924 Power, J. A., Nye, C. J., Coombs, M. L., Wessels, R. L., Cervelli, P. F., Dehn, J., Wallace, K.
1925 L., Freymueller, J. T., Doukas, M. P., 2006. The reawakening of Alaska's Augustine
1926 volcano. *Eos, Transactions American Geophysical Union* 87, 373-377.
- 1927 Pritchard, M. E., Simons, M., 2002. A satellite geodetic survey of large-scale
1928 deformation of volcanic centres in the central Andes. *Nature* 418, 167-171.
- 1929 Pyle, D., 2000. Sizes of volcanic eruptions. In: Sigurdsson, H. (Ed.), *Encyclopaedia of*
1930 *Volcanoes.* Academic, Sydney, pp. 263-269.
- 1931 Ratdomopurbo, A., Beauducel, F., Subandriyo, J., Nandaka, I. M. A., Newhall, C. G.,
1932 Suhama, Sayudi, D. S., Suparwaka, H., Sunarta, 2013. Overview of the 2006 eruption
1933 of Mt. Merapi. *Journal of Volcanology and Geothermal Research* 261, 87-97.

- 1934 Ratdomopurbo, A., Poupinet, G., 2000. An overview of the seismicity of Merapi
1935 volcano (Java, Indonesia), 1983-1994. *Journal of Volcanology and Geothermal*
1936 *Research* 100, 193-214.
- 1937 Reubi, O., Blundy, J., 2009. A dearth of intermediate melts at subduction zone
1938 volcanoes and the petrogenesis of arc andesites . *Nature* 461, 1269-1273.
- 1939 Ripepe, M., Donne, D. D., Genco, R., Maggio, G., Pistolesi, M., Marchetti, E., Lacanna, G.,
1940 Ulivieri, G., Poggi, P., 2015. Volcano seismicity and ground deformation unveil the
1941 gravity-driven magma discharge dynamics of a volcanic eruption. *Nature*
1942 *Communications* 6.
- 1943 Rittmann, A., 1962. *Volcanoes and their Activity*. New York: John Wiley & Sons.
- 1944 Roberge, J., Delgado-Granados, H., Wallace, P. J., 2009. Mafic magma recharge
1945 supplies high CO₂ and SO₂ gas fluxes from Popocatepetl volcano, Mexico. *Geology* 37,
1946 107-110.
- 1947 Robin, C., Camus, G., Gourgaud, A., 1991. Eruptive and magmatic cycles at Fuego de
1948 Colima volcano (Mexico). *Journal of Volcanology and Geothermal Research* 45, 209-
1949 225.
- 1950 Rodríguez, L. A., Watson, I. M., Rose, W. I., Branan, Y. K., Bluth, G. J., Chigna, G., Matías,
1951 O., Escobar, D., Carn, S. A., Fischer, T. P., 2004. SO₂ emissions to the atmosphere from
1952 active volcanoes in Guatemala and El Salvador, 1999-2002. *Journal of Volcanology*
1953 *and Geothermal Research* 138, 325-344.
- 1954 Roman, D. C., Cashman, K. V., Gardner, C. A., Wallace, P. J., Donovan, J. J., 2006.
1955 Storage and interaction of compositionally heterogeneous magmas from the 1986
1956 eruption of Augustine Volcano, Alaska. *Bulletin of Volcanology* 68, 240-254.
- 1957 Roman, D. C., Power, J. A., Moran, S. C., Cashman, K. V., Doukas, M. P., Neal, C. A.,
1958 Gerlach, T. M., 2004. Evidence for dike emplacement beneath Iliamna Volcano,
1959 Alaska in 1996 . *Journal of Volcanology and Geothermal Research* 130, 265-284.
- 1960 Rose, W. I., 1973. Pattern and mechanism of volcanic activity at the Santiaguito
1961 Volcanic Dome, Guatemala. *Bulletin of Volcanology* 37, 73-94.
- 1962 Rose, W.I., J., 1972. Notes on the 1902 eruption of Santa María volcano, Guatemala.
1963 *Bulletin Volcanologique* 36, 29-45.
- 1964 Rose, W. I., Bluth, G. J. S., Ernst, G. G. J., 2000. Integrating retrievals of volcanic cloud
1965 characteristics from satellite remote sensors: a summary. *Philosophical*
1966 *Transactions of the Royal Society of London A: Mathematical, Physical and*
1967 *Engineering Sciences* 358, 1585- 1606.
- 1968 Rose, W. I., Heiken, G., Wohletz, K., Eppler, D., Barr, S., Miller, T., Chuan, R. L.,
1969 Symonds, R. B., 1988. Direct Rate Measurements of Eruption Plumes at Augustine
1970 Volcano: A Problem of Scaling and Uncontrolled Variables. *Journal of Geophysical*
1971 *Research: Solid Earth* 93, 4485-4499.

- 1972 Rutherford, M. J., Sigurdsson, H., Carey, S., Davis, A., 1985. The May 18, 1980,
1973 eruption of Mount St. Helens: 1. Melt composition and experimental phase
1974 equilibria. *Journal of Geophysical Research: Solid Earth* 90, 2929-2947.
- 1975 Samaniego, P., Le Pennec, J-L., Robin, C., Hidalgo, S., 2011. Petrological analysis of the
1976 preeruptive magmatic process prior to the 2006 explosive eruptions at Tungurahua
1977 volcano (Ecuador). *Journal of Volcanology and Geothermal Research* 199, 69-84.
- 1978 Sandri, L., Marzocchi, W., Zaccarelli, L., 2004. A new perspective in identifying the
1979 precursory patterns of eruptions. *Bulletin of Volcanology* 66, 263-275.
- 1980 Savov, I. P., Luhr, J. F., Navarro-Ochoa, C., 2008. Petrology and geochemistry of lava
1981 and ash erupted from Volcán Colima, Mexico, during 1998-2005. *Journal of*
1982 *Volcanology and Geothermal Research* 174, 241-256.
- 1983 Scandone, R., Cashman, K. V., Malone, S. D., 2007. Magma supply, magma ascent and
1984 the style of volcanic eruptions. *Earth and Planetary Science Letters* 253, 513-529.
- 1985 Scandone, R., Malone, S. D., 1985. Magma supply, magma discharge and
1986 readjustment of the feeding system of mount St. Helens during 1980. *Journal of*
1987 *Volcanology and Geothermal Research* 23, 239-262.
- 1988 Schilling, S. P., Thompson, R. A., Messerich, J. A., Iwatsubo, E. Y., 2008. Use of Digital
1989 Aerophotogrammetry to Determine Rates of Lava Dome Growth, Mount St. Helens,
1990 Washington, 2004-2005. In: Sherrod, D. R., Scott, W. E., Stauffer, P. H. (Eds.), *A*
1991 *Volcano Rekindled: The Renewed Eruption of Mount St. Helens, 2004-2006*. U.S.
1992 Geological Survey Professional Paper 1750, Ch. 8, pp. 145-168.
- 1993 Scott, J. A. J., Pyle, D. M., Mather, T. A., Rose, W. I., 2013. Geochemistry and evolution
1994 of the Santiaguito volcanic dome complex, Guatemala. *Journal of Volcanology and*
1995 *Geothermal Research* 252, 92-107.
- 1996 Scott, J. A., Mather, T. A., Pyle, D. M., Rose, W. I., Chigna, G., 2012. The magmatic
1997 plumbing system beneath Santiaguito Volcano, Guatemala. *Journal of Volcanology*
1998 *and Geothermal Research* 237-238, 54-68.
- 1999 Scott, W. E., Sherrod, D. R., Gardner, C. A., 2008. Overview of the 2004 to 2006, and
2000 Continuing, Eruption of Mount St. Helens, Washington. In: Sherrod, D. R., Scott, W. E.,
2001 Stauffer, P. H. (Eds.), *A Volcano Rekindled: The Renewed Eruption of Mount St.*
2002 *Helens, 2004-2006*. U.S. Geological Survey Professional Paper 1750, Ch. 1, pp. 3-26.
- 2003 Sernageomin, 2013. Reporte de Actividad Volcánica (RAV) REGIÓN DE
2004 ANTOFAGASTA Año 2013 Julio-Volumen 9. Tech. rep., Sernageomin.
2005 http://www.sernageomin.cl/reportesVolcanes/20130813010943279RAV_
2006 [Antofagasta_2013_julio_vol_9.pdf](http://www.sernageomin.cl/reportesVolcanes/20130813010943279RAV_)
- 2007 Shcherbakov, V. D., Plechov, P. Y., Izbekov, P. E., Shipman, J. S., 2011. Plagioclase
2008 zoning as an indicator of magma processes at Bezymianny Volcano, Kamchatka.
2009 *Contributions to Mineralogy and Petrology* 162, 83-99.

2010 Sheldrake, T., 2014. Long-term forecasting of eruption hazards: A hierarchical
2011 approach to merge analogous eruptive histories. *Journal of Volcanology and*
2012 *Geothermal Research* 286, 15-23.

2013 Shepherd, J. B., Tomblin, J., Woo, D., 1971. Volcano-Seismic Crisis in Montserrat,
2014 West Indies, 1966-67. *Bulletin Volcanologique* 35, 143-152.

2015 Shinohara, H., 2008. Excess degassing from volcanoes and its role on eruptive and
2016 intrusive activity. *Review of Geophysics* 46.

2017 Sides, I. R., Edmonds, M., MacLennan, J., Swanson, D. A., Houghton, B. F., 2014.
2018 Eruption style at Kilauea Volcano in Hawaii linked to primary melt composition.
2019 *Nature Geoscience* 7, 464- 469.

2020 Siebert, L., Simkin, T., Kimberley, P., 2010. *Volcanoes of the World*. Berkeley:
2021 University of California Press.

2022 Siswowidjoyo, S., Suryo, I., Yokoyama, I., 1995. Magma eruption rates of Merapi
2023 volcano, Central Java, Indonesia during one century (1890-1992). *Bulletin of*
2024 *Volcanology* 57, 111-116.

2025 Smith, A. L., Roobol, M. J., 1990. Mt. Pelée, Martinique; A Study of an Active Island-arc
2026 Volcano. *Geological Society of America Memoirs* 175, 1-110.

2027 Smith, R., Kilburn, C., Sammonds, P., 2007. Rock fracture as a precursor to lava dome
2028 eruptions at Mount St Helens from June 1980 to October 1986. *Bulletin of*
2029 *Volcanology* 69, 681-693.

2030 Sobradelo, R., Martí, J., 2015. Short-term volcanic hazard assessment through
2031 Bayesian inference: retrospective application to the Pinatubo 1991 volcanic crisis.
2032 *Journal of Volcanology and Geothermal Research* 290, 1-11.

2033 Sobradelo, R., Bartolini, S., Martí, J., 2013. HASSET: a probability event tree tool to
2034 evaluate future volcanic scenarios using Bayesian inference. *Bulletin of Volcanology*
2035 76.

2036 Solano, J. M. S., Jackson, M. D., Sparks, R. S. J., Blundy, J. D., Annen, C., 2012. Melt
2037 Segregation in Deep Crustal Hot Zones: a Mechanism for Chemical Differentiation,
2038 Crustal Assimilation and the Formation of Evolved Magmas. *Journal of Petrology*.

2039 Sparks, R. S. J., 2003. Forecasting volcanic eruptions. *Earth and Planetary Science*
2040 *Letters* 210, 1-15.

2041 Sparks, R. S. J., 1997. Causes and consequences of pressurisation in lava dome
2042 eruptions. *Earth and Planetary Science Letters* 150, 177-189.

2043 Sparks, R. S. J., Biggs, J., Neuberg, J. W., 2012. Monitoring Volcanoes. *Science* 335.

2044 Sparks, R. S. J., Aspinall, W. P., 2004. Volcanic Activity: Frontiers and Challenges in
2045 Forecasting, Prediction and Risk Assessment. In: *AGU Geophysical Monograph "State*
2046 *of the Planet"* 150. pp. 359-374.

2047 Sparks, R. S. J., Young, S. R., 2002. The eruption of Soufrière Hills Volcano,
 2048 Montserrat (1995-1999): overview of scientific results. Geological Society, London,
 2049 Memoirs 21, 45-69.

2050 Spiegelhalter, D. J., 1986. Probabilistic prediction in patient management and clinical
 2051 trials. *Statistics in Medicine* 5, 421-433.

2052 Stasiuk, M. V., Jaupart, C., Sparks, R. S. J., 1993. On the variations of flow rate in non-
 2053 explosive lava eruptions. *Earth and Planetary Science Letters* 114, 505-516.

2054 Steffke, A. M., Fee, D., Garces, M., Harris, A., 2010. Eruption chronologies, plume
 2055 heights and eruption styles at Tungurahua Volcano: Integrating remote sensing
 2056 techniques and infrasound. *Journal of Volcanology and Geothermal Research* 193,
 2057 143-160.

2058 Stith, J. L., Hobbs, P. V., Radke, L. F., 1978. Airborne particle and gas measurements in
 2059 the emissions from six volcanoes. *Journal of Geophysical Research: Oceans* 83, 4009-
 2060 4017.

2061 Straub, S. M., Martin-Del Pozzo, A. L., 2001. The significance of phenocryst diversity
 2062 in tephra from recent eruptions at Popocatepetl volcano (central Mexico).
 2063 *Contributions to Mineralogy and Petrology* 140, 487-510.

2064 Surono, Jousset, P., Pallister, J., Boichu, M., Buongiorno, M. F., Budisantoso, A., Costa,
 2065 F., Andreastuti, S., Prata, F., Schneider, D., Clarisse, L., Humaida, H., Sumartí, S.,
 2066 Bignami, C., Griswold, J., Carn, S., Oppenheimer, C., Lavigne, F., 2012. The 2010
 2067 explosive eruption of Java's Merapi volcano-A '100-year' event. *Journal of*
 2068 *Volcanology and Geothermal Research* 241242, 121-135.

2069 Swanson, D., Holcomb, R., 1990. Regularities in Growth of the Mount St. Helens
 2070 Dacite Dome, 1980-1986. In: Fink, J. H. (Ed.), *Lava Flows and Domes*. Vol. 2 of IAVCEI
 2071 *Proceedings in Volcanology*. Springer Berlin Heidelberg, pp. 3-24.

2072 Swanson, D. A., Casadevall, T. J., Dzurisin, D., Malone, S. D., Newhall, C. G., Weaver, C.
 2073 S., 1983. Predicting Eruptions at Mount St. Helens, June 1980 Through December
 2074 1982. *Science* 221, 1369-1376.

2075 Swanson, S. E., Nye, C. J., Miller, T. P., Avery, V. F., 1994. Geochemistry of the 1989-
 2076 1990 eruption of Redoubt volcano: Part II. Evidence from mineral and glass
 2077 chemistry. *Journal of Volcanology and Geothermal Research* 62, 453-468.

2078 Swanson, S. E., Kienle, J., 1988. The 1986 Eruption of Mount St. Augustine: Field Test
 2079 of a Hazard Evaluation. *Journal of Geophysical Research: Solid Earth* 93, 4500-4520.

2080 Symonds, R. B., Rose, W. I., Gerlach, T. M., Briggs, P. H., Harmon, R. S., 1990.
 2081 Evaluation of gases, condensates, and SO₂ emissions from Augustine volcano,
 2082 Alaska: the degassing of a Cl-rich volcanic system. *Bulletin of Volcanology* 52, 355-
 2083 374.

2084 Szakács, A., 2010. From a definition of volcano to conceptual volcanology. *Geological*
 2085 *Society of America Special Papers*, 470, 67-76.

2086 Szakács, A., Cañón-Tapia, E., 2010. Some challenging new perspectives of
2087 volcanology. Geological Society of America Special Papers, 470, 123-140.

2088 Tanguy, J-C., 2004. Rapid dome growth at Montagne Pelée during the early stages of
2089 the 1902-1905 eruption: a reconstruction from Lacroix's data. Bulletin of
2090 Volcanology 66, 615-621.

2091 Tanguy, J. C., 1994. The 1902-1905 eruptions of Montagne Pelée, Martinique:
2092 anatomy and retrospection. Journal of Volcanology and Geothermal Research 60, 87-
2093 107.

2094 Taran, Y., Gavilanes, J. C., Cortés, A., 2002. Chemical and isotopic composition of
2095 fumarolic gases and the SO₂ flux from Volcán de Colima, México, between the 1994
2096 and 1998 eruptions. Journal of Volcanology and Geothermal Research 117, 105-119.

2097 Tarasewicz, J., White, R. S., Woods, A. W., Brandsdóttir, B., Gudmundsson, M. T.,
2098 2012. Magma mobilization by downward-propagating decompression of the
2099 Eyjafjallajökull volcanic plumbing system. Geophysical Research Letters 39.

2100 Thelan, W., West, M., Senyukov, S., 2010. Seismic characterisation of the fall 2007
2101 eruptive sequence at Bezymianny Volcano, Russia. Journal of Volcanology and
2102 Geothermal Research 194, 201-213.

2103 Toramaru, A., Noguchi, S., Oyoshihara, S., Tsune, A., 2008. MND (microlite number
2104 density) water exsolution rate meter. Journal of Volcanology and Geothermal
2105 Research 175, 156-167.

2106 Troll, V. R., Deegan, F. M., Jolis, E. M., Harris, C., Chadwick, J. P., Gertisser, R.,
2107 Schwarzkopf, L. M., Borisova, A. Y., Bindeman, I. N., Sumarti, S., Preece, K., 2013.
2108 Magmatic differentiation processes at Merapi Volcano: inclusion petrology and
2109 oxygen isotopes. Journal of Volcanology and Geothermal Research 261, 38-49.

2110 Turner, S. J., Izbekov, P., Langmuir, C., 2013. The magma plumbing system of
2111 Bezymianny Volcano: Insights from a 54 year time series of trace element whole-
2112 rock geochemistry and amphibole compositions. Journal of Volcanology and
2113 Geothermal Research 263, 108-121.

2114 Ui, T., Takarada, S., Yoshimoto, M., 2000. Debris avalanches. In: Sigurdsson, H. et al.
2115 (Eds.), Encyclopedia of Volcanoes. 1st Edition. Academic Press, pp. 617-626.

2116 Umakoshi, K., Takamura, N., Shinzato, N., Uchida, K., Matsuwo, N., Shimizu, H., 2008.
2117 Seismicity associated with the 1991-1995 dome growth at Unzen Volcano, Japan .
2118 Journal of Volcanology and Geothermal Research 175, 91-99.

2119 USGS, 2014. Cascades Volcano Observatory Information Statement: Wednesday,
2120 April 30, 2014 16:05 UTC. Tech. rep., USGS,
2121 <http://volcanoes.usgs.gov/activity/archiveupdate.php?noticeid=10035>.

2122 van Manen, S. M., Dehn, J., Blake, S., 2010. Satellite thermal observations of the
2123 Bezymianny lava dome 1993-2008: Precursory activity, large explosions, and dome
2124 growth. Journal of Geophysical Research: Solid Earth 115.

- 2125 Varley, N., Arámbula-Mendoza, R., Reyes-Dávila, G., Sanderson, R., Stevenson, J.,
2126 2010. Generation of Vulcanian activity and long-period seismicity at Volcán de
2127 Colima, Mexico. *Journal of Volcanology and Geothermal Research* 198, 45-56.
- 2128 Varley, N. R., Taran, Y., 2003. Degassing processes of Popocatepetl and Volcán de
2129 Colima, Mexico. *Geological Society, London, Special Publications* 213, 263-280.
- 2130 Voight, B., 1988. A method for prediction of volcanic eruptions. *Nature* 332, 125-
2131 130.
- 2132 Voight, B., Constantine, E. K., Siswowidjoyo, S., Torley, R., 2000. Historical eruptions
2133 of Merapi Volcano, Central Java, Indonesia, 1768-1998. *Journal of Volcanology and*
2134 *Geothermal Research* 100, 69-138.
- 2135 Voight, B., Janda, R. J., Glicken, H., Douglass, P. M., 1983. Nature and mechanics of the
2136 Mount St Helens rockslide-avalanche of 18 May 1980. *Géotechnique* 33, 243-273.
- 2137 Volpe, A. M., Hammond, P. E., 1991. ^{238}U - ^{230}Th - ^{226}Ra disequilibria in young Mount St.
2138 Helens rocks: time constraint for magma formation and crystallization. *Earth and*
2139 *Planetary Science Letters* 107, 475-486.
- 2140 Wadge, G., Aspinall, W. P., 2014. A review of volcanic hazard and risk-assessment
2141 praxis at the Soufrière Hills Volcano, Montserrat from 1997 to 2011. *Geological*
2142 *Society, London, Memoirs* 39, 439-456.
- 2143 Wadge, G., Voight, B., Sparks, R. S. J., Cole, P. D., Loughlin, S. C., Robertson, R. E. A.,
2144 2014. An overview of the eruption of Soufrière Hills Volcano, Montserrat from 2000
2145 to 2010. *Geological Society, London, Memoirs* 39, 1-40.
- 2146 Wadge, G., Herd, R., Ryan, G., Calder, E. S., Komorowski, J. C., 2010. Lava production
2147 at Soufrière Hills Volcano, Montserrat: 1995-2009. *Geophysical Research Letters* 37.
- 2148 Wallace, P. J., 2005. Volatiles in subduction zone magmas: concentrations and fluxes
2149 based on melt inclusion and volcanic gas data. *Journal of Volcanology and*
2150 *Geothermal Research* 140, 217-240.
- 2151 Wallace, P. J., 2003. From mantle to atmosphere: magma degassing, explosive
2152 eruptions, and volcanic volatile budgets. In: Vivo, B. D., Bodnar, R. J. (Eds.), *Melt*
2153 *Inclusions in Volcanic Systems Methods, Applications and Problems*. Vol. 5 of
2154 *Developments in Volcanology*. Elsevier, pp. 105-127.
- 2155 Wallace, P. J., Edmonds, M., 2011. The Sulfur Budget in Magmas: Evidence from Melt
2156 Inclusions, Submarine Glasses, and Volcanic Gas Emissions. *Reviews in Mineralogy*
2157 *and Geochemistry* 73, 215-246.
- 2158 Walter, T. R., Wang, R., Zimmer, M., Grosser, H., Lühr, B., Ratdomopurbo, A., 2007.
2159 Volcanic activity influenced by tectonic earthquakes: Static and dynamic stress
2160 triggering at Mt. Merapi. *Geophysical Research Letters* 34.
- 2161 Watts, R. B., Herd, R. A., Sparks, R. S. J., Young, S. R., 2002. Growth patterns and
2162 emplacement of the andesitic lava dome at Soufrière Hills Volcano, Montserrat.
2163 *Geological Society, London, Memoirs* 21, 115-152.

2164 Webster, J. D., Mandeville, C. W., Goldoff, B., Coombs, M. L., Tappen, C., 2010.
 2165 Augustine Volcano-The Influence of Volatile Components in Magmas Erupted A.D.
 2166 2006 to 2,100 Years Before Present. In: Power, J. A., Coombs, M. L., Freymueller, J. T.
 2167 (Eds.), The 2006 Eruption of Augustine Volcano, Alaska. U.S. Geological Survey
 2168 Professional Paper 1769, Ch. 16, p. 383.

2169 Werner, C., Kelly, P. J., Doukas, M., Lopez, T., Pfeffer, M., McGimsey, R., Neal, C., 2013.
 2170 Degassing of CO₂, SO₂, and H₂S associated with the 2009 eruption of Redoubt
 2171 Volcano, Alaska. *Journal of Volcanology and Geothermal Research* 259, 270-284.

2172 West, M. E., 2013. Recent eruptions at Bezymianny volcano-A seismological
 2173 comparison. *Journal of Volcanology and Geothermal Research* 263, 42-57.

2174 Whelley, P. L., Newhall, C.G., Bradley, K. E., 2015, The frequency of explosive volcanic
 2175 eruptions in Southeast Asia. *Bulletin of Volcanology* 77.

2176 White, R., McCausland, W., 2016. Volcano-tectonic earthquakes: A new tool for
 2177 estimating intrusive volumes and forecasting eruptions. *Journal of Volcanology and*
 2178 *Geothermal Research* 309, 139-155.

2179 Witter, J. B., Kress, C. C., Newhall, C. G., 2005. Volcano Popocatepetl, Mexico.
 2180 Petrology, Magma Mixing, and Immediate Sources of Volatiles for the 1994-Present
 2181 Eruption. *Journal of Petrology* 46, 2337-2366.

2182 Wolf, K. J., Eichelberger, J. C., 1997. Syneruptive mixing, degassing, and
 2183 crystallization at Redoubt Volcano, eruption of December, 1989 to May 1990.
 2184 *Journal of Volcanology and Geothermal Research* 75, 19-37.

2185 Wright, H. M., Cashman, K. V., Mothes, P. A., Hall, M. L., Ruiz, A. G., Le Pennec, J-L.,
 2186 2012. Estimating rates of decompression from textures of erupted ash particles
 2187 produced by 1999-2006 eruptions of Tungurahua volcano, Ecuador. *Geology* 40,
 2188 619-622.

2189 Yokoyama, I., 2005. Growth rates of lava domes with respect to viscosity of magmas.
 2190 *Annals Of Geophysics*, 48.

2191 Young, S. R., Sparks, R. S. J., Aspinall, W. P., Lynch, L. L., Miller, A. D., Robertson, R. E.
 2192 A., Shepherd, J. B., 1998. Overview of the eruption of Soufrière Hills Volcano,
 2193 Montserrat, 18 July 1995 to December 1997. *Geophysical Research Letters* 25, 3389-
 2194 3392.

2195 Zellmer, G. F., Sparks, R. S. J., Hawkesworth, C. J., Wiedenbeck, M., 2003a. Magma
 2196 Emplacement and Remobilization Timescales Beneath Montserrat: Insights from Sr
 2197 and Ba Zonation in Plagioclase Phenocrysts. *Journal of Petrology* 44, 1413-1431.

2198 Zellmer, G. F., Hawkesworth, C. J., Sparks, R. S. J., Thomas, L. E., Harford, C. L., Brewer,
 2199 T. S., Loughlin, S. C., 2003b. Geochemical Evolution of the Soufrière Hills Volcano,
 2200 Montserrat, Lesser Antilles Volcanic Arc. *Journal of Petrology* 44, 1349-1374.

2201 Zharinov, N., Demyanchuk, Y., 2008. The growth of an extrusive dome on Shiveluch
 2202 Volcano, Kamchatka in 1980-2007: Geodetic observations and video surveys.
 2203 *Journal of Volcanology and Seismology* 2, 217-227.

Zobin, V. M., Varley, N. R., González, M., Orozco, J., Reyes, G. A., Reyes, C. A., Navarro, C., Bretón, M., 2008. Monitoring the 2004 andesitic block-lava extrusion at Volcán de Colima, México from seismic activity and SO₂ emission. *Journal of Volcanology and Geothermal Research* 177, 367-377.

Zobin, V., Luhr, J., Taran, Y., Bretón, M., Cortés, A., De la Cruz-Reyna, S., Dominguez, T., Galindo, I., Gavilanes, J., Muñiz, J., Navarro, C., Ramírez, J., Reyes, G., Ursúa, M., Velasco, J., Alatorre, E., Santiago, H., 2002. Overview of the 1997-2000 activity of Volcán de Colima, México. *Journal of Volcanology and Geothermal Research* 117, 1-19.

Figure Captions:

(1.5 column)

Figure 1: Locations of the 15 dome-building volcanoes in this study: (a) Augustine; (b) Bezymianny; (c) Colima; (d) Kudryavy; (e) Lascar; (f) Merapi; (g) Mount St. Helens; (h) Mont Pelée; (i) Popocatépetl; (j) Redoubt; (k) Santiaguito; (l) Shiveluch; (m) Soufrière Hills Volcano; (n) Tungurahua; (o) Mount Unzen. They are all found in subduction settings: either oceanic-continental or oceanic-oceanic boundaries.

(1.5 column)

Figure 2: Binary plots indicating whether (magmatic) eruptive activity (ash explosions and lava dome growth) was recording in each year since 1800 C.E, at each of the 15 volcanoes in this study. Importantly, the red bars do not equate to continuous eruptive activity, but instead are meant to indicate the variation in long-term patterns of eruptive activity. Labels are MER - Merapi; LAS - Lascar; COL - Colima; SHI - Shiveluch; SAN - Santiaguito; BEZ - Bezymianny; POP - Popocatépetl; TUN - Tungurahua; SHV - Soufrière Hills Volcano; HEL - Mount St. Helens; AUG - Augustine; RED - Redoubt; UNZ - Unzen; PEL - Pelée; KUD - Kudryavy. Volcanoes with the most persistent behaviour are found towards the top of the figure, and we have highlighted issues with specifically identifying a persistent regime in older records. The record of volcanic activity is based upon the Smithsonian database (Siebert and Simkin, 2002), and references specific to each volcano that can be found in section 3 and the supplementary material.

(Single column)

Figure 3: Representative cartoons for the two different eruptive regimes that are identified in this review; (a) Episodic behaviour, where the duration a volcano remains in an eruptive state is proportionally much shorter than the duration it remains in non-eruptive state. Degassing is temporally correlated with eruptive activity, and the regime is characterised by periods of no eruptive in which degassing is negligible, which we define as inter-eruptive repose; (b) Persistent

behaviour, where the duration a volcano remains in an eruptive state is proportionally similar to the duration it remains in non-eruptive state. Degassing is not necessarily temporally correlated with eruptive activity, and the regime is characterised by periods of no eruptive in which degassing is continuous and sustained, which we define as intra-eruptive repose. (c) A third mixed regime is characterised to identify how a volcano can exhibit both episodic and persistent behaviour in its eruptive record.

(Double column)

Figure 4: A hierarchical construct for historical eruptive activity at dome-building volcanoes. The first sub-level of this construct identifies the two different behaviours, episodic and persistent. The second sub-level of this construct identifies two different styles of episodic and persistent behaviour that are observed in historical records, over identical timescales (i.e. between points a and b). Key characteristics for each behaviour are identified in the boxes below each cartoon.

(Double column)

Figure 5: (a) Episodic behaviour at Augustine between 1970 and 2008, consisting of four eruptive episodes lasting months (red lines represent onsets), adapted from Power and Lalla, (2010). SO₂ degassing (orange) is temporally correlated with the eruptive episodes, as indicated by the data from McGee et al., (2010), overlaid on the lower chart. Black bars represent seismicity, which is elevated prior and during eruptive episodes; (b) Persistent behavior at Merapi between 1990 and 2006, with several phases of dome growth (blue bars) and associated explosions (blue vertical arrows), adapted from Ratdomopurbo et al., (2013). SO₂ degassing (orange) is temporally uncorrelated with eruptive activity, as observed by the overlaid data between 1992 and 1998. Seismicity is correlated with phases of eruptive activity, as indicated by the variation in the cumulative seismic energy (red line).

(Single column)

Figure 6: Estimated effusion rate (blue dots) at Unzen between 1990-1995, from Nakada et al. (1999). This is an example of a single eruptive episode at Unzen that lasted 5 years between 1990-1995 (Fig. 2). The latter stages of the eruptive episode are characterised by crystal-rich lavas and low effusion rates. During the eruptive episode, however, there are periodic increases in effusion rate, such as in 1993.

(Single column)


Figure 7: Estimated extrusion rates (blue dots) for 23 phases of dome growth at Bezymianny volcano between 1993 and 2008, from van Manen et al., (2010). This pattern of activity is an example of a persistent regime, in which frequent periods of dome-growth occur, with a consistent long-term extrusion rate. However, the intensity and frequency of phases of dome growth can vary. The red dashed line indicates the cumulative extruded volume, in which periods of dome growth and repose can be observed.


(Single column)

Figure 8: Example of a conceptual model for eruptive activity associated with the shallow chamber paradigm at La Soufrière, Guadeloupe, adapted from Hincks et al. (2014), where geophysical and geochemical observations at the surface are interpreted in terms of shallow crustal magmatic processes.


(Single column)

Figure 9: Schematic for the interaction of melt layers in a transcrustal magmatic system at lava dome-building volcanoes. Possible scenarios for eruptive activity and volcanic unrest; (a) complete destabilisation of the transcrustal system, involving deeply sourced mafic melts that provide volatiles and heat, resulting in major explosive activity; (b) partial destabilisation of the transcrustal system involving magma stored in shallow crustal regions resulting in effusive and minor explosive activity; (c) partial destabilisation of the magmatic system resulting in volcanic unrest but not eruptive activity. Importantly, this is in no way a true representation of the structure and dimensions of magmatic systems at lava dome-building volcanoes as they are found in subduction zones. Indeed, perpendicular to tectonic plate margins the arc widths of active volcanism are generally very narrow (~5 km or less).








(a) Episodic regime





(b) Persistent regime



(c) Mixed regime




Historical eruptive activity at dome-building volcanoes




Episodic behaviour

- Degassing correlated with eruptive state
- Duration in repose state > duration in eruptive state




- Episodic behaviour, with eruptive episodes lasting several years.
- Magmas are well mingled with homogeneous lavas.




Persistent behaviour

- Degassing uncorrelated with eruptive state
- Duration in repose state \approx duration in eruptive state



- Historical records characterised by continuous persistent behaviour.
- Large-magnitude explosions can occur at any time in the historical record.
- SiO_2 content of lava is consistent throughout the historical record.



- Persistent behaviour following a long-duration in state of repose (\sim millenia).
- Large-magnitude explosions occur with the onset of a persistent regime.
- SiO_2 content of lava decreases throughout the historical record.

