

How dose of biochar and biochar with nitrogen can improve the parameters of soil organic matter and soil structure?

Article

Accepted Version

Simansky, V., Horak, J., Igaz, D., Jonczak, J., Markiewicz, M., Felber, R., Rizhiya, E. Y. and Lukac, M. ORCID: <https://orcid.org/0000-0002-8535-6334> (2016) How dose of biochar and biochar with nitrogen can improve the parameters of soil organic matter and soil structure? *Biologia*, 71 (9). pp. 989-995. ISSN 0006-3088 doi: 10.1515/biolog-2016-0122 Available at <https://centaur.reading.ac.uk/66736/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

To link to this article DOI: <http://dx.doi.org/10.1515/biolog-2016-0122>

Publisher: De Gruyter

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online

How dose of biochar and biochar with nitrogen can improve the parameters of soil organic matter and soil structure?

Vladimír ŠIMANSKÝ^{1*}, Ján HORÁK², Dušan IGAZ², Jerzy JONCZAK³, Maciej MARKIEWICZ⁴,
Raphael FELBER⁵, Elena Y. RIZHIYA⁶ & Martin LUKAC^{7, 8}

¹*Department of Soil Science, Faculty of Agrobiolgy and Food Resources, Slovak University of Agriculture, Tr. A. Hlinku 2, 949 76 Nitra, Slovakia; e-mail: Vladimír.Simansky@uniag.sk*

²*Department of Biometeorology and Hydrology, Horticulture and Landscape Engineering Faculty, Slovak University of Agriculture in Nitra, Hospodárska 7, Nitra 94901, Slovakia*

³*Warsaw University of Life Sciences, Department of Soil Environment Sciences, Nowoursynowska Str. 159, 02-776 Warsaw, Poland*

⁴*Department of Soil Science and Landscape Management, Faculty of Earth Sciences, Nicolaus Copernicus University, Lwowska 1, 87-100 Toruń, Poland*

⁵*Agroscope, Climate & Air Pollution, Reckenholzstrasse 191, 8046 Zürich, Switzerland*

⁶*Agrophysical Research Institute, Grazhdansky pr. 14, St. Petersburg, 195220 Russia*

⁷*School of Agriculture, Policy and Development, University of Reading, Reading RG66AR, UK*

⁸*Department of Forest Management, Faculty of Forestry and Wood Sciences, Czech University of Life Sciences, 16521 Prague, Czech Republic*

*Corresponding author.

Running Title: Soil organic matter and soil structure

Abstract: Biochar application to agricultural soils has a significant potential to influence soil resource availability and thus crop performance. A factorial experiment investigating effects

of different biochar application rates combined with nitrogen fertilizer was conducted in field conditions on Haplic Luvisol. The aim of this study was to evaluate the effects of biochar and biochar combined with fertilization on soil organic matter and soil structure parameters. The treatments comprised combinations of biochar application of 0, 10 and 20 t ha⁻¹ (B0, B10 and B20) and 0, 40 and 80 kg N ha⁻¹ of nitrogen fertilizer (N0, N40, N80) applied in a full-factorial design. Biochar application rate of 20 t ha⁻¹ significantly increased soil organic carbon content (*SOC*) and non-labile carbon content (*C_{NL}*), but decreased carbon lability (*L_C*). The addition of biochar at 10 t ha⁻¹ together with 40 and 80 kg N ha⁻¹ significantly increased the values of *SOC* and *C_{NL}*. On the other hand, B10N80 treatment resulted in a considerable decrease of carbon lability (*L_C*). Overall, the highest average content of water-stable macro-aggregates was found in the B20N80 treatment and then with B10N0 < B20N40 < B20N0 < B10N80 < B0N0 < B10N40. Biochar applied at 20 t ha⁻¹ increased the critical level of soil organic matter and decreased the crusting index.

Key words: Biochar; N fertilization; Soil organic matter; Soil structure; Water-stable aggregates.

Introduction

The growing need for soil management strategies that enhance wider environmental benefits derived from arable soils while maintaining their productive function requires the development of novel approaches which ensure long-term sustainability of crop production. It is crucial to preserve or enhance favorable chemical, physical and biological soil properties, which in most soils are closely correlated with soil organic matter (*SOM*) content. *SOM* plays an important role in maintaining soil quality and ecosystem functionality (Benbi et al. 2015) and is an important aspect of agricultural soil quality and soil ecology (Gaida et al. 2013). Soil

organic carbon (*SOC*) content is one of the qualitative parameters of the soil humus regime (Howard & Howard 1990). Soil structure is one of the soil's most important physical properties, critically important for many soil environmental processes (Czachor & Lichner 2013; Garbout et al. 2013; Kodešová et al. 2015; Leelamanie & Mapa 2015). Soil structure is the key factor regulating soil function, its ability to support plant and animal life, and moderate environmental quality (Bronick & Lal 2005).

Intensive agriculture often leads to a decline of *SOC* content. Organic fertilizers, such as farmyard manure, are often the most important sources of organic compounds in systems with continuous removal of organic crop residue. The last two decades have been characterized by a continuous decline of livestock population in Slovakia, resulting in a decreasing availability of organic fertilizers. At the same time, environmental and regulatory constraints have driven arable agriculture towards lower-input soil management, highlighting the need to maintain optimal soil function and a favourable balance of organic compounds in the soil. could be an Application of biochar to arable soils could constitute an alternative to the historical use of organic fertilizers, acting as an important source of stable organic matter (Lehman 2007; Fischer & Glaser 2012).

Biochar is the product of thermal decomposition of organic materials in the absence of air (pyrolysis), and is distinguished from charcoal by its use as a soil amendment (Zimmerman 2010). Its use in agriculture may be advisable due to the confirmed positive effects on crop yields, mainly in sandy soils (Butnan et al. 2015). Biochar amends soil chemical properties such as *pH* (van Zwieten et al. 2010), reduces pesticide and nutrient leaching to groundwater (Novak et al. 2009), and improves nutrient regime of soils (Purakayastha et al. 2015). Biochar has also been shown to change soil biological community composition and abundance (Lehmann et al. 2011). Applying biochar to soils has had positive effects on soil physical properties, such as soil water holding capacity, bulk density, porosity

(Kammann et al. 2011) inner surface area (van Zwieten et al. 2009) and soil structure (Obia et al. 2016).

The relationship between organic matter and soil structure have been studied previously in different soil types, climate conditions and under varying soil management practices (Leelamanie & Karube 2014; Bartlová et al., 2015; Rajkai et al. 2015; Schacht & Marschner 2015). However, the interaction between biochar and biochar with nitrogen fertilizer applied in commercial setting and in field conditions has not been explored yet.

In this context, we hypothesised that the application of biochar to the soil would (i) increase *SOM* and (ii) improve soil structure. The objective of this study was to determine whether the addition of biochar or biochar together with nitrogen fertilizer has an effect on the soil organic matter and parameters of the soil structure.

Material and methods

Site description and experimental details

The field trial was conducted at an experimental site of SAU-Nitra (Nitra-Malanta) in Nitra region of Slovakia (lat. 48°19'00''; lon. 18°09'00'') during the period from March to July 2014 when a single crop of spring barley was grown. The soil at the site is classified as Haplic Luvisol (WRB 2006). Average annual air temperature was 10.3°C and annual precipitation was 640 mm during 2014. Soil samples from soil depth of 0–20 cm at 10 random locations (experimental field trial) were taken on 4th of March prior to setting up the experiment. On average, the soil contained 360.4 g kg⁻¹ of sand, 488.3 g kg⁻¹ of silt and 151.3 g kg⁻¹ of clay. Soil organic carbon content was 9.13 g kg⁻¹, while the average soil *pH* (KCl) was 5.71.

The experiment was established on 7th March 2014, followed by biochar application (10th of March) and crop drilling (11th of March). The replicated (n=3) spring barley (*Hordeum vulgare* L.) trial plots (4 m x 6 m) were laid out in a randomized block design in an

experimental field that has been used for continuous crop production for the last several years. The experiment consisted of the following treatments, separated by a protection row 0.5 m in width (Fig. 1): 1. B0N0 - no biochar, no N fertilization, 2. B10N0 - biochar (10 t ha⁻¹), 3. B20N0 - biochar (20 t ha⁻¹), 4. B0N40 - no biochar, fertilizer (40 kg N ha⁻¹), 5. B10N40 - biochar (10 t ha⁻¹) + fertilizer (40 kg N ha⁻¹), 6. B20N40 - biochar (20 t ha⁻¹) + fertilizer (40 kg N ha⁻¹), 7. B0N80 - no biochar, fertilizer (80 kg N ha⁻¹), 8. B10N80 - biochar (10 t ha⁻¹) + fertilizer (80 kg N ha⁻¹), 9. B20N80 - biochar (20 t ha⁻¹) + fertilizer (80 kg N ha⁻¹).

The field was ploughed, harrowed and biochar was evenly applied to the soil surface and immediately incorporated into the 0-10 cm soil layer combined with or without N fertilization using a combinator. To maintain consistency, plowing and mixing treatments were also performed in the control plots where no biochar or N fertilization was applied. A standard N fertilizer (Calc-Ammonium nitrate with dolomite, LAD 27) was used in this experiment. Biochar was produced from paper fiber sludge and grain husks (1:1 w/w) (company Sonnenerde, Austria) by pyrolysis at 550 °C for 30 minutes in a Pyreg reactor (Pyreg GmbH, Dörth, Germany) with particle size of 1–5 mm in size when applied (Table 1).

Soil sampling and analytical methods

Soil samples were repeatedly taken from the soil depth of 0–20 cm to cover the whole spring barley growing season (19th March, 17th April, 15th May, 16th Jun, and 13th July). Three different locations within each replicate plot were selected for soil sampling with and samples from a single plot were pooled to produce an average representative sample. Soil samples were always collected from the vicinity of the same sampling points and taken with the aid of a spade to maintain the soil aggregates throughout the experiment. Any roots and large fragments of plant litter were removed. Soil samples were transported to the laboratory where

the large clods were gently broken up along natural fracture lines, followed by air-drying at the lab temperature.

Particle-size distribution was determined by the pipette method (Fiala et al. 1999): dissolution of CaCO_3 with 2 M HCl, decomposition of the organic matter with 6% H_2O_2 , repeated washing, dispersing using $\text{Na}(\text{PO}_3)_6$ and then determination of the particle-size distribution. Soil organic carbon content (SOC) was estimated by the Tyurin wet oxidation method. The reagent mixture used in 0.07 M H_2SO_4 and $\text{K}_2\text{Cr}_2\text{O}_7$, with titration using 0.01M Mohr's salt (Dziadowiec & Gonet 1999). Labile carbon was extracted from 1 g soil samples by shaking them in 50 mL of 0.005 M KMnO_4 for two hours. After centrifugation, labile carbon content (C_L) was determined by oxidation of 0.07 M H_2SO_4 and $\text{K}_2\text{Cr}_2\text{O}_7$ with titration using 0.05 M Mohr's salt (Loginow et al. 1987). On the base of determined SOC and C_L we calculated the following parameters of SOM: carbon lability (L_C) and non-labile carbon content (C_{NL}), as suggested by Blair et al. (1995).

The L_C was calculated according to equation (1):

$$L_C = \frac{C_L}{C_{NL}} \quad (1)$$

where the non-labile carbon content (C_{NL}) is calculated as:

$$C_{NL} = \text{SOC} - C_L \quad (2)$$

Size classes of water-stable aggregates (WSA) were determined using the Baksheev method (Vadjunina & Korchagina 1986). Soil samples were first overflowed with distilled water (water level 1 cm above aggregates). After two hours, each sample was transferred to the top sieve (>5 mm) of a cylindrical container (Baksheev device), which was filled with distilled water. The cylinder was hermetically sealed and the sample was sieved for 12 minutes. The size fractions of WSA were as follows: >5, 5–3, 3–2, 2–1, 1–0.5, 0.5–0.25 mm (macro-aggregates, WSA_{ma}) and <0.25 mm (micro-aggregates, WSA_{mi}). The remaining material except for WSA_{mi} was quantified in each sieve. The micro-aggregate fraction was

calculated as the difference between the total weight of the soil sample and the sums of macro-aggregates. The index of crusting (Lal & Shukla 2004), and the critical level of soil organic matter according to Pieri (1991) were calculated as well.

The index of crusting (I_c) was calculated according to equation (3):

$$I_c = \frac{1.5S_f + 0.75S_c}{Cl + (10 \times SOM)} \quad (3)$$

where S_f is % fine silt, S_c is % coarse silt, Cl is % clay, and SOM is % soil organic matter content.

Critical soil organic matter content (St) was calculated using equation (4):

$$St = \frac{SOM}{(Clay + Silt)} \quad (4)$$

Statistical analysis

Statistical analysis was performed using the Statgraphics Centurion XV.I programme (Statpoint Technologies, Inc., USA). Effects of biochar and biochar combined with N fertilizer on SOM and soil structure parameters were tested using one-way ANOVA and then the least significant difference (LSD) method was used to compare treatment means for the two levels of biochar and two levels of nitrogen application at the significant level of $\alpha = 0.05$. The link between the SOM and soil structure parameters was assessed by a correlation matrix.

Results and discussion

Effects of biochar on the soil organic matter

SOC content was higher in B10N0 ($14.5 \pm 1.34 \text{ g kg}^{-1}$) and B20N0 ($18.9 \pm 3.30 \text{ g kg}^{-1}$) than in B0N0 ($12.2 \pm 0.85 \text{ g kg}^{-1}$). Biochar addition has been shown to increase SOC in soils (Fisher & Glasher 2012; Agegnehu et al. 2016). The same trends were observed in other treatments

when biochar was applied with N (40 and 80 kg N ha⁻¹), however, N addition interacted with biochar addition effects on *SOC* (Table 2). Addition of N at 40 and 80 kg ha⁻¹ together with 10 t ha⁻¹ of biochar significantly increased the value of *SOC* by 10 and 24%, respectively, compared to B10N0. The *C_{NL}* contents reflected the *SOC* status. Overall, the highest average content of *C_{NL}* was found in the B20N0 and B10N80 treatments compared to B0N0 (Table 2). We also evaluated the effects of biochar and biochar with N on changes of *L_C* which is used for the determination of smaller changes and changes over a short period of time (Blair et al. 1995, Bendi et al. 2015). Higher values of *L_C* indicate that the *SOM* is readily degradable by micro-organisms, while lower values of *L_C* indicate that the *SOM* has greater stability and resistance to microbial degradation (Szobathová 1999, Laik et al. 2009). The values of carbon lability (*L_C*) were significantly affected in B20N0 and B10N80 treatments compared to B0N0. Higher doses of biochar with no N fertilization and lower doses of biochar applied with higher doses of N appear to increase *SOM* resistance to microbial degradation.

Effects of biochar on the soil structure parameters

Several authors (Cornelissen et al. 2013; Herath et al. 2013) indicate positive effects of biochar on soil physical properties such as bulk density, porosity and soil structure stability. Our results from biochar-only plots are in general agreement with existing literature, but our study indicates the presence of interactive effects of biochar and nitrogen fertilization (Table 2). Biochar can improve soil physical conditions (Cornelissen et al. 2013; Obia et al. 2016) such as adsorption of cations (Liang et al. 2006), soil aggregate stability (Obia et al. 2016), but the effects on individual fractions of aggregates can differ, as indicated in our study (Fig. 2). For example, biochar (10 t ha⁻¹) applied without N fertilizer increased *WSA_{ma}* 5–2 mm content, but at the same time decreased *WSA_{ma}* 0.5–0.25 mm content. Application of biochar (20 t ha⁻¹) had no remarkable influence on the content of *WSA_{ma}*. Adding lower amounts of

biochar may thus be more beneficial for soil aggregation than higher rates of biochar addition. Secondly, biochar particles are 1–5 mm in size when applied. Conversion to WSA_{ma} 0.5–0.25 mm might therefore be difficult and could occur only after some time. Most of all, the biochar is very stable in the soil compared to the other forms of organic matter applications (Fischer & Glaser 2012). The surface of biochar particles after oxidation may contain hydroxyl and carboxylic groups which are able to adsorb soil particles and clays and form macro-aggregates (Jien & Wang 2013) however this process requires a substantial length of time. In our case, the experiment was established in March 2014 and soil sampling was carried out during the same growing season. Biochar, due to its mostly inert nature, is often applied to soils in conjunction with organic or mineral fertilizers (Fischer & Glaser 2012). In their study, application of N fertilizer together with biochar had a positive effect on the incorporation of biochar into the larger aggregates, which confirmed our results. In case of B20N80 treatment, the values of WSA_{ma} in the size fractions 3–2 mm (75%) and 5–3 mm (149%) were higher, while the size fraction of 0.5–0.25 mm (27%) was lower than in B20N0. We observed considerably lower content of WSA_{ma} 5–2 mm in B10N80. Dose of 40 kg N ha⁻¹ together with 10 and 20 t ha⁻¹ of biochar did not have a noticeable effect on WSA_{ma} (except size fraction 3–2 mm in B10N40). Adding nitrogen to the soil can improve microbial activity (Lehmann et al. 2011), increase the intensity of the biochar mineralization processes and increase CEC and active surface area (Yeboah et al. 2009), which results in higher aggregation (Bronic & Lal 2005).

Numerous reports show positive effects of biochar on aggregate stability (Herath et al. 2013; Sun & Lu 2014) and our study fully confirm these findings (Table 2). Biochar in dose of 20 t ha⁻¹ increased St (55%) compared to B0N0. Addition of manure (Whalen and Chang, 2002), as well as fertilizer application generally improve soil aggregation (Haynes and Naidu, 1998). However, under some conditions fertilizers may also decrease SOC concentration, reduce

aggregation, and reduce microbial communities compared to manured soils. Yet, using chemical fertilizers often improves soil structure in comparison to unfertilized soils (Munkholm et al., 2002). St increased by 24% in B10N80 compared to B10N0. Soil crust is a major structural feature of the surface soil and one of the most important physical properties. Organic fertilizers increase the stock of SOM , through which they positively influence the soil physicochemical properties, such as the formation of favourable structure (Czachor et al. 2015). In our study biochar also affected the crusting index (Table 2). Higher amounts of biochar significantly decreased I_c values compared to B0N0. The increased mean weight diameter of soil aggregates due to biochar application could be attributed to the increase in the amount of oxidized functional groups after mineralization of biochar (Jien & Wang 2013), which allowed flocculation of the soil particles and biochar, which also meant better structural state.

Relationships between SOM and the soil structure parameters

Pieri (1991) proposed the concept of the critical level of soil organic matter concentration (St) for structural stability of the soils. St values ranged between 3.28 and 5.08 and biochar and biochar with N had significant influence on St (Table 2). Similar to the Pieri (1991) evaluation, our results show lower values of St , meaning loss of soil structure and high susceptibility to erosion. The values of St increased only after higher amounts of biochar application, confirming the strong relationship between SOM and St (Table 3). Šimanský & Bajčan (2014) and Rabbi et al. (2015) also present a very strong positive relationship between aggregate stability and organic carbon content. It is a result of the strong linkage between colloidal fractions of soils and SOM (Kirkby & Morgan 1980). A very important parameter for the evaluation of soil structure is index of crusting (I_c). This is influenced by soil texture, SOM concentration and soil management practices (Lal & Shukla 2004). Higher SOM

content resulted in lower I_c values (Table 3). Higher content of labile carbon resulted in lower content of higher size fractions of aggregates (>3 mm). On the other hand, higher C_L content had a positive effect on the increase of WSA_{ma} 2–1 mm.

Conclusion

The results of our study indicate a positive response of *SOM* and soil structure parameters of Haplic Luvisol to biochar application. The most favourable effects on *SOC*, non-labile carbon and soil structure stability were observed when 20 t ha⁻¹ of biochar were applied. The same positive effects were observed after the application of 10 t ha⁻¹ of biochar in combination with nitrogen at rates of 40 and 80 kg ha⁻¹. Biochar improved aggregation and stability of soil structure, especially labile carbon and carbon lability had positive effects in the size fractions 2–1 mm and 5–2 mm of water-stable macro-aggregates, respectively.

The results of our study indicate that the application of biochar increases *SOC* content, in both its labile and non-labile forms, and also improves soil structure parameters. This information is very important for farmers, soil management practices can be optimized to avoid environmental degradation of their soils. Based on our findings, we recommend biochar application to pursue sustainable soil management with respect to carbon sequestration and soil structure preservation.

Acknowledgement

This study was supported by the Slovak Academy of Sciences and the Slovak Research and Development Agency under the contract APVV-15-0160. This study is also the result of the project implementation: Centre of excellence for the integrated river basin management in the changing environmental conditions, ITMS code 26220120062; supported by the Research & Development Operational Programme funded by the ERDF.

References

- Agegnehu G., Bass A. M., Nelson P. N. & Bird M.I. 2016. Benefits of biochar, compost and biochar–compost for soil quality, maize yield and greenhouse gas emissions in a tropical agricultural soil. *Sci. Tot. Environ.* **543**: 295–306
- Bartlová J., Badalíková B., Pospíšilová L., Pokorný E. & Šarapatka B. 2015. Water stability of soil aggregates in different systems of tillage. *Soil & Water Res.*, **10**: 147–154.
- Benbi D. K., Brar K., Toor A.S. & Sharma S. 2015. Sensitivity of labile soil organic carbon pools to long-term fertilizer, straw and manure management in rice-wheat system. *Pedosphere* **25**: 534–545.
- Blair G. J., Lefroy R. D. B. & Lisle L. 1995. Soil carbon fractions based on their degree of oxidation, and the development of a carbon management index for agricultural system. *Aust. J. Agri. Res.* **46**: 1459–1466.
- Bronick C. J. & Lal R. 2005. The soil structure and land management: a review. *Geoderma* **124**: 3–22.
- Butnan S., Deenik J. L., Toomsan B., Antal M. J. & Vityakon P. 2015. Biochar characteristics and application rates affecting corn growth and properties of soils contrasting in texture and mineralogy. *Geoderma* **237-238**: 105–116.
- Cornelissen G., Martinsen V., Shitumbanuma V., Alling V., Breedveld G., Rutherford D., Sparrevik M., Hale S., Obia A. & Mulder J. 2013. Biochar effect on maize yield and soil characteristics in five conservation farming sites in Zambia. *Agronomy* **3**: 256–274.
- Czachor H. & Lichner L. 2013. Temperature influences water sorptivity of soil aggregates. *J. Hydrol. Hydromech.* **61**: 84–87.
- Czachor H., Charytamowicz M., Gonet S., Niewczas J., Józefaciuk G. & Lichner L. 2015. Impact of long-term mineral and organic fertilizer application on the water stability,

- wettability and porosity of aggregates obtained from two loamy soils. *Eur. J. Soil Sci.* **66**: 1–12.
- Dziadowiec H. & Gonet S. S. 1999. Methodical guide-book for soil organic matter studies. Polish Society of Soil Science, Warszawa, 65 p. (in Polish).
- Fiala K., Kobza J., Matušková L., Brečková V., Makovníková J., Barančíková G., Búrik V., Litavec T., Houšková B., Chromaničová A., Váradiová D. & Pechová B. 1999. Valid Methods of Soil Analyses. Partial monitoring system – Soil. Soil Science and Conservation Research Institute, Bratislava, 142 p. (in Slovak).
- Fischer D. & Glaser B. 2012. Synergisms between compost and biochar for sustainable soil amelioration, pp. 167–198. In: Kumar S. (ed), Management of Organic Waste, In Tech Europe, Rijeka.
- Gaida A. M., Przewłoka B. & Gawryjolek K. 2013. Changes in soil quality associated with tillage system applied. *Int. Agrophys.* **27**: 133–141.
- Garbout A., Munkholm L. J. & Hansen S. B. 2013. Temporal dynamics for soil aggregates determined using X-ray CT scanning. *Geoderma* **204–205**: 15–22.
- Herath H. M. S. K., Camps-Arbestain M. & Hedley M. 2013. Effect of biochar on soil physical properties in two contrasting soils: an Alfisol and an Andisol. *Geoderma* **209–210**: 188–197.
- Howard P. J. A. & Howard D. M. 1990. Use of organic carbon and loss-on-ignition to estimate soil organic matter in different soil types and horizons. *Biol. Fertil. Soils* **9**: 306–310.
- IUSS Working Group WRB. 2006. World reference base for soil resources 2006. 2nd edition. World Soil Resources Reports No. 103, FAO, Rome, 112 p.
- Jien S. H. & Wang Ch. S. 2013. Effects of biochar on soil properties and erosion potential in a highly weathered soil. *Catena* **110**: 225–233

- Kammann C., Linsel S., Gößling J. & Koyro H. W. 2011. Influence of biochar on drought tolerance of *Chenopodium quinoa* Willd and on soil–plant relations. *Plant Soil* **345**: 195–210.
- Kirkby M. J. & Morgan R. P. C. 1980. *Soil Erosion*. John Wiley & Sons, London, 312 p.
- Kodešová R., Němeček K., Žigová A., Nikodem A. & Fér M. 2015. Using dye tracer for visualizing roots impact on soil structure and soil porous system. *Biologia* **70**: 1439–1443.
- Laik R., Kumar K., Das D. K. & Chaturvedi O. P. 2009. Labile soil organic matter pools in a calciorthent after 18 years of afforestation by different plantations. *Appl. Soil Ecol.* **42**: 71–78.
- Lal R. & Shukla M. K. 2004. *Principles of Soil Physics*. Marcel Dekker, New York, 719 p.
- Lehmann J., Rillig M. C., Thies J., Masiell, C. A., Hockaday W. C. & Crowley D. 2011. Biochar effects on soil biota, A review. *Soil Biol. Biochem.* **43**: 1812–1836.
- Leelamanie D.A.L. & Karube J. 2014. Water stable aggregates of Japanese Andisol as affected by hydrophobicity and drying temperature. *J. Hydrol. Hydromech.* **62**: 97–100.
- Leelamanie D.A.L. & Mapa R.B. 2015. Alterations in soil aggregate stability of a tropical Ultisol as mediated by changes in land use. *Biologia* **70**: 1444–1449.
- Lehmann J. 2007. Bio-energy in the black. *Front. Ecol. Environ.* **5**: 381–387.
- Liang B., Lehmann J., Solomon D., Kinyang J., Grossman J., O'Neill B., Skjemstard J.O., Thies J., Luiza F. J., Peterson J. & Neves, E. G. 2006. Black carbon increases CEC in soils. *Soil Sci. Soc. Am. J.* **70**: 1719–1730.
- Loginow W., Wisniewski W., Gonet S. S. & Ciescinska B. 1987. Fractionation of organic carbon based on susceptibility to oxidation. *Pol. J. Soil Sci.* **20**: 47–52.

- Munkholm L. J., Schjonning P., Debosz K., Jensen H. E. & Christensen B. T. 2002. Aggregate strength and mechanical behaviour of a sandy loam soil under long-term fertilization treatments. *Eur. J. Soil Sci.* **53**: 129–137.
- Novak J. M., Lima I., Xing B., Gaskin J. W., Steiner C., Das K. C., Ahmedna M., Rehrah D., Watts D. W., Busscher W.J. & Schomberg H. 2009. Characterization of designer biochar produced at different temperatures and their effects on a loamy sand. *Ann. Environ. Sci.* **3**: 195–206.
- Obia A., Mulder J., Martinsen V., Cornelissen G. & Børresen T. 2016. In situ effects of biochar on aggregation, water retention and porosity in light-textured tropical soils. *Soil Till. Res.* **155**: 35–44
- Pieri C. 1991. Fertility of soils: A future for farming in the West African savannah. Springer-Verlag, Berlin, 348 p.
- Purakayastha T. J., Kumari S. & Pathak H. 2015. Characterisation, stability, and microbial effects of four biochars produced from crop residues. *Geoderma* **239-240**: 293–303.
- Rabbi S. M. F., Wilson B. R., Lockwood P. V., Daniel H. & Young I. M. 2015. Aggregate hierarchy and carbon mineralization in two Oxisols of New South Wales, Australia. *Soil Till. Res.* **146**: 193–203.
- Rajkai K., Tóth B., Barna G., Hernádi H., Kocsis M. & Makó A. 2015. Particle-size and organic matter effects on structure and water retention of soils. *Biologia* **70**: 1456–1461.
- Schacht K. & Marschner B. 2015. Treated wastewater irrigation effects on soil hydraulic conductivity and aggregate stability of loamy soils in Israel. *J. Hydrol. Hydromech.* **63**: 47–54.
- Szombathová N. 1999. The comparison of soil carbon susceptibility to oxidation by KMnO_4 solutions in different farming systems. *Hum. Subst. Environ.* **1**: 35 – 39.

- Sun F. & Lu, S. 2014. Biochars improve aggregate stability, water retention, and pore-space properties of clayey soil. *J. Plant Nutr. Soil Sci.* **177**: 26–33.
- Šimanský V. & Bajčan D. 2014. The stability of soil aggregates and their ability of carbon sequestration. *Soil & Water Res.*, **9**: 111–118.
- Vadjunina A. F. & Korchagina Z. A. 1986. *Methods of Study of Soil Physical Properties*. Agropromizdat, Moscow, 415 p. (in Russian).
- van Zwieten L., Kimber S., Morris S., Chan K. Y., Downie A., Rust J., Joseph S. & Cowie A. 2010. Effects of biochar from slow pyrolysis of papermill waste on agronomic performance and soil fertility. *Plant Soil* **327**: 235–246.
- van Zwieten L., Singh B. P., Joseph S., Kimber S., Cowie A. & Chan K. Y. 2009. Biochar and emissions of non-CO₂ greenhouse gases from soil, pp. 227–249. In: Lehmann J. & Joseph S. (eds.), *Biochar for Environmental Management: Science and Technology*, Earthscan, London.
- Yeboah E., Ofori P., Quansah G. W., Dugan, E. & Sohi S. P. 2009. Improving soil productivity through biochar amendments to soils. *Afr. J. Environ. Sci. Technol.* **3**: 34–41.
- Zimmerman A. 2010. Abiotic and microbial oxidation of laboratory-produced black carbon (biochar). *Environ. Sci. Tech.* **44**: 1295–1301.

Table 1. Basic chemical and physical properties of the biochar (biochar certificate Nr.1013069001 provided by company Sonnenerde, Austria).

biochar		
pH	(KCl)	8.8
Ca	g kg ⁻¹	57
Mg	g kg ⁻¹	3.9
K	g kg ⁻¹	15
Na	g kg ⁻¹	0.7
Total C	g kg ⁻¹	53.1
Total N	g kg ⁻¹	14
C/N		3.79
SSA	m ² g ⁻¹	21.7
Ash	%	38.3

Table 2. Statistical evaluation of soil organic matter and parameters of soil structure

	<i>SOC</i>	<i>C_L</i>	<i>C_{NL}</i>	<i>L_C</i>	<i>St</i>	<i>Ic</i>	<i>WSA_{mi}</i>
	g kg ⁻¹						%
treatments							
B0N0	12.2±0.85 ^a	1.54±0.31 ^a	10.6±0.74 ^a	0.145±0.03 ^c	3.28±0.23 ^a	2.88±0.03 ^d	30.2±5.41 ^b
B10N0	14.5±1.34 ^{ab}	1.77±0.24 ^a	12.7±1.44 ^{ab}	0.142±0.03 ^{bc}	3.90±0.36 ^{ab}	2.82±0.04 ^{cd}	20.5±9.77 ^{ab}
B20N0	18.9±3.30 ^c	1.89±0.40 ^a	16.9±2.99 ^c	0.112±0.02 ^{ab}	5.08±0.89 ^d	2.71±0.08 ^a	26.7±11.3 ^{ab}
B10N40	15.9±1.96 ^{bc}	1.65±0.41 ^a	14.3±1.88 ^{bc}	0.116±0.03 ^{abc}	4.30±0.53 ^{bc}	2.78±0.05 ^{bc}	30.4±8.33 ^b
B20N40	17.9±2.90 ^{cd}	1.89±0.36 ^a	16.0±2.70 ^c	0.119±0.02 ^{abc}	4.83±0.78 ^{cd}	2.73±0.07 ^{ab}	23.8±11.6 ^{ab}
B10N80	18.0±2.03 ^{cd}	1.67±0.26 ^a	16.3±2.10 ^c	0.104±0.02 ^a	4.85±0.55 ^{cd}	2.73±0.05 ^{ab}	27.4±8.49 ^{ab}
B20N80	16.4±2.16 ^{bc}	1.78±0.31 ^a	14.7±1.87 ^{bc}	0.121±0.01 ^{abc}	4.43±0.58 ^{bcd}	2.70±0.06 ^{abc}	17.3±5.23 ^a

Table 3. Correlation coefficients between soil organic matter and soil structure parameters

	<i>St</i>	<i>Ic</i>	>5	5–3	3–2	2–1	1–0.5	0.5–0.25	<i>WSA_{mi}</i>
<i>SOC</i>	1.000 ^{***}	–0.999 ^{***}	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.
<i>C_L</i>	0.425 ^{**}	–0.417 ^{**}	–0.442 ^{**}	n.s.	n.s.	0.382 [*]	n.s.	n.s.	n.s.
<i>L_C</i>	–0.673 ^{***}	0.683 ^{***}	–0.502 ^{***}	–0.355 [*]	n.s.	n.s.	n.s.	n.s.	n.s.




Fig. 1. Field site location and an areal view of experimental plots.




Fig. 2. Content of individual size-fraction of water-stable macro-aggregates.

Different letters between columns (a, b, c) indicate that treatment means are significantly different at $P < 0.05$ according to LSD multiple-range test.