

Existence of 1D vectorial absolute minimisers in L infinity under minimal assumptions

Article

Accepted Version

Abugirda, H. and Katzourakis, N. (2017) Existence of 1D vectorial absolute minimisers in L infinity under minimal assumptions. *Proceedings of the American Mathematical Society*, 145 (6). pp. 2567-2575. ISSN 0002-9939 doi: 10.1090/proc/13421 Available at <https://centaur.reading.ac.uk/67059/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

To link to this article DOI: <http://dx.doi.org/10.1090/proc/13421>

Publisher: American Mathematical Society

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online

EXISTENCE OF 1D VECTORIAL ABSOLUTE MINIMISERS IN L^∞ UNDER MINIMAL ASSUMPTIONS

HUSSIEN ABUGIRDA AND NIKOS KATZOURAKIS

ABSTRACT. We prove the existence of vectorial Absolute Minimisers in the sense of Aronsson to the supremal functional $E_\infty(u, \Omega') = \|\mathcal{L}(\cdot, u, Du)\|_{L^\infty(\Omega')}$, $\Omega' \Subset \Omega$, applied to $W^{1,\infty}$ maps $u : \Omega \subseteq \mathbb{R} \rightarrow \mathbb{R}^N$ with given boundary values. The assumptions on \mathcal{L} are minimal, improving earlier existence results previously established by Barron-Jensen-Wang and by the second author.

1. INTRODUCTION

The main goal of this paper is to prove the existence of vectorial Absolute Minimisers with given boundary values to the supremal functional

$$(1.1) \quad E_\infty(u, \Omega') := \operatorname{ess\,sup}_{x \in \Omega'} \mathcal{L}(x, u(x), Du(x)), \quad u \in W_{\operatorname{loc}}^{1,\infty}(\Omega, \mathbb{R}^N), \quad \Omega' \Subset \Omega,$$

applied to maps $u : \Omega \subseteq \mathbb{R} \rightarrow \mathbb{R}^N$, $N \in \mathbb{N}$, where Ω is an open interval and $\mathcal{L} \in C(\Omega \times \mathbb{R}^N \times \mathbb{R}^N)$ is a non-negative continuous function which we call Lagrangian and whose arguments will be denoted by (x, η, P) . By Absolute Minimiser we mean a map $u \in W_{\operatorname{loc}}^{1,\infty}(\Omega, \mathbb{R}^N)$ such that

$$(1.2) \quad E_\infty(u, \Omega') \leq E_\infty(u + \phi, \Omega'),$$

for all $\Omega' \Subset \Omega$ and all $\phi \in W_0^{1,\infty}(\Omega', \mathbb{R}^N)$. This is the appropriate minimality notion for supremal functionals of the form (1.1); requiring at the outset minimality on all subdomains is necessary because of the lack of additivity in the domain argument. The study of (1.1) was pioneered by Aronsson in the 1960s [A1]-[A5] who considered the case $N = 1$. Since then, the (higher dimensional) scalar case of $u : \Omega \subseteq \mathbb{R}^n \rightarrow \mathbb{R}$ has developed massively and there is a vast literature on the topic (see for instance the lecture notes [C, K7]). In the case the Lagrangian is C^1 , of particular interest has been the study of the (single) equation associated to (1.1), which is the equivalent of the Euler-Lagrange equation for supremal functionals and is known as the ‘‘Aronsson equation’’:

$$(1.3) \quad A_\infty u := D(\mathcal{L}(\cdot, u, Du)) \mathcal{L}_P(\cdot, u, Du) = 0.$$

In (1.3) above, the subscript denotes the gradient of $\mathcal{L}(x, \eta, P)$ with respect to P and, as it is customary, the equation is written for smooth solutions. Herein we are interested in the vectorial case $N \geq 2$ but in one spatial dimension. Unlike the scalar case, the literature for $N \geq 2$ is much more sparse and starts much more recently. Perhaps the first most important contributions were by Barron-Jensen-Wang [BJW1, BJW2] who among other deep results proved the existence

2010 *Mathematics Subject Classification.* Primary 35J47, 35J62, 53C24; Secondary 49J99.

Key words and phrases. Vectorial Calculus of Variations in L^∞ ; Vectorial Absolute Minimisers; ∞ -Laplacian.

of Absolute Minimisers for (1.1) under certain assumptions on \mathcal{L} which we recall later. However, their contributions were at the level of the functional and the appropriate (non-obvious) vectorial analogue of the Aronsson equation was not known at the time. The systematic study of the vectorial case of (1.1) (actually in the general case of maps $u : \Omega \subseteq \mathbb{R}^n \rightarrow \mathbb{R}^N$) together with its associated system of equations begun in the early 2010s by the second author in a series of papers, see [K1]-[K6], [K8]-[K12] (and also the joint contributions with Croce, Pisante and Pryer [CKP, KP1, KP2]). The ODE system associated to (1.1) for smooth maps $u : \Omega \subseteq \mathbb{R} \rightarrow \mathbb{R}^N$ turns out to be

$$(1.4) \quad \mathcal{F}_\infty(\cdot, u, Du, D^2u) = 0, \quad \text{on } \Omega,$$

where

$$(1.5) \quad \begin{aligned} \mathcal{F}_\infty(x, \eta, P, X) := & \left[\mathcal{L}_P(x, \eta, P) \otimes \mathcal{L}_P(x, \eta, P) \right. \\ & + \mathcal{L}(x, \eta, P) [\mathcal{L}_P(x, \eta, P)]^\perp \mathcal{L}_{PP}(x, \eta, P) \Big] X \\ & + \left(\mathcal{L}_\eta(x, \eta, P) \cdot P + \mathcal{L}_x(x, \eta, P) \right) \mathcal{L}_P(x, \eta, P) \\ & + \mathcal{L}(x, \eta, P) [\mathcal{L}_P(x, \eta, P)]^\perp \left(\mathcal{L}_{P\eta}(x, \eta, P) P \right. \\ & \left. + \mathcal{L}_{Px}(x, \eta, P) - \mathcal{L}_\eta(x, \eta, P) \right). \end{aligned}$$

Quite unexpectedly, in the case $N \geq 2$ the Lagrangian needs to be C^2 for the equation to make sense, whilst the coefficients of the full system are *discontinuous*; for more details we refer to the papers cited above. In (1.5) the notation of subscripts symbolises derivatives with respect to the respective variables and $[\mathcal{L}_P(x, \eta, P)]^\perp$ is the orthogonal projection to the hyperplane normal to $\mathcal{L}_P(x, \eta, P) \in \mathbb{R}^N$:

$$(1.6) \quad [\mathcal{L}_P(x, \eta, P)]^\perp := I - \text{sgn}(\mathcal{L}_P(x, \eta, P)) \otimes \text{sgn}(\mathcal{L}_P(x, \eta, P)).$$

The system (1.4) reduces to the equation (1.3) when $N = 1$. In the paper [K9] the existence of an absolutely minimising *generalised solution* to (1.4) was proved, together with extra partial regularity and approximation properties. Since (1.4) is a quasilinear non-divergence degenerate system with discontinuous coefficients, a notion of appropriately defined “weak solution” is necessary because in general solutions are non-smooth. To this end, the general new approach of \mathcal{D} -solutions which has recently been proposed in [K8] has proven to be the appropriate setting for vectorial Calculus of Variations in L^∞ (see [K8]-[K10]), replacing to some extent viscosity solutions which essentially apply only in the scalar case.

Herein we are concerned with the existence of absolute minimisers to (1.1) without drawing any connections to the differential system (1.4). Instead, we are interested in obtaining existence under the weakest possible assumptions. Accordingly, we establish the following result.

Theorem 1. *Let $\Omega \subseteq \mathbb{R}$ be a bounded open interval and let also*

$$\mathcal{L} : \overline{\Omega} \times \mathbb{R}^N \times \mathbb{R}^N \rightarrow [0, \infty),$$

be given continuous function with $N \in \mathbb{N}$. We assume that:

- (1) *For each $(x, \eta) \in \overline{\Omega} \times \mathbb{R}^N$, the function $P \mapsto \mathcal{L}(x, \eta, P)$ is level-convex, that is for each $t \geq 0$ the sublevel set*

$$\{P \in \mathbb{R}^N : \mathcal{L}(x, \eta, P) \leq t\}$$

is a convex set in \mathbb{R}^N .

- (2) there exist non-negative constants C_1, C_2, C_3 , and $0 < q \leq r < +\infty$ and a positive locally bounded function $h : \mathbb{R} \times \mathbb{R}^N \rightarrow [0, +\infty)$ such that for all $(x, \eta, P) \in \overline{\Omega} \times \mathbb{R}^N \times \mathbb{R}^N$

$$C_1|P|^q - C_2 \leq \mathcal{L}(x, \eta, P) \leq h(x, \eta)|P|^r + C_3.$$

Then, for any affine map $b : \mathbb{R} \rightarrow \mathbb{R}^N$, there exist a vectorial Absolute Minimiser $u^\infty \in W_b^{1,\infty}(\Omega, \mathbb{R}^N)$ of the supremal functional (1.1) (Definition (1.2)).

Theorem 1 generalises two respective results in the both the papers [BJW1] and [K9]. On the one hand, in [BJW1] Theorem 1 was established under the extra assumption $C_2 = C_3 = 0$ which forces $\mathcal{L}(x, \eta, 0) = 0$, for all $(x, \eta) \in \mathbb{R} \times \mathbb{R}^N$. Unfortunately this requirement is incompatible with important applications of (1.1) to problems of L^∞ -modelling of variational Data Assimilation (4DVar) arising in the Earth Sciences and especially in Meteorology (see [B, BS, K9]). An explicit model of \mathcal{L} is given by

$$(1.7) \quad \mathcal{L}(x, \eta, P) := |k(x) - K(\eta)|^2 + |P - \mathcal{V}(x, \eta)|^2,$$

and describes the “error” in the following sense: consider the problem of finding the solution u to the following ODE coupled by a pointwise constraint:

$$Du(t) = \mathcal{V}(t, u(t)) \quad \& \quad K(u(t)) = k(t), \quad t \in \Omega.$$

Here $\mathcal{V} : \Omega \times \mathbb{R}^N \rightarrow \mathbb{R}^N$ is a time-dependent vector field describing the law of motion of a body moving along the orbit described by $u : \Omega \subseteq \mathbb{R} \rightarrow \mathbb{R}^N$ (e.g. Newtonian forces, Galerkin approximation of the Euler equations, etc), $k : \Omega \subseteq \mathbb{R} \rightarrow \mathbb{R}^M$ is some partial “measurements” in continuous time along the orbit and $K : \mathbb{R}^N \rightarrow \mathbb{R}^M$ is a submersion which corresponds to some component of the orbit that is observed. We interpret the problem as that u should satisfy the law of motion and also be compatible with the measurements along the orbit. Then minimisation of (1.1) with \mathcal{L} as given by (1.7) leads to a uniformly optimal approximate solution without “spikes” of large deviation of the prediction from the actual orbit.

On the other hand, in the paper [K9] Theorem 1 was proved under assumptions allowing to model Data Assimilation but strong convexity, smoothness and structural assumptions were imposed, allowing to obtain stronger results accordingly. In particular, the Lagrangian was assumed to be radial in P , which means it can be written in the form

$$\mathcal{L}(x, \eta, P) := \mathcal{H}\left(x, \eta, \frac{1}{2}|P - \mathcal{V}(x, \eta)|^2\right).$$

In this paper we relax the hypotheses of both the aforementioned results.

2. PROOF OF THE MAIN RESULT

In this section we establish Theorem 1. In the proof we will utilise the following lemma essentially proved in the paper [BJW1] which we recall right below for the convenience of the reader.

Lemma 2. (cf. Lemma 2.2 in [BJW1]) *In the setting of theorem 1 and under the same hypotheses, for a fixed affine map $b : \mathbb{R} \rightarrow \mathbb{R}^N$, set*

$$C_m := \inf \left\{ E_m(u, \Omega) : u \in W_b^{1,qm}(\Omega, \mathbb{R}^N) \right\},$$

$$C_\infty := \inf \left\{ E_\infty(u, \Omega) : u \in W_b^{1,\infty}(\Omega, \mathbb{R}^N) \right\}.$$

where E_∞ is as in (1.1) and

$$(2.1) \quad E_m(u, \Omega) := \int_{\Omega} \mathcal{L}(x, u(x), Du(x))^m dx.$$

Then, there exist $u^\infty \in W_b^{1,\infty}(\Omega, \mathbb{R}^N)$ which is a (mere) minimiser of (1.1) over $W_b^{1,\infty}(\Omega, \mathbb{R}^N)$ and a sequence of approximate minimisers $\{u^m\}_{m=1}^\infty$ of (2.1) in the spaces $W_b^{1,qm}(\Omega, \mathbb{R}^N)$ such that, for any $s \geq 1$,

$$u^m \rightharpoonup u^\infty, \text{ weakly as } m \rightarrow \infty \text{ in } W^{1,s}(\Omega, \mathbb{R}^N)$$

along a subsequence. Moreover,

$$(2.2) \quad E_\infty(u^\infty, \Omega) = C_\infty = \lim_{m \rightarrow \infty} (C_m)^{\frac{1}{m}}.$$

By approximate minimiser we mean that u^m satisfies

$$(2.3) \quad |E_m(u^m, \Omega) - C_m| < 2^{-m^2},$$

Finally, for any $A \subseteq \Omega$ measurable of positive measure the following lower semi-continuity inequality holds

$$(2.4) \quad E_\infty(u^\infty, A) \leq \liminf_{m \rightarrow \infty} E_m(u^m, A)^{\frac{1}{m}}.$$

The idea of the proof of (2.3) is based on the use of Young measures in order to bypass the lack of convexity for the approximating L^m minimisation problems (recall that $\mathcal{L}(x, \eta, \cdot)$ is only assumed to be level-convex); without weak lower-semicontinuity of E_m , the relevant infima of the approximating functionals may not be realised. For details we refer to [BJW1] (this method of [BJW1] has most recently been applied to higher order L^∞ problems, see [KP2]). We also note that (2.4) has been established in p. 264 of [BJW1] in slightly different guises, whilst the scaling of the functionals E_m is also slightly different therein. However, it is completely trivial for the reader to check that their proofs clearly establish our Lemma 2.

Proof of Theorem 1. Our goal now is to prove that the candidate u^∞ of Lemma 2 above is actually an Absolute Minimiser of (1.1), which means we need to prove u^∞ satisfies (1.2).

The method we utilise follows similar lines to those of [K9], although technically has been slightly simplified. The main difference is that due to the weaker assumptions than those of [K9], we invoke the general Jensen's inequality for level-convex functions. In [K9] the Lagrangian was assumed to be radial in the third argument, a condition necessary and sufficient for the symmetry of the coefficient matrix multiplying the second derivatives in (1.4); this special structure of \mathcal{L} led to some technical complications. Also, herein we have reduced the number of auxiliary parameters in the energy comparison map (defined below) by invoking a diagonal argument.

Let us fix $\Omega' \Subset \Omega$. Since Ω' is a countable disjoint union of open intervals, there is no loss of generality in assuming Ω' is itself an open interval. By a simple rescaling argument, it suffices to assume that $\Omega' = (0, 1) \subseteq \mathbb{R}$. For, let $\phi \in W_0^{1,\infty}((0, 1), \mathbb{R}^N)$ be an arbitrary variation and set $\psi^\infty := u^\infty + \phi$. In order to conclude, it suffices to establish

$$E_\infty(u^\infty, (0, 1)) \leq E_\infty(\psi^\infty, (0, 1)).$$

Obviously, $u^\infty(0) = \psi^\infty(0)$ and $u^\infty(1) = \psi^\infty(1)$. We define the energy comparison function $\psi^{m,\delta}$, for any fixed $0 < \delta < 1/3$ as

$$\psi^{m,\delta}(x) := \begin{cases} \left(\frac{\delta-x}{\delta}\right) u^m(0) + \left(\frac{x}{\delta}\right) \psi^\infty(\delta), & x \in (0, \delta], \\ \psi^\infty(x), & x \in (\delta, 1-\delta), \\ \left(\frac{1-x}{\delta}\right) \psi^\infty(1-\delta) + \left(\frac{x-(1-\delta)}{\delta}\right) u^m(1), & x \in [1-\delta, 1), \end{cases}$$

where $m \in \mathbb{N} \cup \{\infty\}$. Then, $\psi^{m,\delta} - u^m \in W_0^{1,\infty}((0, 1), \mathbb{R}^N)$ and

$$D\psi^{m,\delta}(x) = \begin{cases} \frac{\psi^\infty(\delta) - u^m(0)}{\delta}, & \text{on } (0, \delta), \\ D\psi^\infty, & \text{on } (\delta, 1-\delta), \\ \frac{\psi^\infty(1-\delta) - u^m(1)}{-\delta}, & \text{on } (1-\delta, 1). \end{cases}$$

Now, note that

$$(2.5) \quad \psi^{m,\delta} \longrightarrow \psi^{\infty,\delta} \text{ in } W^{1,\infty}((0, 1), \mathbb{R}^N), \text{ as } m \rightarrow \infty,$$

because $\psi^{m,\delta} \longrightarrow \psi^{\infty,\delta}$ in $L^\infty((0, 1), \mathbb{R}^N)$ and for a.e. $x \in (0, 1)$ we have

$$\begin{aligned} |D\psi^{m,\delta}(x) - D\psi^{\infty,\delta}(x)| &= \chi_{(0,\delta)} \frac{|u^\infty(0) - u^m(0)|}{\delta} + \chi_{(1-\delta,1)} \frac{|u^\infty(1) - u^m(1)|}{\delta} \\ &\leq \left(\frac{1}{\delta} + \frac{1}{\delta}\right) \|u^m - u^\infty\|_{L^\infty(\Omega)} \\ &= o(1), \end{aligned}$$

as $m \rightarrow \infty$ along a subsequence.

Now, recall that $\psi^{m,\delta} = u^m$ at the endpoints $\{0, 1\}$. Let us also remind to the reader that after the rescaling simplification, $(0, 1)$ is a subinterval of $\Omega \subseteq \mathbb{R}$ whilst (2.3) holds only for the whole of Ω . Since u^m is an approximate minimiser of (2.1) over $W_b^{1,m}(\Omega, \mathbb{R}^N)$ for each $m \in \mathbb{N}$, by utilising the approximate minimality of u^m (given by (2.3)), the additivity of E_m with respect to its second argument, we obtain the estimate

$$E_m(u^m, (0, 1)) \leq E_m(\psi^{m,\delta}, (0, 1)) + 2^{-m^2}.$$

Hence, by Hölder inequality

$$(2.6) \quad \begin{aligned} E_m(u^m, (0, 1))^{\frac{1}{m}} &\leq E_m(\psi^{m,\delta}, (0, 1))^{\frac{1}{m}} + 2^{-m} \\ &\leq E_\infty(\psi^{m,\delta}, (0, 1)) + 2^{-m}. \end{aligned}$$

On the other hand, we have

$$E_\infty(\psi^{m,\delta}, (0, 1)) = \max \left\{ E_\infty(\psi^{m,\delta}, (0, \delta)), \right. \\ \left. E_\infty(\psi^{m,\delta}, (\delta, 1 - \delta)), \right. \\ \left. E_\infty(\psi^{m,\delta}, (1 - \delta, 1)) \right\}$$

and since $\psi^{m,\delta} = \psi^\infty$ on $(\delta, 1 - \delta)$, we have

$$(2.7) \quad E_\infty(\psi^{m,\delta}, (0, 1)) \leq \max \left\{ E_\infty(\psi^{m,\delta}, (0, \delta)), E_\infty(\psi^\infty, (0, 1)), \right. \\ \left. E_\infty(\psi^{m,\delta}, (1 - \delta, 1)) \right\}.$$

Combining (2.5)-(2.7) and (2.4), we get

$$(2.8) \quad E_\infty(u^\infty, (0, 1)) \leq \liminf_{m \rightarrow \infty} \left(\max \left\{ E_\infty(\psi^{m,\delta}, (0, \delta)), E_\infty(\psi^\infty, (0, 1)), \right. \right. \\ \left. \left. E_\infty(\psi^{m,\delta}, (1 - \delta, 1)) \right\} \right) \\ \leq \max \left\{ E_\infty(\psi^\infty, (0, 1)), E_\infty(\psi^{\infty,\delta}, (0, \delta)), \right. \\ \left. E_\infty(\psi^{\infty,\delta}, (1 - \delta, 1)) \right\}.$$

Let us now denote the difference quotient of a function $v : \mathbb{R} \rightarrow \mathbb{R}^N$ as $D^{1,t}v(x) := \frac{1}{t}[v(x+t) - v(x)]$. Then, we may write

$$D\psi^{\infty,\delta}(x) = D^{1,\delta}\psi^\infty(0), \quad x \in (0, \delta), \\ D\psi^{\infty,\delta}(x) = D^{1,-\delta}\psi^\infty(1), \quad x \in (1 - \delta, 1),$$

Note now that

$$(2.9) \quad \begin{cases} E_\infty(\psi^{\infty,\delta}, (0, \delta)) = \max_{0 \leq x \leq \delta} \mathcal{L}(x, \psi^{\infty,\delta}(x), D^{1,\delta}\psi^\infty(0)), \\ E_\infty(\psi^{\infty,\delta}, (1 - \delta, 1)) = \max_{1 - \delta \leq x \leq 1} \mathcal{L}(x, \psi^{\infty,\delta}(x), D^{1,-\delta}\psi^\infty(1)). \end{cases}$$

In view of (2.8)-(2.9), it suffices to prove that there exist an infinitesimal sequence $(\delta_i)_{i=1}^\infty$ such that

$$(2.10) \quad E_\infty(\psi^\infty, (0, 1)) \geq \max \left\{ \limsup_{i \rightarrow \infty} \max_{[0, \delta_i]} \mathcal{L}(\cdot, \psi^{\infty,\delta_i}, D^{1,\delta_i}\psi^\infty(0)), \right. \\ \left. \limsup_{i \rightarrow \infty} \max_{[1 - \delta_i, 1]} \mathcal{L}(\cdot, \psi^{\infty,\delta_i}, D^{1,-\delta_i}\psi^\infty(1)) \right\}.$$

The rest of the proof is devoted to establishing (2.10). Let us begin by recording for later use that

$$(2.11) \quad \begin{cases} \max_{0 \leq x \leq \delta} |\psi^{\infty,\delta}(x) - \psi^\infty(0)| \rightarrow 0, \quad \text{as } \delta \rightarrow 0, \\ \max_{1 - \delta \leq x \leq 1} |\psi^{\infty,\delta}(x) - \psi^\infty(1)| \rightarrow 0, \quad \text{as } \delta \rightarrow 0. \end{cases}$$

Fix a generic $u \in W^{1,\infty}(\Omega, \mathbb{R}^N)$, $x \in [0, 1]$ and $0 < \varepsilon < 1/3$ and define

$$A_\varepsilon(x) := [x - \varepsilon, x + \varepsilon] \cap [0, 1].$$

We claim that there exist an increasing modulus of continuity $\omega \in C(0, \infty)$ with $\omega(0^+) = 0$ such that

$$(2.12) \quad E_\infty(u, A_\varepsilon(x)) \geq \operatorname{ess\,sup}_{y \in A_\varepsilon(x)} \mathcal{L}(x, u(x), Du(y)) - \omega(\varepsilon).$$

Indeed for a.e. $y \in A_\varepsilon(x)$ we have $|x - y| \leq \varepsilon$ and by the continuity of \mathcal{L} and the essential boundedness of the derivative Du , there exist ω such that

$$\left| \mathcal{L}(x, u(x), Du(y)) - \mathcal{L}(y, u(y), Du(y)) \right| \leq \omega(\varepsilon)$$

for a.e. $y \in A_\varepsilon(x)$, leading directly to (2.12). Now, we show that

$$(2.13) \quad \sup_{A_\varepsilon(x)} \left\{ \limsup_{t \rightarrow 0} \mathcal{L}(x, u(x), D^{1,t}u(y)) \right\} \leq \operatorname{ess\,sup}_{A_\varepsilon(x)} \mathcal{L}(x, u(x), Du(y)).$$

Indeed, for any Lipschitz function u , we have

$$(2.14) \quad D^{1,t}u(y) = \frac{u(y+t) - u(y)}{t} = \int_0^1 Du(y + \lambda t) d\lambda,$$

when $y, y+t \in A_\varepsilon(x)$, $t \neq 0$. Further, for any $x \in \Omega$ the function $\mathcal{L}(x, u(x), \cdot)$ is level-convex and the Lebesgue measure on $[0, 1]$ is a probability measure, thus Jensen's inequality for level-convex functions (see e.g. [BJW1, BJW2]) yields

$$\begin{aligned} \mathcal{L}(x, u(x), D^{1,t}u(y)) &= \mathcal{L}\left(x, u(x), \int_0^1 Du(y + \lambda t) d\lambda\right) \\ &\leq \operatorname{ess\,sup}_{0 \leq \lambda \leq 1} \mathcal{L}(x, u(x), Du(y + \lambda t)), \end{aligned}$$

when $y \in A_\varepsilon(x)$, $0 < t < 1$. Consequently,

$$\begin{aligned} \sup_{A_\varepsilon(x)} \left\{ \limsup_{t \rightarrow 0} \mathcal{L}(x, u(x), D^{1,t}u(y)) \right\} &\leq \sup_{A_\varepsilon(x)} \left\{ \limsup_{t \rightarrow 0} \left[\operatorname{ess\,sup}_{0 \leq \lambda \leq 1} \mathcal{L}(x, u(x), Du(y + \lambda t)) \right] \right\} \\ &\leq \sup_{A_\varepsilon(x)} \left\{ \lim_{s \rightarrow 0} \left[\operatorname{ess\,sup}_{y-s \leq z \leq y+s} \mathcal{L}(x, u(x), Du(z)) \right] \right\} \\ &= \operatorname{ess\,sup}_{A_\varepsilon(x)} \mathcal{L}(x, u(x), Du(y)), \end{aligned}$$

as desired. Above we have used the following known property of the L^∞ norm (see e.g. [C])

$$\|f\|_{L^\infty(\Omega)} = \sup_{x \in \Omega} \left(\lim_{\varepsilon \rightarrow 0} \left\{ \operatorname{ess\,sup}_{(x-\varepsilon, x+\varepsilon)} |f| \right\} \right).$$

Note now that by (2.12) we have

$$E_\infty(u, (0, 1)) \geq E_\infty(u, A_\varepsilon(x)) \geq \operatorname{ess\,sup}_{A_\varepsilon(x)} \mathcal{L}(x, u(x), Du(y)) - \omega(\varepsilon)$$

which combined with (2.13) leads to

$$\begin{aligned} E_\infty(u, (0, 1)) &\geq \sup_{A_\varepsilon(x)} \left(\limsup_{t \rightarrow 0} \mathcal{L}(x, u(x), D^{1,t}u(y)) \right) - \omega(\varepsilon) \\ &\geq \limsup_{t \rightarrow 0} \left(\mathcal{L}(x, u(x), D^{1,t}u(x)) \right) - \omega(\varepsilon). \end{aligned}$$

By passing to the limit as $\varepsilon \rightarrow 0$ we get

$$(2.15) \quad E_\infty(u, (0, 1)) \geq \limsup_{t \rightarrow 0} \left(\mathcal{L}(x, u(x), D^{1,t}u(x)) \right),$$

for any fixed $u \in W^{1,\infty}(\Omega, \mathbb{R}^N)$ and $x \in [0, 1]$. Now, since

$$|D^{1,t}u(x)| \leq \|Du\|_{L^\infty(\Omega)}, \quad x \in (0, 1), \quad t \neq 0,$$

for any *finite* set of points $x \in (0, 1)$, there is a common infinitesimal sequence $(t_i(x))_{i=1}^\infty$ such that

$$(2.16) \quad \text{the limit vectors } \lim_{i \rightarrow \infty} D^{1,t_i(x)}u(x) \text{ exists in } \mathbb{R}^N.$$

Utilising the continuity of \mathcal{L} together with (2.15)-(2.16) we obtain

$$(2.17) \quad \begin{aligned} E_\infty(u, (0, 1)) &\geq \limsup_{i \rightarrow \infty} \mathcal{L}(x, u(x), D^{1,t_i(x)}u(x)) \\ &= \mathcal{L}\left(x, u(x), \lim_{i \rightarrow \infty} D^{1,t_i(x)}u(x)\right). \end{aligned}$$

Now we apply (2.17) to $u = \psi^\infty$ and $x \in \{0, 1\}$ to deduce the existence of a sequence $(\delta_i)_{i=1}^\infty$ along which

$$(2.18) \quad \text{the limit vectors } \lim_{i \rightarrow \infty} D^{1,\delta_i}\psi^\infty(0), \quad \lim_{i \rightarrow \infty} D^{1,-\delta_i}\psi^\infty(1) \text{ exist in } \mathbb{R}^N$$

and also

$$(2.19) \quad \begin{aligned} E_\infty(\psi^\infty, (0, 1)) &\geq \max \left\{ \mathcal{L}\left(0, \psi^\infty(0), \lim_{i \rightarrow \infty} D^{1,\delta_i}\psi^\infty(0)\right), \right. \\ &\quad \left. \mathcal{L}\left(1, \psi^\infty(1), \lim_{i \rightarrow \infty} D^{1,-\delta_i}\psi^\infty(1)\right) \right\}. \end{aligned}$$

By recalling (2.9), (2.11) and (2.18), for $\delta = \delta_i$ we obtain

$$(2.20) \quad \begin{aligned} \lim_{i \rightarrow \infty} E_\infty(\psi^{\infty,\delta_i}, (0, \delta_i)) &= \lim_{i \rightarrow \infty} \max_{[0, \delta_i]} \mathcal{L}\left(\cdot, \psi^{\infty,\delta_i}, D^{1,\delta_i}\psi^\infty(0)\right) \\ &= \mathcal{L}\left(0, \psi^\infty(0), \lim_{i \rightarrow \infty} D^{1,\delta_i}\psi^\infty(0)\right), \end{aligned}$$

and also

$$(2.21) \quad \begin{aligned} \lim_{i \rightarrow \infty} E_\infty(\psi^{\infty,\delta_i}, (1 - \delta_i, 1)) &= \lim_{i \rightarrow \infty} \max_{[1 - \delta_i, 1]} \mathcal{L}\left(\cdot, \psi^{\infty,\delta_i}, D^{1,-\delta_i}\psi^\infty(1)\right) \\ &= \mathcal{L}\left(1, \psi^\infty(1), \lim_{i \rightarrow \infty} D^{1,-\delta_i}\psi^\infty(1)\right). \end{aligned}$$

By putting together (2.19)-(2.21), (2.10) ensues and we conclude the proof. \square

REFERENCES

- [A1] G. Aronsson, *Minimization problems for the functional $\sup_x \mathcal{F}(x, f(x), f'(x))$* , Arkiv für Mat. 6 (1965), 33 - 53.
- [A2] G. Aronsson, *Minimization problems for the functional $\sup_x \mathcal{F}(x, f(x), f'(x))$ II*, Arkiv für Mat. 6 (1966), 409 - 431.
- [A3] G. Aronsson, *Extension of functions satisfying Lipschitz conditions*, Arkiv für Mat. 6 (1967), 551 - 561.
- [A4] G. Aronsson, *On the partial differential equation $u_x^2 u_{xx} + 2u_x u_y u_{xy} + u_y^2 u_{yy} = 0$* , Arkiv für Mat. 7 (1968), 395 - 425.
- [A5] G. Aronsson, *Minimization problems for the functional $\sup_x \mathcal{F}(x, f(x), f'(x))$ III*, Arkiv für Mat. (1969), 509 - 512.

- [BJW1] E. N. Barron, R. Jensen and C. Wang, *The Euler equation and absolute minimizers of L^∞ functionals*, Arch. Rational Mech. Analysis 157 (2001), 255 - 283.
- [BJW2] E. N. Barron, R. Jensen, C. Wang, *Lower Semicontinuity of L^∞ Functionals* Ann. I. H. Poincaré AN 18, 4 (2001) 495 - 517.
- [B] J. Broecker, *On variational data assimilation in continuous time*, Quarterly Journal of the Royal Meteorological Society 136, 1906 - 1919 (2010).
- [BS] J. Broecker, I. Szendro, *Sensitivity and Out-Of-Sample Error in Continuous Time Data Assimilation*, Quarterly Journal of the Royal Meteorological Society 138, 785 - 801 (2011).
- [C] M. G. Crandall, *A visit with the ∞ -Laplacian*, in *Calculus of Variations and Nonlinear PDE*, Springer Lecture notes in Mathematics 1927, Cetraro 2005.
- [CKP] G. Croce, N. Katzourakis, G. Pisante, *\mathcal{D} -solutions to the system of vectorial Calculus of Variations in L^∞ via the Baire Category method for the singular values*, preprint.
- [K1] N. Katzourakis, *L^∞ -Variational Problems for Maps and the Aronsson PDE system*, J. Differential Equations, Volume 253, Issue 7 (2012), 2123 - 2139.
- [K2] N. Katzourakis, *∞ -Minimal Submanifolds*, Proceedings of the AMS, 142 (2014) 2797-2811.
- [K3] N. Katzourakis, *On the Structure of ∞ -Harmonic Maps*, Communications in PDE, Volume 39, Issue 11 (2014), 2091 - 2124.
- [K4] N. Katzourakis, *Explicit 2D ∞ -Harmonic Maps whose Interfaces have Junctions and Corners*, Comptes Rendus Acad. Sci. Paris, Ser.I, 351 (2013) 677 - 680.
- [K5] N. Katzourakis, *Optimal ∞ -Quasiconformal Immersions*, ESAIM Control, Opt. and Calc. Var., to appear (2015) DOI: <http://dx.doi.org/10.1051/cocv/2014038>.
- [K6] N. Katzourakis, *Nonuniqueness in Vector-valued Calculus of Variations in L^∞ and some Linear Elliptic Systems*, Comm. on Pure and Appl. Anal., Vol. 14, 1, 313 - 327 (2015).
- [K7] N. Katzourakis, *An Introduction to viscosity Solutions for Fully Nonlinear PDE with Applications to Calculus of Variations in L^∞* , Springer Briefs in Mathematics, 2015, DOI 10.1007/978-3-319-12829-0.
- [K8] N. Katzourakis, *Generalised solutions for fully nonlinear PDE systems and existence-uniqueness theorems*, ArXiv preprint, <http://arxiv.org/pdf/1501.06164.pdf>.
- [K9] N. Katzourakis, *Absolutely minimising generalised solutions to the equations of vectorial Calculus of Variations in L^∞* , ArXiv preprint, <http://arxiv.org/pdf/1502.01179.pdf>.
- [K10] N. Katzourakis, *A new characterisation of ∞ -Harmonic and p -Harmonic maps via affine variations in L^∞* , ArXiv preprint, <http://arxiv.org/pdf/1509.01811.pdf>.
- [K11] N. Katzourakis, *Mollification of \mathcal{D} -solutions to fully nonlinear PDE systems*, ArXiv preprint, <http://arxiv.org/pdf/1508.05519.pdf>.
- [K12] N. Katzourakis, *Solutions of vectorial Hamilton-Jacobi equations are rank-one Absolute Minimisers in L^∞* , ArXiv preprint, <http://arxiv.org/pdf/1604.00802.pdf>.
- [KP1] N. Katzourakis, T. Pryer, *On the numerical approximation of ∞ -Harmonic mappings*, ArXiv preprint, <http://arxiv.org/pdf/1511.01308.pdf>.
- [KP2] N. Katzourakis, T. Pryer, *Second order L^∞ variational problems and the ∞ -Polylaplacian*, ArXiv preprint, <http://arxiv.org/pdf/1605.07880.pdf>.

DEPARTMENT OF MATHEMATICS , COLLEGE OF SCIENCE, UNIVERSITY OF BASRA, BASRA, IRAQ
& DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF READING, WHITEKNIGHTS, PO
Box 220, READING RG6 6AX, UK

E-mail address: h.a.h.abugirda@student.reading.ac.uk

DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF READING, WHITEKNIGHTS,
PO Box 220, READING RG6 6AX, UK

E-mail address: n.katzourakis@reading.ac.uk