

Potential ecological effects of Piriformospora indica, a possible biocontrol agent, in UK agricultural systems

Article

Accepted Version

Creative Commons: Attribution-Noncommercial-No Derivative Works 4.0

Rabiey, M., Ullah, I., Shaw, L. J. and Shaw, M. W. (2017) Potential ecological effects of Piriformospora indica, a possible biocontrol agent, in UK agricultural systems. *Biological Control*, 104. pp. 1-9. ISSN 1049-9644 doi: <https://doi.org/10.1016/j.biocontrol.2016.10.005> Available at <https://centaur.reading.ac.uk/67733/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

To link to this article DOI: <http://dx.doi.org/10.1016/j.biocontrol.2016.10.005>

Publisher: Elsevier

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online

1 **Potential ecological effects of *Piriformospora indica*, a possible biocontrol agent, in**
2 **UK agricultural systems**

3 Mojgan Rabiey^{a*}, Ihsan Ullah^b, Liz J Shaw^c, and Michael W Shaw^b

4 ^a School of Biological Sciences, Knight Building, University of Reading, RG6 6AJ

5 ^b School of Agriculture, Policy and Development, University of Reading, RG6 6AR,
6 UK;

7 ^c Soil Research Centre, School of Archaeology, Geography and Environmental Science,
8 University of Reading, RG6 6AB, UK.

9

10 *Corresponding author: Mojgan Rabiey, m.rabiey@reading.ac.uk, Tel: +447415414119

11

12

13

14

15

16

17

18

19

20

21

22

23 **Abstract**

24 *Piriformospora indica* (Sebacinaceae), a root endophytic fungus, was originally isolated
25 from an arid sub-tropical soil. *P. indica* forms mutualistic symbioses with a broad range
26 of host plants, increases biomass production, resistance and tolerance to fungal
27 pathogens and abiotic stresses. These characteristics make it a very attractive
28 component of more sustainable agriculture. So, it is desirable to understand its wider
29 ecosystem effects. We determined how long *P. indica* could survive in the soil and how
30 it interacts with other soil microorganisms and some important arable weeds.

31 Survival of *P. indica* in the soil, under winter and summer conditions in the UK was
32 tested by isolating DNA and RNA of *P. indica* from pots of soil which had been left
33 open to winter-summer weather conditions without host plants, followed by PCR and
34 reverse transcription-PCR (RT-PCR) with *P. indica*-specific primers. *P. indica* effects
35 on other soil and root microorganisms were tested by PCR-denaturing gradient gel
36 electrophoresis analysis of DNA extracted from soil and roots from pots in which *P.*
37 *indica*-infected wheat had been grown. The effect of *P. indica* on growth of black-grass
38 (*Alopecurus myosuroides*), wild-oat (*Avena fatua*) and cleavers (*Galium aparine*) was
39 tested alone and in competition with wheat.

40 In soil *P. indica*-mRNA and DNA could still be detected after eight months, but not
41 after 15 months. Soils from *P. indica*-inoculated pots had distinct fungal and bacterial
42 species communities which were more diverse than non-inoculated controls. *P. indica*
43 infected *A. myosuroides* and *A. fatua* but was not detected in *G. aparine*. The average
44 above-ground competitiveness of the weeds with wheat was decreased.

45 If applied to field crops in the UK, *P. indica* would be persistent for up to 15 months
46 and likely to alter competitive relations within vegetation. Increased soil microbial
47 diversity during the first eight weeks after inoculation, although usually desirable, could
48 alter soil composition or functioning.

49 **Keywords:** denaturing gradient gel electrophoresis, microbial diversity, reverses
50 transcription-PCR, root endophytic fungus, soil microorganisms, sustainable
51 agriculture.

52

53 **1. Introduction**

54 *Piriformospora indica*, a root endophytic fungus (Sebacinales: Basidiomycota), was
55 first found in the Thar dessert of India (Verma et al., 1998), an arid region which
56 experiences extreme day-time heat and diurnal temperature fluctuations as well as
57 extended drought. *P. indica* promotes plant growth, increases root and above ground
58 biomass and final yield during its mutualistic relationship with a wide variety of plants
59 (Shrivastava & Varma, 2014). It increases resistance of several hosts from diverse
60 families to many biotic stresses under glasshouse and field conditions (Waller et al.,
61 2005, Deshmukh & Kogel, 2007, Ghahfarokhy et al., 2011, Harrach et al., 2013).
62 Tolerance to abiotic stresses is also increased in a variety of hosts (Zarea et al., 2012,
63 Alikhani et al., 2013, Ghabooli et al., 2013, Varma et al., 2013). Our previous studies
64 showed that *P. indica* protected wheat from Fusarium crown rot in pot experiments
65 (Rabiey et al., 2015). Our work also suggests that Fusarium head blight and mycotoxin
66 deoxynivalenol contamination are reduced in pot experiments under UK weather
67 conditions (Rabiey & Shaw, 2015). Tests on a field scale are therefore attractive, but as

68 the organism is an alien, it would first be necessary to understand its ecosystem effects
69 and persistence.

70 How *P. indica* interacts with other soil microorganisms is still unclear. Endophytic
71 fungal symbionts can have profound effects on plant ecology, fitness, and evolution
72 (Brundrett, 2006), shaping plant communities (Clay & Holah, 1999), increasing plant
73 tolerance to abiotic stresses (Murphy et al., 2015), increasing plant resistance to
74 pathogens (Rodriguez et al., 2009, Murphy et al., 2014) and manifesting strong effects
75 on the community structure and diversity of associated organisms (e.g. bacteria,
76 nematodes and insects; Omacini et al. (2001)). Studies on the effects of arbuscular
77 mycorrhizal fungi (AMF) on rhizosphere bacteria have shown variable results, with
78 both negative (decreasing the population of bacteria) (Christensen & Jakobsen, 1993,
79 Amora-Lazcano et al., 1998) and positive (increasing the population of bacteria)
80 (Andrade et al., 1997, Abdel-Fattah & Mohamedin, 2000) effects. The variable results
81 could be due to the fact that some bacteria are being stimulated and others being
82 repressed by AMF (Wamberg et al., 2003). Söderberg et al. (2002) suggested that the
83 effect of AMF differed between plant species; the strength of the effect on the bacterial
84 community in the rizosphere depended more on the plant species than on AMF
85 colonisation. If *P. indica* is going to be applied to crops, a clear picture of how it affects
86 other soil microorganisms would be needed, as the soil microflora plays a major role in
87 the availability of nutrients to plants and has a strong influence on plant health and
88 productivity.

89 Weed competition can threaten crop quality and quantity and ultimately the farmer's
90 profitability (Bockus et al., 2010); it is usually managed by herbicide application. The

91 key herbicide-resistant weed species of arable crops in the UK are: black-grass
92 (*Alopecurus myosuroides*), wild-oats (*Avena fatua*), cleavers (*Gallium aparine*), Italian
93 rye-grass (*Lolium multiflorum*), common poppy (*Papaver rhoeas*), common chickweed
94 (*Stellaria media*), and scentless mayweed (*Tripleurospermum inodorum*) (Bond et al.,
95 2007, Moss et al., 2011, Hull et al., 2014). These are also important world-wide and in
96 other crops (Yu et al., 2013). Herbicide resistance in the UK is an important and
97 increasing problem, as in other parts of the world including western, central and
98 northern Europe (Mennan & Isik, 2004, Moss et al., 2007, Bertholdsson, 2012). *P.*
99 *indica* has a wide range of hosts which might include weeds as well. If *P. indica* was as
100 beneficial to weeds as to wheat, it could make weed control more difficult, or increase
101 the damage done by weeds; alternatively, it might increase the competitiveness of wheat
102 against some species or in some settings, which would be useful in managing herbicide
103 resistant weeds. Also, the spread of *P. indica* might have side-effects outside arable
104 fields.

105 In this study the following hypotheses were tested: *P. indica* would survive the UK
106 weather and soil conditions; *P. indica* would not affect the composition of the bulk soil
107 or root-zone microflora; and *P. indica* would be as beneficial to weeds as to wheat.

108 **2. Materials and methods**

109 **2.1. *P. indica* survival and viability experiment**

110 The utility of mRNA and DNA measurements as indicators of viability of *P. indica* was
111 determined by performing RT-PCR and PCR on heat and cold treated pure cultures of
112 *P. indica*. For this purpose, *P. indica* was obtained from Dr. Patrick Schafer, Warwick
113 University, UK (originally from German Collection of Microorganisms and Cell

114 Culture, strain number DSM 11827) and mycelia was grown on complex
115 modified *Aspergillus* liquid medium (CM medium) (Pham *et al.*, 2004) and incubated
116 on an orbital shaker at 140 rpm at room temperature (21 ± 1 °C) for two weeks. Samples
117 were then kept at 80 °C in a hot water bath for 6 hours, then stored at -80 °C for 6 hours,
118 one and four weeks. After storage, separate samples of mycelia were transferred to
119 potato dextrose agar to check whether they would grow and used for RNA and DNA
120 extraction followed by RT-PCR and PCR respectively. This experiment was repeated to
121 confirm the results.

122 *P. indica* survival in the soil under UK weather conditions was tested in different soil
123 types based on either the soil series or textural classification and each soil was under a
124 different crop management. The soils were collected from the Reading University Farm
125 at Sonning (grid ref: SU76187547). These were (1) a Clay Loam (CL) of the Neville
126 series, from an area under winter barley which had previously been under winter wheat;
127 (2) a Sandy Clay Loam (SCL) of the Sonning series from an area under ryegrass at the
128 time and for the previous two years; (3) a Loamy Sand (LSO) of the Rowland series,
129 under organic management, from an area under faba bean cultivation; (4) a Loamy Sand
130 (LS) of the Rowland series, under non-organic management, from an area under
131 ryegrass cultivation. The experiment was carried out between December 2013 and
132 March 2015 at the University of Reading, under outdoor weather conditions. Six pots (3
133 L, top diameter: 18 cm, bottom diameter: 14 cm, depth: 15 cm) were filled with each
134 soil. Five out of six pots received 4 g of liquid culture of *P. indica* inoculum containing
135 an unquantified mixture of chlamydospores and mycelium and mixed thoroughly with
136 the soil. The control pot received 4 g of sterilised liquid culture of *P. indica* inoculum.

137 The pots were placed in holes with the tops level with the surrounding soil level to
138 make temperature fluctuations realistic. Around 50 g of each soil type was collected,
139 with a small core (diameter: 12 mm, depth: 8 cm) from the middle of pots, at three and
140 half months (mid-March 2014), 8 months (end of July 2014) and 15 months (end of
141 March 2015) after inoculation with *P. indica*. When collecting the samples, they were
142 kept in a cool box on ice and transferred immediately to -20 °C before DNA and RNA
143 were extracted and PCR or RT-PCR performed. Maximum and minimum temperatures
144 of soil in the pots were recorded every 2 days by a digital thermometer placed in the
145 centre of one of the pots.

146 **2.2. Soil community composition**

147 To examine whether *P. indica* affects other soil microorganisms, wheat was grown in 3
148 L pots containing one of two soil types, SCL or LSO, as above. Winter wheat seeds, cv.
149 Battalion, were surface disinfected by rinsing for 2 mins in 20 mL L⁻¹ sodium
150 hypochlorite (Fisher Scientific UK Ltd, UK), followed by three rinses in sterilized
151 distilled water, and germinated on damp filter paper in a Petri dish at room temperature
152 (21 ± 1 °C) under natural indoor light for 48 hours. No micro-organisms grew from a
153 sample of seeds so treated and placed on PDA plates for one week. Pre-germinated
154 seeds were planted into 3 L pots (one seed per pot). This experiment had a 2×2×4
155 factorial combinations of ±*P. indica* × two soil types × four harvesting points, with two
156 replications completely randomised. The pots were incubated at temperatures ranging
157 between 15 and 25 °C; humidity and light were not controlled. Inoculation with 4 g
158 liquid culture of *P. indica* mixed with soil was done at the time of sowing. Root and soil
159 samples were collected at 2, 4, 6 and 8 weeks after inoculation (wai) for DNA

160 extraction, PCR and DGGE analysis, as below. Samples were transferred and stored as
161 described above.

162 **2.2.1. DNA and RNA isolation**

163 Total genomic DNA from *P. indica* and root samples was isolated using a DNeasy plant
164 mini kit (QIAGEN, UK), and from soil samples by using a PowerLyzer™ PowerSoil®
165 DNA Isolation kit (CAMBIO Ltd, UK) following the manufacturer's instructions. Total
166 RNA from *P. indica* was isolated using a RNeasy Plant Mini Kit (QIAGEN, UK), and
167 from soil samples by using a RNA PowerSoil® Total RNA Isolation kit (CAMBIO Ltd,
168 UK). Samples were stored at -20 °C until required. Bulk DNA concentration was
169 measured using a NanoDrop-lite spectrophotometer (Thermo Scientific, Life
170 Technologies Ltd, UK). The extent of shearing of DNA and RNA was determined by
171 electrophoresis of an aliquot of DNA in a 1 % agarose gel in 1x TAE buffer.

172 **2.2.2. Primer development and PCR condition for RT-PCR study**

173 The gene-specific primer for the RT-PCR study was designed using the PRIMER
174 BLAST tool from NCBI (<http://www.ncbi.nlm.nih.gov/tools/primer-blast>) to amplify
175 fragments of the *P. indica* mRNA for EF-1-alpha (*TEF* gene, forward: 5-
176 CCACCATCACTGAAGTCCCTC-3 and reverse: 5-TCAATGCCACCGCACTTGTA-
177 3, 148 bp, accession number AJ249912.1, <http://www.ncbi.nlm.nih.gov>). The primers
178 were supplied by Invitrogen (Thermo Scientific, Life Technologies Ltd, UK). To assess
179 specificity of the primers for the targeted gene, RT-PCR was done using RNA isolated
180 from a pure culture of *P. indica*. The PCR products of the selected primer were sent to
181 Source Bioscience (<http://www.sourcebioscience.com/>) for sequencing to verify their
182 specificity.

183 EF (EF-1-alpha (*TEF* gene)) primer amplified cDNA of 148 bp and gDNA of 227 bp.
184 The PCR amplicon sequence corresponded to genomic sequence from 1547 to 1756 bp
185 of the *P. indica TEF* gene, GenBank: accession number AJ249911.2, as expected.
186 PCR was performed in 0.2 mL PCR tubes (Fisher Scientific UK Ltd, UK) with 20 µL
187 final reaction volume containing 2x Biomix PCR master mix, 0.25 µM forward and
188 reverse primer, and template genomic DNA. Amplification was performed in a thermal
189 cyclor (Applied Biosystems® GeneAmp® PCR System 9700, Thermo Scientific, Life
190 Technologies Ltd, UK) programmed as: 94 °C for 5 min followed by 35 cycles of 94 °C
191 for 30 s, 56 °C for 45s and 72 °C for 30 s, followed by incubation at 72 °C for 7 min.
192 Amplification was confirmed by electrophoresis of an aliquot of the PCR products in 2
193 % agarose gel in 1x TAE buffer.

194 **2.2.3. Reverse Transcription-PCR (RT-PCR)**

195 RT-PCR for *P. indica* was performed by using Invitrogen SuperScript® III First-Strand
196 Synthesis SuperMix (Life Technologies Ltd, UK) in a 20 µL final reaction volume
197 using 10 µL 2× RT Reaction Mix, 2 µL RT Enzyme Mix, RNase-free water and 4 µL *P.*
198 *indica* RNA. Reverse transcription was done in a thermal cyclor. Samples were first
199 incubated at 50 °C for 30 minutes, then held at 85 °C for 5 minutes and then chilled on
200 ice for 5 min. Thereafter, 1 µL *E. coli* RNase H was added to the tube which was then
201 incubated at 37 °C for 20 minutes. PCR was then performed using the complementary
202 DNA (cDNA) obtained from the reverse transcription.

203 RT-PCR for soil samples was performed by using a One-Step RT-PCR Kit (QIAGEN,
204 UK), in a 25 µL final reaction volume using 5 µL 5x QIAGEN OneStep RT-PCR
205 Buffer, 1 µL dNTP Mix, 1 µL of Enzyme Mix, 0.6 µM of each primer, RNase-free

206 water and 4 μ L *P. indica* and samples RNA. Thermal cycler was set up at 30 min 50 °C,
207 15 min 95 °C, 35 cycles of 94 °C for 30 s, 56 °C for 45 s, 72 °C for 30 s, followed by
208 incubation at 72 °C for 7 min.

209 **2.2.4. Primer and PCR condition for DGGE study**

210 Bacterial 16S rRNA genes, from the extracted DNA, were amplified using the primer
211 341F-

212 CGCCCGCCGCGCGCGGGCGGGGCGGGGGCACGGGGGGCCTACGGGAG

213 GCAGCAG and 534R-ATTACCGCGGCTGCTGG (Muyzer et al., 1993). Fungal 18S

214 rRNA genes were amplified using the primer NS1F-GTAGTCATATGCTTGTCTC and

215 GCFung-R-

216 CGCCCGCCGCGCCCCGCGCCCGGCCCGCCCGCCCCCGCCCCATTCCCCGTTAC

217 CCGTTG (Hoshino & Morimoto, 2008).

218 The PCR was performed in a 20 μ L final reaction volume using 2 \times Biomix PCR master

219 mix, 50 pmol μ L⁻¹ (for bacterial study) and 0.3 pmol μ L⁻¹ (for fungal study) of forward

220 and reverse primer, and sample DNA. Touchdown PCR for the bacterial study was

221 performed in a thermal cycler set up at 94 °C for 10 min, denaturation at 94 °C for 1

222 min, an annealing temperature which was set at 65 °C initially, then decreased by 1 °C

223 after each 2 cycles until it reached 55 °C. Primer extension was performed at 72 °C for

224 2 min. The above reaction was performed for 20 cycles, followed by 15 cycles of 94 °C

225 for 1 min, 55 °C for 1 min and 72 °C for 2 min. A final extension step was performed

226 for 10 min at 72 °C (Sasaki et al., 2009).

227 For the fungal primers, amplification was set at 94°C for 2 min, 30 cycles of 94°C for
228 15 s, 50 °C for 30 s and 68 °C for 30 s with a final extension of 72 °C for 5 minutes
229 (Hoshino & Morimoto, 2008).

230 **2.2.5. Denaturing gradient gel electrophoresis of fungi and bacteria**

231 Denaturing gradient gel electrophoresis was performed according to the method
232 described by Muyzer et al. (1993) (for bacterial study) and Hoshino & Morimoto (2008)
233 (for fungal study) using the Bio-Rad DCode™ Universal Mutation Detection System.
234 PCR samples (20 µL+loading dye) were applied directly onto 8 % (wt/vol)
235 polyacrylamide gels (40 % acrylamide 37.5:1) with denaturing gradients of 40-60 %
236 (for bacteria) and denaturing gradients of 20-40 % (for fungi), where 60 % denaturant
237 comprised 24 mL 100 mL⁻¹ Formamide and 25.2 g 100 mL⁻¹ Urea (Sigma Aldrich
238 Company Ltd, UK). Electrophoresis was performed at a constant voltage of 75 V and a
239 temperature of 60 °C for 17 hours for bacteria and voltage of 50 V and a temperature of
240 60 °C for 20 hours for fungi. After electrophoresis, the gels were fixed (0.5 % glacial
241 acetic acid and 10 % ethanol) and silver-stained (1 g L⁻¹ silver nitrate), scanned, and the
242 images analysed.

243 **2.2.5.1. Statistical analysis of DGGE banding patterns**

244 The DNA bands that migrated within each gel to the same relative distance were each
245 ascribed the same label. In each lane, corresponding to a sample, the presence of a band
246 with that label was scored 1 and absence scored 0. The band corresponding to *P. indica*
247 band (which had the same position in all *P. indica*-inoculated samples) was not included
248 in the scoring. These data were then analysed by two methods:

249 (i) Canonical variates analysis (CVA, GenStat 17th ed, VSN) was used to evaluate
250 differences in community structure and allow the comparison of community profiles
251 between groups of samples. CVA differentiate between groups variation, using a trace
252 statistic as a summary of differentiation. CVA will produce a visualization of the data
253 that shows groups as clearly separated, whether the differences are genuine or the result
254 of chance sampling effects. The natural measure of how separate the groups found are is
255 the trace of the matrix ratio $W^{-1}B$, where B is the matrix of between-group sums of
256 squares and products and W is the matrix of within-group sums of squares and products.
257 This measure and a randomization test (10,000 replicates) were used. The significance
258 of the observed separation between groups, to determine whether groups were more
259 distinct than expected by chance, was assessed by randomisation tests of 10,000
260 replicates (Rajaguru & Shaw, 2010).

261 (ii) Shannon-Wiener diversity index (H' , GenStat 17th ed, VSN) was used to quantify
262 the diversity of species (bands) present in a group of samples. This index was calculated
263 by the following equation:

$$264 \quad H' = - \sum_i (R_i / R) \times \log (R_i / R)$$

265 where R_i is the total number of occurrences of band i in a group of observations, and R
266 is total number of bands of any type observed in the group. Confidence intervals for the
267 index were obtained by randomly re-sampling band abundances from a multinomial
268 with the observed probabilities of each band type, and re-calculating the index.

269 **2.3. *P. indica* interaction with weeds**

270 Black-grass (*Alopecurus myosuroides*, 16 seeds per pot), wild-oat (*Avena fatua*, 6 seeds
271 per pot), cleavers (*Galium aparine*, 3 seeds per pot) with and without wheat (6 seeds per

272 pot) were planted in 5 L pots (top diameter: 22.5 cm, bottom diameter: 16.5 cm, depth:
273 17.5 cm) at a depth of 1 cm in one part non-sterilised vermiculite (Medium, Sinclair,
274 UK) and one part sand, mixed with 4 g L⁻¹ of slow release fertiliser (3-4 months,
275 Osmocote® Pro), with and without 4 g pot⁻¹ of liquid *P. indica* inoculum mixed into the
276 soil. Four replicates, distributed in four randomised blocks, were used with the
277 following factorial combinations of treatments: ± *P. indica*, ± wheat, and three weed
278 species. Wheat alone with and without *P. indica* was included as a control.

279 The pots were placed outside under natural conditions in the first two weeks of
280 November/2014 for vernalisation, and then incubated in the glasshouse at temperatures
281 ranging between 5 °C and 18 °C; humidity and light were not controlled. All pots were
282 harvested, when wheat flag leaf was fully emerged (Zadoks Growth Stage (GS) 39;
283 Zadoks et al. (1974)), and roots teased apart, washed and separated from the above
284 ground parts before drying and weighing. This experiment was repeated to confirm the
285 results.

286 In a separate experiment, to confirm the colonisation of weed roots with *P. indica*
287 microscopically, seeds of black-grass, wild-oat and cleavers were planted separately in
288 1 L pots (top diameter: 13 cm, bottom diameter: 10 cm, depth: 11 cm) in one part non-
289 sterilised vermiculite (Medium; Sinclair) and one part sand, and inoculated with *P.*
290 *indica* at sowing. The roots were harvested at one and four weeks after inoculation,
291 stained according to the method described in Rabiey et al., 2015, and viewed under a
292 microscope with 10x and 40x objectives.

293 Competitiveness of each weed species with wheat was quantified as log (wheat
294 biomass/weed biomass).

295 **2.4. Statistical analysis of pot experiments**

296 ANOVA was used to analyse all data using Genstat 17th ed, (VSN, UK) with
297 appropriate blocking.

298 **3. Results**

299 **3.1. Weather conditions during 2013-15**

300 Winter 2013-14 was an “exceptionally” stormy season, with at least 12 major winter
301 storms affecting the UK. Mean temperatures and total rainfall were 2 °C and 211 mm
302 respectively, above the long-term average over Reading (Nov-Mar 2013-14 average 7.1
303 °C). Soil temperature was 1 °C above average. Soil froze on only five occasions (Fig.
304 1).

305 Following this, the mean air and soil temperature of spring and summer 2014 was near
306 the average (April-June 2014 average 13.5 °C); total rainfall was, 55 mm and 31 mm
307 respectively, above the long term average (Fig. 1).

308 The weather of autumn 2014 was warm (Sep-Nov 2014 average 12.7 °C), 1.6 °C above
309 the average with the number of air frosts well below average. Rainfall totals and soil
310 temperature were above average, 11 mm and 1.5 °C respectively. Winter 2014-15 was
311 sunny with mean air (Dec-Mar 2014/15 average 5.6 °C) and soil temperature near
312 average. Soil froze on 20 occasions. Rainfall totals were 13 mm below average (Fig. 1.
313 www.met.reading.ac.uk/weatherdata).

314 **3.2. *P. indica* viability under UK winter weather conditions**

315 The viability of *P. indica* mycelia was tested under laboratory conditions. Exposure of
316 mycelia to 80 °C for 6 hours, then to -80 °C for 6 hours, one or four weeks killed them:
317 plates showed no growth of fungus after one month. RT-PCR detected *P. indica* mRNA

318 after 6 hours exposure to 80 °C then 6 hours at -80 °C, but did not detect *P. indica*
319 mRNA after exposure to 80 °C followed by one or four weeks storage at -80 °C. PCR
320 detected DNA in all treatments (Table 1).

321 RNA and DNA of *P. indica* were successfully isolated from all four soil after winter
322 2013 (collected mid March 2014) (Table 2). DNA of *P. indica* was successfully isolated
323 from all different soil types following a UK spring and summer (collected end of July
324 2014), but RNA could be detected in only six of the pots. After 15 months (collected
325 mid March 2015), neither RNA, nor DNA of *P. indica* could be detected from any of
326 the soils (Table 2). *P. indica* could not be detected in the controls that was not
327 inoculated with *P. indica*, which shows the primers could only detect *P. indica* mRNA.

328 **3.3. *P. indica* effect on other soil microorganisms**

329 **3.3.1. Canonical variate analysis**

330 Canonical variate analysis was used to differentiation between groups variation, using a
331 trace statistic as a summary of differentiation. Canonical variate analysis of band
332 patterns (Fig. 2), including both bacteria and fungi separated the four different harvested
333 time points (trace: 1.9, $P < 0.0001$), mainly because the first sample was distinct (Fig. 3
334 a). Root samples were clearly distinguishable from soil samples (trace: 3.9, $P < 0.0001$,
335 Fig. 3 b), and soil types were clearly distinct (trace: 1.6, $P < 0.0001$, Fig. 3 c). *P. indica*-
336 inoculated and non-inoculated samples were distinct (trace: 0.6, $P = 0.001$, Fig. 3 d), *P.*
337 *indica*-inoculated were distinguishable from non-inoculated samples by CVA when
338 restricted to either fungal (trace: 1.1, $P < 0.03$, Fig. 3 e), or bacterial primers (trace: 1.2,
339 $P < 0.02$, Fig. 3 f) or soil samples (trace: 2.9, $P < 0.0001$, Fig. 3 g) but not root samples
340 (trace: 0.6, $P = 0.6$, Fig. 3 h).

341 To check the interaction between the effects of *P. indica* and soil-root zones combined
342 factors were created. CVA of groups of samples classified by both *P. indica* inoculation
343 and root-soil zone, including both bacterial and fungal bands, separated *P. indica*-
344 inoculated from non-inoculated samples (trace: 5.5, $P < 0.0001$, Fig. 3 i).

345 **3.3.2. Shannon-Wiener diversity index**

346 Samples harvested at different time points did not differ in diversity. Rowland series
347 soils (LSO) had more fungal and bacterial band diversity than Sonning series (SCL).
348 Both types of soil had more fungal and bacterial band diversity in the presence of *P.*
349 *indica* (Fig. 4) and samples inoculated with *P. indica* had more bands of all types than
350 non-inoculated samples. Root samples had more fungal species diversity when *P. indica*
351 was present, but slightly fewer bacterial species diversity (Fig. 5).

352 **3.4. *P. indica* interaction with weeds**

353 Two *Avena fatua* root samples out of 10 were colonised by *P. indica* at two wai and
354 three out of ten at four wai. Two *Alopecurus myosuroides* root samples out of ten were
355 colonised at four wai. No *Galium aparine* root samples (of ten samples) were colonised.
356 *P. indica* application at sowing time increased wheat shoot and root biomass by 33 %
357 (main effect $P = 0.05$) and 100 % (main effect $P = 0.02$) respectively, as expected (Table
358 3; supporting information Table 1).

359 *P. indica* increased root biomass, averaged over *Avena fatua*, *Alopecurus myosuroides*
360 and *G. aparine*, by 35 % ($P = 0.04$). As expected, competition reduced root biomass (by
361 about 26 %, $P = 0.05$) and there were differences between species ($P = 0.03$; *A. fatua* was
362 about 50 % heavier than the other two species). All interactions were non-significant
363 ($P > 0.4$). In particular, the effect of inoculation did not differ between weed species, and

364 the effect of inoculation did not differ in the competition pots (Table 3; supporting
365 information Table 1).

366 Shoot biomass of all plants was decreased about 24 % ($P=0.005$) by competition and
367 differed greatly between the species ($P=0.001$) because *G. aparine* had a lower biomass.
368 The effect of *P. indica* was slight (a 12 % increase; $P=0.2$) and no interactions were
369 significant ($P>0.2$ in all cases) (Table 3; supporting information Table 1).

370 The average competitiveness with wheat of *Avena fatua*, *Alopecurus myosuroides* and
371 *G. aparine*, measured by the ratio of shoot weights, was reduced by 40 %
372 (backtransformed from the \log_{10} scale; $P=0.02$) when *P. indica* was present in the soil
373 (Table 4; supporting information Table 2). Although the competitiveness differed
374 significantly between species, no interaction terms were significant ($P>0.5$). There were
375 no significant differences in competitiveness measured by the ratio of root weights
376 ($P>0.13$ for all main and interaction terms). This experiment was repeated (supporting
377 information Table 3, 4) and the results were confirmed.

378

379 **4. Discussion**

380 This study indicates (1) that *P. indica* can survive the UK weather and soil conditions
381 for period of months, even when there is no host present (Table 2); (2) that the
382 inoculation of *P. indica* to soil has a substantial effect on soil and wheat root-associated
383 microflora at least during the first eight weeks (Fig. 3,4,5); (3) that *P. indica* affects at
384 least two of three tested native arable weeds, and alters their competitive relations with
385 wheat, and with each other and at least some other species (Table 3,4; and supporting
386 information Table 3,4).

387 If it were used in field applications in England, *P. indica* would probably remain active
388 in the soil for up to 15 months and there might be no need to re-apply it within season.
389 However, in the event of adverse side-effects, it would be hard or impossible to
390 eradicate. The longevity of *P. indica* inoculum in the soil, coupled with its strong
391 growth promotional effects on some species might alter the competitive relations
392 between existing native species. It also might affect other methods of disease
393 management as the altered soil microflora could influence crop physiology in
394 undetermined ways. The longevity of inoculum in soil might be specifically due to the
395 mild weather of 2013-15 compared with the climatic average. However, the UK is
396 predicted to experience milder winter conditions over the next decades (UKCIP;
397 www.ukcip.org.uk/).

398 Exposure of *P. indica* to heat (80 °C) then immediately to -80 °C, killed the mycelia
399 (Table 1). mRNA can be used as an indicator of *P. indica* viability, as it could not be
400 detected a few hours after mycelia of *P. indica* were killed, while DNA of *P. indica*
401 could be detected even four weeks after mycelium was killed (Table 1). This agrees
402 with other studies. Herdina et al. (2004) concluded that mycelium of *Gaeumannomyces*
403 *graminis* var. *tritici* killed by heating to 55 °C for 1 hour and DNA could still be
404 detected by PCR after eight days. Chimento et al. (2012) killed *Phytophthora ramorum*
405 mycelia by rapid lyophilisation and could detect DNA three months later while mRNA
406 was only detected up to one week after the treatment, despite its relatively mild nature.

407 The DGGE analysis showed detectable changes in the microbial community structure
408 and increased diversity in the fungal and bacterial community of both root and soil
409 samples inoculated with *P. indica*, which are reflected in increases in Shannon diversity

410 indices (Fig. 3,4,5). How this might affect soil function is unknown. There is lots of
411 debate about the importance of microbial community structure and diversity for soil
412 function, plant productivity, resilience and stability. Changes in the composition of the
413 soil microbial community can change ecosystem process rates, specifically
414 decomposition, and affect plant productivity (positively, negatively or not at all)
415 depending on the composition of the initial microbial community (McGuire & Treseder,
416 2010, Gera Hol et al., 2015). The two soils tested differed in their initial diversity, but
417 responded similarly to inoculation with *P. indica*. The increase in microbial diversity
418 might be due to *P. indica* causing changes in root exudate (composition and quantity)
419 patterns, or directly through fungal exudates, as reported for AMF (Barea, 2000,
420 Gryndler, 2000, Jeffries et al., 2003).

421 The primer sets 341/534 and NS1/GCFung for the bacterial and fungal community
422 study were used as Muyzer et al. (1993) and Hoshino and Morimoto (2008) suggested
423 these primer sets could most clearly discriminate bacterial and fungal communities in
424 the soil. To obtain more specific results from DGGE, PCR primers must amplify only
425 specific groups of fungi and bacteria (Jumpponen, 2007, Hoshino, 2012). The DGGE
426 gave an overview of *P. indica*-induced changes in bacterial and fungal community
427 structure but next generation sequencing approaches could be employed in the future for
428 in depth study of the effects of *P. indica* on community structure and composition
429 (Rincon-Florez et al., 2013).

430 *P. indica* has a very wide host range, and may be able to interact with and improve
431 growth of economically-damaging weeds as well as crops. The effect of *P. indica* on
432 *Alopecurus myosuroides*, *Avena fatua* and *Galium aparine*, three of the most important

433 weeds in UK wheat production was evaluated. As expected, the weeds reduced wheat's
434 root and shoot biomass significantly. *P. indica* did not colonise *G. aparine*, but did
435 colonise *A. fatua* and *A. myosuroides*, though less than wheat (Table 3,4; supporting
436 information Table 3,4). The average root biomass of the three species was nonetheless
437 increased by inoculation with *P. indica*, but less than that of wheat. The ratio of wheat
438 shoot biomass to weed shoot biomass was increased in pots inoculated with *P. indica* so
439 the effect on wheat had outweighed the effect on the weeds. This suggests that wheat
440 might be a favourable host for *P. indica* and that field application of *P. indica* might not
441 make weed control more difficult. However, since only three species were tested, on a
442 small scale, the main conclusion is that the fungus can alter competitive relations among
443 both host and non-host species. The survival time and wide host range suggests that the
444 fungus would escape into natural communities and might alter their composition or
445 functioning. Changes would not necessarily be detrimental but these results do imply a
446 need for extensive assessments on an ecosystem scale.

447 Previous studies (Rabiey & Shaw, 2015, Rabiey et al., 2015) show that *P. indica* could
448 be extremely useful in stabilising and increasing wheat yields and quality in the UK;
449 other studies in northern Europe suggest it might also benefit other crops (Achatz et al.,
450 2010, Fakhro et al., 2010, Sun et al., 2010). The present results suggest *P. indica* effect
451 on both weeds and soil function should be studied further. Considering a close relative
452 of *P. indica*, *Serendipita herbamans*, has recently been discovered in Germany (Riess et
453 al 2014), a search for native organisms with similar characteristics might be a better
454 direction to go in (Hodkinson & Murphy, 2015).

455 **4.1. Conclusion**

456 In the case of field application, *P. indica* would probably remain active in the soil
457 within season for up to 15 months. *P. indica* increased root and soil fungal and bacterial
458 diversity. Although usually desirable, this indicates substantial effects on soil
459 composition or functioning during the first eight weeks after inoculation. The organism
460 would be likely to alter competitive relations among both host and non-host species.
461 The wider effects of *P. indica* and similar organisms need to be better understood before
462 agricultural deployment.

463

464 **Acknowledgment**

465 This work was funded by the Sir Halley Stewart Trust.

466 **5. References**

467 Abdel-Fattah GM, Mohamedin AH, 2000. Interactions between a vesicular-arbuscular
468 mycorrhizal fungus (*Glomus intraradices*) and *Streptomyces coelicolor* and their effects
469 on sorghum plants grown in soil amended with chitin of prawn scales. *Biology and*
470 *Fertility of Soils* **32**, 401-9.

471 Achatz B, Von Ruden S, Andrade D, Neumann E, Pons-Kuhnemann J, Kogel KH,
472 Franken P, Waller F, 2010. Root colonization by *Piriformospora indica* enhances grain
473 yield in barley under diverse nutrient regimes by accelerating plant development. *Plant*
474 *and Soil* **333**, 59-70.

475 Alikhani M, Khatabi B, Sepehri M, Nekouei MK, Mardi M, Salekdeh GH, 2013. A
476 proteomics approach to study the molecular basis of enhanced salt tolerance in barley
477 (*Hordeum vulgare* L.) conferred by the root mutualistic fungus *Piriformospora indica*.
478 *Molecular Biosystems* **9**, 1498-510.

479 Amora-Lazcano E, Vazques MM, Azcon R, 1998. Response of nitrogen-transforming
480 microorganisms to arbuscular mycorrhizal fungi. *Biology and Fertility of Soils* **27**, 65–
481 70.

- 482 Andrade G, Mihara KL, Linderman RG, Bethlenfalvay GJ, 1997. Bacteria from
483 rhizosphere and hyphosphere soils of different arbuscular-mycorrhizal fungi. *Plant and*
484 *Soil* **192**, 71–9.
- 485 Barea JM, 2000. Rhizosphere and mycorrhiza of field crops. In: Toutant Jp. BE,
486 Galante E., Lynch J.M., Schepers J.S., Werner D., Werry P.A., ed. *Biological resource*
487 *management: connecting science and policy*. (OECD) INRA Editions and Springer,
488 Berlin Heidelberg New York, pp 110-125.
- 489 Bertholdsson NO, 2012. Allelopathy-a tool to improve the weed competitive ability of
490 wheat with herbicide-resistant black-grass (*Alopecurus myosuroides* Huds.) *Agronomy*
491 **2**, 284-94.
- 492 Bockus WW, Bowden RL, Hunger RM, Morrill WL, Murray TD, Smiley RW, 2010.
493 Compendium of Wheat Diseases and Pests, Third Edition. *American Phytopathological*
494 *Society*.
- 495 Bond W, Davies G, Turner R, 2007. The biology and non-chemical control of Cleavers
496 (*Galium aparine* L.). *HDRA The Organic Organisation*.
497 <http://www.gardenorganic.org.uk/organicweeds>.
- 498 Brundrett MC, 2006. Understanding the roles of multifunctional mycorrhizal and
499 endophytic fungi. In: Schulz B.J.E. BCJC, Sieber T.N., ed. *Microbial root endophytes*.
500 Berlin, Germany: Springer-Verlag, 281-293 pp.
- 501 Chimento A, Cacciola SO, Garbelotto M, 2012. Detection of mRNA by reverse-
502 transcription PCR as an indicator of viability in *Phytophthora ramorum*. *Forest*
503 *Pathology* **42**, 14-21.
- 504 Christensen H, Jakobsen I, 1993. Reduction of bacterial growth by a vesicular-
505 arbuscular mycorrhizal fungus in the rhizosphere of cucumber (*Cucumis sativus* L.).
506 *Biology and Fertility of Soils* **15**, 253-8.
- 507 Clay K, Holah J, 1999. Fungal endophyte symbiosis and plant diversity in successional
508 fields. *Science* **285**, 1742-4.
- 509 Deshmukh SD, Kogel KH, 2007. *Piriformospora indica* protects barley from root rot
510 caused by *Fusarium graminearum*. *Journal of Plant Diseases and Protection* **114**, 263-
511 8.

- 512 Fakhro A, Andrade-Linares DR, Von Bargen S, Bandte M, Buttner C, Grosch R,
513 Schwarz D, Franken P, 2010. Impact of *Piriformospora indica* on tomato growth and on
514 interaction with fungal and viral pathogens. *Mycorrhiza* **20**, 191-200.
- 515 Gera Hol WH, De Boer W, De Hollander M, Kuramae EE, Meisner A, Van Der Putten
516 W, 2015. Context dependency and saturating effects of loss of rare soil microbes on
517 plant productivity. *Frontiers in Plant Science* **6**, 485, doi: 10.3389/fpls.2015.00485.
- 518 Ghabooli M, Khatabi B, Ahmadi FS, Sepehri M, Mirzaei M, Amirkhani A, Jorrin-Novo
519 JV, Salekdeh GH, 2013. Proteomics study reveals the molecular mechanisms
520 underlying water stress tolerance induced by *Piriformospora indica* in barley. *Journal*
521 *of Proteomics* **94**, 289-301.
- 522 Ghahfarokhy MR, Goltapeh EM, Purjam E, Pakdaman, B.S., Modarres Sanavy SaM,
523 Varma A, 2011. Potential of mycorrhiza-like fungi and Trichoderma species in
524 biocontrol of Take-all Disease of wheat under greenhouse condition *Journal of*
525 *Agricultural Technology* **7(1)**, 185-95. <http://www.ijat-aatsea.com/>.
- 526 Gryndler M, 2000. Interactions of arbuscular mycorrhizal fungi with other soil
527 organisms. In: Kapulnik Y, And Douds D.D., ed. *Arbuscular mycorrhizas: physiology*
528 *and function*. Kluwer Academic, Dordrecht, The Netherlands, pp 239-262.
- 529 Harrach BD, Baltruschat H, Barna B, Fodor J, Kogel K-H, 2013. The mutualistic fungus
530 *Piriformospora indica* protects Barley roots from a loss of antioxidant capacity caused
531 by the necrotrophic pathogen *Fusarium culmorum*. *Molecular Plant-Microbe*
532 *Interactions* **26**, 599-605.
- 533 Herdina, Neate S, Jabaji-Hare S, Ophel-Keller K, 2004. Persistence of DNA of
534 *Gaeumannomyces graminis* var. *tritici* in soil as measured by a DNA-based assay.
535 *FEMS Microbiol Ecology* **47**, 143-52.
- 536 Hodkinson TR, Murphy BR, 2015. Native fungal microorganisms enhance important
537 agronomic traits in barley. . *Graduate Students' Union of the University of Dublin,*
538 *Trinity College. Journal of Postgraduate Research* **14**.
- 539 Hoshino Y, T., 2012. Molecular analyses of soil fungal community. In: S. MCH, ed.
540 *Methods and Applications, Soil Health and Land Use Management*. InTech, pp. 279-
541 304.

- 542 Hoshino YT, Morimoto S, 2008. Comparison of 18S rDNA primers for estimating
543 fungal diversity in agricultural soils using polymerase chain reaction-denaturing
544 gradient gel electrophoresis. *Soil Science and Plant Nutrition* **54**, 701-10.
- 545 Hull R, Tatnell LV, Cook SK, Beffa R, Moss SR, 2014 Current status of herbicide-
546 resistant weeds in the UK. Crop Production in Southern Britain: Precision Decisions for
547 Profitable Cropping *Aspects of Applied Biology* **127**, 261-72.
- 548 Jeffries P, Gianinazzi S, Perotto S, Turnau K, Barea J-M, 2003. The contribution of
549 arbuscular mycorrhizal fungi in sustainable maintenance of plant health and soil
550 fertility. *Biology and Fertility of Soils* **37**, 1-16.
- 551 Jumpponen A, 2007. Soil fungal communities underneath willow canopies on a primary
552 successional glacier forefront: rDNA sequence results can be affected by primer
553 selection and chimeric data. *Microbial Ecology* **53**, 233-46.
- 554 Mcguire KL, Treseder KK, 2010. Microbial communities and their relevance for
555 ecosystem models: decomposition as a case study. *Soil Biology and Biochemistry* **42**,
556 529-35.
- 557 Mennan H, Isik D, 2004. The competitive ability of *Avena* spp. and *Alopecurus*
558 *myosuroides* Huds. influenced by different wheat (*Triticum aestivum*) cultivars. *Turkish*
559 *Journal of Agriculture and Forestry* **28**, 245-51.
- 560 Moss SR, Marshall R, Hull R, Alarcon-Reverte R, 2011. Current status of herbicide-
561 resistant weeds in the United Kingdom. Crop Protection in Southern Britain. *Aspects of*
562 *Applied Biology* **106**, 1-10.
- 563 Moss SR, Perryman SaM, Tatnell LV, 2007. Managing herbicide-resistant blackgrass
564 (*Alopecurus myosuroides*): Theory and practice. *Weed Technology* **21**, 300-9.
- 565 Murphy BR, Doohan FM, Hodkinson TR, 2014. Fungal endophytes of barley roots. *The*
566 *Journal of Agricultural Science* **152**, 602-15.
- 567 Murphy BR, Martin Nieto L, Doohan FM, Hodkinson TR, 2015. Fungal endophytes
568 enhance agronomically important traits in severely drought-stressed barley. *Journal of*
569 *Agronomy and Crop Science* **201** 419-27.

- 570 Muyzer G, Dewaal EC, Uitterlinden AG, 1993. Profiling of Complex Microbial-
571 Populations by Denaturing Gradient Gel-Electrophoresis Analysis of Polymerase Chain
572 Reaction-Amplified Genes-Coding for 16s Ribosomal-Rna. *Applied and Environmental*
573 *Microbiology* **59**, 695-700.
- 574 Omacini M, Chaneton EJ, Ghera CM, Müller CB, 2001. Symbiotic fungal endophytes
575 control insect host-parasite interaction webs. *Nature* **409**, 78-81.
- 576 Rabiey M, Shaw MW, 2015. *Piriformospora indica* reduces fusarium head blight
577 disease severity and mycotoxin DON contamination in wheat under UK weather
578 conditions. *Plant Pathology*, DOI: 10.1111/ppa.12483.
- 579 Rabiey M, Ullah I, Shaw MW, 2015. The endophytic fungus *Piriformospora indica*
580 protects wheat from fusarium crown rot disease in simulated UK autumn conditions.
581 *Plant Pathology* **64**, 1029–40. DOI: 10.1111/ppa.12335.
- 582 Rajaguru BaP, Shaw MW, 2010. Genetic differentiation between hosts and locations in
583 populations of latent *Botrytis cinerea* in southern England. *Plant Pathology* **59**, 1081-
584 90.
- 585 Rincon-Florez VA, Carvalhais LC, Schenk PM, 2013. Culture-independent molecular
586 tools for soil and Rhizosphere microbiology *Diversity* **5**, 581-612.
- 587 Rodriguez RJ, White JF, Jr., Arnold AE, Redman RS, 2009. Fungal endophytes:
588 diversity and functional roles. *New Phytologist* **182**, 314-30.
- 589 Shrivastava S, Varma A, 2014. From *Piriformospora indica* to rootonic: a review.
590 *African Journal of Microbiology Research* **8**, 2984-92.
- 591 Söderberg KH, Olsson PA, Bååth E, 2002. Structure and activity of the bacterial
592 community in the rhizosphere of different plant species and the effect of arbuscular
593 mycorrhizal colonisation. *FEMS Microbiology Ecology* **40**, 223-31.
- 594 Sun C, Johnson JM, Cai D, Sherameti I, Oelmüller R, Lou B, 2010. *Piriformospora*
595 *indica* confers drought tolerance in Chinese cabbage leaves by stimulating antioxidant
596 enzymes, the expression of drought-related genes and the plastid-localized CAS protein.
597 *Journal of Plant Physiology* **167**, 1009-17.

- 598 Varma A, Kost G, Oelmuller R, Eds., 2013. *Piriformospora indica; Sebaciniales and*
599 *their biotechnological applications*. Soil Biology 33. Berlin-Heidelberg: Springer-
600 Verlag. pp 17.
- 601 Verma S, Varma A, Rexer KH, Hassel A, Kost G, Sarbhoy A, Bisen P, Butehorn B,
602 Franken P, 1998. *Piriformospora indica*, gen. et sp. nov., a new root-colonizing fungus.
603 *Mycologia* **90**, 896-903.
- 604 Waller F, Achatz B, Baltruschat H, Fodor J, Becker K, Fischer M, Heier T,
605 Huckelhoven R, Neumann C, Von Wettstein D, Franken P, Kogel KH, 2005. The
606 endophytic fungus *Piriformospora indica* reprograms barley to salt-stress tolerance,
607 disease resistance, and higher yield. *Proceedings of the National Academy of Sciences*
608 *of the United States of America* **102**, 13386-91.
- 609 Wamberg C, Christensen S, Jakobsen I, Müller AK, Sørensen SJ, 2003. The
610 mycorrhizal fungus (*Glomus intraradices*) affects microbial activity in the rhizosphere
611 of pea plants (*Pisum sativum*). *Soil Biology and Biochemistry* **35**, 1349-57.
- 612 Yu Q, Ahmad-Hamdani MS, Han H, Christoffers MJ, Powles SB, 2013. Herbicide
613 resistance-endowing ACCase gene mutations in hexaploid wild oat (*Avena fatua*):
614 insights into resistance evolution in a hexaploid species. *Heredity* **110**, 220-31.
- 615 Zadoks JC, Chang TT, Konzak CF, 1974. A decimal code for the growth stages of
616 cereals. *Weed Research* **14**, 415-21.
- 617 Zarea MJ, Hajinia S, Karimi N, Goltapeh EM, Rejali F, Varma A, 2012. Effect of
618 *Piriformospora indica* and *Azospirillum* strains from saline or non-saline soil on
619 mitigation of the effects of NaCl. *Soil Biology & Biochemistry* **45**, 139-46.

620

621

622

623

624

625

626

627

628

629

630

631 **Figure Legends**

632 **Fig 1.** Reading mean air temperature, mean 10 cm soil temperature, and total rainfall
633 between winter 2013-14 and winter 2014-15, compared with 1981-2010 average
634 (source: www.met.reading.ac.uk/weatherdata).

635

636 **Fig 2.** Denaturing gradient gel electrophoresis profiles of the wheat (a) root fungal and
637 (b) bacterial community in Sonning series (SCL) or Rowland series (LSO) soil
638 inoculated with (+) or without (-) *Piriformospora indica*, harvested at 2 weeks after
639 inoculation (wai) (T1), 4 wai (T2), 6 wai (T3) and 8 wai (T4), (first lane: Hyper Ladder
640 I-100 lanes (Bioline)).

641

642 **Fig 3.** Canonical variates analysis of bands from denaturing gradient gel electrophoresis
643 using universal fungal and bacterial primers for wheat root samples grown in Sonning
644 series (SCL) or Rowland series (LSO) soils, inoculated with/without *Piriformospora*
645 *indica*, (Pi: *P. indica*). First or first and second canonical axes are shown for data
646 classified by (a) the four time points of harvest; (b) Root and soil source; (c) soil types;
647 (d) *P. indica*-inoculation status; (e-h) *P. indica*-inoculation status using but restricted to

648 fungal (e), or bacterial primers (f) or to soil samples (g) or root samples (h); (i) both *P.*
649 *indica* inoculation and root or soil source. In (a) and (i) the circles indicate 95%
650 confidence intervals for the centre of the group in the canonical variate space under
651 normality assumptions.

652

653 **Fig 4.** Shannon-Weiner diversity index for Sonning (SCL) and Rowland series (LSO)
654 soil samples inoculated or not with *Piriformospora indica*. Based on denaturing gel
655 electrophoresis of DNA extracts amplified using universal fungal and bacterial primers.
656 Each bar shows mean \pm 95% bootstrap confidence interval.

657

658 **Fig 5.** Shannon-Weiner diversity index for wheat root and soil samples inoculated or
659 not with *Piriformospora indica* (Pi), based on denaturing gel electrophoresis of DNA
660 extracts amplified using universal fungal and bacterial primers. Both soil types
661 (Sonning series (SCL) and Rowland series (LSO)) are combined. Each bar shows mean
662 \pm 95% bootstrap confidence interval.

663

664

665 **Table Legends**

666 **Table 1.** Recovery of *Piriformospora indica* DNA and RNA after the mycelia were
667 killed by exposure to heat and cold or grown in covered petri dishes of potato
668 dextrose agar (n=3 for each condition).

669

670 **Table 2.** Recovery of *Piriformospora indica* DNA and RNA from four soil types, left in
671 pots under prevailing weather conditions without plant roots present from December
672 2013 with sample collections at mid-March 2014, end-July 2014 and mid-March 2015,
673 n=5 (S.e.d.: Standard error of the difference of means).

674

675 **Table 3.** Dry weights (g) of root and shoot of *Alopecurus myosuroides*, *Avena fatua* and
676 *Galium aparine* alone and in competition with wheat, with and without inoculation with
677 *Piriformospora indica* (error d.f.: 33; S.e.d.: Standard error of the difference of means).

678

679 **Table 4.** Competitiveness of *Alopecurus myosuroides*, *Avena fatua*, and *Galium aparine*
680 with wheat, measured as \log_{10} (weed dry weight/wheat dry weight), in the presence and
681 absence of inoculum of *Piriformospora indica* in the soil (d.f.: 15).

682

683

684 **Supporting Information**

685 **Table 1.** ANOVA P-value for Table 3. for dry weights (g) of root and shoot of weed
686 species (*Alopecurus myosuroides*, *Avena fatua* and *Galium aparine*) alone and in
687 competition with wheat, with and without inoculation with *Piriformospora indica*.

688

689

690 **Table 2.** ANOVA P-value for Table 4. for competitiveness of weed species (*Alopecurus*
691 *myosuroides*, *Avena fatua*, and *Galium aparine*) with wheat, in the presence and
692 absence of inoculum of *Piriformospora indica* in the soil.

693

694 **Table 3a.** Dry weights (g) of root and shoot of *Alopecurus myosuroides*, *Avena fatua*
695 and *Galium aparine* alone and in competition with wheat, with and without inoculation
696 with *Piriformospora indica* (error d.f.: 33; S.e.d.: Standard error of the difference of
697 means) (second repeat).

698

699 **Table 3b.** ANOVA P-value for Table 3a. for dry weights (g) of root and shoot of weed
700 species (*Alopecurus myosuroides*, *Avena fatua* and *Galium aparine*) alone and in
701 competition with wheat, with and without inoculation with *Piriformospora indica*.

702

703 **Table 4a.** Competitiveness of *Alopecurus myosuroides*, *Avena fatua*, and *Galium*
704 *aparine* with wheat, measured as \log_{10} (weed dry weight/wheat dry weight), in the
705 presence and absence of inoculum of *Piriformospora indica* in the soil (d.f.:15; S.e.d.:
706 Standard error of the difference of means) (second repeat).

707

708 **Table 4b.** ANOVA P-value for Table 4a. for competitiveness of weed species
709 (*Alopecurus myosuroides*, *Avena fatua*, and *Galium aparine*) with wheat, in the presence
710 and absence of inoculum of *Piriformospora indica* in the soil.
711
712
713