

The Lancet Countdown: tracking progress on health and climate change

Article

Accepted Version

Creative Commons: Attribution-Noncommercial-No Derivative Works 4.0

Watts, N., Adger, W. N., Ayeb-Karlsson, S., Bai, Y., Byass, P., Campbell-Lendrum, D., Colbourn, T., Cox, P., Davies, M., Depledge, M., Depoux, A., Dominguez-Salas, P., Drummond, P., Ekins, P., Flahault, A., Grace, D., Graham, H., Haines, A., Hamilton, I., Johnson, A., Kelman, I., Kovats, S., Liang, L., Lott, M., Lowe, R., Luo, Y., Mace, G., Maslin, M., Morrissey, K., Murray, K., Neville, T., Nilsson, M., Oreszczyn, T., Parthemore, C., Pencheon, D., Robinson, E. J. Z. ORCID: https://orcid.org/0000-0002-4950-0183, Schütte, S., Shumake-Guillemot, J., Vineis, P., Wilkinson, P., Wheeler, N., Xu, B., Yang, J., Yin, Y., Yu, C., Gong, P., Montgomery, H. and Costello, A. (2017) The Lancet Countdown: tracking progress on health and climate change. Lancet, 389 (10074). pp. 1151-1164. ISSN 0140-6736 doi: 10.1016/S0140-6736(16)32124-9 Available at https://centaur.reading.ac.uk/68063/

It is advisable to refer to the publisher's version if you intend to cite from the work. See <u>Guidance on citing</u>.

Published version at: http://dx.doi.org/10.1016/S0140-6736(16)32124-9

To link to this article DOI: http://dx.doi.org/10.1016/S0140-6736(16)32124-9

Publisher: Elsevier



All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the End User Agreement.

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online

The Lancet Countdown: Tracking Progress on Health and Climate Change

3 4 Nick Watts, W. Neil Adger, Sonja Ayeb-Karlsson, Yuqi Bai, Peter Byass, Diarmid Campbell-Lendrum, 5 Tim Colbourn, Peter Cox, Michael Davies, Michael Depledge, Anneliese Depoux, Paula Dominguez-Salas, Paul Drummond, Paul Ekins, Antoine Flahault, Delia Grace, Hilary Graham, Andy Haines, Ian 6 7 Hamilton, Anne Johnson, Ilan Kelman, Sari Kovats, Lu Liang, Melissa Lott, Robert Lowe, Yong Luo, 8 Georgina Mace, Mark Maslin, Karyn Morrissey, Kris Murray, Tara Neville, Maria Nilsson, Tadj 9 Oreszczyn, Christine Parthemore, David Pencheon, Elizabeth Robinson, Stefanie Schütte, Joy 10 Shumake-Guillemot, Paolo Vineis, Paul Wilkinson, Nicola Wheeler, Bing Xu, Jun Yang, Yongyuan Yin, 11 Chaoqing Yu, Peng Gong, Hugh Montgomery, Anthony Costello 12 13 Abstract 14 15 The Lancet Countdown: Tracking Progress on Health and Climate Change is an international, multi-16 disciplinary research collaboration between academic institutions and practitioners across the world. 17 It follows on from the work of the 2015 Lancet Commission, which concluded that the response to 18 climate change could be "the greatest global health opportunity of the 21st century". 19 The Lancet Countdown aims to track the health effects of climate change; health resilience and 20 adaptation; health co-benefits of mitigation; climate finance and economics; and political and 21 broader engagement. These focus areas form the five thematic working groups of the Lancet 22 Countdown and represent different aspects of the complex relationships between health and 23 climate change. These thematic groups will provide indicators for a global overview of health and 24 climate change; national case studies highlighting countries leading the way or going against the 25 trend; and engagement with a range of stakeholders. 26 The Lancet Countdown ultimately aims to report annually on a series of indicators across these five 27 working groups. This paper outlines these potential indicators and indicator domains to be tracked 28 by the collaboration, with suggestions on the methodologies, and data sets available to achieve this 29 end. The proposed indicator domains require further refinement, and mark the beginning of an 30 ongoing consultation process – from November 2016 to early 2017 – to develop these domains, 31 identify key areas not currently covered, and change indicators where necessary. It will actively seek 32 to engage with existing monitoring processes, such as the UN Sustainable Development Goals, and 33 the World Health Organization's Climate and Health Country Profiles. Additionally, the indicators will 34 evolve throughout their lifetime through ongoing collaboration with experts and a range of

stakeholders, and dependent on the emergence of new evidence and knowledge. During the course

of its work, the Lancet Countdown will adopt a collaborative and iterative process, which aims to

complement existing initiatives, welcome engagement with new partners, and be open to

developing new research projects on health and climate change.

39

35

36

37

ır	١Ŧ	rc	\sim		ГΙ	\cap	n
	H.	LU	u	LU	L I	U	

- 41 The World Health Organization (WHO) estimated that in 2012, 12.6 million deaths (23% of all
- deaths) were attributable to modifiable environmental factors, many of which could be influenced
- by climate change, or are related to the driving forces of climate change. The 2009 UCL-Lancet
- 44 Commission: Managing the health effects of climate change described the ways in which climate
- change acts as a force multiplier for threats to global health.² These initiatives have drawn on long-
- 46 standing expertise and leadership in the health and climate space from institutions such as the WHO
- and the Intergovernmental Panel on Climate Change's (IPCC).^{3,4}
- 48 The 2015 Lancet Commission on Health and Climate Change: policy responses to protect public
- 49 health built on these foundations and explored the health benefits of climate change mitigation and
- adaptation policies.⁵ As first described in the 2009 Lancet series, greenhouse gas (GHG) mitigation
- across a range of sectors can result in significant improvements in public health. ⁶⁻¹⁰ Taken together,
- 52 the potential to avoid significant impacts of climate change and the potential co-benefits of climate
- 53 mitigation and adaptation led the 2015 Lancet Commission to conclude that "tackling climate
- change could be the greatest global health opportunity of the 21st century".5
- 55 Direct impacts of climate change result from rising temperatures, heatwaves, and increases in the
- 56 frequency of complex extreme weather events such as windstorms, floods and droughts. 11 The
- 57 health and social consequences of these events are far-reaching, ranging from reduced labour
- 58 productivity and heat-related deaths, through to direct injury, the spread of infectious disease, and
- 59 mental health impacts following wide-spread flooding. The effects of climate change will also be
- 60 heterogeneously mediated across different environmental and social systems, resulting in changing
- patterns of the burden and distribution of infectious diseases, changes in food productivity and
- 62 potential effects on food and water shortages, population displacement, and conflict.³ Climate
- 63 change places undue burden on countries least responsible and least able to respond, with low- and
- 64 middle-income countries experiencing multiple impacts simultaneously (see Figure 1). 12




Figure 1. The health impacts of climate change (source: Lancet Commission, 2015⁵)

The Rockefeller Foundation-Lancet Commission on Planetary Health described how sustained human health and development depend on flourishing natural systems. This Commission and others have drawn attention to the fact that human activities are breaching environmental limits across a range of areas, driving terrestrial and marine biodiversity loss, ocean acidification, stratospheric ozone depletion, soil degradation, and other potentially irreversible processes. 13,14

At the international level, the Paris Agreement provides the framework for future international cooperation and national action on climate change. Modelling suggests that the full implementation of all national government pledged mitigation actions would limit global average warming to around 2.7°C by 2100; an improvement on the high-end 4.8°C or more scenario, but substantially higher than the agreed United Nations target of "well below 2.0°C". 15

Responsibility for the implementation of the Paris Agreement now falls to national governments. The next 15 years, from 2016 to 2030, are a crucial window that will determine the trajectory of climate change and human development for the coming century. As part of this transition, countries will have to shift from an understanding of climate change solely as a threat, to one which embraces the response to climate change as an opportunity for human health and wellbeing. Tracking and communicating this shift will be the central focus of the *Lancet Countdown: Tracking Progress on Health and Climate Change*.

103

Aims of the Lancet Countdown on Health and Climate Change

- 86 The Lancet Countdown aims to: track the impacts of climate change and the speed of the transition
- 87 to a decarbonised global economy, a transition that is already underway; analyse and demonstrate
- 88 the health benefits available; provide a global picture of successes and obstructions in this shift;
- 89 draw out exemplary case-studies for shared learning; and engage with policymakers and the broader
- 90 health community to better communicate the opportunities available in responding to climate
- 91 change both for health and more broadly.
- To do this, the Lancet Countdown will report annually on key indicators that reflect progress on
- 93 health and climate change. Published each year, before the UN Framework Convention on Climate
- 94 Change's (UNFCCC) international negotiations, the annual Lancet paper will consider global, regional,
- 95 national, and, on a selective basis, city level trends. Five interrelated thematic working groups will
- cover different aspects of the relationship between health and climate change, including: the health
- 97 impacts of climate change; health resilience and adaptation; the health co-benefits of mitigation;
- 98 finance and economics; and policy, politics and public opinion.
- 99 The Lancet Countdown is an international, multi-disciplinary research collaboration between
- academic institutions and experts across the world. Where the 2015 Lancet Commission existed as a
- partnership primarily between European and Chinese academics, the Lancet Countdown will build
- on these foundations to be more global, both in expertise and outlook.

104 Indicators of Progress: A Call for Input and Engagement

- The work of the Lancet Countdown is divided into five working groups, each of which will be
- responsible for their own set of indicators. Table 1 and the rest of this paper outlines proposed
- indicator domains for these working groups. These are presented for consultation with varying
- degrees of certainty, ranging from the presentation of a specific indicator, through to the description
- of a broad domain within which a number of indicators might function.
- 110 The ongoing framing and selection of indicators to mark the progress to a low-carbon and climate
- resilient society could take a number of forms, such as focusing on the interactions between society
- and environment (as seen in the DPSEEA framework, adapted in Appendix 1); or vulnerability, risk
- and adaptive capacity to climate change. 16-18 The selected indicators will need to address the
- 114 challenges of representing spatial and temporal factors in a summarized form. The framing that will
- be used to select indicators as part of the Lancet Countdown is primarily from the health
- 116 perspective. In turn, the focus is on those indicators that capture the greatest effects that climate
- change has on health; the anthropogenic drivers that have the greatest contribution to climate
- change and the measures and actions that would substantially reduce the effects climate change or
- 119 yield health co-benefits of mitigation policies.
- 120 The proposed indicator domains reflect a pragmatic need to capture markers of progress in the key
- interactions between health and climate change using the best available data, and those that can be
- translated to the health community and more widely. They were developed through an iterative
- 123 process, following an initial, broad consultation process. This sought input from a wide variety of
- 124 experts working in the field. These were further discussed and refined by the Lancet Countdown's
- academic working groups at a series of multidisciplinary meetings throughout 2016. Each proposed

indicator domain was assessed for its ability to cover a unique part of the relationship between health and climate change; potential data availability; feasibility given current resource constraints; applicability to countries across a variety of resource settings; and policy relevance.

This collaborative process is intended to work closely with other monitoring initiatives, such as the information being collected under the WHO's Climate and Health Country Profiles, the SHUE (Sustainable Healthy Urban Environments) project, the Sendai Framework, and the ClimateWorks Foundation's Carbon Transparency Initiative. ¹⁹⁻²² It will also look to draw on the UN SDGs where appropriate. The potential links between these initiatives and the Lancet Countdown's indicator domains have been summarised in Appendix 2. Where relevant cross-over exists, the Lancet Countdown will look to incorporate this data into its work to provide a more complete, and standardised overview.

This paper marks the beginning of an external consultation process, to further refine the suggestions below. Indeed, it is expected that, the indicators and metrics used will continuously evolve to make use of emerging evidence and data availability. To this end, the Lancet Countdown is committed to maintaining an open approach to further developing its work programme, inviting external input and actively consulting over the coming months. The research collaboration welcomes engagement with new academic partners, with the expertise and capacity to make substantial contributions to the final indicator process. The collaboration is also open to developing new research projects on as-yet neglected areas of health and climate change, with the possibility to jointly seek additional funding and capacity for this work in future. We invite direct input on the content, methods, and data for the proposed indicators and indicator domains, as well as proposals for new partnerships, through the website here: www.LancetCountdown.org/IndicatorConsultation

Thematic Working Group	Indicator Domains		
	1.1 Exposure to temperature change		
	1.2 Change in labour productivity		
	1.3 Exposure to heatwave		
1. Health Impacts of Climate Hazards	1.4 Exposure to flood		
	1.5 Exposure to drought		
	1.6 Spread of infectious disease across sentinel sites		
	1.7 Food security and undernutrition		
	2.1 Integration of health in national adaptation plans		
2. Health Resilience and Adaptation	2.2 Climate services for health		
	2.3 Adaptation finance for health		
	3.1. Coal phase-out		
	3.2 Growth in renewable energy resources		
	3.3 Energy access		
	3.4. Energy access for health facilities		
3. Health Co-Benefits of Mitigation	3.5 Ambient and household air pollution exposure		
	3.6 Deployment of low-emission vehicles		
	3.7 Active transport infrastructure and uptake		
	3.8 Food consumption, production, and waste		
	3.9 Carbon footprint of healthcare systems		
	4.1 Change in annual investment in renewable energy		
4. Finance and Economics	4.2 Change in annual investment in energy efficiency		
	4.3 Low-carbon technology patent generation and		

	innovation
	4.4 Valuing the health co-benefits of climate change
	mitigation
	4.5 Direct and indirect fossil fuel subsidies
	4.6 Coverage and strength of carbon pricing
	4.7 Equity of the low-carbon transition
	5.1 Public engagement in health and climate change
	5.2 Academic publications on health and climate change
	5.3 Health and climate change in the UNFCCC and UNGA
5. Political and Broader Engagement	high-level statements
3. Folitical and broader Engagement	5.4 Inclusion of health and climate change within medical
	and public health curricula
	5.5 Implementation and estimated health benefits of the
	Nationally Determined Contributions

Table 1. Proposed indicator domains for the Lancet Countdown: Tracking Progress on Health and Climate Change. Importantly, these indicator domains are heterogeneous in nature – some reflect outcomes (GHG emissions or health) and others reflect process indicators with both direct and indirect links to climate change. Additionally, some can be modelled at a global or national level, whereas others reflect location-specific issues and would depend on data collection at sentinel sites.

Health impacts of climate change

The health effects of climate change are projected to become increasingly severe in the future, and threaten to undermine the last half-century of gains in public health and development.^{3,5} These effects are unevenly distributed within and between countries, with all risks having important social, economic, and geographical mediating factors.¹⁹ The first working group of the Lancet Countdown proposes seven indicator domains to be considered and then tracked.

1.1-1.5: Human exposure to extreme weather

These domains will use observed meteorological data to create indicators of exposure to extremes of weather across five areas: annual mean temperature change; heat index relevant for outdoor labour productivity; heatwave; drought; and flood risk. This data will be complemented by a review of the detection and attribution studies linking climate change to specific extreme weather events that have affected human health. These indices are deliberately similar to those presented in the Lancet Commission for consistency and transparency. For the Lancet Countdown, we will focus on metrics calculated from observational data rather than climate model projections. However, we aim to maintain comparability between these metrics for the monitoring of progress, and the exposure metrics implied by the future projections presented in the Lancet Commission report.

Mean temperature increase and changes in the severity and frequency of heatwaves bring substantial and potentially fatal health risks to most populations. These include particularly exposed individuals (ie. those engaged in outdoor physical labour), and individuals with reduced capacity to maintain physiological homeostasis, such as the infirm, neonates, or the elderly. The direct relationship between extremes of heat and heat-related morbidity and mortality is well established, as is the relationship between indicators of thermal stress such as wet bulb globe

temperature (WBGT), and reductions in outdoor labour productivity, to a lesser extent.²⁴ Examples of direct and indirect health effects were seen in the 2010 Russian heatwave, which resulted in approximately 11,000 excess deaths from heat and poor air quality from subsequent forest fires. 25 The Lancet Countdown will utilise the population related metrics developed from the 2015 Lancet Commission to calculate the mean warming experienced by people.⁵ It will also make use of the index proposed by Jacob et al., defining a heatwave as more than three consecutive days where minimum temperature exceeds the 99th percentile for the recent past.²⁶ In addition, changes in labour productivity will be modelled using WBGT, which has been used to identify thresholds of heat

187 stress.²⁷

The fourth and fifth indicator domains will follow human exposure to flood and drought. Here, flood refers to meteorological floods, related to rain, storm surges, and sea level rise, rather than due to tsunamis or volcanic eruptions, melting snow and ice. Drought refers to meteorological drought, so a deficit of precipitation, rather than other forms of drought, such as water depletion caused by increasing demand. Observational data suggests that many regions demonstrating rising frequency of meteorological drought over the past 60 years overlap with crucial agricultural zones and regions expecting rapid population expansion in sub-Saharan Africa and South Asia. Analysis in the 2015 Commission projected an additional 1.4 billion person drought exposure events per year by 2100, as a result of population change and climate change.

Of particular importance, is the effect that climate change will have on mental health and wellbeing, an issue which is often amplified in low-resource settings which lack the protective social and public health institutions. The Lancet Countdown is currently exploring options to track the mental health impacts of climate change.

1.6: Human Exposure to Infectious Disease

Infectious diseases make significant contributions to the global burden of disease, and many infectious diseases, their vectors and/or reservoirs, are influenced directly or indirectly by climate.³² Distributions and impacts of infectious diseases are already responding to the various dimensions of climate change so far observed, with projections that this will worsen for many infectious diseases in future.³³⁻³⁵

Given existing information about climate-sensitive infectious diseases, we will derive a shortlist of 'sentinel' diseases or disease groups to road-test the indicator protocol, and then expand the list to include other relevant infectious diseases and following wider input and consultation with infectious disease experts. Examples from three key groups will be tracked: food-borne diseases, and vector-borne and parasitic diseases/zoonotic diseases. Each of these groups, and specific diseases within each group, are likely to be affected by climate change in diverse ways. Our aim is to place a finger on the pulse of these impacts at a global scale and facilitate trend tracking through time. For this indicator domain, we will thus leverage surveillance and research networks that monitor and synthesise existing data to model changes in climate change relevant infectious disease impacts and/or risks and exposure. This process will identify sentinel sites (as comprehensive monitoring is not feasible) across a range of geographies. We would welcome suggestions of suitable sites and diseases.

A number of sub-indicators will be derived for this purpose, broadly covering the areas of outbreaks, occurrence and spread (of disease, causative agents, or vector or reservoir species), and prevalence and incidence, providing a picture of changing trends in exposure to, and impacts from infectious

diseases due to climate change. Four focal metrics proposed for each of these indicators include 1) change in cases observed and predicted in the human population, 2) changes observed or predicted in geographical or temporal extent 3) observed or predicted changes in environmental suitability for sentinel pathogens, vectors or reservoirs, 4) changes in other environmental exposures and confounding factors.

1.7: Food security and undernutrition

Having reliable access to sufficient, affordable and nutritious food can be negatively affected by climate change in many ways. This ranges from the direct impact of drought, flood and heat on harvest yields, through to the health and social impacts of climate change, resulting in unhealthy populations unable to farm or work enough to earn money to purchase food. Furthermore, food trade may be disrupted due to infrastructure damaged by climate shocks. ⁴⁰ Populations in low-income countries reliant on rain-fed agriculture are often particularly vulnerable to climate change and weather shocks. These disproportionately affect the availability and cost of staples, as a result of unreliable access to international markets and low food stocks being unable to buffer price spikes. ⁴⁰

Whilst the health implications of food insecurity are local, international and national drivers are important. ⁴¹ Further, measures to ensure climate-resilient food systems also improve food security, public health and community development. ⁴⁰ Climate-related food security indicators can address direct availability of food (agricultural production), households' ability to purchase food (rural and urban poverty relative to food prices), and resilience to shocks (food stocks and international trade in grains). The seventh proposed indicator domain will focus on food price indices and food stocks as a proxy for food affordability and availability. Other environmental and socioeconomic factors are likely to be key to understanding food security and undernutrition. To this end, the Lancet Countdown will seek partnership with external activities to fully address this interaction.

Health Resilience and Adaptation

Adaptation interventions designed to minimise the health impacts of climate change are already required, today. The second working group of the Lancet Countdown will focus on the design and deployment of adaptation and resilience interventions. It will particularly draw on data collected for the WHO/UNFCCC Climate and Health Country Profiles, including responses to surveys from national Ministries of Health.⁴²

2.1: Health adaptation planning

Past and ongoing human influence on the atmosphere means we are now committed to climate change for several decades to come. Health and related systems, such as water, sanitation and nutrition, will need to become more resilient and adapt to changing climate conditions, in order to continue to protect and promote health in a changing climate. WHO, UNFCCC and other international agencies are supporting countries to develop the health components of national adaptation plans, and promoting a comprehensive approach to build resilience into the building blocks of health and other relevant systems. ^{43,44} This eventual indicator will use the monitoring systems established for SDG indicator 13.2.1, monitoring submissions to the UNFCCC, and survey

responses from national Ministries of Health, to track the number of countries that have developed a health adaptation plan, the range of functions covered, and the extent of implementation.

267

268

265

266

2.2: Climate services for health

- 269 Informed adaptation and sustainable development requires the use of climate information for
- 270 evidence-based decision-making in the health sector. This depends fundamentally upon the
- availability of relevant, high quality climate and environmental observations, institutional and
- 272 human capacity to transform climate data into reliable, and relevant climate products and services.
- The availability, access to, and use of climate services are thus a cornerstone for health adaptation.
- This should be monitored as an indicator of health sector capacity to help anticipate and prepare for
- 275 climate risks, appropriately target long- and short-term investments, and avoid potentially
- 276 maladaptive choices.
- 277 For the purposes of the Lancet Countdown, we plan to collaborate with the WMO to conduct a
- 278 periodic survey of National Hydrological and Meteorological Services, to monitor the demand,
- availability and provision of climate information services provided to the public and national health
- authorities. Categories of services surveyed may include: sharing of historical climate and
- 281 hydrological observations, tailored forecasts or monitoring for exposure to hazardous air quality,
- pollen, extreme heat, floods and storms; or provision of tailored climate scenarios and impact
- projections. This will also be cross-referenced with WHO surveys of national Ministries of Health to
- 284 measure the extent to which countries use this information to inform health surveillance and to
- develop early warning and response systems. An additional dimension to this indicator domain could
- involve analysing national expenditure on climate information services.

287

288

2.3: Adaptation finance for health

- Health is widely recognized as a priority for adaptation. For example, over 95% of Least Developed
- 290 Countries (LDCs) identified health as a priority in their National Adaptation Programmes for Action.⁴⁵
- 291 However, this is not yet reflected in financial flows, with less than 1.5% of international climate
- 292 finance for adaptation has been directed to projects specifically addressing health. 42 This proposed
- 293 indicator domain will use information from monitoring systems of multilateral and bilateral climate
- finance, including SDG indicator 13.a.1, as well as survey responses from health ministries, to
- 295 measure investment of international and domestic resources in health adaptation to climate change.

296

297

Health co-benefits of mitigation

- The existence of ancillary health benefits (co-benefits) of climate change mitigation policies provides
- a powerful incentive to accelerate policy change, since these benefits are experienced in the near-
- term, as opposed to the long-term benefits of climate change mitigation. As noted, however, such
- 301 benefits are not automatic, and care is needed to avoid unintended adverse consequences for
- 302 health. In order to assess progress in climate change mitigation and the potential resultant ancillary
- 303 health effects (mainly co-benefits), the third working group envisions tracking nine indicators across
- four systems energy, transport, food, and healthcare.
- Here, relevant categories of data include trends in GHG and short lived climate pollutant (SLCP)
- emissions, indicators relevant to the pathways by which health co-benefits are achieved (air

307 pollution exposures, transport-related physical activity patterns, and dietary survey data), and 308 regulations (e.g. restrictions on polluting vehicles, energy sources, and energy performance) in 309 sectors that are also responsible for GHG/SLCP emissions. Trends in GHG emissions by country can be assessed through the UNFCCC reporting mechanisms and notification is subject to new reporting 310 311 requirements. 312 3.1-3.5: The Energy Sector 313 The energy sector (both production and use) represents the largest single source of anthropogenic 314 GHG emissions globally, producing an estimated two-thirds of such emissions. 15,46 It is also the 315 predominant source of air pollution, with almost all globally produced sulphur dioxide and nitrogen 316 317 oxide emissions, as well as around 85% of particulate matter, being produced by energy production 318 and energy use in buildings, industry and transport.⁴⁷ 319 320 3.1: Coal phase-out Coal use comprises 29% of total global fuel use. 48 Globally, coal is used to generate 40% of electricity 321 and, among all energy sources for electricity production, coal-fired generation contributes most 322 323 (50%) to ambient air pollution (and consequently to adverse impacts on health) as well as to CO₂ emissions. ⁴⁹ Coal is responsible for approximately 60% of global sulphur dioxide emissions. Until 324 325 recently, coal use grew steadily through to 2014, with China being the major user (over 80% of global growth since 2000 and approximately 50% of total global use). 49 326 Counts of the number and capacity of coal-fired plants, their use of coal, and their emissions, can be 327 328 monitored, but more informative would be estimates of the loss of life expectancy attributable to 329 the contribution of coal-fired combustion to ambient air pollution. The estimation of such burdens is 330 theoretically possible, but depends on high quality emissions inventory data, and air pollution 331 modelling of source contributions to human exposure. This is feasible in data rich (primarily OECD) settings, but not universally. The International Energy Agency (IEA) produces Market Reports on coal 332 use and forecasts for both the OECD and non-OECD countries.⁴⁹ The data are derived from country 333 level estimates of installed capacity, fuel consumption or power generation; the fuel mix of coal and 334 335 emission standards will be used to derive estimates of coal-related air pollution. Initially, this will be 336 feasible in specific geographical locations, with ambition to expand the work globally. 337 338 3.2: Growth in renewable energy use 339 Globally, renewable energy from wind, solar thermal, photovoltaic, hydro, tidal, geothermal, 340 biofuels and waste comprised 14% of total primary energy supply, 22% of global electricity generation, and accounted for nearly half of the new generation capacity added in 2014. 48,50 341 342 Renewable energy offers a number of important potential mechanisms for addressing climate 343 change and improving health. Most forms of renewable energy produce no direct emissions related 344 to electricity generation (with the exception of biomass) and therefore help alleviate air pollution 345 exposure. Renewables can also scale and be deployed as decentralised systems, providing greater 346 penetration and provision of modern energy to hard-to-reach populations and health facilities.

Renewable energy growth is primarily measured in terms of capacity and total final energy

consumption (TFEC). The Lancet Countdown plans to use this as an indicator of growth in renewable

347

348

energy, using the IEA and International Renewable Energy Agency's (IRENA) regularly published estimates.

3.3: Energy access

- Access to adequate and clean energy supplies in the household offers numerous benefits to health, and improved life expectancy. 8,51 In 2013, the IEA estimated that around 1.2 billion people do not have access to electricity and around 2.7 billion people rely on burning unsustainable and inefficient solid fuels for cooking and heating. 48 The household air pollution that results from these fuels and other sources has an attributable impact of around 4.3 million deaths annually, related to pneumonia, stroke, lung cancer, stroke, heart disease, and COPD. 52 Although access to electricity is growing, with the current average national electrification rate being 83%, there is enormous variability, with urban access to electricity as low as 1-4% in South Sudan, Liberia, and Central African Republic. 53,54
 - For the purposes of the Lancet Countdown on Health and Climate Change, the IEA and World Bank produce national statistics on metrics of energy use based on surveys and data provided by member countries and their own research. These metrics include: energy use per capita, percentage of population with access to non-solid fuels, and percentage of population with access to electricity. The SDG indicators focus on the latter two metrics of energy access. The Lancet Countdown is also exploring the feasibility of monitoring the expansion of micro-grids in low-resource settings, as an important component of the expansion of renewable energy.

3.4: Energy access for health facilities

Access to energy is crucial for the delivery of healthcare. Providing adequate lighting, cooling of medicines, controlling indoor thermal exposure, and hot water for washing and sterilization and clinical procedures rely on a consistent delivery of energy. Among low-income countries, healthcare facilities struggle to ensure access to consistent and affordable energy. A review of healthcare facilities in a number of sub-Saharan countries showed on average 26% of those facilities having no access to electricity; 28% had reliable access to electricity, while 7% relied solely on generators. The WHO have proposed a 'multi-tier metric' for assessing electricity access among healthcare facilities and include peak power capacity, daily energy capacity, duration of supply, evening peak hours supply, affordability, quality, reliability, operational sustainability, and environmental sustainability and health. This metric has yet to be operationalised, but the Lancet Countdown will draw on this measure for reporting on the healthcare sector.


3.5: Ambient and household air pollution exposure

An estimated 18,000 people die every day due to air pollution exposure, making it the world's largest single environmental health risk. This public health emergency is particularly pertinent in urban areas, but also affects non-urban populationshere. As Figure 2 shows, the vast majority (~90%) of Europeans living in urban areas and almost all (98%) of those living in cities in low- and middle-income countries are exposed to air pollution levels in excess of WHO guidelines. Moreover, current evidence suggests health effects at concentrations below guideline levels,

without threshold, so that health benefits can be expected the more pollution is reduced regardless of initial concentrations.

Figure 1.7

Average annual outdoor PM_{2.5} concentrations in selected urban areas



Sources: WHO (2016) Global Urban Ambient Air Pollution Database; Demographia (2015) for population; country groups per income based on World Bank (2016).

Figure 2 - Average annual outdoor PM2.5 concentrations in selected urban areas (IEA, 2016, Energy and Air Pollution)⁴⁷

 The WHO's Urban Ambient Air Pollution Database now has annual mean outdoor concentrations of PM_{10} and $PM_{2.5}$ for almost 3,000 cities. As part of the Lancet Countdown, this database will provide an important data source for cities. We propose to track a number of indicators of (urban) air pollution, primarily annual mean $PM_{2.5}$ and/or PM_{10} together with selected other pollutants (e.g. NO_2), using data for individual cities and, where feasible, population weighted averages for other geographic scales.

3.6-3.7: The Transport Sector

Transportation systems – including road vehicles, rail, shipping, and aviation – are a key source of GHG emissions, contributing 14% of global GHG emissions in 2010. 47,48,59 It is also a major source of air pollutants, including particulate matter (PM), nitrogen oxides (NO_x), particulate matter (PM), sulphur dioxide, carbon monoxide, and volatile organic compounds, and, indirectly, ozone (O₃). The IEA estimates over half of global NO_x emissions are produced by the transport sector. 47

3.6: Deployment of low-emission vehicles

Switching to low-emission transport systems is an important component of climate mitigation and will help reduce concentrations of most ambient air pollutants (though counterintuitively in some

settings, may lead to greater concentrations of ozone because of the 'titration effect' of NO₂).⁵ Personal exposure to traffic-related air pollution is a function of both ambient concentrations (a function of vehicle technology and other factors) and time activity patterns.⁶⁰⁻⁶⁴ The IEA maintains a technical-economic database that includes detailed transport activity, vehicle activity, energy demand, and well-to-wheel GHG and pollutant emissions.⁶⁵ The Global Electric Vehicle Outlook report tracks sales of electric vehicles, and the International Council on Clean Transportation maintains a set of data tables, comparison charts, and a conversion tool for comparing passenger vehicle fuel standards. These types of databases and methods will provide the quantitative backing to this indicator domain for the Lancet Countdown, which will track clean transport technology (e.g. electric vehicles) deployment and sector-specific emission factor trends at a variety of geographic scales.

3.7: Active transport infrastructure and uptake

- While decarbonizing the vehicle fleet is essential for meeting climate mitigation targets and improving urban air quality, in most settings encouraging the greater uptake of active travel (walking and cycling) for shorter journeys offers the greater opportunity for public health benefit. This is because active travel can lead to appreciable improvement in the levels of physical activity at population level, with all the attendant benefits in terms of reduced risk of cardiovascular disease, selected cancers, dementia, and diabetes, and improvement in mental well-being. Whilst these benefits may be partly offset by increases in exposure to road danger and, in some settings, higher exposure to ambient air pollution, the injury risks can be moderated by policies to improve road safety.⁷
- Indicators to be considered include (where available) proportion of journeys and distance taken on foot and bicycle in major urban areas. Such data require travel surveys, which are routinely implemented and with comparable methods only in selected (mainly high income) settings. In these cases, data on the duration of active travel, and the number of road crashes that occur could be collected. The contribution of change in active travel to population health could be computed given other population health and activity data, but to date, has seldom been assessed outside research studies.

3.8: The Food System

Sub-indicators under this heading will consider how food consumption and production affects and is affected by climate change, and the additional impacts this has on health. There is growing evidence on the benefits to health from more sustainable food systems and dietary change.

Consumption

Although all foods contribute to the emission of climate pollutants to different extents, among those with the greatest GHG footprints are meat and dairy products. These also contribute to water scarcity, land use change and erosion, and ultimately increase risks for cardiovascular disease, and some cancers in the case of red and processed meats. Emissions per unit protein produced are especially high for ruminants (cattle, goat and sheep), as compared to pork and poultry production. Although desirable to track trends in consumption patterns – especially of meat, dairy and

vegetables – the calculation of associated GHG emissions in particular, is complex, and so too the computation of health effects. In certain countries, the contribution of livestock to GHG emissions may be less substantial than in most high-income countries. Additionally, in many low- and middle-income countries, animal products may be important, providing an invaluable source of nutrients particularly to children and pregnant women. We propose measures of consumption of meat, dairy, vegetables and legumes, derived from standardized population dietary surveys and food diaries could be used to track consumption changes. By converting these datasets into *per capita* quantities, these trends could be tracked annually. However, data from (nationally) representative surveys is not widely available. Case-studies may therefore be a useful tool for highlighting the health and climate benefits of reduced meat and dairy consumption.

Production

- Agricultural production can be a significant contributor to GHG emissions. As such, there is clear room for improvement in water, carbon and nitrogen management in over-fertilized regions (e.g. China and India) to reduce GHG emissions and water pollution from agricultural lands and enhance environmental sustainability. For example, it is estimated that in the farmlands of China, a nitrogen use efficiency improvement from 31% to 50% would cut synthetic nitrogen use by 41%, and GHG emissions by 39%. The second element of this indicator domain will track changes in food production and food waste over time and the consequent impacts upon GHG emissions and health. GHG emissions associated with agriculture (including livestock systems and biogeochemical processes) will be quantified using existing models (e.g. DAYCENT or DNDC). To DNDC).
- An understanding of the potential for multiple environmental factors to affect food systems, and the nuances within and between countries and cultures is needed. Further work will be required to refine indicators for this area, and the scientific community is invited to suggest potential metrics and data sources.

3.9: Emissions from the Healthcare System

The health sector is a major contributor to GHG emissions, and has both a special responsibility and significant measurable opportunity to lead by example in reducing its carbon footprint. Pharmaceuticals, for example, are associated with high levels of avoidable GHG emissions, and there is nearly always scope for savings in transport and procuring goods needed to support the health system. Such actions have already been demonstrated to deliver health, social, environmental, and economic benefits, both immediate and long term. Calculation of the carbon intensity and emissions of the health sector has been achieved in England and the USA (serially in the former), despite complexities in capturing all inputs to provide comparable data over time, place and sub-sector. For the purposes of the Lancet Countdown, we will collect purposive samples from countries where data are available initially to raise the profile of the topic within the health community locally, nationally, and globally.

Economics and Finance

Article 2 of the Paris Agreement establishes the importance of ensuring financial flows consistent with a pathway towards a low-carbon economy. The focus of the Lancet Countdown's fourth working group is on the ways in which flows of finance and economic incentives are developing to

accelerate progress on health and climate change. Indicators fall into three broad 'themes': investment in the low-carbon economy; valuing the health co-benefits of mitigation; and pricing the health externalities of fossil fuels.

4.1 to 4.3: Investing in the low-carbon economy

Having made the case for a comprehensive response to climate change and the resultant health benefits, three of the proposed indicator domains in working group 4 will track investment in the low-carbon economy; specifically in renewable energy, energy efficiency and innovation.

The first two are closely linked, measuring changes in annual investments in renewable energy and in energy efficiency. In order to decarbonise the global energy system in order to meet the global climate change commitments in the Paris Agreement, whilst simultaneously managing a rise in demand for energy over the coming decades, low-carbon technologies and energy efficiency must account for around 90% of the \$2.5 trillion global annual investment required by 2035. In 2014, this value stood at 23%. Data for annual global investment in renewable energy is compiled and reported by Bloomberg New Energy Finance, which may be presented by the Lancet Countdown. Investment in energy efficiency, however, may be more difficult to track, as there is no standard agreed definition on what constitutes energy efficiency investment, which is carried out by a multitude of agents (often without the use of external finance), and is difficult to disaggregate from other activities. One approach is to estimate energy efficiency investment through modelling techniques. Further work will include discussions with the IEA and other organizations to determine the most appropriate definition to employ and how estimates of investment may be calculated. Estimates of total energy system investment are also published frequently by the IEA, allowing a proportional value for low-carbon technologies and energy efficiency to be calculated.

The third indicator domain here would track innovation in the low-carbon sector, by measuring annual changes in the generation of patents for low-carbon and energy efficient technologies. The data for such calculations may be found in various databases¹ and would capture the results of a substantial proportion of research and development efforts and funding by both public and private sector actors.

4.4: Valuing the health co-benefits of climate change mitigation

Building on work from working group 3, this indicator domain will aim to capture the costs and savings resulting from the health co-benefits of mitigation, across a variety of sectors. In particular, the health-related economic benefits (or costs) of changes in coal-based electricity generation, conventional car sales (i.e. petrol and diesel), and a rise in active transport. It is estimated that the annual value of the health impacts of ambient air pollution, principally caused by coal-based electricity generation and conventional vehicles, is as high as \$3.5 trillion (~5% Gross World Product) in the OECD (plus India and China).⁷⁸ Estimates of health-related economic benefits that result from mitigation policies would draw on indicators compiled and reported in working group 3 (e.g. coal phase out rates, low-emission vehicle sales and investment in active transport). Depending on the final form of the indicators presented in working group 3, these indicators may be produced either by relatively simple calculations, or through the use of energy system models that consider air pollution aspects of system developments. If the latter approach is required, further development of

¹ For example, the European Patent Office Worldwide Database (PATSTAT).

in-house air pollution modelling capabilities, or collaboration with other institutions, would be sought. Furthermore, this indicator domain closely relates to the estimating of health benefits of National Determined Contributions (NDCs) under working group 5; these two indicator domains will be jointly refined to ensure they complement each other.

4.5-4.7: Pricing the health externalities of fossil fuels

The third and final indicator theme within this working group would measure whether we are 'getting the prices right' to encourage the development of a low-carbon economy, and the heath-related benefits this brings, including ensuring that that inequities are addressed. Three areas of work would fall under this theme. The first concerns the presence of subsidies (such as tax breaks) for fossil fuel production and consumption, which incentivise their use and increase relative costs of renewable alternatives. In 2014, global fossil fuel subsidies stood at around \$490 billion – around four times the level of subsidy afforded to the deployment of renewable energy. Although the reform of such subsidies between 2009 and 2014 means that current subsidy levels are around \$117 billion lower than they would otherwise have been, much more needs to be done. This is recognised by SDG 12c, the indicator for which, when developed, may be employed here. However, further work will be conducted to determine which definition of fossil fuel subsidies may be suitability employed for the purposes of the Countdown.

The second indicator domain would cover the spread and strength of carbon pricing, which seeks to internalise the 'market externality' of CO₂ (and other GHG) emissions globally. Carbon pricing instruments currently cover around 12% of global GHG emissions, although with wide ranging values (from under \$1/tCO₂e to around \$130/tCO₂e).⁷⁹ This indicator may consist of two elements; the change in (and absolute level of) the proportion of global GHG emissions to which carbon pricing is applied, and the change in (and absolute value of) the weighted-average global carbon price. These data may be drawn directly from, or calculated based on, the World Bank's annual *State and Trends of Carbon Pricing* report.

The development of such indicators interact with and complement Indicator domains 4.1-4.3. Whilst the reduction of fossil fuel subsidies and increase in the spread and strength of carbon pricing 'pushes' the flow of finances towards the deployment and development of low-carbon and energy efficient technologies and measures, other policies, such as renewable energy subsidies, help to 'pull' finance towards such investments. Indicator domains 4.1 - 4.3 implicitly measure the impact generated by both influences.

What has been presented thus far do not address potential concerns surrounding the equity of the low-carbon transition, with carbon pricing on fuels having potentially regressive impacts. These impacts may be dampened or avoided with the appropriate public policies, such as environmental tax reform (ETR). ETR involves shifting the burden of tax from 'goods', such as labour or environmentally beneficial products or actions, to 'bads', such as pollution. Such a shift in economic incentives may, when well designed, produce a 'double dividend' of environmental improvement with social and economic benefit. As such, the third indicator domain under this theme concerns the use of revenue generated by carbon pricing instruments, with qualitative consideration for the intended end-use of this revenue. Further work is required to determine whether revenue from the reduction of fossil fuel subsidies may also be included in this indicator.

Political and Broader Engagement

The fifth working group will focus on the broader context within which progress on health and climate change is being made. These indicator domains will track the implementation of political commitments within the UNFCCC, alongside analysis of scientific and public engagement with health and climate change, which both provide background and context for policy implementation.

5.1: Public engagement in health and climate change

Globally, public engagement with climate change is mixed. In two surveys of a range of high-income, middle-income and low-income countries, most people considered that climate change was a 'very serious problem' and 'a major threat', and would impact directly on the conditions for health. ^{81,82} Asked when they thought climate change would start to substantially harm people in their country (now, in 10 years, in 25 years, in 50 years, in 100 years, never), the majority of respondents in most countries thought that their citizens were being substantially harmed now. Figure 3 presents the proportions responding 'now' and 'in 10 years'. Given that lack of understanding is one of the largest perceived barriers to individual engagement, the finding that framing climate change as a public health issue enhances engagement, is significant. ⁸³⁻⁸⁵




Figure 3. Proportions of the population who regard climate as substantially harming people in their country now or within 10 years (World Bank, 2009⁸²)

The Lancet Countdown will bring together evidence on changes in public understanding relating to health and climate change – more specifically, in terms of perceptions of threats and opportunities

in the response. One possible long-term, but resource-intensive approach would be using phone-and online-based public opinion polling techniques across a range of countries and settings. In the interim, the Lancet Countdown proposes to undertake an annual analysis of social media to build a broad understanding of public perceptions, and track the evolution of public engagement and knowledge. The Lancet Countdown's social media analysis will track levels of public engagement with health and climate change over time and identify key events that cause spikes in engagement; whether such spikes result in longer-term engagement; and countries where engagement is particularly high or low.

- 5.2 and 5.3: Academic publications on health and climate change & its inclusion within the medical curricula
- The Lancet Countdown will also track scientific engagement with health and climate change. Annual reviews of published scientific articles, using a bibliometric search relating to "climate" and "health" provides a potentially useful indicator, demonstrating research trends and coverage. Historical trends and research gaps including disciplinary and geographical focus could also be explored. This study will provide a more extensive and inclusive overview than previously published reports and reviews, and may also be a useful resource for informing future research funding by gaps and priorities. A study protocol for this scoping review has been written and accepted for publication. 86
 - To accompany an analysis of the academic literature, the Lancet Countdown will also follow the extent to which health and climate change is incorporated into the educational curricula of health professionals (initially, medical professionals), around the world. This analysis would determine not just the inclusion of climate change in these curricula, but also the emphasis on these issues (for example, whether this subject matter is mandatory or optional). This may be used to provide background context, rather than as an indicator in its own right.

- 5.4 and 5.5: Health and climate change in the UNFCCC and UNGA and implementation and estimated health benefits of the NDCs
- At the international level, the UNFCCC negotiations and the Paris Agreement provide an important framework for mitigation and adaptation policies. Indicator domains 5.4 and 5.5 would both look to provide an overview of the extent to which human health and wellbeing is considered within these political processes. Indicator domain 5.4 would examine the inclusion of health within the transcripts of the high-level statements delivered at the UNFCCC's annual Conference of the Parties, and the UN General Assembly. The former has previously been compiled on an ad hoc basis and would require additional work, but a database of the latter is readily available. The Lancet Countdown would analyse high-level statements to monitor how the inclusion and framing of health and climate change evolves over time. This work could be back-dated to extend this analysis to historical high-level statements, thus providing a longer time series to analyse.
- The final indicator domain for the Lancet Countdown would estimate the health benefits or disadvantages of the NDCs. Initially, the NDCs and subsequent communiques to the UNFCCC will be explored for substantive references and considerations of the relationship between public health and climate change. Over time, it is hoped that the potential health co-benefits of mitigation, from a reduction in air pollution, may be modelled. This would be conducted in a similar way to the analysis conducted by the Day, Höhne, and Gonzales in their 2015 assessment.⁸⁷ Many of these changes may

650 651	be captured by the UNFCCC's NAZCA (non-state actor zone for climate action) process — a potential source for future indicators and monitoring.
652	
653 654 655 656	Conclusion The Lancet Countdown: Tracking Progress on Health and Climate Change is an international, multi-disciplinary research collaboration, dedicated to tracking progress on health and climate change from 2016 to 2030.
657 658 659 660 661 662	The Lancet Countdown will be governed by a board comprising the research leads for each working group, and coordinated by a smaller executive team responsible for supporting the working groups to deliver and communicate the academic content. Over the coming months, it will work to establish an international advisory board, to provide strategic direction to the process and assist with policy and stakeholder engagement. This advisory board will be made up of academics and senior health and climate change experts from a broad range of geographies.
663 664 665 666	The indicators and indicator domains proposed in this paper are intended to form the foundation of our process, and will be refined and developed further over the coming months and throughout the Lancet Countdown's process. We invite ongoing direct input on the content, methods, and data of each of these, through the forms available at www.LancetCoundown.org/IndicatorConsultation
667	
668	
669	
670	
671 672	[Contributors, Declaration of Interests, Acknowledgements]

References

- World Health Organization. Preventing disease through healthy environments: a global assessment of the burden of disease from environmental risks. Geneva, 2016.
- 677 2. Costello A, Abbas M, Allen A, et al. Managing the health effects of climate change. *The*
- 678 *Lancet* 2009; **373**(9676): 1693-733.
- 679 3. Smith KR, Woodward A, Campell-Lendrum D, et al. Human Health Impacts, adaptation and
- 680 co-benefits. Climate Change 2014: Impacts, Adaptation, and Vulnerability Working Group II
- Contribution to the IPCC 5th Assessment Report. Cambridge, UK and New York, NY, USA: Cambridge
- University Press; 2014.
- 4. Haines A, McMichael A, Epstein P. Global Health Watch: Monitoring Impacts of
- 684 Environmental Change. *The Lancet* 1993; **342**: 1464-9.
- 685 5. Watts N, Adger N, Agnolucci P, et al. Lancet Commission on Health and Climate Change:
- Policy Responses to Protect Public Health. *The Lancet* 2015; **386**(10006): 1861-914.
- 687 6. Friel S, Dangour A, Garnett T, et al. Public health benefits of strategies to reduce
- greenhouse-gas emissions: food and agriculture. *The Lancet* 2009; **374**: 2016-25.
- Woodcock J, Edwards P, Tonne C, et al. Public health benefits of strategies to reduce
- 690 greenhouse-gas emissions: urban land transport. The Lancet 2009; **374**(9705): 1930-43.
- 691 8. Wilkinson P, Smith KR, Davies M, et al. Public health benefits of strategies to reduce
- 692 greenhouse-gas emissions: household energy. The Lancet 2009; 374(9705): 1917-29.
- 693 9. Markandya A, Armstrong BG, Hales S, et al. Public health benefits of strategies to reduce
- greenhouse-gas emissions: low-carbon electricity generation. *The Lancet* 2009; **374**(9706): 2006-15.
- 695 10. Haines A, McMichael AJ, Smith KR, et al. Public health benefits of strategies to reduce
- 696 greenhouse-gas emissions: overview and implications for policy makers. The Lancet 2009; **374**(9707):
- 697 2104-14.
- 698 11. IPCC. Summary for Policymakers. In: Field CB, V.R. Barros, D.J. Dokken, K.J. Mach, M.D.
- 699 Mastrandrea, T.E. Bilir, M. Chatterjee, K.L. Ebi, Y.O. Estrada, R.C. Genova, B. Girma, E.S. Kissel, A.N.
- 700 Levy, S. MacCracken, P.R. Mastrandrea, and L.L. White ed. Climate Change 2014: Impacts,
- 701 Adaptation, and Vulnerability Part A: Global and Sectoral Aspects Contribution of Working Group II
- to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change. Cambridge,
- 703 United Kingdom and New York, NY, USA: Cambridge University Press; 2014: 1-32.
- 704 12. Patz J, Gibbs H, Foley J, Rogers J, Smith K. Climate Change and Global Health: Quantifying a
- 705 Growing Ethical Crisis. *EcoHealth* 2007; **4**(4): 397-405.

- 706 13. Whitmee S, Haines A, Beyrer C, et al. The Rockefeller Foundation Lancet Commission on
- 707 Planetary Health: Safeguarding human health in the Anthropocene epoch. *The Lancet* 2015;
- 708 **386**(10007): 1973-2028.
- 709 14. Rockström J, Steffen W, Noone K, et al. Planetary boundaries: exploring the safe operating
- space for humanity. *Ecology and Society* 2009; **14**(2).
- 711 15. International Energy Agency. Energy and Climate Change: World Energy Outlook Special
- 712 Briefing for COP21. Paris, 2015.
- 713 16. Organisation for Economic Co-operation and Development. A Core Set of Indicators for
- 714 Environmental Performance Reviews Paris, 1993.
- 715 17. Brooks N. Vulnerability, risk and adaptation: A conceptual framework. Norwich: Tyndall
- 716 Centre for Climate Change Research, 2003.
- 717 18. Hambling T, Weinstein P, Slaney D. A review of frameworks for developing environmental
- 718 health indicators for climate change and health. *International journal of environmental research and*
- 719 *public health* 2011; **8**(7): 2854-75.
- 720 19. World Health Organization, UN Framework Convention on Climate Change. Climate and
- Health Profiles A Global Overview. Geneva, 2015.
- 722 20. United Nations. Transforming Our World: The 2030 Agenda for Sustainable Development.
- 723 A/Res/70/1. New York, 2015.
- 724 21. UCL Institute for Environmental Design and Engineering. Sustainable Healthy Urban
- 725 Environments. 2016. https://www.bartlett.ucl.ac.uk/iede/research/project-
- 726 <u>directory/projects/sustainable-healthy-urban-environments</u> (accessed 05 Oct 2016).
- 727 22. Reduction UOfDR. Indicators to Monitor Global Targets of the Sendai Framework for Disaster
- 728 Risk Reduction 2015-2030: A Technical Review. Geneva, 2015.
- 729 23. Åström C, Orru H, Rocklöv J, Strandberg G, Ebi KL, Forsberg B. Heat-related respiratory
- hospital admissions in Europe in a changing climate: a health impact assessment. BMJ Open 2013;
- 731 **3**(1).
- 732 24. Kjellstrom T, Briggs D, Freyberg C, Lemke B, Otto M, Hyatt O. Heat, Human Performance, and
- 733 Occupational Health: A Key Issue for the Assessment of Global Climate Change Impacts. Annu Rev
- 734 *Public Health* 2016; (37): 97-112.
- 735 25. Revitch B, Shaposhnikov D. Climate change, heat and cold waves as risk factors of increased
- 736 mortality in Russia. *Ecoforum* 2012; **2**(10): 122-38.
- 737 26. Jacob D, Petersen J, Eggert B, et al. EURO-CORDEX: new high-resolution climate change
- 738 projections for European impact research. Reg Environ Change 2013; (1-16).

- 739 27. Dunne JP, Stouffer RJ, John JG. Reductions in labour capacity from heat stress under climate
- 740 warming. *Nature Clim Change* 2013; **3**(6): 563-6.
- 741 28. Glantz M, Katz R. When is a drought a drought? *Nature* 1977; **267**: 192-3.
- 742 29. Wilhite D, Glantz M. Understanding the Drought Phenomenon: The Role of Definitions.
- 743 *Watern International* 1985; **10**(3): 111-20.
- 744 30. Dai A. Increasing drought under global warming in observations and models. Nature Clim
- 745 *Change* 2013; **3**: 52-8.
- 31. Bongaarts J. Development: Slow down population growth. *Nature* 2016; **530**(7591): 409-12.
- 747 32. McMichael A, Woodruff R. Climate change and infectious diseases. In: Mayer KH, Pizer HF,
- 748 eds. The Social Ecology of Infectious Diseases: Academic Press; 2011.
- 749 33. Altizer S, Ostfeld RS, Johnson PTJ, Kutz S, Harvell CD. Climate Change and Infectious
- 750 Diseases: From Evidence to a Predictive Framework. Science 2013; 341(6145): 514-9.
- 751 34. Siraj A, Bouma M, Santos-Vega M, Pascual M. Temperature and population density
- determine reservoir regions of seasonal persistence in highland malaria. *Proceedings of the Royal*
- 753 *Society B: Biological Sciences* 2015; **282**(1820).
- 754 35. Lafferty KD. Calling for an ecological approach to studying climate change and infectious
- 755 diseases. *Ecology* 2009; **90**(4): 932-3.
- 756 36. Australian Academy of Science. Climate change challenges to health: Risks and
- 757 opportunities. Canberra, 2014.
- 758 37. Victor LY, Edberg SC. Global Infectious Diseases and Epidemiology Network (GIDEON): a
- 759 world wide Web-based program for diagnosis and informatics in infectious diseases. Clin Infect Dis
- 760 2005; **40**(1): 123-6.
- 761 38. Brownstein JS, Freifeld CC, Reis BY, Mandl KD. Surveillance Sans Frontieres: Internet-based
- 762 emerging infectious disease intelligence and the HealthMap project. PLoS Med 2008; 5(7): e151.
- 763 39. Victor LY, Madoff LC. ProMED-mail: an early warning system for emerging diseases. *Clin*
- 764 Infect Dis 2004; **39**(2): 227-32.
- 765 40. United Nations Environment Programme. Avoiding Future Famines: Strengthening the
- Ecological Foundation of Food Security through Sustainable Food Systems. Nairobi, 2012.
- 767 41. Food and Agricultural Organization, International Fund for Agricultural Development, World
- 768 Food Programme. The State of Food Insecurity in the World: How does international price volatility
- affect domestic economies and food security? Rome, 2011.

- 770 42. WHO. Climate and Health Country Profiles 2015: A Global Overview. Geneva: World Health
- 771 Organization, 2015.
- 772 43. WHO. WHO guidance to protect health from climate change through health adaptation
- 773 planning. Geneva: World Health Organization, 2014.
- 774 44. WHO. Health in the Intended Nationally Determined Contributions (INDCs) to the United
- 775 Nations Framework Convention on Climate Change, 2015. Geneva: World Health Organization, 2016.
- 776 45. Manga L, Bagayoko M, Meredith T, Neira M. Overview of health considerations within
- National Adaptation Programmes of Action for climate change in least developed countries
- and small island states, 2010.
- 779 46. Bruckner T, Bashmakov I, Mulugetta Y, et al. Energy Systems. In: Edenhofer O, Pichs-
- 780 Madruga R, Sokona Y, et al., eds. Climate Change 2014: Mitigation of Climate Change Contribution of
- 781 Working Group III to the Fifth Assessment Report of the Intergovernmental Panel on Climate
- 782 Change. Cambridge, United Kingdom and New York, NY, USA: Cambridge University Press; 2014.
- 783 47. Agency IE. Energy and Air Pollution. World Energy Outlook Special Report 2016.
- 784 48. International Energy Agency. World Energy Outlook 2014. Paris, 2015.
- 785 49. International Energy Agency. Medium-Term Coal Market Report 2015: Market Analysis and
- 786 Forecasts to 2020. Paris, 2015.
- 787 50. International Energy Agency. Renewables Information 2016. Paris, 2016.
- 788 51. Wilkinson P, Smith KR, Beevers S, Tonne C, Oreszczyn T. Energy, energy efficiency, and the
- 789 built environment. *Lancet* 2007; **370**: 1175-87.
- 790 52. World Health Organization. Burden on Disease from Air Pollution in 2012; 2014.
- 791 http://www.who.int/phe/health_topics/outdoorair/databases/FINAL_HAP_AAP_BoD_24March2014
- 792 <u>.pdf</u> (accessed 7 Oct 2014).
- 793 53. Hancock K. The expanding horizon of renewable energy in sub-Saharan Africa:Leading
- research in the social sciences. *Energy Research & Social Science* 2015; **5**: 1-8.
- 795 54. International Energy Agency. World Energy Outlook 2015 Electricity Access Database. Paris,
- 796 2015.
- 797 55. Adair-Rohani H, Zukor K, Bonjour S, et al. Limited electricity access in health facilities of sub-
- 798 Saharan Africa: a systematic review of data on electricity access, sources, and reliability. Glob Heal
- 799 *Sci Pract* 2013; **1**(2): 249-61.
- 800 56. Bhatia M, Angelou N, Soni R, et al. Access to Modern Energy Services for Health Facilities in
- 801 Resource-Constrained Settings: The World Bank and World Health Organization, 2015.

- 802 57. Sokhi RS, Kitwiroon N. Air Pollution in Urban Areas. In: Sokhi RS, ed. World Atlas of
- 803 Atmospheric Pollution: Anthem Press; 2011: 19-34.
- 804 58. Health Effects Institute. Traffic-Related Air Pollution: A Critical Review of the Literature on
- 805 Emissions, Exposure and Health Effects. 2010; **Special Report**: 386.
- 806 59. Daly HE, Ramea K, Chiodi A, Yeh S, Gargiulo M, Gallachóir BÓ. Incorporating travel behaviour
- and travel time into TIMES energy system models. APPLIED ENERGY 2014; 135: 429-39.
- 808 60. Yim SHL, Stettler MEJ, Barrett SRH. Air quality and public health impacts of UK airports. Part
- 809 II: Impacts and policy assessment. *Atmospheric Environment* 2013; **67**: 184-92.
- 810 61. Yim SHL, Barrett SRH. Public health impacts of combustion emissions in the United Kingdom.
- 811 Environmental Science & Technology 2012; **46**: 4291-6.
- 812 62. Walton BH, Dajnak D, Beevers S, Williams M, Watkiss P, Hunt A. Understanding the Health
- 813 Impacts of Air Pollution in London. 2015; 2015.
- 814 63. Stettler MEJ, Eastham S, Barrett SRH. Air quality and public health impacts of UK airports.
- Part I: Emissions. *Atmospheric Environment* 2011; **45**: 5415-24.
- 816 64. Caiazzo F, Ashok A, Waitz Ia, Yim SHL, Barrett SRH. Air pollution and early deaths in the
- 817 United States. Part I: Quantifying the impact of major sectors in 2005. Atmospheric Environment
- 818 2013; **79**: 198-208.
- 819 65. International Energy Agency. Modelling of the transport sector in the Mobility Model
- 820 (MoMo). 2016. https://www.iea.org/etp/etpmodel/transport/ (accessed 12 Aug 2016).
- 821 66. World Cancer Research Fund. Colorectal Cancer Report: Food, nutrition, physical activity,
- and the prevention of colorectal cancer. London, 2011.
- 823 67. Pelster D, Gisore B, Goopy J, et al. Methane and nitrous oxide emissions from cattle excreta
- on an East African grassland. *Journal of Environmental Quality* 2016; **45**(5): 1531-9.
- 825 68. Zhang X, Davidson E, Mauzerall D, Searchinger T, Dumas P. Managing nitrogen for
- sustainable development. *Nature* 2015; **528**(7580): 51-9.
- 827 69. Huang Y, Tang Y. An estimate of greenhouse gas (N2O and CO2) mitigation potential under
- various scenarios of nitrogen use efficiency in Chinese croplands. *Global Change Biology* 2010;
- **11**(16): 2958-70.
- 830 70. Grosso S, Parton W, Mosier A, Walsh M, Ojima D, Thornton P. DAYCENT: National-Scale
- Simulations of Nitrous Oxide Emissions from Cropped Soils in the United States. *Journal of*
- 832 *Environmental Quality* 2006; **35**(4): 1451-60.
- 833 71. Gilhespy S, Anthony S, Cardenas L, et al. First 20 years of DNDC (DeNitrification
- DeComposition): Model evolution. *Ecological Modelling* 2014; **292**: 51-62.

- 835 72. Li C, Salas W, Zhang R, Krauter C, Rotz A, Mitloehner F. Manure-DNDC: a biogeochemical
- 836 process model for quantifying greenhouse gas and ammonia emissions from livestock manure
- systems. Nutrient Cycling in Agroecosystems 2012; 93(2): 163-200.
- 838 73. UK Sustainable Development Unit. Module on Carbon Hotspots. Sustainable Development
- 839 Strategy for the Health and Care System 2014 2020. London; 2014.
- 840 74. NHS Sustainable Development Unit. Sustainable Development in Health and Care Health
- 841 Check 2016: NHS England & Public Health England, 2016.
- 842 75. Chung J, Meltzer D. Estimate of the Carbon Footprint of the US Healthcare Sector. Journal of
- 843 the American Medical Association 2009; **302**(18): 1970-2.
- 844 76. Eckelman M, Sherman J. Environmental Impacts of the US Health Care System and Effects on
- 845 Public Health. *PLoS ONE* 2016; **11**(6).
- 846 77. International Energy Agency. World Energy Investment Outlook. Paris; 2014.
- 78. Organisation of Economic Co-operation and Development. The Cost of Air Pollution: Health
- 848 Impacts of Road Transport. Paris, 2014.
- 849 79. World Bank Group. State and Trends of Carbon Pricing. Washinton D.C., 2015.
- 850 80. Patuelli R, Nijkamp P, Pels E. Environmental tax reform and the double dividend: A meta-
- analytical performance assessment. *Ecological Economics* 2005; **55**(4): 564-83.
- 852 81. Pew Research Center. Climate Change and Financial Instability Seen as Top Global Threats
- Washington DC: Pew Research Center; 2013.
- 854 82. World Bank Group. Public attitudes toward climate change: findings from a multi-country
- poll. Background note to the world development report 2010. Washington, D.C., 2009.
- 856 83. Lorenzoni I, Nicholson-Cole S, Whitmarsh L. Barriers perceived to engaging with climate
- change among the UK public and their policy implications. *Global Environmental Change* 2007; **17**(3-
- 858 4): 445–59.
- 859 84. Myers T, Nisbet M, Maibach E, Leiserowitz A. A public health frame arouses hopeful
- emotions about climate change. *Climate Change* 2012; **113**(3-4): 1105–12.
- 85. Maibach E, Nisbet M, Baldwin P, Akerlof K, Diao G. Reframing climate change as a public
- health issue: an exploratory study of public reactions. *BMC Public Health* 2010; **10**(299).
- 86. Herlihy N, Bar-Hen A, Verner G, et al. Climate Change and Human Health: What are the
- Research Trends? A scoping review protocol. . *BMJ Open (accepted, not published)* 2016.

865 87. Day T, Höhne N, Gonzales S. Assessing the missed benefits of countries' national contributions: quantifying potential co-benefits. Cologne: New Climate Institute, 2015.
867
868
869
870
871

Appendices

 Appendix 1: Re-considering the health and climate indicators using an adaptation of the DPSEEA Framework:

Red indicates available data, orange data which can be obtained by data processing or are available only for selected sites and blue markers which require modelling or special data gathering. Related Sustainable Development Goals are indicated where relevant.

Duining Fores	Duccessures	States/Functions	Effects	Actions (I	Responses)
Driving Forces	Pressures	States/Exposures	Effects	Specific	General
Dependence on energy from combustion of fossil fuels; other human activities leading to emissions of climate active pollutants (CAPs), including agriculture and land use change	Emission of CAPs → altered concentration in the atmosphere	Increased radiative forcing → global warming, with regional variations	Impacts on: • health • productivity	(1) Measures to a effects of climate (2) Measures to n change	change
<u>MARKERS</u>					
 Per capita use of energy (kw.hr.person⁻¹) [national, reported data] Per capita energy use per US\$ GDP (kw.hr.US\$M⁻¹) 	 Annual total and per capita emissions of climate active pollutants (Gtonne CO₂.e) [country,] CO₂ concentrations 	Mean of (warm season) daily maximum or mean temperatures [city, observed series]	Heat- (and cold-) related mortality/ morbidity [city, requires epi modelling] Reduction in	 Implementati on of heatwave plans [national] Building regulation for protection against heat risks [national] 	Health and climate change in the UNFCCC and UNGA high-level statements Academic publication and funding on health and climate change
(kw.hr.US\$M ⁻) [national, reported data]	in atmosphere (ppmv) [global, monitored data Mauna Loa]	wet bulb globe temperatures for working hours [city, derived variable]	labour productivity from excess heat [national, model- based from WBGT]		 Integration of health in national adaptation plans
		population and proportion affected by flooding	Flood-related mortality and morbidity (including mental		Direct and indirect fossil fuel subsidies

	T	T	L	T	
		[national]	health)		[national]
			[national,		
			immediate		Adaptation
			observed deaths		finance for
			estimable only]		health [
		Periods of low	Nutrition-related		definition]
		rainfall resulting in	growth and		
		reduced crop yields	mortality impacts		 Change in
		[national]	in children		annual
			[national,		investment in
			requires		energy
			modelling]		efficiency
		Warm season mean	Deaths/		
		of ozone	morbidity from		 Public opinion
		concentrations?	ozone		on health and
		[city, not specific to	concentrations		climate
		climate change]	[city, modelled]		change
		Transmissions	Burdens (cases)		[national,
		potential for	of specific vector-		needs
		specific vector-	borne diseases		definition]
		borne diseases	[national, sentinel		definition
		(malaria, dengue)	sites]		<u>Note</u>
		[sentinel sites,	sitesj		SDG 12.c.1
		modelled from			Fossil-fuel
					subsidy per unit
		weather data only]			GDP
					GDP
					SDG 13.2.1
					Operationalizati on of
					adaptation plan
Fossil fuel	Number (net	Ambient	Mortality/	Growth in	
combustion for	,	concentration	morbidity	renewable	
	capacity in	(annual mean) of	attributable to		
electricity	Gigawatts) of	,		energy	 Cost-savings
generation	coal fired	PM _{2.5} from coal	ambient PM _{2.5}	resources	from the
(Tera-joule per	power stations	fired power	derived from coal	[national,	health co-
million population)	[national,	generation	fired generation	reported	benefits
[national, reported	reported data]	[city, requires	[city, model	data]	
data]		modelling or	based]		Coverage and
	 Emissions of 	method of source		 Renewable 	strength of
	CAPs from coal	apportionment]		energy patent	carbon pricing
	fired electricity			generation	carbon pricing
	production			and	
	[national]			innovation	
				[national,	Noto
				requires	Note
				definition]	SDG 13.a.1
1					Mobilized
			l	Notes	US\$/year
			Note:	Note:	toward- (100
			Note: SDG 3.9.1	SDG 7.1.2	towards \$100
				SDG 7.1.2	billion
			SDG 3.9.1 mortality	SDG 7.1.2 proportion of	billion commitment for
			SDG 3.9.1	SDG 7.1.2 proportion of population with	billion
			SDG 3.9.1 mortality attributable to	SDG 7.1.2 proportion of population with primary reliance	billion commitment for
			SDG 3.9.1 mortality attributable to household and	SDG 7.1.2 proportion of population with	billion commitment for
			SDG 3.9.1 mortality attributable to household and	SDG 7.1.2 proportion of population with primary reliance	billion commitment for

Per capita energy consumption, housing sector [national, reported data] Proportion of housing which is energy inefficient [national, requires definition]	Energy efficiency of housing stock (mean energy requirement to maintain standardized heating and/or cooling regime, as well as cooking and other household needs) [national, housing survey + modelling]		annual mean PM _{2.5} in cities Ancillary effects on deaths/ morbidity relating to exposures of the indoor environment [national, model-based in selected locations only]	SDG 7.2.1 renewable energy as share of final energy consumption Building regulation for energy efficiency [national]
Mean per capita energy use for transport (kj.person ⁻¹ .year ⁻¹) [national, reported data] Transport: per capita distance travelled by motorized transport [national or city, survey data]	Transport-related emissions of Climate Active Pollutants and ambient air pollutants [city, emissions inventories (where available)]	Ambient air PM _{2.5} concentrations attributable to transport-related emissions [city, requires modelling or method of source apportionment]	Reduction in deaths/morbidity from (transport-related contribution to) physical activity [national or city, definition of counterfactual] Deaths from transport-related PM _{2.5} exposure [city, model-based]	 Deployment of low-emission vehicles Active transport infrastructure and uptake [city, definition]
Per capita energy consumption, food and agriculture sector [national, reported data]		Per capita consumption of red meat & dairy products (kilojoules per person) [national, survey based]	Mortality/ morbidity attributable to consumption of red meat and dairy products [national, modelled and setting specific]	Note SDG 12.3.1 Global food loss index
[Carbon footprint of healthcare systems]				Implementation and health benefits of the NDCs [national, requires special data gathering and modelling]

Appendix 2: Indicators from other monitoring processes relevant to the Lancet Countdown

The table below maps the Lancet Countdown's indicators with those used for the Sustainable Development Goals, Sendai Framework for Disaster Risk Reduction, ClimateWorks Foundation Carbon Transparency Initiative, and WHO Climate and Health Country Profiles. It is important to note that while the intent of these initiatives is for eventual global coverage, they are still in development and so currently their implementation is limited to some countries. For example, the ClimateWorks Foundation Carbon Transparency Initiative has currently used their indicators for China, the EU, India, Mexico, and the US and WHO have Climate and Health Country Profiles for 40 countries.

Working Group	Lancet Countdown Indicator Domains	Sustainable Development Goals	Sendai Framework for Disaster Risk Reduction	ClimateWorks Foundation Carbon Transparency Initiative	WHO Climate and Health Country Profiles
Health Impacts and Climate Change	Track populations' exposure to heat Track changes in labour productivity Track populations' exposure to heatwaves	1.5.1 Number of deaths, missing persons and persons affected by disaster per 100,000 people 1.5.3 Number of countries with national and local disaster risk reduction strategies 6.4.2 Level of water stress: freshwater withdrawal as a proportion of available freshwater resources	A-1. Number of deaths and missing due to hazardous events per 100,000. A-2. Number of deaths due to hazardous events. A-3. Number of missing due to hazardous events. B-1. Number of affected people per 100,000. B-2. Number of injured or ill people due to hazardous events. B-3. Number of people who left their	Illitrative	Warmer and/or fewer cold days and nights over most land areas. Warmer and/or more frequent hot days and nights over most land areas. Heat-related mortality. Heat stress and work productivity. Warm spells/heatwaves. Frequency and/or duration increases over most land areas. Heat-related mortality. Heat-related mortality. Heavy precipitation events.
	Track populations' exposure to floods		places of residence due to hazardous events.		Increase in the frequency, intensity, and/or amount of heavy precipitation.

Track popu		B-3a. Number of evacuated people due to hazardous events. B-3b. Number of relocated people due to hazardous events. B-4. Number of people whose houses were damaged due to hazardous events. B-5. Number of people whose houses were destroyed due to hazardous events.	Increases in intense tropical cyclone activity. Increased incidence and/or magnitude of extreme high sea level. River flooding. Exposure to flooding due to sea level rise. Increases in intensity and/or
exposure to droughts		B-6. Number of people who received food relief aid due to hazardous events.	duration of drought.
Track the s			Populations at risk of infectious and vector-borne diseases for malaria and dengue fever.
Track popu food securi	2.1.1 Prevalence of undernourishment 2.1.2 Prevalence of moderate or severe food insecurity in the population, based on the Food Insecurity Experience Scale (FIES)	C-2. Direct agricultural loss due to hazardous events.	

		standard deviation from the median of the WHO Child Growth Standards) among children under 5 years of age, by type (wasting and overweight) 2.4.1 Proportion of agricultural area under productive and sustainable agriculture		
	Track the integration of health in National Adaptation Plans	3.d.1 International Health Regulations (IHR) capacity and health emergency preparedness 11.b.1 Proportion of local governments that adopt and implement local disaster risk reduction strategies in line with the Sendai Framework for Disaster Risk Reduction 2015-2030	E-3. Number of countries that integrate climate and disaster risk into development planning.	Governance and policy. Vulnerability, impact and
Health Resilience and Adaptation	Track climate services for health	13.2.1 Number of countries that have communicated the establishment or operationalization of an integrated policy/strategy/plan which increases their ability to adapt to the adverse impacts of climate change, and foster climate resilience and low greenhouse gas emissions development in a manner that does not threaten food production	D-2. Number of health facilities destroyed or damaged by hazardous events.	adaptation (health) assessments. Health adaptation strategies and action plans. Preparedness, risk management and integrated risk monitoring. Awareness raising and capacity building.
	Track adaptation finance for health	 (including a national adaptation plan, nationally determined contribution, national communication, biennial update report or other) 13.3.2 Number of countries that have communicated the strengthening of institutional, systemic and individual capacity-building to implement adaptation, mitigation and 		Financing.

		technology transfer, and development actions		
	Track the phase-out of coal		Share amount of coal in total final energy consumption—that is, the share of an economy's energy derived from coal. Share of electricity from coal generation.	
	Track the growth in renewable energy resources	7.1.2 Proportion of population with primary reliance on clean fuels and technology 7.2.1 Renewable energy share in the total final energy consumption	Share amount of renewable energy in total final energy consumption—that is, an economy's share of energy derived from renewable sources. Share of electricity from renewable energy generation.	
Health Co-	Track energy access	7.1.1 Proportion of population with access to electricity		
Benefits of Mitigation	Track energy access for health facilities	3.8.1 Coverage of essential health services (defined as the average coverage of essential services based on tracer interventions that include reproductive, maternal, newborn and child health, infectious diseases, non-communicable diseases and service capacity and access, among the general and the most disadvantaged population) 3.b.1 Proportion of the population with access to affordable medicines and vaccines on a sustainable basis		
	Track ambient air pollution exposure	3.9.1 Mortality rate attributed to household and ambient air pollution 11.6.2 Annual mean levels of fine particulate matter (e.g. PM2.5 and		Current exposures and health risks due to air pollution, including outdoor air pollution exposure, short-lived climate pollutants, and household air

		PM10) in cities (population weighted)			pollution.
-		weighted)	Sha	are of new vehicles in a	
			par	rticular geography that are	
	Track the		elec	ectric drive rather than internal	
	deployment of low		con	mbustion engine vehicles.	
	emission vehicles				
	CITIISSIOTI VETIICIES			are of electric drive vehicles for	
				e light-duty fleet in a particular	
			yea		
		11.2.1 Proportion of population that		tal terrestrial passenger	
		has convenient access to public		ometers, meaning the total	
		transport, by sex, age and persons		imber of kilometers that a	
		with disabilities		pulation travels, including on	
				ivate, public, and passenger	
			rail	II.	
			Nur	umber of kilometers travelled in	
				rrestrial modes—private,	
				iblic, and rail—on a per capita	
			bas		
			543		
			Tot	otal number of kilometers	
	Track active			avelled in private modes—light-	
	transport			ity vehicles, two wheelers, and	
	infrastructure and			ree-wheelers.	
	uptake				
			Tot	tal number of kilometers	
				avelled in private modes—light-	
				ity vehicles, two wheelers, and	
			thre	ree-wheelers—on a per capita	
			bas	isis.	
				otal number of kilometers	
				avelled in public modes—bus	
			and	d rail.	
				tal accordance of hill-residence	
				tal number of kilometers	
				avelled in public modes—bus	
			and	d rail—on a per capita basis.	

		Total number of vehicle kilometers travelled in private modes—light-duty vehicles, twowheelers, and three-wheelers. Share of passenger kilometers associated with public transport—bus and rail. Total share of electricity in the energy mix for all terrestrial transport—private, public, and freight modes. Share of kilometers associated with private modes of	
		transport—light-duty, two wheelers, and three-wheelers. Total amount of greenhouse gas	
		emissions associated with the Agriculture Sector.	
Track food	11.3.1 Ratio of land consumption rate to population growth rate 12.1.1 Number of countries with sustainable consumption and	Total amount of greenhouse gas emissions associated with the Agriculture Sector from direct sources in production and onsite energy use.	
consumption and production	production (SCP) national action plans or SCP mainstreamed as a priority or a target into national policies	Total amount of greenhouse gas emissions associated with the Agriculture Sector from electricity.	
	12.3.1 Global food loss index	Size of a herd of cattle in a given geography and year on a per capita basis. This metric does not include dairy cattle.	

			Share of agricultural emissions	
			associated with non-dairy cattle.	
			Share of agricultural emissions	
			associated with fertilizers.	
			Greenhouse gas emissions intensity associated with agriculture on a per capita basis.	
	Track the carbon footprint of healthcare systems			Annual greenhouse gas emissions by sector (metric tonnes in CO2-equivalent) – although not for healthcare.
	Track change in annual investment in renewable energy	7.2.1 Renewable energy share in the total final energy consumption		
	Track change in annual investment in energy efficiency	7.a.1 Mobilized amount of United States dollars per year starting in 2020 accountable towards the \$100		
	Track low-carbon technology patent generation and innovation	billion commitment for climate finance		
Finance and Economics	Track the value the health co-benefits of climate change mitigation			
	Track direct and indirect fossil fuel subsidies	12.c.1 Amount of fossil-fuel subsidies per unit of GDP (production and consumption) and as a proportion of total national expenditure on fossil fuels		
	Track the coverage and strength of carbon pricing			
	Equity of the low- carbon transition			

	Track public, civil society and community mobilisation on health and climate change Track academic publications on health and climate change			
	Track health and climate change in the UNFCCC and UNGA high-level statements			Governance and policy.
Political and Broader Engagement	Track the inclusion of health and climate change within medical and public health curricula	12.8.1 Extent to which (i) global citizenship education and (ii) education for sustainable development (including climate change education) are mainstreamed in (a) national education policies; (b) curricula; (c) teacher education; and (d) student assessment 13.3.1 Number of countries that have integrated mitigation, adaptation, impact reduction and early warning into primary, secondary and tertiary curricula		
	Track the implementation and estimated health benefits of the Nationally Determined Contributions	13.2.1 Number of countries that have communicated the establishment or operationalization of an integrated policy/strategy/plan which increases their ability to adapt to the adverse impacts of climate change, and foster climate resilience and low greenhouse gas emissions development in a manner that does not threaten food production (including a national adaptation plan,		Governance and policy.

	nationally determined contribution,		
	national communication, biennial		
	update report or other)		