

Flavanone-rich citrus beverages counteract the transient decline in postprandial endothelial function in humans: a randomised, controlled, double-masked, cross-over intervention study

Article

Accepted Version

Rendeiro, C., Dong, H., Saunders, C., Harkness, L., Blaze, M., Hou, Y., Belanger, R. L., Corona, G., Lovegrove, J. A. and Spencer, J. P. E. (2016) Flavanone-rich citrus beverages counteract the transient decline in postprandial endothelial function in humans: a randomised, controlled, double-masked, cross-over intervention study. *British Journal of Nutrition*, 116 (12). pp. 1999-2010. ISSN 0007-1145 doi: <https://doi.org/10.1017/S0007114516004219> Available at <https://centaur.reading.ac.uk/68614/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

Published version at: <http://dx.doi.org/10.1017/S0007114516004219>

To link to this article DOI: <http://dx.doi.org/10.1017/S0007114516004219>

Publisher: Cambridge University Press

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online

1 **Flavanone-rich citrus beverages counteract the transient decline in postprandial endothelial**
2 **function in humans: a randomized, controlled, double-masked, crossover intervention study**

3

4 **Catarina Rendeiro^{1,5}, Honglin Dong^{1,6}, Caroline Saunders², Laura Harkness³, Melvin Blaze⁴,**
5 **Yanpeng Hou⁴, Ronald L. Belanger⁴, Giulia Corona^{1,7}, Julie A. Lovegrove¹ and Jeremy P. E.**
6 **Spencer^{1*}.**

7 ¹ Department of Food and Nutritional Sciences, School of Chemistry, Food and Pharmacy, University of
8 Reading, PO Box 226, RG2 6AP, Reading, UK. ⁵*Current Address:* Beckman Institute for Advanced
9 Technology, University of Illinois UC, 405 Mathews Av, 61802, Urbana, USA; ⁶ *Current Address:*
10 School of Life Sciences, Faculty of Health and Life Sciences, Coventry University, Coventry, UK.
11 ⁷*Current Address:* Life Sciences Department, Whitelands College, University of Roehampton,
12 Holybourne Avenue, London, UK

13 ² PepsiCo R+D Nutrition, PepsiCo Inc., UK;

14 ³ Global R+D Nutrition, PepsiCo Inc., USA;

15 ⁴ PepsiCo R+D Biological & Discovery Analytics, PepsiCo Inc., USA

16

17 **Study registration:** The National Institutes of Health (NIH)-randomized trial records held on the NIH
18 ClinicalTrials.gov website (Registration number: NCT01963416; 10-10-2013).

19

20 **Funding:** This work was supported by PepsiCo Inc. (PEP-1122).

21

22 **Address Correspondence:** Jeremy PE Spencer, Molecular Nutrition Group, School of Chemistry, Food
23 and Pharmacy, University of Reading, Reading RG2 6AP, UK; e-mail: j.p.e.spencer@reading.ac.uk; tel:
24 +44 118 378 8724.

25

26 **Running head:** Flavanones and endothelial function

27 **ABSTRACT**

28

29 Specific flavonoid-rich foods/beverages are reported to exert positive effects on vascular function;
30 however, data relating to effects in the postprandial state are limited. The present study investigates the
31 postprandial, time-dependent (0 -7 h) impact of citrus flavanone intake on vascular function. An acute,
32 randomized, controlled, double-masked, crossover intervention study was conducted in middle aged
33 healthy men (30-65 yrs, n=28) to assess the impact of flavanone intake (Orange Juice: 128.9 mg;
34 flavanone-rich Orange Juice: 272.1 mg; homogenized whole orange: 452.8 mg; isocaloric control: 0 mg
35 flavanones) on postprandial (double meal delivering a total of 81 g of fat) endothelial function.
36 Endothelial function was assessed by flow-mediated dilatation (FMD) of the brachial artery at 0, 2, 5 and
37 7 h. Plasma levels of naringenin / hesperetin metabolites (sulphates and glucuronides) and nitric oxide
38 species (NO_x) were also measured. All flavanone interventions were effective at attenuating transient
39 impairments in FMD induced by the double meal (7 h post intake; $P<0.05$), but no dose response effects
40 were observed. The effects on FMD coincided with the peak of naringenin/hesperetin metabolites in
41 circulation (7 h) and sustained levels of plasma nitrite. In summary, citrus flavanones are effective at
42 counteracting the negative impact of a sequential double meal on human vascular function, potentially
43 through the actions of flavanone metabolites on NO.

44

45 **Key words:** citrus flavanones, endothelial function, high-fat meal, nitric oxide, postprandial

46 **Introduction**

47 A transient impairment of vascular function is known to occur in the postprandial or fed state ⁽¹⁻⁴⁾ and is
48 widely believed to impact on endothelial dysfunction and lifetime cardiovascular disease risk ⁽⁵⁻⁸⁾. In
49 particular, endothelial function has been shown to be transiently impaired (2-8 h) following ingestion of
50 moderate to high fat meals (36-80 g of fat) ⁽⁹⁻¹³⁾, potentially driven by hyperglycemia and
51 hypertriglyceridemia which occurs during the postprandial state ^(1, 14, 15). Observational data have
52 highlighted that the consumption of diets rich in flavonoids might lead to an improved cardiovascular
53 prognosis ⁽¹⁶⁻²⁰⁾. Indeed, flavonoid-rich foods and beverages are well reported to improve endothelial
54 function in humans both acutely ⁽²¹⁻²⁴⁾, short-term ⁽²⁵⁻²⁷⁾ and long-term ⁽²⁸⁻³¹⁾. However, most of the acute
55 interventions were undertaken with volunteers in the fasted state, which is considered less representative
56 of the free living state, whilst data relating to flavonoid potential to ameliorate acutely postprandial
57 endothelial impairments are more scarce ⁽³²⁻³⁴⁾.

58

59 Although, the precise mechanisms by which absorbed flavonoids and their circulating metabolites
60 mediate beneficial vascular effects remain unclear, there is evidence to suggest that the modulation of
61 circulating NO levels might be involved ^(22-24, 35-38). Notably, flavanol-rich cocoa has been consistently
62 shown to improve endothelium-dependent vasodilation in healthy individuals ^(24, 27, 31, 39, 40), smokers ^{(23,}
63 ²⁶⁾, patients with coronary artery disease (CAD) ⁽⁴¹⁾, hypertension ⁽⁴²⁾, or diabetes ⁽⁴³⁾. Particularly, acute
64 vascular improvements have been shown to coincide with the appearance of flavanol metabolites in the
65 circulation and with peak plasma NO levels ^(23, 24, 26, 42). Furthermore, flavanol-induced improvement in
66 vascular function are inhibited following co-administration of an eNOS inhibitor, suggesting a cause-
67 and-effect relationship between flavonoid intake, plasma NO levels and vascular function ^(23, 24).
68 Although less studied, flavanones from citrus, have also been shown to exert beneficial effects on human
69 vascular function ^(28, 44). In particular, chronic interventions with orange juice, or the pure flavanone,

70 hesperedin, resulted in a decrease in blood pressure in overweight volunteers and acute (6 h)
71 improvements in micro-vascular reactivity ⁽⁴⁵⁾. Short-term intake of pure hesperedin also resulted in
72 significant improvements in endothelial function (as measured by brachial artery FMD) in volunteers
73 with metabolic syndrome ⁽⁴⁶⁾.

74

75 In the present study, we assessed the impact of an acute intervention with increasing doses of orange
76 flavanones (sourced from differently processed orange beverages) on human vascular function in the
77 postprandial state. A sequential double meal (breakfast and lunch, delivering a total of 81 g of fat) was
78 used to simulate the fed state and investigate the postprandial time-dependent effects of flavanone intake
79 on endothelial function as measured by brachial artery FMD.

80

81 **SUBJECTS AND METHODS**

82 **Ethics**

83 The clinical trial was registered at clinicaltrials.gov (NCT01963416) and conducted according to the
84 Declaration of Helsinki following Good Clinical Practice (GCP). It was approved for conduct by the
85 University of Reading's Research Ethics Committee (ethics reference number 12/06). All volunteers
86 signed an informed consent form before commencing the study.

87

88 **Intervention study volunteers**

89 Volunteers were recruited from the University of Reading and surrounding area by use of the Hugh
90 Sinclair Unit volunteers' database, poster advertisement within the university and local community via
91 local websites (April-Dec 2012). Fifty-nine male healthy volunteers, aged 30-65 years old were assessed
92 for screening and selected according to the following inclusion criteria: 1) fasting lipids in the upper half

93 of the normal range (triacylglycerol 0.8-3.2 mmol/l and total cholesterol 6.0-8.0 mmol/l) or BMI 25-32
94 kg/m²; 2) Non-smoker; 3) Not diabetic (diagnosed or fasting glucose > 7 mmol/l) or suffer from
95 endocrine disorders; 4) hemoglobin and liver enzymes levels within the normal range; 5) Not having
96 suffered a myocardial infarction/stroke in the past 12 months; 6) Not suffering from renal or bowel
97 disease or have a history of choleostatic liver or pancreatitis; 7) Not on drug treatment for hyperlipidemia,
98 hypertension, inflammation or hyper-coagulation; 8) Not taking any fish oil, fatty acid or vitamin and
99 mineral supplements; 9) No history of alcohol misuse; 10) Not planning or on a weight reducing regime;
100 11) Not having taken antibiotics in the 6 months prior to the study; 12) Not being able to consume the
101 study meals. Those selected for the study were instructed not to alter their usual dietary or fluid intake.
102 Volunteers were asked for 24 h prior to, and during, the study to refrain from the following: 1)
103 consumption of polyphenol-rich foods including fruits (including citrus fruits) vegetables, cocoa,
104 chocolate, coffee, tea, fruit juices and wine; 2) consumption of foods-rich in nitrates, including beetroot,
105 spinach, lettuce, rocket, celery, parsley, cabbage (defined as containing more than 50 mg nitrates/ 100 g
106 fresh weight ⁽⁴⁷⁾; 3) participating in vigorous exercise and 4) consuming any of alcohol beverage.
107 Volunteers were further asked to fast for 12 hours before each study visit and during that period only
108 consume low-nitrate water provided. The same standard meal, low in polyphenols and nitrates, was also
109 provided for dinner for the day before each visit. Written informed consent was obtained from all eligible
110 volunteers prior to their participation in the study.

111

112 **Study Design**

113 The study design consisted of an acute, randomized, placebo controlled, double-masked postprandial
114 crossover study (Figure 1). After the initial screening visit to assess the eligibility of volunteers to
115 participate in the study, volunteers were enrolled in the study (by researchers CR and HD) and visited

116 the Hugh Sinclair Unit at the University of Reading on four separate occasions separated by a two week
117 period (June-Dec 2012). Volunteers were asked to consume either a A) Control drink (C); B) Orange
118 juice beverage (OJ); C) Flavanone-rich orange juice (FROJ) or D) Whole Orange beverage (WO),
119 together with a high fat breakfast (at baseline, $t = 0$ h), followed by a medium-fat lunch ($t = 5.5$ h). HD
120 assigned participants to the 3 digit coded drink interventions for their 4 visits according to a random
121 allocation sequence generated by a third party. Details on the flavonoid composition of the interventions,
122 as well as micro and macronutrient composition, can be found on Table 1. Compliance to a 24 hour low-
123 polyphenol intake period and 12 hour fasting was monitored by a 24 hour dietary recall conducted in
124 each study visit. On each visit day, volunteers rested for 30 minutes in a quiet, temperature controlled
125 room before they were cannulated by a qualified research nurse and blood samples were collected in the
126 fasted state (0 h) and at 2, 5, 7 and 24 h after consumption of each intervention drink. Flow mediated-
127 dilation of the brachial artery (FMD) was the primary outcome and it was measured at 0, 2, 5 and 7 h
128 post consumption. Secondary outcomes of the study included systolic and diastolic blood pressure (0, 2,
129 5, 7 h), plasma flavanone levels (0, 2, 5, 7, 24 h) and Nitric Oxide (NO) plasma levels (0, 2, 5, 7 h). After
130 baseline measurements were taken, the high fat breakfast (Table 2) was consumed with one of the clinical
131 products (C, OJ, FROJ, WO). Volunteers were asked to consume the intervention drink and the high fat
132 meal in 10-15 min. At 5.5 hours from baseline, a medium fat lunch was provided (Table 2). Last
133 measurement of the day was performed at 7 hours and the volunteers were asked to return to the clinical
134 unit the following morning to provide a 24 h blood sample (fasted). From 7 to 24 h, volunteers were
135 asked to consume the free-polyphenol dinner provided by the research team and to continue on the low-
136 polyphenol diet, as well as refrain from exercise and consuming alcohol. Blood samples for flavonoid
137 analysis were collected in EDTA-containing tubes (Greiner Bio-One Ltd, Stonehouse, UK), immediately
138 centrifuged for 15 min at 4 °C (4000 x g) and the plasma spiked formic acid (1.5% of a 50% water
139 solution) and ascorbic acid (5% of a 10 mM solution) and stored at -80 °C. Blood samples for Nitric

140 Oxide analysis were collected in heparin-containing tubes, immediately (within 3 min of collection)
141 centrifuged for 15 min at 4 °C (4000 x g) and plasma rapidly collected, aliquoted and stored at -80 °C to
142 reduce inactivation of nitroso species. All procedures involving human volunteers were approved by the
143 University of Reading Research Ethics Committee. The clinical trial was registered at clinicaltrials.gov
144 as NCT01963416.

145 **Sequential double-meal**

146 The sequential double meal protocol was based on the department's extensive experience on postprandial
147 studies, which have been collated into the DISRUPT database ⁽⁴⁸⁾. It consisted of two meals 1) high fat
148 breakfast (51 g fat; 14 g protein; 64 g carbohydrates; 777 kcal) administered with the intervention drink
149 and 2) medium-fat lunch (30 g fat; 15 g protein; 80 g carbohydrates; 628 kcal) (Table 2) administered
150 5.5 hours after the intervention drink. The high fat meal consisted of two butter croissants (47 g of fat)
151 and 5 g of butter (4 g of fat). The medium fat meal consisted of 2 slices of white bread (2 g of fat); 42 g
152 of Philadelphia soft cheese (13 g of fat); a small bag of salted crisps (9 g of fat) and two shortbread
153 biscuits (6 g of fat) (Table 2). The volunteers were asked to consume meals within 10-15 min.

154

155 **Flavanone-containing interventions**

156 The preparations of the intervention drinks were carried out in accordance to good manufacturing
157 practice as described in HACCP. The control (C) drink was matched for sugars found in the orange
158 beverages and 0.67% citric acid and orange flavoring was added for flavor purposes. The levels of total
159 β -carotenes present in the flavanone-treatments are considered negligible (~ 0.25 mg; 2-RSD: 15%) with
160 regards to endothelial function effects; with a 15 mg dose (6 weeks intervention, in combination with
161 150 mg of vitamin C) resulting in no significant changes on endothelial biomarkers ⁽⁴⁹⁾. (Table 1). The
162 levels of folate present (~ 60 μ g; 2-RSD: 16%) can also be considered insignificant in regards to its

163 potential to impact on endothelial function; folate has been shown to drive small improvements in
164 endothelial function only in long-term interventions (1- 4 months) of at least 5000 – 10000 μg /day of
165 folate, but not with lower doses in the ranges of 400 – 800 μg /day⁽⁵⁰⁾.

166 Orange juice intervention (OJ) was a 100% commercial pure orange juice (Tropicana Pure Premium).
167 The flavanone-rich orange juice intervention (FROJ) was Tropicana Pure Premium with added orange
168 pomace. Pomace comprised the edible part of a whole orange which is leftover during the production of
169 Tropicana pure premium orange juice and subjected to particle size reduction. Orange pomace is rich in
170 fiber (40:60 ratio of soluble to insoluble) and contains small amounts of micronutrients and a high
171 proportion of the polyphenols found in whole orange. The whole orange intervention (WO) consisted of
172 lightly blended whole table orange, without the peel. Drinks displayed slightly different viscosities, but
173 specific measurements were no undertaken to assess this. All drinks were stored in individual portions
174 (255 g/ 240 ml) in aluminum canisters, frozen at -20 °C and labeled with a 3-digit code to ensure double-
175 masking. Drinks were defrosted overnight in the fridge (4 °C) just before being used for each study day.
176 Participants, care-providers and all researchers assessing outcomes were blind until all the data was
177 analyzed. Quantification of flavanones from orange beverages (OJ, FROJ and WO) was performed by
178 UHPLC-MS. Sample preparation was performed by diluting the juice sample with DMSO, the internal
179 standard (IS) solution (10 μg /mL d4-Dimethylphthalate in 50% Acetonitrile/water) and 50%
180 Acetonitrile/water, followed by vortexing and centrifugation (10 min, 2500 rpm). The supernatant was
181 filtered prior to analysis in an Agilent 1290 UHPLC, with a Zorbax Eclipse Plus C18 column (1.8 μm ,
182 2.1 mm x 100 mm; linear gradient starting at 100% (A) containing 2% Acetonitrile in water with 0.1%
183 Formic acid, to 90% (B) containing Acetonitrile with 0.1% Formic acid, followed by 100% B). MS
184 detection was performed in ESI positive ion mode, on an Agilent 6530A Q-ToF MS with MassHunter
185 Software for instrument control and data processing. Calibration standards were prepared from analytical
186 grade materials purchased from Indofine Chemical Chromadex or LKT Laboratories. The levels of

187 flavanones in the test products are presented in Table 1. Briefly, the total levels of flavanones in a) OJ is
188 128.88 mg, b) FROJ is 272.14 mg, c) WO is 452.80 mg (Table 1). The flavanone hesperedin is the main
189 flavonoid present in the intervention beverages, ranging from 107.30 mg (OJ) to 352.80 mg (WO).

190

191 **Flow-mediated dilation (FMD)**

192 FMD of the brachial artery was the primary end point measure of the study and measurements were taken
193 following standard guidelines ⁽⁵¹⁾ using an ALT Ultrasound HDI5000 system (ATL Ultrasound, UK) in
194 combination with a semi-automated computerized analysis system (Brachial Analyzer, Medical Imaging
195 Applications-llc, IL, US). Briefly, after 15 minutes supine rest in a quiet air-conditioned room the
196 brachial artery was imaged longitudinally at 2-10 cm proximal to the antecubital fossa. After baseline
197 images were recorded for 60 seconds, a blood pressure cuff placed around the forearm was inflated to
198 220 mmHg. After 5 min of occlusion, the pressure was rapidly released to allow reactive hyperemia,
199 with image collection continuing for 5 min post release. A single researcher, who was blinded to the
200 measurement details, analyzed all image files and peak diameter was defined as the largest diameter
201 obtained after the occlusion was released. FMD response was calculated as relative diastolic diameter
202 change from baseline as compared to peak diastolic diameter. A total of 28 volunteers were analyzed for
203 their FMD response. Data from 8 volunteers was not analyzed or was excluded due to *i*) measurement of
204 FMD from non-dominant arm (rather than dominant) due to limitations with blood collection (n=2); *ii*)
205 absence of FMD response (n=3); *iii*) technical problems during recording of ultrasound FMD
206 measurements rendered non-analyzable data (n=3).

207

208 **Blood pressure**

209 Systolic and diastolic blood pressure were measured using an Omron MX2 automatic digital upper arm
210 blood pressure monitor (Omron Healthcare UK Ltd, Milton Keynes, UK). All measurements were taken
211 according to standard practice and by a qualified research nurse, prior to and following each intervention
212 period. Before starting blood pressure measurements the volunteers were seated or laying down quietly
213 for at least 20 min. Measurements were taken in the right arm, before FMD procedure for each time
214 point. The subject's right arm was placed resting on a pillow (on a side table positioned at heart level),
215 slightly flexed with palm upward. Volunteers were asked to refrain from speaking during blood pressure
216 measurements. The measurement were repeated 3 times and blood pressure was considered as the
217 average of these measurements.

218

219 **Plasma flavanone analysis**

220 Blood samples were collected in EDTA blood tubes and centrifuge at 4 °C for 10-15 min at 4000g.
221 Formic acid (1.5% of a 50% solution) and ascorbic acid (5% of a 10 mM solution prepared fresh
222 everyday) were added to the plasma samples to preserve flavanones before freezing at -80 °C. A subset
223 of 20 volunteers were selected randomly for analysis of their flavanone content. A high throughput
224 analytical method using Ultra-high Performance Liquid Chromatography coupled with tandem mass
225 spectrometry (UHPLC-MS/MS) was developed and validated to measure simultaneously naringenin and
226 hesperetin in human plasma. Enzymatic hydrolysis and methanol extraction was applied as described
227 before ⁽⁵²⁾ with modifications to accommodate *in situ* monitoring of enzyme efficiency and automated
228 sample preparation using a Hamilton Microlab Star liquid handling system (Hamilton, UK). Plasma
229 samples (45 µL) were incubated after addition of β-glucuronidase type VII-A (Sigma, USA) and sulfatase
230 type H-5 (Sigma, USA) for 90 min and 60 min at 37 °C, respectively. To monitor the enzyme activity,
231 every individual sample was spiked with a known concentration of phenolphthalein glucuronide and

232 potassium 4-nitrophenyl sulfate (Sigma, USA) as enzyme substrates in addition to caffeine-(trimethyl-
233 d9) (Sigma, USA) as internal standard (IS) prior to incubation. The enzyme hydrolyzed samples were
234 subsequently extracted with methanol and centrifuged. The supernatant (6 μ L) was analyzed using an
235 Agilent 1290 UHPLC coupled with an Agilent 6490 triple quadrupole mass spectrometer. Naringenin
236 and hesperetin were separated in a Waters BEH C18 (100 x 2.1 mm, 1.7 micrometre particle size) at a
237 flow rate of 0.6 mL/min in a 6.5 min gradient 99% solvent A (water containing 0.1% formic acid) and
238 1% solvent B (acetonitrile containing 0.1% formic acid) initially; 70% solvent A at 0.5 minutes; 55%
239 solvent A at 2.5 min; 2% solvent A at 3.0 min; 2% solvent A at 4.0 min; 99% solvent A at 4.5 min
240 followed by post equilibration for 2 minutes). The mass spectrometer was operated in ESI positive
241 ionization mode and Multiple Reaction Monitoring (MRM) mode by monitoring quantifier and qualifier
242 ions for both naringenin and hesperetin. MRM transitions were determined as 204.1/144.0 (m/z)
243 corresponding to caffeine, 495.1/319.1 (m/z) corresponding to phenolphthalein-glucuronide and
244 217.9/137.9 (m/z) corresponding to potassium 4-nitrophenyl sulfate, as Quantifier ions. MRM transitions
245 were determined as 303.1/153.1 (m/z) corresponding to hesperetin and 273.1/147.1 (m/z) corresponding
246 to naringenin as qualifier ions. Concentrations of hesperetin and naringenin were then calculated based
247 on ratios of their integrated peak area for the quantifier ions to that of IS using two sets of eight point
248 calibration curves. Accuracy of the analysis was monitored by systematic counter-balancing between
249 plasma samples and quality control samples spiked with a known concentration of hesperetin and
250 naringenin. The method was validated for a linear calibration range of 0.0313 μ M to 8.02 μ M for
251 naringenin and 0.0282 μ M to 7.22 μ M for hesperetin, respectively. Additionally, limits of detection for
252 naringenin and hesperetin were determined as 2 nM and 7 nM, respectively.

253

254 **Biochemical analysis**

255 The blood samples collected in pre-chilled lithium or heparin tubes were spun (4000 x g; 10-15 min;
256 4°C) immediately after collection (within 3 min). Samples were also collected in serum separation tubes
257 (SST) and allowed to stand for 30 min prior to centrifugation (1300 x g; 10 min; 21°C). All samples
258 were aliquoted and frozen at -80°C until analysis. **Plasma NO analysis:** Plasma samples for
259 measurement of total nitroso species (NOx) were aliquoted in 150 µl aliquots to avoid freeze-thawing of
260 the samples for each measurement. Samples were defrosted just before the measurements took place
261 (within 10 min), these were kept on ice throughout. Plasma samples (n=28) were analysed for nitrite,
262 nitrate and other nitroso species (RXNO, including nitrosothiols, nitrosamines, iron-nitrosylhemoglobin
263 and nitrosohemoglobin) by ozone-based chemiluminescence (model 88 AM, Eco Physics) as previously
264 described ⁽⁵³⁾. In brief, for total NOx measurement (NO derived from nitrate, nitrite and RXNO), one
265 aliquot of plasma was injected in airtight microreaction vessel containing a solution of vanadium (III)
266 chloride (50 mM) dissolved in 1 M HCl, connected to a chemiluminescence analyser. For measurement
267 of nitrite and other RXNO, *i*) one aliquot of plasma was injected in the same apparatus into a glacial acid
268 acetic solution containing 45 mM of potassium iodide and 10 mM of iodide, at 60 °C actively purged by
269 inert helium, which allowed the detection of NO from both nitrite and RXNO (but no nitrate). *ii*)
270 Subsequently, the plasma sample was treated with acidic sulphanilamide (1 M HCL) to scavenge nitrite,
271 before injection, allowing for quantification of RNXO alone. Nitrite levels in the plasma samples was
272 determined by the difference between these two measurements (*i* and *ii*). Nitrate concentration was
273 determined by subtracting Nitrite + RXNO from total NOx. Samples used for calibration curves were
274 prepared fresh every day and displayed consistent values across days. **Plasma baseline lipids and**
275 **glucose:** Plasma levels of total cholesterol, LDL cholesterol, HDL cholesterol, glucose and
276 triacylglycerol (TAG) were assayed on an ILAB 600 chemistry analyzer (Instrumentation Laboratory,
277 Warrington, UK) using enzyme based colorimetric tests supplied by Instrumentation Laboratory.

279 **Power calculation and statistical analysis**

280 Power calculations were performed for the primary endpoint, change in FMD response. Power was based
281 on the intra-individual variability of the operator that performed the FMD analysis (5% CV, SD=0.3%).
282 Previous measures of variability in a control group estimated the standard deviation within subjects to
283 be 2.3%. At 90% power and 0.05 significance, the number of volunteers required to detect a difference
284 of 1.5% in the response of matched pairs in a crossover study is 25. The statistical analysis was performed
285 using the SPSS Statistics 21 (IBM) package. FMD, blood pressure, plasma levels of Nitric Oxide species
286 (Nitrate, Nitrite and Nitroso species) and plasma levels of flavanones were analyzed using a two-way
287 repeated measures ANOVA within subjects with Time (0, 2, 5, 7 hours) and Treatment (C, OJ, FROJ,
288 WO) as main factors. Post-hoc and Pairwise comparisons were carried out using the Bonferroni
289 correction for multiple comparisons. Significance was defined as $P < 0.05$ (95 % confidence interval)
290 for all outcome measures, with p-values represented in the figures as follows: * $P = 0.01-0.05$, ** $P =$
291 $0.001-0.01$, *** $P < 0.001$. Pharmacokinetics parameters were calculated as follows: a) the maximum
292 plasma concentration (C_{max}) and b) the time to reach the maximum plasma concentration (T_{max}) were
293 determined from the individual data obtained from each participant; c) the area under the plasma
294 concentration versus time curve (AUC) was calculated using the trapezoidal method. Multiple regression
295 analysis was used to predict the value of FMD (dependent variable) based on the value of hesperetin and
296 naringenin plasma levels (independent variables). Random allocation sequence was generated by a third
297 party statistician using SAS version 9.1 (procedure plan and seed = 122700). The randomized block
298 design contained 4 blocks and 9 randomized sequences within each block.

299 **RESULTS**

300 **Baseline characteristics and tolerance of intervention**

301 The baseline characteristics of volunteers recruited were within the desired ranges, with either
302 triacylglycerol ranging from 0.8-3.2 mmol/l and total cholesterol from 6.0-8.0 mmol/l or/and BMI from
303 25-32 kg/m² (Table 3). All intervention beverages were well tolerated by all volunteers, as well as the
304 high and medium fat meals administered throughout the study. No adverse events were reported.

305

306 **Flavanone modulation of postprandial FMD**

307 A 2-factor repeated-measures ANOVA for endothelium-dependent brachial artery vasodilation,
308 measured FMD response, revealed a highly significant interaction between the interventions (C, OJ,
309 AOJ, WO) and time of the day (0, 2, 5, 7 h) [F(9,243)=3.27, P<0.0001], as well as significant main
310 effects of time of the day [F(3,81)=12.062, P<0.0001] and intervention [F(3,81)=2.78, P<0.05]. At
311 baseline (t = 0 h) there were no significant differences in brachial artery FMD between visits with the
312 average baseline levels of FMD for the study population being 4.80 ± 0.03 FMD units. Two hours after
313 intake of the high-fat meal, a significant decrease in % FMD was detected for both control ($P < 0.0001$)
314 and the 3 flavanone interventions ($P < 0.05$) (Figure 2). In particular, in the control group, the % FMD
315 decreased by 0.99 ± 0.17 % FMD after 2 h ($P < 0.0001$) and remained significantly suppressed 5 h ($P <$
316 0.05) and 7 h ($P < 0.0001$) after intake, relative to baseline levels. In contrast, all orange flavanone
317 interventions resulted in a recovery in % FMD to that of baseline levels between 5-7 h (OJ : $4.51 \pm 0.23\%$
318 FMD; FROJ: $4.74 \pm 0.25\%$ FMD and WO: $4.75 \pm 0.23\%$ FMD) (Figure 2). At 5 h post intervention,
319 there were no significant differences in % FMD between control and each of the flavanone interventions
320 (OJ, FROJ, WO), whereas at 7 h (following intake of the medium-fat meal at 5.5 h), we observed a
321 significantly higher % FMD for OJ ($P < 0.05$), FROJ ($P < 0.01$) and WO ($P < 0.01$) in comparison to
322 control. There were no significant differences between the flavanone interventions at 7 h, with all three
323 doses of flavanones administered (OJ: 128.8 mg; FROJ: 272.1 mg and WO: 452.7 mg) counteracting the
324 deleterious effect of the double meal challenge on % FMD response to a similar extent. Blood pressure

325 was not significantly altered following consumption of any of the flavanone interventions, relative to
326 baseline or to the control beverage (Table 4).

327

328 **Modulation of plasma flavanones**

329 Total flavanones, naringenin and hesperetin (including glucuronides and sulfates), were not detected in
330 the plasma of volunteers at baseline, indicating compliance to the 24 h low flavonoid diet prior to the
331 study visits. Flavanone metabolites were not detected in the circulation of individuals following intake
332 of the control drink (Figure 3). Significant increases in plasma levels of hesperetin/metabolites were
333 detected at 5 and 7 h ($P < 0.0001$) (Figure 3A), and at 2, 5 and 7 h for naringenin/metabolites ($P < 0.0001$)
334 (Figure 3B). The time to reach T_{max} for hesperetin and naringenin were not significantly different
335 between treatments and coincided with the timeframe of FMD effects (Table 5). At 2, 5 and 7 h, plasma
336 levels of naringenin were significantly higher following WO intake compared to OJ ($P < 0.01$) (Figure
337 3B). Similarly, at 7 h, both FROJ and WO showed a trend towards higher plasma concentrations of
338 hesperetin, relative to OJ ($P < 0.1$) (Figure 3A). With respect to the C_{max} and AUC (0-24 h) for plasma
339 hesperetin, both FROJ and WO were significantly higher than after OJ ($P < 0.05$), whilst for naringenin,
340 both the C_{max} and AUC were only significantly higher for WO in relation to OJ ($P < 0.005$) (Table 5).
341 No significant differences in plasma flavanone levels were detected between FROJ and WO despite the
342 different levels present in the treatment drinks. At 24 h, the levels of flavanones were not significantly
343 different from baseline, indicating the flavanone metabolites have been cleared from circulation ($P =$
344 0.13) (Figure 3). A multivariate regression analysis, including both plasma naringenin and hesperetin,
345 showed that hesperetin ($P = 0.001$), but not naringenin ($P = 0.092$), predicted changes in % FMD over
346 the course of 0-7 h. Specifically, at 7 h, at the peak of FMD response, hesperetin significantly predicted
347 the magnitude of FMD increase ($r = 0.32$, $P = 0.005$) following flavanone intake.

348

349 **Modulation of plasma nitrite, nitrate and RXNO**

350 Levels of nitrate, nitrite and other RXNO (nitrosothiols, nitrosamines, iron-nitrosylhemoglobin and
351 nitrosohemoglobin) were determined in plasma at baseline and 2, 5 and 7 h post treatment (Figure 4).
352 Nitrite plasma levels are known to reflect more accurately endogenous NO production in humans
353 (estimate of 70-80% of plasma nitrite deriving from endothelial nitric oxide synthase activity), whilst the
354 other major source is diet-derived nitrate (by reduction to nitrite). This was the rationale for detecting
355 separately levels of nitrite, nitrate and other RXNO species. The average levels of plasma nitrate, nitrite
356 and other RXNO at baseline were 32.1 $\mu\text{mol/L}$, 68.3 nmol/L and 0.4 nmol/L , respectively, which is in
357 agreement with the values reported in the literature^(37,54). A significant decrease in plasma nitrate levels
358 was observed at 2, 5 and 7 h for all the interventions, including control ($P < 0.0001$) (Figure 4B). In
359 contrast, plasma nitrite levels remained constant up to 7 h post-treatment (not significantly different from
360 baseline) following OJ, FROJ and WO intake whilst the control group nitrite levels decreased
361 significantly ($P < 0.01$) (Figure 4A). No significant changes were detected in RXNO levels in plasma
362 (NS) (Figure 4C).

363

364 **DISCUSSION**

365 Considerable evidence suggests that dysregulation of endothelial function in the postprandial state is an
366 important contributing factor for cardiovascular disease risk^(2, 5, 7, 8), whilst intake of
367 flavonoid/polyphenol-rich foods, such as cocoa, tea and berries have been shown to exert positive effects
368 on vascular function. In support of this, clinical trial data has indicated that intake of such
369 foods/beverages may lower CVD disease risk, at least partially, through their actions in mitigating fed-
370 state metabolic and vascular disturbances (reviewed in⁽³⁴⁾). In the present study, we showed that
371 intervention with orange flavanones, both in juice or whole orange homogenised form, counteracts

372 impairments in vascular function evoked by a sequential double meal challenge, which reflects a regular
373 eating pattern and a typical dietary intake in the population ⁽⁴⁸⁾. Each flavanone intervention tested was
374 effective in reversing vascular impairments, to a physiologically similar degree, despite them containing
375 different levels of flavanones (ranging from 128 mg to 452 mg) and resulting in different concentrations
376 of plasma flavanone metabolites. No changes in blood pressure were observed. The rescue of transient
377 impairments in vascular function, as assessed using brachial artery FMD, coincided with the peak of
378 flavanone metabolites (total sulphates and glucuronides) in the circulation (7 h) and with sustained levels
379 of plasma Nitrite, the latter of which was significantly reduced by the double meal challenge. Thus, our
380 data support the concept that the observed postprandial vascular benefits may be linked to the actions of
381 circulating flavanone metabolites on NO bioavailability.

382 Our findings are consistent with previous RCT datasets indicating that cocoa flavanols partially
383 counteract the decrease in FMD following high fat meal loading ⁽³²⁾. Furthermore, pure quercetin has
384 also been shown to ameliorate postprandial FMD following maltose overload ⁽³³⁾. To our knowledge this
385 is the first data indicating that citrus flavanones are also capable of attenuating postprandial impairments
386 in endothelial function following a sequential high-medium fat double meal in individuals displaying
387 mild cardio-metabolic risk factors. Furthermore and in support of our findings, previous studies
388 conducted in the fasted state report that chronic interventions with flavanones in at risk groups (e.g.
389 hypertensive, overweight or metabolic syndrome patients) induce positive effects on blood pressure and
390 endothelial function (FMD) ^(45,46,55) and improvements in microvascular reactivity ⁽⁴⁵⁾. Most importantly,
391 in the present study, no dose-dependent effects on brachial artery FMD were observed, despite the
392 interventions containing different amounts of flavanones (WO: 3.5 x OJ). This may indicate that at these
393 intake levels a threshold plateau may be reached, similarly to what has been shown previously with other
394 flavonoid-rich interventions ⁽²²⁾. It also further suggests that lower doses of flavanones (approx. 130 mg)
395 can be efficacious at modulating postprandial endothelial function. No changes in blood pressure were

396 observed in the present study, which is in agreement with previous human intervention trials reporting
397 modulation of blood pressure only after chronic interventions with flavonoid-rich foods, but not in an
398 acute manner ^(27, 31).

399

400 We observed concurrent modulation of FMD, nitrite, and circulating flavanone metabolites (total
401 sulphates and glucuronides) suggesting that the latter may be linked to NO availability and subsequent
402 improvements in vascular function, although we cannot establish a causal relationship at this time.
403 Specifically, both naringenin and hesperetin metabolites peak plasma levels for all three interventions
404 occurs at approximately 7 h, which coincide with significant improvements in endothelial function (at 7
405 h) and sustained levels of circulating nitrite after flavanone interventions in comparison to control.
406 Plasma levels of flavanone metabolites peaked slightly later than previously reported (4 - 6 h) ^(45, 56, 57),
407 most likely due to the concomitant intake of fat, which is thought to interfere with flavonoid absorption
408 ⁽⁵⁸⁾. On the other hand, no significant differences in time of absorption were detected between the
409 flavanone treatment groups (peak occurs at approx. 7 h for all three treatments).

410 In support of the link between flavanone intake and human vascular function, we observe that when low
411 or no levels of flavanone metabolites are detected in circulation (e.g 2 h), no differences in postprandial
412 brachial FMD are observed between control and flavanone-rich beverages. Further multiple regression
413 analysis suggests that mainly hesperetin metabolites seem to predict significantly the magnitude of
414 changes in FMD ($r = 0.32$, $P = 0.005$), suggesting an important role of this flavanone in the effects
415 observed. This is corroborated by previous studies showing that pure hesperidin can trigger both acute
416 and chronic improvements in vascular function in humans ^(45, 46). It is important to further note that only
417 the sulphated and glucuronidated portion of the flavanone metabolites were quantified in our study and
418 these are likely to account for a fraction (approx. 16%) of the total flavanone metabolites absorbed ⁽⁵⁹⁾.
419 As such, we anticipate that gut-derived phenolic compounds might also contribute to the improvements

420 in endothelial function observed. This is supported by our observation that hesperetin metabolites can
421 only significantly predict a small percentage (approx. 30%) of the FMD response observed; therefore it
422 is likely that stronger correlations might be apparent once gut derived small phenolic metabolites are
423 taken into consideration, however such extensive analysis was outside of the scope of our study.

424

425 Our study also indicates that the impairment in postprandial FMD induced by the sequential high fat
426 meal might be linked to decreases in circulating levels of NO species, in particular nitrite and nitrate.
427 Although, the precise mechanisms underlying postprandial endothelium impairments are not established,
428 mechanistic animal studies suggest a role for NO signalling, showing, for example, that endothelial
429 dysfunction induced by fat intake also results in decreases in NO production ⁽⁶⁰⁻⁶²⁾. Importantly, the
430 flavanone interventions only prevented the decrease in nitrite, but not nitrate. Numerous evidence suggest
431 that nitrite reflects more accurately endogenous NO production in humans, with an estimate of 70-80%
432 of plasma nitrite deriving from endothelial nitric oxide synthase (eNOS) activity ^(63, 64) and also better
433 reflects the degree of endothelial dysfunction in humans ⁽⁶⁵⁾. In agreement with our data, previous human
434 clinical data suggests an ability of some flavonoid-rich foods to modulate NO bioavailability ^(22, 24, 35, 37).
435 In particular, cocoa flavanols induced improvements in FMD have been causally linked to NO production
436 in humans ⁽²³⁾. More recently Bondonno *et al.*, 2012, also showed that apples containing (-) epicatechin
437 and quercetin increased levels of nitrite along with FMD response after 2 h of intake ⁽³⁷⁾. Additionally
438 and in agreement with the present data, both pure (-) epicatechin and quercetin were shown to specifically
439 increase plasma nitrite, but not nitrate in healthy humans ⁽³⁵⁾. Supporting *in vitro* mechanistic studies (in
440 endothelial cells) have demonstrated the flavanone hesperetin and some of its *in vivo* metabolites (e.g.
441 7-O- β -D glucuronide) can stimulate NO production *via* activation/expression of eNOS ^(46, 66) or by
442 decreasing NO degradation through inhibition of nicotinamide adenine dinucleotide phosphate-oxidase
443 (NADPH) ⁽⁶⁷⁾ and these are possible mechanistic pathways by which flavanone metabolites might

444 modulate postprandial FMD. Although the specific modulation of nitrite by flavanone-containing
445 interventions is an interesting observation in the present study, the interpretation of the temporal
446 dynamics (time course) of flavanone appearance in plasma and levels of plasma nitrite is not
447 straightforward in regards to explaining the effects of nitrite on FMD. This seems to suggest that the
448 impact of flavanone metabolites on FMD response cannot be explained completely by modulation of
449 nitrite (as a measure of NO). Since, the present study was not designed (or powered) to detect changes
450 in NO species, we are limited in our ability to establish a clear-cut link between FMD modulation and
451 NO at this time. However, we believe this preliminary data is very novel and valuable for future, more
452 mechanism-focused, human RCT.

453

454 Interestingly, FROJ intake resulted in similar levels of plasma flavanones to WO, despite lower initial
455 concentrations, which might be related to characteristics of food matrix itself, such as viscosity, which
456 is known to influence the bioavailability of polyphenols (reviewed in ⁽⁶⁸⁾). It is possible that the reduced
457 particle size of the pomace in FROJ aided the release of polyphenols from the fiber matrix, making these
458 more accessible for gut microbiota metabolism ⁽⁶⁹⁾. It is known that dietary fiber can physically trap
459 polyphenols within the fiber matrix in the fruit tissue reducing the accessibility to enzymes and the gut
460 microbiota ⁽⁷⁰⁾. On the other hand, the rate of release of polyphenols from fibrous particles is inversely
461 proportional to the fiber particle size ⁽⁷¹⁾, therefore by reducing particle size in the pomace, we are likely
462 to increase the bioavailability of flavanones in FROJ. In order to confirm that this is the case, future
463 studies will focus on measuring accurately total urine excretion (e.g. over a 24 h period). Nonetheless,
464 our study seems to suggest that particle size reduction of fiber-rich orange pomace and the re-introduction
465 of this product into orange juice might be an effective strategy to increase the bioavailability of
466 polyphenols *in vivo*. Importantly, the increased bioavailability of flavanones after FROJ intake did not
467 enhance significantly postprandial FMD in comparison to lower flavanone-containing OJ, again

468 suggesting that perhaps a certain level of flavanone metabolites is necessary in circulation to trigger
469 postprandial FMD improvements but further increases in flavanone levels may not produce additional
470 benefits ⁽²²⁾.

471

472 One of the limitations in the design of the present study is related to the composition of the control
473 intervention, which did not take into account the levels of ascorbic acid present in the citrus beverages.
474 Clinical studies suggest that doses up to 500 mg of vitamin C do not impact on biomarkers of endothelial
475 function (e.g. ^(49, 72)). More specifically, it has been shown in a recent stratified meta-analysis that doses
476 ranging from 90 to 500 mg of ascorbic acid do not produce improvements in endothelial function, both
477 acutely or chronically ^(73, 74), especially in populations with normal vitamin C status ^(75, 76). Furthermore,
478 previous studies reporting acute beneficial effects of ascorbic acid on endothelium dependent
479 vasodilation (within 2- 4 h of intake), deliver doses of at least 2000 mg (e.g. ⁽⁷⁷⁻⁷⁹⁾) and in many cases
480 positive outcomes are achieved by delivering ascorbic acid intravenously, resulting in supra-
481 physiological plasma levels of vitamin C, which cannot be achieved by oral ingestion ^(79, 80). Since, the
482 levels of vitamin C in the present study were approx. 80-120 mg, we are confident that these can be
483 considered negligible with respect to acute effects on endothelial function, as measured by FMD.
484 Therefore, despite these limitations, we can safely argue that our conclusions are reasonable when
485 attributing the FMD response to circulating flavanone metabolites (at 7 h post intake) and that our data
486 are relevant in furthering the understanding of flavonoid-rich foods/beverages impact on postprandial
487 endothelial function.

488

489 In summary, our results suggest that acute intake of a beverage containing at least 128 mg of flavanones
490 can be an effective dietary strategy to blunt the acute transient impairment in endothelial function induced
491 by a sequential double meal that reflects a typical intake in the population. Although we cannot draw

492 firm conclusions regarding the mechanisms by which flavanones elicit vascular responses, our results
493 suggest that these might be linked to an ability of flavanone metabolites to sustain basal circulating NO
494 levels. Collectively these observations have important implications considering that most individuals
495 spend the majority of the day in the postprandial state and such temporary vascular changes repeated on
496 a daily basis can critically impact on long-term vascular health and overall chronic disease risk.

497

498 **DISCLOSURES**

499 CS works as a Senior Scientist at PepsiCo Inc, LH works as a Senior Director at Global R&D Nutrition
500 at PepsiCo Inc, RLB, MB and YH work as Principle Scientists at PepsiCo Inc. The other authors declare
501 no conflicts of interest.

502 The views expressed in this manuscript are those of the authors and do not necessarily reflect the position
503 or policy of PepsiCo Inc.

504 **ACKNOWLEDGMENTS**

505 This work was supported by PepsiCo Inc. and is greatly appreciated.

506 **AUTHORS' CONTRIBUTIONS**

507 CR: coordinated and conducted the study, undertook all FMD measurements, did data analysis and wrote
508 the manuscript; HD: coordinated and conducted the study; CS: design and coordination of the study,
509 writing of the manuscript; LH: design of the study; RLB: conducted study drinks analysis MB, YH:
510 conducted plasma flavanone analysis; GC: conducted NO species measurements and analysis; JL: co-
511 investigator in the study, involved in experimental design; JPES: principal investigator, involved in
512 experimental design and writing of the manuscript. All authors reviewed the manuscript.

REFERENCES

1. Ceriello A, Taboga C, Tonutti L, Quagliario L, Piconi L, Bais B, et al. Evidence for an independent and cumulative effect of postprandial hypertriglyceridemia and hyperglycemia on endothelial dysfunction and oxidative stress generation: effects of short- and long-term simvastatin treatment. *Circulation*. 2002;106(10):1211-8. Epub 2002/09/05.
2. de Koning EJ, Rabelink TJ. Endothelial function in the post-prandial state. *Atherosclerosis*. 2002;Suppl. 3:11-6.
3. Nappo F, Esposito K, Cioffi M, Giugliano G, Molinari AM, Paolisso G, et al. Postprandial endothelial activation in healthy subjects and in type 2 diabetic patients: role of fat and carbohydrate meals. *J Am Coll Cardiol*. 2002;39(7):1145-50. Epub 2002/03/30.
4. Alipour A, Elte JW, van Zaanen HC, Rietveld AP, Cabezas MC. Postprandial inflammation and endothelial dysfunction. *Biochemical Society transactions*. 2007;35(Pt 3):466-9. Epub 2007/05/22.
5. Ebenbichler CF, Kirchmair R, Egger C, Patsch JR. Postprandial state and atherosclerosis. *Current opinion in lipidology*. 1995;6(5):286-90. Epub 1995/10/01.
6. Karpe F. Postprandial lipid metabolism in relation to coronary heart disease. *The Proceedings of the Nutrition Society*. 1997;56(2):671-8. Epub 1997/07/01.
7. Ansar S, Koska J, Reaven PD. Postprandial hyperlipidemia, endothelial dysfunction and cardiovascular risk: focus on incretins. *Cardiovascular diabetology*. 2011;10:61. Epub 2011/07/09.
8. Ceriello A. The post-prandial state and cardiovascular disease: relevance to diabetes mellitus. *Diabetes/metabolism research and reviews*. 2000;16(2):125-32. Epub 2000/04/07.
9. Newens KJ, Thompson AK, Jackson KG, Wright J, Williams CM. DHA-rich fish oil reverses the detrimental effects of saturated fatty acids on postprandial vascular reactivity. *The American journal of clinical nutrition*. 2011;94(3):742-8. Epub 2011/08/13.
10. Jackson KG, Armah CK, Minihane AM. Meal fatty acids and postprandial vascular reactivity. *Biochemical Society transactions*. 2007;35(Pt 3):451-3. Epub 2007/05/22.

11. Vogel RA, Corretti MC, Plotnick GD. Effect of a single high-fat meal on endothelial function in healthy subjects. *The American journal of cardiology*. 1997;79(3):350-4. Epub 1997/02/01.
12. Keogh JB, Grieger JA, Noakes M, Clifton PM. Flow-mediated dilatation is impaired by a high-saturated fat diet but not by a high-carbohydrate diet. *Arteriosclerosis, thrombosis, and vascular biology*. 2005;25(6):1274-9. Epub 2005/03/19.
13. Steer P, Sarabi DM, Karlstrom B, Basu S, Berne C, Vessby B, et al. The effect of a mixed meal on endothelium-dependent vasodilation is dependent on fat content in healthy humans. *Clin Sci (Lond)*. 2003;105(1):81-7. Epub 2003/03/11.
14. Steinberg HO, Tarshoby M, Monestel R, Hook G, Cronin J, Johnson A, et al. Elevated circulating free fatty acid levels impair endothelium-dependent vasodilation. *The Journal of clinical investigation*. 1997;100(5):1230-9. Epub 1997/09/01.
15. Bae JH, Bassenge E, Kim KB, Kim YN, Kim KS, Lee HJ, et al. Postprandial hypertriglyceridemia impairs endothelial function by enhanced oxidant stress. *Atherosclerosis*. 2001;155(2):517-23. Epub 2001/03/20.
16. Cassidy A, Mukamal KJ, Liu L, Franz M, Eliassen AH, Rimm EB. High anthocyanin intake is associated with a reduced risk of myocardial infarction in young and middle-aged women. *Circulation*. 2013;127(2):188-96. Epub 2013/01/16.
17. Mink PJ, Scrafford CG, Barraj LM, Harnack L, Hong CP, Nettleton JA, et al. Flavonoid intake and cardiovascular disease mortality: a prospective study in postmenopausal women. *The American journal of clinical nutrition*. 2007;85(3):895-909. Epub 2007/03/09.
18. McCullough ML, Peterson JJ, Patel R, Jacques PF, Shah R, Dwyer JT. Flavonoid intake and cardiovascular disease mortality in a prospective cohort of US adults. *The American journal of clinical nutrition*. 2012;95(2):454-64.
19. Zamora-Ros R, Jimenez C, Cleries R, Agudo A, Sanchez MJ, Sanchez-Cantalejo E, et al. Dietary flavonoid and lignan intake and mortality in a Spanish cohort. *Epidemiology*. 2013;24(5):726-33. Epub 2013/07/25.
20. Wang X, Ouyang YY, Liu J, Zhao G. Flavonoid intake and risk of CVD: a systematic review and meta-analysis of prospective cohort studies. *The British journal of nutrition*. 2014;111(1):1-11. Epub 2013/08/21.

21. Rodriguez-Mateos A, Del Pino-Garcia R, George TW, Vidal-Diez A, Heiss C, Spencer JP. Impact of processing on the bioavailability and vascular effects of blueberry (poly)phenols. *Molecular nutrition & food research*. 2014;58(10):1952-61. Epub 2014/07/22.
22. Rodriguez-Mateos A, Rendeiro C, Bergillos-Meca T, Tabatabaee S, George TW, Heiss C, et al. Intake and time dependence of blueberry flavonoid-induced improvements in vascular function: a randomized, controlled, double-blind, crossover intervention study with mechanistic insights into biological activity. *The American journal of clinical nutrition*. 2013;98(5):1179-91. Epub 2013/09/06.
23. Heiss C, Kleinbongard P, Dejam A, Perre S, Schroeter H, Sies H, et al. Acute consumption of flavanol-rich cocoa and the reversal of endothelial dysfunction in smokers. *J Am Coll Cardiol*. 2005;46(7):1276-83. Epub 2005/10/04.
24. Schroeter H, Heiss C, Balzer J, Kleinbongard P, Keen CL, Hollenberg NK, et al. (-)-Epicatechin mediates beneficial effects of flavanol-rich cocoa on vascular function in humans. *Proceedings of the National Academy of Sciences of the United States of America*. 2006;103(4):1024-9.
25. Fisher ND, Hughes M, Gerhard-Herman M, Hollenberg NK. Flavanol-rich cocoa induces nitric-oxide-dependent vasodilation in healthy humans. *Journal of hypertension*. 2003;21(12):2281-6.
26. Heiss C, Finis D, Kleinbongard P, Hoffmann A, Rassaf T, Kelm M, et al. Sustained increase in flow-mediated dilation after daily intake of high-flavanol cocoa drink over 1 week. *Journal of cardiovascular pharmacology*. 2007;49(2):74-80.
27. Heiss C, Sansone R, Karimi H, Krabbe M, Schuler D, Rodriguez-Mateos A, et al. Impact of cocoa flavanol intake on age-dependent vascular stiffness in healthy men: a randomized, controlled, double-masked trial. *Age (Dordr)*. 2015;37(3):9794. Epub 2015/05/28.
28. Chanet A, Milenkovic D, Manach C, Mazur A, Morand C. Citrus flavanones: what is their role in cardiovascular protection? *Journal of agricultural and food chemistry*. 2012;60(36):8809-22. Epub 2012/05/12.
29. Hooper L, Kay C, Abdelhamid A, Kroon PA, Cohn JS, Rimm EB, et al. Effects of chocolate, cocoa, and flavan-3-ols on cardiovascular health: a systematic review and meta-analysis of randomized trials. *The American journal of clinical nutrition*. 2012;95(3):740-51. Epub 2012/02/04.
30. Shrime MG, Bauer SR, McDonald AC, Chowdhury NH, Coltart CE, Ding EL. Flavonoid-rich cocoa consumption affects multiple cardiovascular risk factors in a meta-analysis of short-term studies. *The Journal of nutrition*. 2011;141(11):1982-8. Epub 2011/10/01.

31. Sansone R, Rodriguez-Mateos A, Heuel J, Falk D, Schuler D, Wagstaff R, et al. Cocoa flavanol intake improves endothelial function and Framingham Risk Score in healthy men and women: a randomised, controlled, double-masked trial: the Flaviola Health Study. *The British journal of nutrition*. 2015;114(8):1246-55. Epub 2015/09/09.
32. Westphal S, Luley C. Flavanol-rich cocoa ameliorates lipemia-induced endothelial dysfunction. *Heart and vessels*. 2011;26(5):511-5. Epub 2010/12/09.
33. Nakayama H, Tsuge N, Sawada H, Higashi Y. Chronic intake of onion extract containing quercetin improved postprandial endothelial dysfunction in healthy men. *J Am Coll Nutr*. 2013;32(3):160-4. Epub 2013/07/28.
34. Burton-Freeman B. Postprandial metabolic events and fruit-derived phenolics: a review of the science. *The British journal of nutrition*. 2010;104 Suppl 3:S1-14. Epub 2010/10/20.
35. Loke WM, Hodgson JM, Proudfoot JM, McKinley AJ, Puddey IB, Croft KD. Pure dietary flavonoids quercetin and (-)-epicatechin augment nitric oxide products and reduce endothelin-1 acutely in healthy men. *The American journal of clinical nutrition*. 2008;88(4):1018-25. Epub 2008/10/10.
36. Zhu Y, Xia M, Yang Y, Liu F, Li Z, Hao Y, et al. Purified anthocyanin supplementation improves endothelial function via NO-cGMP activation in hypercholesterolemic individuals. *Clinical chemistry*. 2011;57(11):1524-33. Epub 2011/09/20.
37. Bondonno CP, Yang X, Croft KD, Considine MJ, Ward NC, Rich L, et al. Flavonoid-rich apples and nitrate-rich spinach augment nitric oxide status and improve endothelial function in healthy men and women: a randomized controlled trial. *Free radical biology & medicine*. 2012;52(1):95-102. Epub 2011/10/25.
38. Duarte J, Francisco V, Perez-Vizcaino F. Modulation of nitric oxide by flavonoids. *Food & function*. 2014;5(8):1653-68. Epub 2014/06/06.
39. Engler MB, Engler MM, Chen CY, Malloy MJ, Browne A, Chiu EY, et al. Flavonoid-rich dark chocolate improves endothelial function and increases plasma epicatechin concentrations in healthy adults. *J Am Coll Nutr*. 2004;23(3):197-204. Epub 2004/06/11.
40. Monahan KD, Feehan RP, Kunselman AR, Preston AG, Miller DL, Lott ME. Dose-dependent increases in flow-mediated dilation following acute cocoa ingestion in healthy older adults. *J Appl Physiol*. 2011;111(6):1568-74. Epub 2011/09/10.
41. Heiss C, Jahn S, Taylor M, Real WM, Angeli FS, Wong ML, et al. Improvement of endothelial function with dietary flavanols is associated with mobilization of circulating angiogenic cells in patients

with coronary artery disease. *Journal of the American College of Cardiology*. 2010;56(3):218-24. Epub 2010/07/14.

42. Heiss C, Dejam A, Kleinbongard P, Schewe T, Sies H, Kelm M. Vascular effects of cocoa rich in flavan-3-ols. *Jama*. 2003;290(8):1030-1. Epub 2003/08/28.

43. Balzer J, Rassaf T, Heiss C, Kleinbongard P, Lauer T, Merx M, et al. Sustained benefits in vascular function through flavanol-containing cocoa in medicated diabetic patients a double-masked, randomized, controlled trial. *J Am Coll Cardiol*. 2008;51(22):2141-9. Epub 2008/05/31.

44. Del Rio D, Rodriguez-Mateos A, Spencer JP, Tognolini M, Borges G, Crozier A. Dietary (poly)phenolics in human health: structures, bioavailability, and evidence of protective effects against chronic diseases. *Antioxidants & redox signaling*. 2013;18(14):1818-92.

45. Morand C, Dubray C, Milenkovic D, Lioger D, Martin JF, Scalbert A, et al. Hesperidin contributes to the vascular protective effects of orange juice: a randomized crossover study in healthy volunteers. *The American journal of clinical nutrition*. 2011;93(1):73-80. Epub 2010/11/12.

46. Rizza S, Muniyappa R, Iantorno M, Kim JA, Chen H, Pullikotil P, et al. Citrus polyphenol hesperidin stimulates production of nitric oxide in endothelial cells while improving endothelial function and reducing inflammatory markers in patients with metabolic syndrome. *The Journal of clinical endocrinology and metabolism*. 2011;96(5):E782-92. Epub 2011/02/25.

47. Santamaria P. Nitrate in vegetables: toxicity, content, intake and EC regulation. *J Sci Food Agric* 2006;86:10-7.

48. Jackson KG, Clarke DT, Murray P, Lovegrove JA, O'Malley B, Minihane AM, et al. Introduction to the DISRUPT postprandial database: subjects, studies and methodologies. *Genes & nutrition*. 2010;5(1):39-48. Epub 2009/09/30.

49. Seljeflot I, Arnesen H, Brude IR, Nenseter MS, Drevon CA, Hjerermann I. Effects of omega-3 fatty acids and/or antioxidants on endothelial cell markers. *European journal of clinical investigation*. 1998;28(8):629-35. Epub 1998/10/10.

50. McRae MP. High-dose folic acid supplementation effects on endothelial function and blood pressure in hypertensive patients: a meta-analysis of randomized controlled clinical trials. *Journal of chiropractic medicine*. 2009;8(1):15-24. Epub 2009/08/04.

51. Corretti MC, Anderson TJ, Benjamin EJ, Celermajer D, Charbonneau F, Creager MA, et al. Guidelines for the ultrasound assessment of endothelial-dependent flow-mediated vasodilation of the

brachial artery: a report of the International Brachial Artery Reactivity Task Force. *Journal of the American College of Cardiology*. 2002;39(2):257-65. Epub 2002/01/15.

52. Matsumoto H, Ikoma Y, Sugiura M, Yano M, Hasegawa Y. Identification and quantification of the conjugated metabolites derived from orally administered hesperidin in rat plasma. *Journal of agricultural and food chemistry*. 2004;52(21):6653-9. Epub 2004/10/14.

53. Ignarro LJ, Fukuto JM, Griscavage JM, Rogers NE, Byrns RE. Oxidation of nitric oxide in aqueous solution to nitrite but not nitrate: comparison with enzymatically formed nitric oxide from L-arginine. *Proceedings of the National Academy of Sciences of the United States of America*. 1993;90(17):8103-7. Epub 1993/09/01.

54. Rassaf T, Heiss C, Hendgen-Cotta U, Balzer J, Matern S, Kleinbongard P, et al. Plasma nitrite reserve and endothelial function in the human forearm circulation. *Free radical biology & medicine*. 2006;41(2):295-301. Epub 2006/07/04.

55. Reshef N, Hayari Y, Goren C, Boaz M, Madar Z, Knobler H. Antihypertensive effect of sweetie fruit in patients with stage I hypertension. *American journal of hypertension*. 2005;18(10):1360-3. Epub 2005/10/06.

56. Mullen W, Archeveque MA, Edwards CA, Matsumoto H, Crozier A. Bioavailability and metabolism of orange juice flavanones in humans: impact of a full-fat yogurt. *Journal of agricultural and food chemistry*. 2008;56(23):11157-64. Epub 2008/11/15.

57. Erlund I, Meririnne E, Alfthan G, Aro A. Plasma kinetics and urinary excretion of the flavanones naringenin and hesperetin in humans after ingestion of orange juice and grapefruit juice. *The Journal of nutrition*. 2001;131(2):235-41. Epub 2001/02/13.

58. Mullen W, Edwards CA, Serafini M, Crozier A. Bioavailability of pelargonidin-3-O-glucoside and its metabolites in humans following the ingestion of strawberries with and without cream. *Journal of agricultural and food chemistry*. 2008;56(3):713-9. Epub 2008/01/24.

59. Pereira-Caro G, Borges G, van der Hooft J, Clifford MN, Del Rio D, Lean ME, et al. Orange juice (poly)phenols are highly bioavailable in humans. *The American journal of clinical nutrition*. 2014;100(5):1378-84. Epub 2014/10/22.

60. Magne J, Huneau JF, Delemasure S, Rochette L, Tome D, Mariotti F. Whole-body basal nitric oxide production is impaired in postprandial endothelial dysfunction in healthy rats. *Nitric oxide : biology and chemistry / official journal of the Nitric Oxide Society*. 2009;21(1):37-43. Epub 2009/05/07.

61. Erdei N, Toth A, Pasztor ET, Papp Z, Edes I, Koller A, et al. High-fat diet-induced reduction in nitric oxide-dependent arteriolar dilation in rats: role of xanthine oxidase-derived superoxide anion. *American journal of physiology Heart and circulatory physiology*. 2006;291(5):H2107-15. Epub 2006/06/27.
62. Razny U, Kiec-Wilk B, Wator L, Polus A, Dyduch G, Solnica B, et al. Increased nitric oxide availability attenuates high fat diet metabolic alterations and gene expression associated with insulin resistance. *Cardiovascular diabetology*. 2011;10:68. Epub 2011/07/26.
63. Lauer T, Preik M, Rassaf T, Strauer BE, Deussen A, Feelisch M, et al. Plasma nitrite rather than nitrate reflects regional endothelial nitric oxide synthase activity but lacks intrinsic vasodilator action. *Proceedings of the National Academy of Sciences of the United States of America*. 2001;98(22):12814-9. Epub 2001/10/19.
64. Kleinbongard P, Dejam A, Lauer T, Rassaf T, Schindler A, Picker O, et al. Plasma nitrite reflects constitutive nitric oxide synthase activity in mammals. *Free radical biology & medicine*. 2003;35(7):790-6. Epub 2003/10/30.
65. Kleinbongard P, Dejam A, Lauer T, Jax T, Kerber S, Gharini P, et al. Plasma nitrite concentrations reflect the degree of endothelial dysfunction in humans. *Free radical biology & medicine*. 2006;40(2):295-302. Epub 2006/01/18.
66. Liu L, Xu DM, Cheng YY. Distinct effects of naringenin and hesperetin on nitric oxide production from endothelial cells. *Journal of agricultural and food chemistry*. 2008;56(3):824-9. Epub 2008/01/17.
67. Takumi H, Nakamura H, Simizu T, Harada R, Kometani T, Nadamoto T, et al. Bioavailability of orally administered water-dispersible hesperetin and its effect on peripheral vasodilatation in human subjects: implication of endothelial functions of plasma conjugated metabolites. *Food & function*. 2012;3(4):389-98. Epub 2012/02/07.
68. Bohn T. Dietary factors affecting polyphenol bioavailability. *Nutrition reviews*. 2014;72(7):429-52. Epub 2014/05/16.
69. Palafox-Carlos H, Ayala-Zavala JF, Gonzalez-Aguilar GA. The role of dietary fiber in the bioaccessibility and bioavailability of fruit and vegetable antioxidants. *Journal of food science*. 2011;76(1):R6-R15. Epub 2011/05/04.
70. Montagne L, Pluske JR, Hampson DJ. A review of interactions between dietary fibre and the intestinal mucosa, and their consequences on digestive health in young non-ruminant animals. *Animal Feed Science and Technology*. 2003;108(1-4):95-117.

71. Cummings JH, Edmond LM, Magee EA. Dietary carbohydrates and health: do we still need the concept of fibre? *Clin Nutr Suppl.* 2004;1(2):5-17.
72. Ward NC, Hodgson JM, Croft KD, Burke V, Beilin LJ, Puddey IB. The combination of vitamin C and grape-seed polyphenols increases blood pressure: a randomized, double-blind, placebo-controlled trial. *Journal of hypertension.* 2005;23(2):427-34. Epub 2005/01/22.
73. Ashor AW, Lara J, Mathers JC, Siervo M. Effect of vitamin C on endothelial function in health and disease: a systematic review and meta-analysis of randomised controlled trials. *Atherosclerosis.* 2014;235(1):9-20. Epub 2014/05/06.
74. Ashor AW, Siervo M, Lara J, Oggioni C, Afshar S, Mathers JC. Effect of vitamin C and vitamin E supplementation on endothelial function: a systematic review and meta-analysis of randomised controlled trials. *The British journal of nutrition.* 2015;113(8):1182-94. Epub 2015/04/29.
75. Frikke-Schmidt H, Lykkesfeldt J. Role of marginal vitamin C deficiency in atherogenesis: in vivo models and clinical studies. *Basic & clinical pharmacology & toxicology.* 2009;104(6):419-33. Epub 2009/06/06.
76. Harris RA, Nishiyama SK, Wray DW, Tedjasaputra V, Bailey DM, Richardson RS. The effect of oral antioxidants on brachial artery flow-mediated dilation following 5 and 10 min of ischemia. *European journal of applied physiology.* 2009;107(4):445-53. Epub 2009/08/12.
77. Raitakari OT, Adams MR, McCredie RJ, Griffiths KA, Stocker R, Celermajer DS. Oral vitamin C and endothelial function in smokers: short-term improvement, but no sustained beneficial effect. *J Am Coll Cardiol.* 2000;35(6):1616-21. Epub 2000/05/12.
78. Pullin CH, Bonham JR, McDowell IF, Lee PJ, Powers HJ, Wilson JF, et al. Vitamin C therapy ameliorates vascular endothelial dysfunction in treated patients with homocystinuria. *Journal of inherited metabolic disease.* 2002;25(2):107-18. Epub 2002/07/18.
79. Eskurza I, Monahan KD, Robinson JA, Seals DR. Effect of acute and chronic ascorbic acid on flow-mediated dilatation with sedentary and physically active human ageing. *The Journal of physiology.* 2004;556(Pt 1):315-24. Epub 2004/02/03.
80. Taddei S, Virdis A, Ghiadoni L, Magagna A, Salvetti A. Vitamin C improves endothelium-dependent vasodilation by restoring nitric oxide activity in essential hypertension. *Circulation.* 1998;97(22):2222-9. Epub 1998/06/19.

TABLES

Table 1: Compositional analysis of orange flavanone beverages and control beverage used in the acute postprandial study.

Compounds	2-RSD (%)	Intervention drink (240 ml)			
		Control ¹	OJ ²	FROJ ³	WO ⁴
Hesperidin (mg)	5.0	-	107.30	220.46	352.80
Narirutin (mg)	5.0	0.08	15.41	34.54	76.58
Others* (mg)		0.02	6.17	17.14	23.33
Total Flavonoids (mg)		0.10	128.88	272.14	452.71
Fructose (g)	4.0	6.38	6.63	6.12	6.89
Glucose (g)	4.0	5.36	5.36	5.10	5.87
Sucrose (g)	2.0	10.20	10.97	11.99	11.48
Total Sugars (g)		21.93	22.95	23.21	24.23
Fiber (total) (g)	12.0	-	0.66	5.36	6.30
Ascorbic acid (mg)	8.0	-	105.57	80.17	123.01
Folate (µg)	16.0	-	54.06	65.28	64.77
Total β carotenes (mg)	15.0	-	0.13	0.26	0.35

¹ Control, sugar matched control; ² OJ, Tropicana pure premium orange juice without pulp; ³ FROJ, Flavanone-rich orange juice: Tropicana pure premium orange juice with added orange pomace; ⁴ WO, juice made from lightly blended fresh whole orange. * Includes Diosmin, Didymin, Nobiletin, Tangeretin, Sinensetin, Me4-Scutellarein. 2-RSD: Relative standard deviation of the measurement (expressed in %)

Table 2: Macronutrient composition of double-meal protocol

		Macronutrient breakdown			Energy (kcal)
	Foods	Fat (g)	Protein (g)	Carbohydrates (g)	
High-fat breakfast	Butter Croissant (2x)	47	14	64	740
	Butter (5 g)	4	n/a	n/a	37
	Total	51	14	64	777
Medium-fat Lunch	2 Slices of sliced white bread (108 g)	2	8.5	50	237
	Philadelphia soft cheese (42 g)	13	3.6	n/a	131
	Crisps (25 g)	9	1.5	13	133
	Shortbread biscuit (22 g)	6	1.4	16	127
	Total	30	15	80	628

Table 3: Baseline clinical characteristics of the study population.

Baseline characteristics	Mean \pm SEM
Age (y)	48 \pm 1
BMI (kg/m ²)	28.4 \pm 0.4
Total cholesterol (mmol/L)	5.6 \pm 0.2
HDL cholesterol (mmol/L)	1.3 \pm 0.3
Triacylglycerol (mmol/L)	1.5 \pm 0.1
Fasting Glucose (mmol/L)	4.8 \pm 0.1
Haemoglobin (g/dL)	14.9 \pm 0.1
Systolic blood pressure (mm Hg)	124.0 \pm 1.7
Diastolic blood pressure (mm Hg)	74.9 \pm 1.5
<i>Liver Enzymes</i>	
Alanine Aminotransferase (ALT) (IU/L)	42.1 \pm 2.1
Gamma-glutamyltransferase (IU/L)	41.7 \pm 5.3

Table 4: Acute postprandial effects of orange flavanone beverages on static blood pressure.

Blood Pressure (mm Hg)		Baseline	2 h	5 h	7 h	<i>P</i>
<i>Systolic</i>	Control ¹	125.5 ± 1.9	126.7 ± 1.9	126.7 ± 1.7	128.6 ± 1.5	NS
	OJ ²	124.8 ± 1.5	125.7 ± 1.2	125.72 ± 1.5	126.4 ± 1.5	
	FROJ ³	126.1 ± 2.1	125.7 ± 2.0	127.3 ± 2.1	127.1 ± 1.6	
	WO ⁴	126.1 ± 1.6	125.1 ± 1.4	124.8 ± 1.5	126.6 ± 1.5	
<i>Diastolic</i>	Control	75.6 ± 1.6	71.5 ± 1.4	74.6 ± 1.4	72.1 ± 1.4	NS
	OJ	74.9 ± 1.5	70.1 ± 1.2	73.7 ± 1.5	70.4 ± 1.5	
	FROJ	75.5 ± 1.5	70.9 ± 1.6	74.6 ± 1.5	72.2 ± 1.4	
	WO	76.0 ± 1.6	69.9 ± 1.4	73.9 ± 1.5	70.9 ± 1.5	

¹ Control, sugar matched control; ² OJ, Tropicana pure premium orange juice without pulp; ³ FROJ, Flavanone-rich orange juice: Tropicana pure premium orange juice with added orange pomace; ⁴ WO, juice made from lightly blended fresh whole orange; Results are presented as mean ± SEM (n=36). Baseline levels did not differ between groups. NS: Non significant differences between treatments

Table 5: Pharmacokinetics of the major plasma flavanones, naringenin and hesperetin, after consumption of beverages containing either 128.88 mg (OJ), 272.14 mg (FROJ) or 452.80 mg (WO) of total orange flavanones in healthy middle-aged men.

		<i>C_{max}</i> (μ M)			<i>T_{max}</i> (h)			<i>AUC</i> (0 - 24 h)		
		OJ	AOJ	WO	OJ	AOJ	WO	OJ	AOJ	WO
<i>Flavanones</i>	Hesperetin	0.1 \pm 0.0	0.3 \pm 0.1 *	0.2 \pm 0.0 *	7.9 \pm 1.3	7.6 \pm 0.9	9.2 \pm 1.4	1.4 \pm 0.2	3.5 \pm 0.8 *	2.4 \pm 0.4 *
	Naringenin	0.05 \pm 0.0	0.2 \pm 0.0	0.1 \pm 0.0 *	6.5 \pm 1.0	6.5 \pm 0.2	6.3 \pm 0.3	0.8 \pm 0.1	1.9 \pm 0.5 #	1.8 \pm 0.2 *

OJ, Tropicana pure premium orange juice without pulp; FROJ, Flavanone-rich orange juice: Tropicana pure premium orange juice with added orange pomace; WO, juice made from lightly blended fresh whole orange. Results are presented as mean \pm SEM (n=20). * $P < 0.05$, indicates a significant difference in C_{max} / AUC in WO or FROJ in relation to OJ. # $P = 0.058$, indicates a trend in AUC in FROJ in relation to OJ. No significant differences in T_{max} were detected between treatments.

FIGURES

Figure 1. CONSORT flow diagram for the postprandial study. CONSORT, Consolidated Standards of Reporting Trials.

Figure 2. Time-course of postprandial FMD following consumption of flavanone beverages containing either 128.88 mg of flavanones (OJ); 272.14 mg of flavanones (FROJ); 452.80 mg of flavanones (WO) or a macronutrient and micronutrient-matched control in middle aged healthy men (n=28). A high-fat breakfast (51 g of fat) was administered at t = 0 h, and a medium-fat lunch (30 g of fat) was administered at t = 5.5 h. Data are presented as mean \pm SEM and analyzed using a 2-factor repeated measures ANOVA with time and treatment as the 2 factors [significant main effects of time x treatment ($P < 0.0001$), time ($P < 0.0001$) and treatment ($P < 0.05$)]. Post hoc analysis were conducted using Bonferroni multiple comparisons test. * $P < 0.05$ OJ significantly different from control at the 7 h; ** $P < 0.01$ FROJ and WO significantly different from control at 7 h. # Significant decrease in FMD response in relation to baseline levels for both control (at 2, 5 and 7 h; $P < 0.0001$, $P < 0.05$, $P < 0.0001$ respectively) and all three flavanone interventions (at 2 h; $P < 0.05$). FMD, flow-mediated dilatation; OJ, orange juice; FROJ, flavanone-rich orange juice; WO, whole blended orange.

Figure 3: Plasma flavanone profile following postprandial consumption of flavanone beverages containing either 128.88 mg of flavanones (OJ); 272.14 mg of flavanones (FROJ); 452.80 mg of flavanones (WO) or a macronutrient and micronutrient-matched control in middle aged healthy men (n=20). A) Hesperetin, B) Naringenin. Data are presented as mean \pm SEM and analyzed using a 2-factor repeated measures ANOVA with time and treatment as the 2 factors [significant main effects of time x treatment ($P < 0.0001$), time ($P < 0.0001$) and treatment ($P < 0.001$)]. Post hoc analysis were conducted using Bonferroni multiple comparisons test. Hesperetin levels are significantly higher in all treatments in comparison to control at 5 and 7 h ($^{\$} 0.00 < P < 0.02$), whilst Naringenin levels are significantly

higher in all treatments in comparison to control at 2, 5 and 7 h ($0.00 < P < 0.03$). *** $P < 0.001$, ** $P < 0.01$: levels of plasma Naringenin are significantly higher in OJ in comparison to WO. # $P < 0.1$: levels of plasma Hesperetin in FROJ and WO show a trend towards higher values than OJ. OJ, orange juice; FROJ, flavanone-rich orange juice; WO, whole blended orange.

Figure 4: Plasma Nitric Oxide levels following postprandial consumption of flavanone beverages containing either 128.88 mg of flavanones (OJ); 272.14 mg of flavanones (FROJ); 452.80 mg of flavanones (WO) or a macronutrient and micronutrient-matched control in middle aged healthy men (n=28). A) Nitrite levels (nmol/L), B) Nitrate levels ($\mu\text{mol/L}$), C) Nitroso species (RN XO) including nitrosothiols, nitrosamines, iron-nitrosylhemoglobin and nitrosohemoglobin (nmol/L). Data are presented as mean \pm SEM and expressed as change from baseline. Data were analyzed using a 2-factor repeated measures ANOVA with time and treatment as the 2 factors [significant main effects of time ($P < 0.0001$)]. Post hoc analysis were conducted using Bonferroni multiple comparisons test. ** $P < 0.01$ Nitrite levels are significantly different from baseline only for the control group at the specified time points. # Nitrate levels are significantly different from baseline for both control and all three flavanone treatments ($P < 0.0001$) at the specified time points. OJ, orange juice; FROJ, flavanone-rich orange juice; WO, whole blended orange.