

The effect of drought on dissolved organic carbon (DOC) release from peatland soil and vegetation sources

Article

Accepted Version

Creative Commons: Attribution 3.0 (CC-BY)

Open Access

Ritson, J. P., Brazier, R. E., Graham, N. J. D., Freeman, C., Templeton, M. R. and Clark, J. ORCID: <https://orcid.org/0000-0002-0412-8824> (2017) The effect of drought on dissolved organic carbon (DOC) release from peatland soil and vegetation sources. *Biogeosciences*, 14 (11). pp. 2891-2902. ISSN 1726-4170 doi: 10.5194/bg-14-2891-2017 Available at <https://centaur.reading.ac.uk/68650/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

Published version at: <http://dx.doi.org/10.5194/bg-2016-517>

To link to this article DOI: <http://dx.doi.org/10.5194/bg-14-2891-2017>

Publisher: Copernicus Publications

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online



The effect of drought on dissolved organic carbon (DOC) release from peatland soil and vegetation sources

Jonathan P. Ritson¹, Richard E. Brazier², Nigel J.D. Graham¹, Chris Freeman³, Michael R. Templeton¹ and Joanna M. Clark⁴.

¹ Department of Civil and Environmental Engineering, Imperial College London, South Kensington, London, SW7 2AZ, UK

² Geography, College of Life and Environmental Sciences, University of Exeter, EX4 4RJ, UK

³ Wolfson Carbon Capture Laboratory, School of Biological Sciences, Bangor University, Bangor, Gwynedd, LL57 2UW, UK

⁴ Department of Geography and Environmental Science; School of Archaeology, Geography and Environmental Science; The University of Reading, Whiteknights campus, PO Box 227, Reading, RG6 6AB, UK.

Correspondence to: Jonathan Ritson (j.ritson12@imperial.ac.uk)

Abstract: Drought conditions are expected to increase in frequency and severity as the climate changes, representing a threat to carbon sequestered in peat soils. Downstream water treatment works are also at risk of regulatory compliance failures and higher treatment costs due to the increase in riverine dissolved organic carbon (DOC) often observed after droughts. More frequent droughts may also shift dominant vegetation in peatlands from *Sphagnum* moss to more drought tolerant species. This paper examines the impact of drought on the production and treatability of DOC from four vegetation litters (*Calluna vulgaris*, *Juncus effusus*, *Molinia caerulea* and *Sphagnum* spp.) and a peat soil. We found that mild droughts caused a 39.6% increase in DOC production from peat and that this DOC was harder to remove by conventional water treatment processes (coagulation/flocculation). Drought had no effect on DOC production from vegetation litters, however large variation was observed between typical peatland species (*Sphagnum* and *Calluna*) and drought tolerant grassland species (*Juncus* and *Molinia*), with the latter producing more DOC per unit weight. This would therefore suggest the increase in riverine DOC often observed post-drought is due entirely to soil microbial processes and DOC solubility rather than litter-layer effects. Long term shifts in species diversity may, therefore, be the most important impact of drought on litter layer DOC flux, whereas more immediate effects are observed in peat soils. These results provide evidence in support of catchment management which increases the resilience of peat soils to drought, such as ditch-blocking to raise water-tables.

Keywords: Dissolved organic carbon, DOC, drought, peat, drinking water treatment

1.0 Introduction

Organic rich peat soils are a major global carbon sink (Limpens et al., 2008) which have formed due to the limited decay of recalcitrant plant litter found in peatland areas, coupled with anoxic conditions created by high



38 water-tables slowing decay (Billett et al., 2010; van Breemen, 1995). The locations in which these conditions
39 exist are threatened by climate change (Clark et al., 2010; Gallego-Sala and Prentice, 2012), and future climate
40 may also destabilise sequestered carbon (Evans and Warburton, 2010; Fenner and Freeman, 2011; Freeman et
41 al., 2001a).

42 Dissolved organic carbon (DOC) represents a significant flux of carbon from peatlands (Dinsmore et al., 2010)
43 and can also lead to difficulties for downstream drinking water treatment plants. DOC can cause colour, odour
44 and taste problems in drinking water and so must be removed as best as possible during treatment, commonly by
45 coagulation, flocculation and sedimentation/flotation. Any DOC which remains may act as a substrate for
46 microbial growth in the distribution system (Rodriguez and Sérodes, 2001) and can react during disinfection to
47 form disinfection by-products (DBPs) (Rook, 1974) which may have human health implications due to their
48 potential genotoxicity and carcinogenicity (Nieuwenhuijsen et al., 2009).

49 Droughts are projected to become more common under future climate conditions in the UK (Jenkins et al.,
50 2009). Droughts can have drastic consequences for peatland carbon storage and riverine DOC concentrations
51 due to the 'enzymatic latch' mechanism, whereby decomposition is suppressed due to the inhibitory effect of
52 phenolic compounds. Under drought conditions, the water table is lowered, creating oxic conditions which
53 stimulates phenol oxidase enzymes, thereby reducing the concentration of phenolics and their inhibitory effect
54 on hydrolase enzymes (Fenner and Freeman, 2011; Freeman et al., 2001a). Altered redox conditions can also
55 change the controls on DOC solubility, meaning organic carbon is not solubilised during the drought but instead
56 flushed from the system once redox conditions return to normal (Clark et al., 2006, 2005; Clark et al., 2011).

57 These processes have led to numerous observations of increased riverine DOC after droughts which may remain
58 elevated for years after the event (Evans et al., 2005; Scott et al., 1998; Watts et al., 2001; Worrall and Burt,
59 2004). How drought effects the treatability of DOC is less well understood although some authors have noted an
60 increase in the hydrophilic component during droughts and more hydrophobic character post-drought (Clark et
61 al., 2011; Scott et al., 1998; Watts et al., 2001). Hydrophobic DOC is commonly regarded as being easier to
62 remove via coagulation than the hydrophilic fraction (Bond et al., 2011; Matilainen et al., 2010).

63 The impact of climate change on DOC production and drinking water treatment is complex and involves a
64 number of biogeochemical cycles (Ritson et al., 2014b). Vegetative change in peatlands has occurred in the
65 recent past (Chambers et al., 2007b) and is projected to continue with *Sphagnum* mosses, which are favoured for
66 peat formation, giving way to vascular plants (Fenner et al., 2007; Weltzin et al., 2003). Many grassland species
67 (*Juncus effusus*, *Molinia caerulea*) have encroached on peatland areas as a result of anthropogenic pressures
68 such as nutrient deposition and management practices (Berendse, 1994; Chambers et al., 2007a; McCorry and
69 Renou, 2003; Shaw et al., 1996). These species are adapted to higher nutrient availability (Aerts, 1999) and thus
70 can out-compete peatland species if nutrient levels are elevated through, for example, nitrogen deposition
71 (Berendse et al., 2001).

72 Vegetative change has implications for carbon storage in peatlands, as *Sphagnum* is responsible for a number of
73 mechanisms (e.g. the production of recalcitrant litter) which allow carbon to be stored over long time periods
74 (van Breemen, 1995). Conversely, many vascular plants can destabilise colonised peat, stimulating
75 decomposition by adding labile carbon at the surface and through their root systems (Fenner et al. 2007; Gogo et
76 al. 2010). As such, a number of programmes have aimed to promote *Sphagnum* dominance for carbon storage



and other ecosystem services (Grand-Clement et al., 2013). However, further evidence is needed on the water quality outcomes of such interventions and the implications for water treatment. Previous work has highlighted both the vegetative source and climate controls on production affecting the ease of removal of DOC and the formation of DBPs (Gough et al., 2012; Reckhow et al., 2007; Ritson et al., 2014a; Tang et al., 2013). The present research sought to quantify the effect of drought on peatland DOC flux and any interaction with projected changes in litter input. To this end, climate simulations of varying drought severities defined in terms of percentiles of mean monthly rainfall were performed on four typical peatland vegetation types (*Calluna vulgaris*, *Juncus effusus*, *Molinia caerulea* and *Sphagnum* spp.) and a peat soil. After a six-week drought simulation, the DOC released upon rewetting was analysed in terms of optical properties and coagulation removal efficiency with ferric sulphate to determine: (a) whether drought conditions affect DOC production from peatland litter and soil types and (b) whether peatland species and invasive, drought tolerant vegetation produce different quantities and quality of DOC with respect to drinking water treatment.

2.0 Methodology

2.1 Field site and sample collection

Samples were collected from the Spooners site (51° 07'23.3'' N 3° 45'11.8'' W) in Exmoor National Park, UK at approximately 400 m elevation. Further site details can be found in Ritson et al., (2014a). The site is part of the MIRE project (Arnott, 2010) and was chosen as this area has been highlighted as a marginal peatland which may be vulnerable to climate change (Clark et al., 2010).

Samples of vegetation and peat soil were collected in one day in May 2014 and were sealed in airtight bags in a chilled container for transport from the field and stored in the dark at 4°C before use. For vascular plants, litter was collected as standing dead biomass. As the decomposition of *Sphagnum* is a continuum process, the section 2–4 cm below the capitulum was taken as equivalent to freshly senesced “litter”, as in other studies (e.g. Bragazza et al., 2007). Samples were sorted to remove any vegetation not belonging to the target species and then cut to 2 cm length and homogenised. Peat samples were collected using a screw auger and peat from 10–30 cm depth was used in the experiments. Peat samples were sorted to remove as many roots as possible but in sites where *Molinia* was present some fine roots remained.

The start times of the drought simulations for different DOC sources were staggered by up to two weeks to allow prompt analysis of water extracts at the end of the experiments. Preliminary work suggested chilled storage gave no significant difference in the amount of water extractable DOC or UV absorbance properties after three weeks of storage in the dark at 4°C.

2.2 Experimental Design

The vegetation and peat samples were homogenised by hand and randomly assigned a drought treatment in a five (vegetation types) x four (drought treatments) design with five replicates per treatment, giving 100 samples in total.

Data were obtained from regional historic climate records of the UK Meteorological Office for the south west of England for the period 1910–2013 (UK Met Office 2014) and these values were used to define three severities of drought and a control value. Data for the months of June, July and August (310 months in total) were used to



find the 50th, 25th, 10th and 5th percentile for total monthly rainfall and these values (Table 1) have been used to set control, mild, moderate and severe droughts, respectively.

Table 1: Monthly rainfall for control group and three severities of drought

Drought Treatment	Monthly rainfall total (mm)
Control (50 th percentile)	79.0
Mild (25 th percentile)	51.5
Moderate (10 th percentile)	34.7
Severe (5 th percentile)	23.3

The number of days of rain per month was fixed at a baseline value of eleven (regional average for June, July and August) and temperature ranged between the mean daily maximum of 18.9 for twelve hours and then and the mean daily minimum of 10.7 °C for twelve hours, calculated using the same historical UK Meteorological Office datasets for the south west of England.

2.3 Experimental procedure and laboratory methods

As in other decomposition studies, vegetation samples were air-dried to constant weight then mixed before subsampling (e.g. Latter et al., 1998). Five subsamples of each vegetation type were then oven-dried at 70 °C until constant weight, to determine the air-dry to oven-dry conversion factor. The peat samples were not air-dried before use as this would have changed the redox conditions within the peat and created a hydrophobic layer which can cause problems for re-wetting (Worrall et al., 2003). This will mean less accuracy in determining the starting weight of the peat sample as some variation in water content may exist, however this was minimised by effective homogenisation.

Buchner funnels fitted into amber-glass bottles were used to hold the sample and collect the simulated rainfall. Approximately 2 g dry-weight of air-dried vegetation/peat was used, however a lower weight of sample was used for *Sphagnum* (~0.65 g) and *Molinia* (~1.5 g) as this was enough to fill the Buchner funnel. The peat samples were spread over the area of the funnel so that a seal was created and the simulated rainwater infiltrated the peat rather than draining directly into the funnel.

The samples were then placed in an incubator for six weeks with simulated rainfall applied eleven times per month using high purity reverse osmosis (RO) treated water as per Table 1, following the methodology of Ritson et al. (2016).

As the samples were collected from the field and had been in contact with litter and soil, no inoculation with microorganisms was required as a suitable decomposer community was likely to be present (Van Meeteren et al., 2007). In this experiment the action of invertebrates and other microfauna was excluded, however their role in the decay of peatland litter is minimal (Dickinson and Maggs, 1974), although their role in DOC production from peat soils may be more significant (Cole et al., 2002).

At the end of the six week simulation the samples were air-dried and weighed. Water extractable DOC from the air dried sample was taken to simulate re-wetting following the end of the drought. DOC was extracted from soil and vegetation samples using approximately 20:1 ratio of RO treated water to sample. Previous work has shown that the amount of water used to extract DOC and whether one extraction is performed or sequential extractions



to simulate multiple rainfall events gives no significant variation in DOC quality (Don and Kalbitz, 2005, Soong et al., 2014), only changes in the total amount of carbon. DOC was measured as non-purgeable organic carbon (NPOC) via a UV/persulphate oxidation method on a Shimadzu TOC-V instrument. The method detection limit was determined by running five blank samples and using the value of three times the standard deviation. This was found to be 0.05 mg l^{-1} .

UV and fluorescence analysis was undertaken before coagulation/flocculation jar testing. UV absorbance was measured on a Perkin Elma Lambda 3 using a 1-cm pathlength quartz cuvette and the specific absorbance, SUVA, was calculated as the absorbance at 254 nm in units of m^{-1} divided by the NPOC content (mgC l^{-1}). Fluorescence analysis was completed using a Vary Eclipse fluorescence spectrophotometer where samples were scanned at excitation wavelengths between 220 and 450 nm at 5 nm intervals and the resulting emission recorded between 300 and 600 nm at 2 nm intervals. An R script was produced based on existing scripts (Lapworth and Kinniburgh, 2009) which performed a blank subtraction, masked out Rayleigh and Raman scattering, visualised the data and calculated fluorescence indices. Data were normalised to the Raman scattering peak of a RO water sample to allow comparison to other laboratories (Lawaetz and Stedmon, 2009). The 'peak C' measure, related to humic-like character, and the tryptophan-like peak, 'peak T' were defined as in Beggs et al., (2013).

Coagulation was performed on 350 ml of sample diluted to 3 mg l^{-1} DOC using a Phipps and Bird PB-700 paddled jar-tester (Phipps and Bird Ltd., Virginia, USA). After settling, the sample was filtered by Whatman qualitative grade 2 filters to remove flocs before NPOC analysis. Preliminary work indicated the following conditions gave effective DOC removal of similar samples: pH 5.5, 30.0 mg l^{-1} ferric sulphate dosed with 28.5 mg l^{-1} calcium hydroxide for pH control during a flash mix of one minute at 175 rpm, followed by a slow mix of 30 minutes at 60 rpm and then one hour of settling. Assessment of DBP formation was attempted, however analysis within the two week period specified in the method was not possible due to instrument failure so data quality could not be assured.

2.4 Data analysis and statistical methods

Statistical analysis was performed in the open source programming language, R, and SPSS version 21 (IBM). Due to problems with normality and heteroscedasticity a Box-Cox transform (Box and Cox, 1964) was applied to the variables before testing with a factorial ANOVA. A Tukey HSD post-hoc procedure was used for pairwise comparisons between the DOC sources and drought conditions. Estimates of effect sizes were made using ω^2 as this is suitable for small samples sizes (Keselman, 1975). Interactive effects from the omnibus ANOVA were followed up using multiple one-way ANOVAs with a Holm-Šidák correction to control the inflation of type one error (Holm, 1979; Šidák, 1967).

2.5 Repetition of the control group

To further investigate the effect of oxygenation of peat on DOC production and treatability, the control condition of this experiment was repeated in August 2015 using peat samples collected from similar ombrotrophic peatland sites in Dartmoor National Park (site details available in Ritson et al., 2016). Water extractable DOC was taken from a subsample before the climate simulation began and analysed for fluorescence and UV properties. Approximately 3.5 g dry weight of peat was then incubated using the same temperature and



rainfall as the control samples of the drought experiment with three replicates. After six weeks water extractable DOC was again taken for fluorescence and UV analysis to assess any changes in DOC quality.

3.0 Results

3.1 Omnibus ANOVA

A factorial ANOVA was performed exploring the source, drought and interactive effects on DOC, SUVA, DOC removal efficiency and the removal of SUVA (Table 2). Extractable DOC and SUVA had significant source, drought and source*drought effects suggesting that there is variation in the sensitivity of the sources to drought. No drought effects were observed for DOC removal or SUVA removal, although the source had strong effects on these parameters. For all significant results the effect size for the source was much greater than that for the drought treatment.

Table 2: p-values from factorial ANOVA (significant values have been highlighted in bold and displayed with ω^2 estimate of effect size in brackets)

Variable	Water extractable DOC	SUVA	DOC removal	SUVA removal
Factor				
DOC source	<0.001 (0.945)	<0.001 (0.422)	<0.001 (0.396)	<0.001 (0.331)
Drought	0.007 (0.004)	0.007 (0.034)	0.418	0.475
DOC source*Drought	0.050 (0.004)	0.005 (0.054)	0.234	0.951

3.2 Water extractable DOC

The omnibus ANOVA suggests both significant source and drought effects as well as an interaction, suggesting the effect of drought varies between the sources. The mean DOC extracted for all samples from each source is shown in Figure 1. The vegetation samples produced more DOC than the peat soil ($0.58 \pm 0.02 \text{ mg g}^{-1}$) with the peatland species, *Sphagnum* and *Calluna*, producing 3.47 ± 0.30 and $6.86 \pm 0.37 \text{ mg g}^{-1}$, respectively whereas the grassland species, *Juncus* and *Molinia*, produced much more at 9.21 ± 0.62 and $16.52 \pm 1.17 \text{ mg g}^{-1}$, respectively. A Tukey HSD test suggested that all DOC sources have significantly different means at the $p < 0.01$ level except the *Calluna* - *Juncus* comparison which was significantly different at the $p < 0.05$ level.




Figure 1: Water extractable DOC of all samples across the different DOC sources (n=20 per source). Error bars at one standard error.

To investigate the source*drought interaction one-way ANOVAs were performed for drought effects on each of the sources (Table 3) using a Holm-Šidák correction to control the inflation of type one error. This method changes the value used for alpha, the significance level, based on how many comparisons have been performed starting with the source with lowest p value and moving to the next lowest until an insignificant comparison is found.

Table 3: ANOVA results testing the effect of drought on water extractable DOC from different sources. Significant effects (Holm-Šidák correction) are highlighted in bold with the ω^2 estimate of effect size in brackets.


DOC Source	p value (DOC extraction)	Alpha used for comparison
Peat	0.010 (0.393)	0.010
Juncus	0.038	0.013
Sphagnum	0.097	-
Calluna	0.418	-
Molinia	0.550	-

Due to the decrease in the level of significance of the p value in the Holm-Šidák method only the peat source was found to have a drought effect on water extractable DOC. The mean values were 0.48, 0.67, 0.61 and 0.58 mg g⁻¹ for the control, mild, moderate and severe treatments, respectively, and this is shown in Figure 2. The mild drought treatment gave a significant increase in extractable DOC, indicated by a Tukey test for comparison to the control group (p=0.007). This corresponded to a 39.6% increase in DOC production for the mild drought



234 treatment. Taken together, the main effects and interaction and ω^2 values suggest that the source of DOC is the
 235 most important factor on extractable DOC and that the effect of drought is significant only for the peat soil and
 236 not for the vegetation.

237
 238



239
 240 **Figure 2: DOC extracted from peat on rewetting following different severities of drought (n=5 per**
 241 **treatment). Letters indicate statistically similar groups from the Tukey test. Error bars at one standard**
 242 **error.**

243
 244 A larger standard error in the moderate and severe drought treatments meant that these were not significantly
 245 different from the control ($p=0.060$ and $p=0.204$, respectively). Observations made throughout the experiment
 246 suggested that in the severe treatment there was a large variation in the extent to which each replicate dried out.
 247 Once peat becomes dry, a hydrophobic layer forms (Spaccini et al. 2002; Worrall et al. 2003), meaning that less
 248 water will infiltrate the sample, therefore possibly increasing the severity of the drought beyond the
 249 experimental design.

250 Variation in peat water content during the experiment was not recorded; however the water content of the peat
 251 samples was measured at the end of the experiment. This averaged 16.11, 14.14, 15.11 and 5.95 g with standard
 252 errors of 7.7, 3.0, 15.9 and 28.1% for the peat control, mild, medium and severe drought treatments respectively.
 253 The much larger standard error in final water content agrees with observations during the experiment and could
 254 perhaps explain some of the increased variation in extractable DOC for the severe drought treatment. This
 255 hypothesis was tested by comparing the variation from group mean in final water content for each sample with
 256 the variation from group mean in extractable DOC. These two measures of variance were found to correlate
 257 (Spearman's ρ coefficient 0.484, $p=0.031$) suggesting some of the variation in DOC extracted may be explained
 258 by different water contents between the samples in each treatment. This could have been caused by small
 259 variations in the way rain was applied over the area of the sample or because shrinkage of the peat mass allowed



260 water to pass through the funnel rather than infiltrate the peat, again possibly increasing the severity of drought
261 beyond the experimental design.

262

263 3.3 SUVA

264 Mean values of SUVA in $\text{L mg}^{-1} \text{m}^{-1}$ for the different sources were in the order *Molinia* (3.03 ± 0.38), peat (3.01
265 ± 0.15), *Juncus* (2.04 ± 0.06), *Calluna* (1.66 ± 0.14) and then *Sphagnum* (1.34 ± 0.13). The Tukey HSD test
266 suggested that the mean values for SUVA formed three subsets with peat and *Molinia* > group two *Calluna* and
267 *Juncus* > *Calluna* and *Sphagnum*.

268

269 To investigate the source*drought interaction one-way ANOVAs were performed for drought effects on SUVA
270 from each of the sources (Table 4) using a Holm-Šidák correction.

271

272 **Table 4: ANOVA results testing the effect of drought on SUVA for different DOC sources. Significant**
273 **effects (Holm-Šidák correction) are highlighted in bold with the ω^2 estimate of effect size in brackets**
274

DOC Source	p value (SUVA)	Alpha used for comparison
<i>Molinia</i>	0.001 (0.546)	0.010
<i>Sphagnum</i>	0.278	0.013
<i>Calluna</i>	0.436	-
Peat	0.696	-
<i>Juncus</i>	0.741	-


275

276 Tukey's test suggested that both the moderate and severe drought treatments were significantly different than
277 the control ($p=0.045$ and 0.026 , respectively) with means of 2.15 , 4.09 and $4.27 \text{ L mg}^{-1} \text{m}^{-1}$ for the control,
278 medium and severe treatment, respectively. Figure 3 shows a graph of SUVA for *Molinia* DOC from the
279 different treatment groups. The SUVA value approximately doubles between the control and the moderate and
280 severe droughts suggesting a large climatic control on the production of aromatic DOC from *Molinia* litter.
281 Taken together, the main effects and interaction and ω^2 values suggest that the source of DOC is the most
282 important factor on SUVA and that the effect of drought is significant only for *Molinia* litter and not for the
283 other vegetation types or the peat soil.

284



285



286

287 **Figure 3: SUVA value of *Molinia caerulea* derived DOC produced under differing severities of drought**
 288 **(n=5 per treatment) with error bars at one standard error. Letters indicate statistically similar groups**
 289 **from the Tukey test.**

290

291


292 3.4 DOC removal efficiency

293 Mean values for DOC removal by coagulation with ferric sulphate were in the order of *Juncus* (54.7 ± 2.3 %),
 294 *Molinia* (37.5 ± 2.6 %), peat (37.0 ± 2.9 %), *Calluna* (35.1 ± 2.0 %) and then *Sphagnum* (26.0 ± 2.9 %). The
 295 Tukey HSD test suggested that the mean values for DOC removal efficiency fell into three subsets with similar
 296 means in the order *Juncus* > *Molinia*, peat and *Calluna* > *Sphagnum*. The factorial ANOVA suggested no drought
 297 effects on removal efficiency ($p=0.418$). The removal efficiency for all samples from each DOC source is
 298 shown in Figure 4. *Juncus* DOC proved to be the easiest to remove via coagulation/flocculation with peat,
 299 *Calluna* and *Molinia* all relatively easily removed at just under 40%. Comparatively poor removal was achieved
 300 for *Sphagnum* DOC (<30%) which may be attributable to the low SUVA and peak C measure also found.

301



302



303

304 **Figure 4: DOC removal efficiency by coagulation/flocculation for different DOC sources (n=20 for each**
 305 **source, error bars at one standard error, letters indicate statistical subset according to Tukey test).**

306

307 3.5 SUVA removal efficiency

308 The removal of aromaticity, measured by SUVA, is of interest in drinking water treatment as aromatic
 309 compounds have a high propensity to form some of the regulated DBPs on chlorination (Bond et al., 2011).
 310 Large, aromatic compounds are selectively removed by coagulation/flocculation and as expected good removal
 311 (>70%) was observed for most of the samples. The mean values for the reduction in SUVA value following
 312 coagulation with ferric sulphate was in the order of peat (76.6 ± 1.8 %), *Sphagnum* (76.3 ± 2.5 %), *Molinia*
 313 (67.7 ± 4.7 %), *Calluna* (49.6 ± 5.3 %) and then *Juncus* (44.5 ± 2.3 %). The Tukey HSD test suggested that
 314 there were two subsets of DOC sources with similar means with peat, *Sphagnum* and *Molinia* > *Juncus* and
 315 *Calluna*. As with the overall DOC removal efficiency, there were no drought effects on SUVA removal
 316 ($p=0.475$). *Sphagnum* DOC showed good removal of SUVA despite relatively poor removal of total DOC,
 317 suggesting the aromatic compounds present in the sample are easily removed but that a large pool of aliphatic
 318 compounds are also present and these are more difficult to treat by conventional means.

319

320 3.6 Correlations between measures of DOC quality and treatability

321 A number of DOC quality indices based on absorbance and fluorescence measures were tested. The correlation
 322 coefficients for the different quality and treatability parameters are shown in Table 5. Peak C, a humic-like
 323 fluorescence peak, showed the best correlation with removal efficiency while the ratio of humic-like to protein-
 324 like fluorescence (Peak C/T) gave a lower but still significant correlation coefficient. The magnitude of peak C
 325 values were in the order *Juncus*>*Molinia*>*Calluna*>peat>*Sphagnum* which is consistent with data on removal
 326 efficiency. The SUVA value showed the best correlation with SUVA removal efficiency, suggesting that DOC
 327 with a lower proportion of aromatic compounds (low SUVA value) contains aromatic compounds which are



harder to remove by coagulation, possibly meaning they are either low molecular weight and/or also contain hydrophilic groups.

Table 5: Spearman's ρ for different DOC quality and treatability measures

DOC quality measure	Treatability measure	Spearman's ρ
Peak C	DOC removal %	0.578, $p < 0.001$
Peak C/T	DOC removal %	0.268, $p = 0.007$
SUVA	SUVA removal %	0.445, $p < 0.001$
Specific Peak C	SUVA removal %	0.235, $p = 0.019$

3.7 Repetition of control group

The data obtained from DOC extracted before and after the repeated simulation were analysed using student's t-test (equal variances assumed, confirmed using Levene's test) to assess whether the DOC extracted was significantly different following six weeks of exposure to oxygen without any experimental treatment. The results of this analysis are shown in Table 6.

Table 6: t-tests for pre and post-incubation peat samples (significant differences highlighted in bold)

Variable	t test	p value	% change
Extractable DOC	5.685	0.005	+41.6
Fluorescence peak C	8.168	0.011	-29.2
Fluorescence C/T	0.180	0.866	Not significant
SUVA	3.195	0.033	-23.0

Water extractable DOC increased significantly from 0.19 to 0.27 mg g⁻¹, an increase of 41.6%. The SUVA value decreased at the end of the simulation from 3.62 to 2.85 L mg m⁻¹, as did the fluorescence Peak C measure, which suggests a decrease in the level of aromaticity and humification of the DOC, respectively. This result may explain why poorer DOC removal for peat DOC was observed in this experiment than in our previous work (Ritson et al., 2016) as exposure to oxygen reduces the aromaticity of peat DOC and therefore its amenability to removal via coagulation.

4.0 Discussion

4.1 Water extractable DOC

The peat soil was affected by the drought treatment with higher extractable DOC observed at the mild severity. This finding is consistent with the 'enzymatic latch' hypothesis that increased oxygenation of peat engages a biogeochemical cascade whereby increased phenol oxidase activity ends the phenol-induced inhibition of hydrolase enzymes, thus increasing overall organic matter decomposition (Freeman et al., 2001a). This is also confirmed by the replication of the control treatment which showed exposure to oxygen even in the absence of drought increased DOC production and decreased DOC aromaticity. This finding has implications for all



laboratory studies which remove peat from anoxic conditions as these may not be representative of in-situ conditions.

No effect was observed with the moderate and severe drought treatments which may be explained by water scarcity limiting microbial activity (Toberman et al., 2008) and/or increased hydrophobic protection decreasing the extractable DOC on rewetting. The very low final water content of the severe treatment and observations of drying out and shrinkage of the peat mass throughout the experiment add weight to these possible explanations, although actual rates of microbial respiration were not monitored during the experiment.

The lack of a drought effect on DOC production from any of the vegetation types suggest the pulse in DOC observed post-drought elsewhere in catchment scale studies (Evans et al., 2005; Scott et al., 1998; Watts et al., 2001; Worrall and Burt, 2004) is likely to be due to the oxygenation of peat soils rather than any litter layer effects. This increase in peat-derived DOC is significant for downstream water treatment as our previous work showed this has more environmental persistence than vegetation sources (Ritson et al., 2016) and the UV and fluorescence data suggested DOC from peat exposed to oxygen may be more difficult to remove by conventional treatment measures. High DOC production was noted for the vascular plants, suggesting they may be an important source of DOC within peatland catchments during the period of their senescence, although drought does not affect the amount they produce. Drought conditions may, however, precipitate a change in vegetation type favouring more drought-tolerant species (Bragazza, 2008), which may have longer term effects for peatland biogeochemistry.

The amount of DOC extracted from *Sphagnum* was low, which may be due to the fact that its litter is recalcitrant to decay due to its high polyphenol content and numerous compounds with antimicrobial and antifungal properties (van Breemen, 1995). The other typically upland species, *Calluna*, produced the second least amount of DOC of the vegetation types, which also agrees with literature surrounding the recalcitrance of its litter (Aerts, 1995; Huang et al., 1998) and field studies suggesting areas of *Calluna* produce more porewater DOC than *Sphagnum* (Armstrong et al., 2012). The two grassland species, *Molinia* and *Juncus*, produced much larger amounts of DOC per g of dry weight. This is in keeping with the growth strategy of these species, whereby they rapidly produce a large amount of above-ground biomass and produce litter which decays readily, providing a positive feedback to its strategy of rapid growth and fast nutrient cycling (Aerts, 1999; Mann and Wetzel, 2000). This growth strategy is in contrast to that of the upland species *Calluna* and *Sphagnum*, which have adapted to low nutrient availability and therefore grow slowly, have nutrient poor litter and invest fewer resources in material which cycles rapidly (Aerts, 1999). Correlations between litter C:N ratio, suggesting nutrient availability, and amount of extractable DOC have been found in our previous work (Ritson et al., 2016) and elsewhere in the literature (Soong et al, 2014).

Molinia encroachment is a well acknowledged problem in Europe (Chambers et al., 2007b; Heil and Diemont, 1983; Hughes et al., 2007; Milligan et al., 2004) and nitrogen deposition and drier summers may mean more grassland species in the UK uplands in the future. The results of this study suggest the transition from *Sphagnum* to *Calluna* and *Molinia* observed in a paleoecological study of the area nearby our Exmoor site (Chambers, 1999) may have increased the amount of extractable DOC in the litter layer on g per g basis, as well as increased the seasonality of its export (Ritson et al., 2016). The much greater effect sizes for DOC source versus drought controls in this study and temperature and rainfall controls in previous work (Ritson et al., 2014a) suggest that the source of the DOC may be the primary driver of DOC quantity and quality in peatland



litters, consistent with litter decomposition studies in boreal peatlands (Straková et al., 2011). This has important implications for overall soil carbon stability in peatlands as the addition of labile carbon from litter can stimulate the decomposition of older carbon (Fontaine et al., 2007).

Studies concerning vegetation control of pore-water DOC are limited, but are reviewed in Ritson et al. (2016). Fenner et al. (2007) found elevated CO₂ caused a transition from *Sphagnum* to *Juncus* dominance on monoliths from flush peat which gave a 66% rise in DOC, attributed to an increase in above-ground biomass, more labile litter and stimulation of peat decomposition through root exudation. Vestgarden et al., (2010) found DOC in pore-waters beneath different vegetation types to be in the order *Molinia*>*Calluna*>*Sphagnum* in shallow samples but *Sphagnum* had higher concentrations than the vascular plants at depth and showed less seasonal variation. This has been linked to the seasonal growth cycles of vascular plants in peatlands which provide litter which decomposes rapidly and produces a large amount of DOC on a mg per g basis creating greater seasonality in DOC export (Ritson et al., 2016).

4.2 SUVA

The SUVA value has been linked to the aromaticity of DOC (Weishaar et al., 2003) and is of interest as a predictor of coagulation removal efficiency and DBP formation (Matilainen et al., 2011) in water treatment. The highest SUVA value was observed for the peat soil and *Molinia* litter, and the lowest value for the statistical subset of *Sphagnum* and *Calluna*. In a similar trend to DOC production, it appears that the grassland species produce DOC of greater aromaticity than the peatland species. *Molinia* also showed an interactive effect with the drought treatment, with a greater flux of aromatic compounds at the moderate and severe treatments, suggesting dry conditions are favourable for the breakdown and/or solubilisation of aromatic compounds in *Molinia* litter. *Molinia* DOC may, therefore, contribute to the increase in the aromaticity of peatland DOC observed after droughts at the catchment scale (Scott et al., 1998; Watts et al., 2001), although solubility controls on peat-derived DOC may be more important (Clark et al., 2006, 2005; Clark et al., 2011).

No drought effect was found for the SUVA value of peat which is in contrast to field studies which have shown a decrease in aromaticity of DOC during drought due to solubility controls and an increase in aromaticity on rewetting (Evans et al., 2005; Scott et al., 1998; Watts et al., 2001; Worrall et al., 2004). This may be explained by the fact that field studies have shown an increase in DOC aromaticity over many years, whereas this study examined a single rewetting event following drought, so the altered biogeochemical controls on DOC aromaticity may not have had enough time to exert a significant effect. The laboratory conditions may also have played a part, as the control sample is likely to have been exposed to more oxygenation through sample collection and setup of the experiment than undisturbed peat in the field, therefore increasing its similarity to the treatment conditions. The changes in DOC properties when the control group was repeated would appear to confirm this hypothesis.

These results suggest encroachment of grassland species into the uplands will increase seasonal DOC flux from the litter layer and increase the aromaticity of exported DOC and create a drought effect where *Molinia* litter is present. The lack of a drought effect for peat suggests that the long-term effects caused by water table drawdown identified elsewhere in the literature will likely be more important for DOC flux than the short-term effects studied here.



437 **4.3 DOC and SUVA removal**

438 DOC removal for all sources were typical of literature values (Matilainen et al., 2010), with *Juncus* DOC
439 proving the easiest to remove and *Sphagnum* DOC the hardest. Repeating the control condition and measuring
440 DOC production and quality parameters allowed an estimate of the effect of oxygen exposure for peat samples.
441 This showed a decrease in SUVA value and humic-like character (fluorescence Peak C) as well as a large
442 increase in extractable DOC. These changes in quality parameters may provide an explanation of why poorer
443 removal by coagulation was achieved for peat following this drought experiment than had been observed in our
444 previous work (Ritson et al., 2016) as less aromatic/humified material is likely to be harder to remove by
445 coagulation (Bond et al. 2011). Poorer removal was observed for *Sphagnum* than in our previous work; the
446 effect of more oxygenated conditions on vegetation decomposition remains an area for further research,
447 particularly as climate change may increase the likelihood of water table draw down in peatlands.
448 The coagulation removal efficiency could best be explained by the Peak C fluorescence index, suggesting humic
449 substances content was the strongest predictor of DOC removal. This is in contrast to our previous work which
450 found the ratio of humic to protein-like DOC to be the most important predictor (Ritson et al. 2014b). Our
451 previous work used DOC collected throughout a two-month simulation rather than a single re-wetting event at
452 the end. The samples will, therefore, have likely undergone microbial processing during this simulation and
453 consequently an increase in the amount of autochthonous DOC, hence the greater importance of the
454 fluorescence measure of protein-like DOC.

455

456 **5.0 Conclusions**

457 Climate projections for the UK vary, however most agree the likelihood of droughts in the future is set to
458 increase. The results of this research suggest the dominant effect of drought on peatland DOC sources is to
459 increase the amount and decrease the treatability of DOC from peat soils. This is likely due to the ‘enzymatic
460 latch’ mechanism increasing decomposition when oxic conditions prevail. No drought effect on different
461 vegetation litters was found, suggesting that the greatest effect of drought for vegetation may be facilitating
462 shifts to drought-tolerant species dominance rather than altering decomposition processes in the short term.
463 Oxygenation of peat appears to greatly increase extractable DOC whilst also decreasing the aromaticity and
464 humification, which may mean it is more difficult to remove at the treatment works. These results provide
465 support for catchment management programmes seeking to increase resilience to drought by raising peatland
466 water tables as a strategy for mitigating against high riverine DOC concentrations following droughts.

467

468 **Author contributions**

469 All authors developed the experimental design and advised on the subsequent analysis. Ritson performed the
470 experiments and data analysis. The manuscript was written by Ritson with contributions from all co-authors.

471

472 **Acknowledgements**

473 This work was supported by the Engineering and Physical Sciences Research Council [grant number
474 EP/N010124/1]. The authors would also like to thank the Grantham Institute: Climate and Environment and
475 Climate-KIC for the financial support of Jonathan Ritson. The authors would also like to thank South West



- Water's Mires project for access to sites as well as Exmoor and Dartmoor National Park Authorities, Natural England and Duchy of Cornwall. Freeman acknowledges NERC Grant NE/K01093X/1.
- References**
- Aerts, R.: The advantages of being evergreen, *Trends Ecol. Evol.*, 10(10), 402–407 [online] Available from: <http://www.sciencedirect.com/science/article/pii/S0169534700891569> (Accessed 29 September 2014), 1995.
- Aerts, R.: Interspecific competition in natural plant communities: mechanisms, trade-offs and plant-soil feedbacks, *J. Exp. Bot.*, 50(330), 29–37 [online] Available from: <http://jxb.oxfordjournals.org/content/50/330/29.short> (Accessed 11 September 2014), 1999.
- Armstrong, A., Holden, J., Luxton, K. and Quinton, J. N.: Multi-scale relationship between peatland vegetation type and dissolved organic carbon concentration, *Ecol. Eng.*, 47, 182–188, 2012.
- Arnott, S.: Exmoor Hydrological and Hydrogeological monitoring plan for the Mires-on-the-Moors project, UK., 2010.
- Beggs, K. M. H., Billica, J. A., Korak, J. A., Rosario-Ortiz, F. L., McKnight, D. M. and Summers, R. S.: Spectral evaluation of watershed DOM and DBP precursors, *J. Am. Water Works Assoc.*, 105, E173–E188, doi:10.5942/jawwa.2013.105.0063, 2013.
- Berendse, F.: Litter decomposability--a neglected component of plant fitness, *J. Ecol.*, 82(1), 187–190 [online] Available from: <http://www.jstor.org/stable/2261398> (Accessed 11 September 2014), 1994.
- Berendse, F., Van Breemen, N., Rydin, H., Buttler, A., Heijmans, M., Hoosbeek, M. R., Lee, J. A., Mitchell, E., Saarinen, T., Vasander, H. and Wallén, B.: Raised atmospheric CO₂ levels and increased N deposition cause shifts in plant species composition and production in Sphagnum bogs, *Glob. Chang. Biol.*, 7(5), 591–598, doi:10.1046/j.1365-2486.2001.00433.x, 2001.
- Billett, M. F., Charman, D. J., Clark, J. M., Evans, C. D., Evans, M. G., Ostle, N. J., Worrall, F., Burden, A., Dinsmore, K. J., Jones, T., McNamara, N. P., Parry, L., Rowson, J. G. and Rose, R.: Carbon balance of UK peatlands: current state of knowledge and future research challenges, *Clim. Res.*, 45(Table 1), 13–29 [online] Available from: <http://www.int-res.com/abstracts/cr/v45/p13-29/>, 2010.
- Bond, T., Goslan, E. H., Parsons, S. A. and Jefferson, B.: Treatment of disinfection by-product precursors., *Environ. Technol.*, 32(1-2), 1–25, doi:10.1080/09593330.2010.495138, 2011.
- Box, G. and Cox, D.: An analysis of transformations, *J. R. Stat. Soc. Ser. B (...)*, 26(2), 211–252 [online] Available from: <http://www.jstor.org/stable/2984418> (Accessed 18 August 2014), 1964.
- Bragazza, L.: A climatic threshold triggers the die-off of peat mosses during an extreme heat wave, *Glob. Chang. Biol.*, 2688–2695, doi:10.1111/j.1365-2486.2008.01699.x, 2008.
- van Breemen, N.: How Sphagnum bogs down other plants, *Trends Ecol. Evol.*, 10(7) [online] Available from: <http://www.sciencedirect.com/science/article/pii/S0169534795900071> (Accessed 27 August 2013), 1995.
- Chambers, F.: Recent rise to dominance of *Molinia caerulea* in environmentally sensitive areas: new perspectives from palaeoecological data, *J. Appl. ...*, 44, 719–733 [online] Available from: <http://onlinelibrary.wiley.com/doi/10.1046/j.1365-2664.1999.00435.x/full> (Accessed 26 September 2014), 1999.
- Chambers, F., Mauquoy, D., Gent, A., Pearson, F., Daniell, J. R. G. and Jones, P. S.: Palaeoecology of degraded blanket mire in South Wales: Data to inform conservation management, *Biol. Conserv.*, 137(2), 197–209, doi:10.1016/j.biocon.2007.02.002, 2007a.



- 517 Chambers, F., Mauquoy, D., Cloutman, E. W., Daniell, J. R. G. and Jones, P. S.: Recent vegetation history of
518 Drygarn Fawr (Elenydd SSSI), Cambrian Mountains, Wales: Implications for conservation management of
519 degraded blanket mires, *Biodivers. Conserv.*, 16, 2821–2846, doi:10.1007/s10531-007-9169-3, 2007b.
- 520 Clark, J., Chapman, P., Adamson, J. and Lane, S.: Influence of drought-induced acidification on the mobility of
521 dissolved organic carbon in peat soils, *Glob. Chang. Biol.*, 11(5), 791–809 [online] Available from:
522 <http://centaur.reading.ac.uk/15575/>, 2005.
- 523 Clark, J., Chapman, P., Heathwaite, A. and Adamson, J.: Suppression of dissolved organic carbon by sulfate
524 induced acidification during simulated droughts, *Environ. Sci. Technol.*, 40(6), 1776–1783 [online] Available
525 from: <http://eprints.lancs.ac.uk/53153/>, 2006.
- 526 Clark, J., Gallego-Sala, A., Allott, T., Chapman, S., Farewell, T., Freeman, C., House, J., Orr, H., Prentice, I.
527 and Smith, P.: Assessing the vulnerability of blanket peat to climate change using an ensemble of statistical
528 bioclimatic envelope models, *Clim. Res.*, 45, 131–150, doi:10.3354/cr00929, 2010.
- 529 Clark, J. M., Heinemeyer, A., Martin, P. and Bottrell, S. H.: Processes controlling DOC in pore water during
530 simulated drought cycles in six different UK peats, *Biogeochemistry*, 109(1-3), 253–270 [online] Available
531 from: <http://centaur.reading.ac.uk/22846/>, 2011.
- 532 Cole, L., Bardgett, R. D., Ineson, P. and Adamson, J. K.: Relationships between enchytraeid worms
533 (Oligochaeta), climate change, and the release of dissolved organic carbon from blanket peat in northern
534 England, *Soil Biol. Biochem.*, 34(5), 599–607, doi:10.1016/S0038-0717(01)00216-4, 2002.
- 535 Dickinson, C. and Maggs, G.: Aspects of the decomposition of *Sphagnum* leaves in an ombrophilous mire, *New*
536 *Phytol.*, (73), 1249–1257 [online] Available from: [http://onlinelibrary.wiley.com/doi/10.1111/j.1469-](http://onlinelibrary.wiley.com/doi/10.1111/j.1469-8137.1974.tb02154.x/abstract)
537 [8137.1974.tb02154.x/abstract](http://onlinelibrary.wiley.com/doi/10.1111/j.1469-8137.1974.tb02154.x/abstract) (Accessed 5 June 2014), 1974.
- 538 Dinsmore, K. J., Billett, M. F., Skiba, U. M., Rees, R. M., Drewer, J. and Helfter, C.: Role of the aquatic
539 pathway in the carbon and greenhouse gas budgets of a peatland catchment, *Glob. Chang. Biol.*, 16(10), 2750–
540 2762, doi:10.1111/j.1365-2486.2009.02119.x, 2010.
- 541 Evans, C. D., Monteith, D. T. and Cooper, D. M.: Long-term increases in surface water dissolved organic
542 carbon: observations, possible causes and environmental impacts., *Environ. Pollut.*, 137(1), 55–71,
543 doi:10.1016/j.envpol.2004.12.031, 2005.
- 544 Evans, C. D., Jones, T. G., Burden, A., Ostle, N., Zieliński, P., Cooper, M. D. A., Peacock, M., Clark, J. M.,
545 Oulehle, F., Cooper, D. and Freeman, C.: Acidity controls on dissolved organic carbon mobility in organic soils,
546 *Glob. Chang. Biol.*, 18(11), 3317–3331, doi:10.1111/j.1365-2486.2012.02794.x, 2012.
- 547 Evans, M. G. and Warburton, J.: Peatland Geomorphology and Carbon Cycling, *Geogr. Compass*, 4(10), 1513–
548 1531, 2010.
- 549 Fenner, N. and Freeman, C.: Drought-induced carbon loss in peatlands, *Nat. Geosci.*, 4(12), 895–900,
550 doi:10.1038/ngeo1323, 2011.
- 551 Fenner, N., Ostle, N. J., McNamara, N., Sparks, T., Harmens, H., Reynolds, B. and Freeman, C.: Elevated CO₂
552 Effects on Peatland Plant Community Carbon Dynamics and DOC Production, *Ecosystems*, 10(4), 635–647,
553 doi:10.1007/s10021-007-9051-x, 2007.
- 554 Fontaine, S., Barot, S., Barré, P., Bdioui, N., Mary, B. and Rumpel, C.: Stability of organic carbon in deep soil
555 layers controlled by fresh carbon supply., *Nature*, 450(7167), 277–280, doi:10.1038/nature06275, 2007.
- 556 Freeman, C., Ostle, N. and Kang, H.: An enzymic “latch” on a global carbon store., *Nature*, 409(6817), 149,
557 doi:10.1038/35051650, 2001a.



- 558 Freeman, C., Evans, C. D., Monteith, D. T., Reynolds, B. and Fenner, N.: Export of organic carbon from peat
559 soils., *Nature*, 412(6849), 785, doi:10.1038/35090628, 2001b.
- 560 Gallego-Sala, A. V. and Prentice, I. C.: Blanket peat biome endangered by climate change, *Nat. Clim. Chang.*,
561 3(2), 152–155, doi:10.1038/nclimate1672, 2012.
- 562 Gogo, S., Laggoun-Défarge, F., Delarue, F. and Lottier, N.: Invasion of a Sphagnum-peatland by *Betula* spp and
563 *Molinia caerulea* impacts organic matter biochemistry. Implications for carbon and nutrient cycling,
564 *Biogeochemistry*, 106(1), 53–69, doi:10.1007/s10533-010-9433-6, 2010.
- 565 Gough, R., Holliman, P., Willis, N., Jones, T. and Freeman, C.: Influence of habitat on the quantity and
566 composition of leachable carbon in the O2 horizon: Potential implications for potable water treatment, *Lake*
567 *Reserv. Manag.*, 28(4), 282–292, doi:10.1080/07438141.2012.741187, 2012.
- 568 Grand-Clement, E., Anderson, K., Smith, D., Luscombe, D., Gatis, N., Ross, M. and Brazier, R. E.: Evaluating
569 ecosystem goods and services after restoration of marginal upland peatlands in South-West England, edited by
570 S. Wan, *J. Appl. Ecol.*, 50(2), 324–334, doi:10.1111/1365-2664.12039, 2013.
- 571 Heil, G. and Diemont, W.: Raised nutrient levels change heathland into grassland, *Vegetatio*, 53, 113–120
572 [online] Available from: <http://link.springer.com/article/10.1007/BF00043031> (Accessed 26 September 2014),
573 1983.
- 574 Holm, S.: A simple sequentially rejective multiple test procedure, *Scand. J. Stat.*, 6(2), 65–70 [online] Available
575 from: <http://www.jstor.org/stable/4615733> (Accessed 15 September 2014), 1979.
- 576 Howson, G., Howard, D. M. and Scott, W. A.: Long term study of litter decomposition on a Pennine peat bog :
577 which regression ?, *Oecologia*, 113(1), 94–103, 1998.
- 578 Huang, Y., Stankiewicz, B. A., Snape, C. E., Evans, B., Latter, P. M. and Ineson, P.: Monitoring
579 biomacromolecular degradation of *Calluna vulgaris* in a 23 year field experiment using solid state ¹³C-NMR and
580 pyrolysis-GC/MS, *Soil Biol. Biochem.*, 30(12), 1517–1528, 1998.
- 581 Hughes, P., Lomas-Clarke, S., Schulz, J. and Jones, P.: The declining quality of late-Holocene ombrotrophic
582 communities and the loss of *Sphagnum austini* (Sull. ex Aust.) on raised bogs in Wales, *The Holocene*, 17(5),
583 613–625, 2007.
- 584 Jenkins, G., Murphy, J., Sexton, D., Lowe, J., Jones, P. and Kilsby, C.: UK climate projections: briefing report,
585 Exeter, UK., 2009.
- 586 Keselman, H.: A Monte Carlo investigation of three estimates of treatment magnitude: Epsilon squared, eta
587 squared, and omega squared., *Can. Psychol. Rev.*, 16, 44–48, 1975.
- 588 Lapworth, D. J. and Kinniburgh, D. G.: An R script for visualising and analysing fluorescence excitation–
589 emission matrices (EEMs), *Comput. Geosci.*, 35(10), 2160–2163, doi:10.1016/j.cageo.2008.10.013, 2009.
- 590 Lawaetz, A. J. and Stedmon, C. A.: Fluorescence intensity calibration using the Raman scatter peak of water.,
591 *Appl. Spectrosc.*, 63(8), 936–40, doi:10.1366/000370209788964548, 2009.
- 592 Limpens, J., Berendse, F., Blodau, C., Canadell, J. G., Freeman, C., Holden, J., Roulet, N., Rydin, H. and
593 Schaepman-Strub, G.: Peatlands and the carbon cycle: from local processes to global implications - a synthesis.,
594 *Biogeosciences*, 5, 1475–1491, doi:10.5194/bg-5-1379-2008, 2008.
- 595 Mann, C. and Wetzel, R.: Effects of the emergent macrophyte *Juncus effusus* L. on the chemical composition of
596 interstitial water and bacterial productivity, *Biogeochemistry*, 48, 307–322 [online] Available from:
597 <http://link.springer.com/article/10.1023/A:1006208213821> (Accessed 19 September 2014), 2000.



- 598 Matilainen, A., Vepsäläinen, M. and Sillanpää, M.: Natural organic matter removal by coagulation during
599 drinking water treatment: a review., *Adv. Colloid Interface Sci.*, 159(2), 189–97, doi:10.1016/j.cis.2010.06.007,
600 2010.
- 601 Matilainen, A., Gjessing, E. T., Lahtinen, T., Hed, L., Bhatnagar, A. and Sillanpää, M.: An overview of the
602 methods used in the characterisation of natural organic matter (NOM) in relation to drinking water treatment.,
603 *Chemosphere*, 83(11), 1431–42, doi:10.1016/j.chemosphere.2011.01.018, 2011.
- 604 McCorry, M. J. and Renou, F.: Ecology and management of *Juncus effusus* (soft rush) on cutaway peatlands,
605 *For. Ecosyst. Res. Gr. Rep. Number 69*, Dublin, Irel., (69), 2003.
- 606 Van Meeteren, M. J. M., Tietema, A. and Westerveld, J. W.: Regulation of microbial carbon, nitrogen, and
607 phosphorus transformations by temperature and moisture during decomposition of *Calluna vulgaris* litter, *Biol.*
608 *Fertil. Soils*, 44(1), 103–112, doi:10.1007/s00374-007-0184-z, 2007.
- 609 Milligan, A. L., Putwain, P. D., Cox, E. S., Ghorbani, J., Le Duc, M. G. and Marrs, R. H.: Developing an
610 integrated land management strategy for the restoration of moorland vegetation on *Molinia caerulea*-dominated
611 vegetation for conservation purposes in upland Britain, *Biol. Conserv.*, 119, 371–385,
612 doi:10.1016/j.biocon.2003.12.002, 2004.
- 613 Nieuwenhuijsen, M. J., Grellier, J., Smith, R., Iszatt, N., Bennett, J., Best, N. and Toledano, M.: The
614 epidemiology and possible mechanisms of disinfection by-products in drinking water., *Philos. Trans. A. Math.*
615 *Phys. Eng. Sci.*, 367(1904), 4043–76, doi:10.1098/rsta.2009.0116, 2009.
- 616 Reckhow, D., Rees, P., Nüsslein, K., Makdissy, G., Devine, G., Conneely, T., Boutin, A. and Bryan, D.: Long-
617 term Variability of BDOM and NOM as Precursors in Watershed Sources, *AwwaRF* [online] Available from:
618 [http://scholar.google.com/scholar?hl=en&btnG=Search&q=intitle:Long-](http://scholar.google.com/scholar?hl=en&btnG=Search&q=intitle:Long-Term+Variability+of+BDOM+and+NOM+as+Precursors+in+Watershed+Sources#0)
619 [Term+Variability+of+BDOM+and+NOM+as+Precursors+in+Watershed+Sources#0](http://scholar.google.com/scholar?hl=en&btnG=Search&q=intitle:Long-Term+Variability+of+BDOM+and+NOM+as+Precursors+in+Watershed+Sources#0) (Accessed 3 January
620 2014), 2007.
- 621 Ritson, J., Bell, M., Graham, N. J. D., Templeton, M. R., Brazier, R. E., Verhoef, A., Freeman, C. and Clark, J.
622 M.: Simulated climate change impact on summer dissolved organic carbon release from peat and surface
623 vegetation: Implications for drinking water treatment, *Water Res.*, 67(0), 66–76,
624 doi:10.1016/j.watres.2014.09.015, 2014a.
- 625 Ritson, J., Graham, N., Templeton, M. R., Clark, J. M., Gough, R. and Freeman, C.: The impact of climate
626 change on the treatability of dissolved organic matter (DOM) in upland water supplies: A UK perspective, *Sci.*
627 *Total ...*, 473–474, 714–730 [online] Available from:
628 <http://www.sciencedirect.com/science/article/pii/S0048969713015787> (Accessed 5 February 2014b), 2014.
- 629 Ritson, J. P., Bell, M., Brazier, R. E., Grand-clement, E., Graham, N. J. D., Freeman, C., Smith, D., Templeton,
630 M. R. and Clark, J. M.: Managing peatland vegetation for drinking water treatment, *Sci. Rep.*, 6:36751,
631 doi:10.1038/srep36751, 2016.
- 632 Rodriguez, M. J. and Sérodes, J. B.: Spatial and temporal evolution of trihalomethanes in three water
633 distribution systems., *Water Res.*, 35(6), 1572–86 [online] Available from:
634 <http://www.ncbi.nlm.nih.gov/pubmed/11317905>, 2001.
- 635 Rook, J. J.: Formation of haloforms during chlorination of natural water, *Water Treat. Exam.*, 23(2), 234–243,
636 1974.
- 637 Scott, M., Jones, M., Woof, C. and Tipping, E.: Concentrations and fluxes of dissolved organic carbon in
638 drainage water from an upland peat system, *Environ. Int.*, 537–546 [online] Available from:
639 <http://www.sciencedirect.com/science/article/pii/S0160412098000439> (Accessed 15 November 2012), 1998.



- 640 Shaw, S. C., Wheeler, B. D., Kirby, P., Philipson, P. and Edmunds, R.: Literature review of the historical effects
641 of burning and grazing of blanket bog and upland wet heath. English Nature Research Reports and Countryside
642 Council for Wales, English Nat. Res. Rep. 172, 1996.
- 643 Šidák, Z.: Rectangular Confidence Regions for the Means of Multivariate Normal Distributions, *J. Am. Stat.*
644 *Assoc.*, 62(318), 626–633, 1967.
- 645 Spaccini, R., Piccolo, A. and Conte, P.: Increased soil organic carbon sequestration through hydrophobic
646 protection by humic substances, *Soil Biol. Biochem.*, 34, 1839–1851 [online] Available from:
647 <http://www.sciencedirect.com/science/article/pii/S0038071702001979> (Accessed 5 September 2014), 2002.
- 648 Straková, P., Niemi, R. M., Freeman, C., Peltoniemi, K., Toberman, H., Heiskanen, I., Fritze, H. and Laiho, R.:
649 Litter type affects the activity of aerobic decomposers in a boreal peatland more than site nutrient and water
650 table regimes, *Biogeosciences*, 8(9), 2741–2755, doi:10.5194/bg-8-2741-2011, 2011.
- 651 Tang, R., Clark, J. M., Bond, T., Graham, N., Hughes, D. and Freeman, C.: Assessment of potential climate
652 change impacts on peatland dissolved organic carbon release and drinking water treatment from laboratory
653 experiments, *Environ. Pollut.*, 173, 270–277, 2013.
- 654 Toberman, H., Freeman, C., Artz, R. R. E., Evans, C. D. and Fenner, N.: Impeded drainage stimulates
655 extracellular phenol oxidase activity in riparian peat cores, *Soil Use Manag.*, 24(December), 357–365,
656 doi:10.1111/j.1475-2743.2008.00174.x, 2008.
- 657 Vestgarden, L. S., Austnes, K. and Strand, L. T.: Vegetation control on DOC, DON and DIN concentrations in
658 soil water from a montane system, southern Norway, *Boreal Environ. Res.*, 15(6), 565–578, 2010.
- 659 Watts, C. D., Naden, P. S., Machell, J. and Banks, J.: Long term variation in water colour from Yorkshire
660 catchments., *Sci. Total Environ.*, 278(1-3), 57–72 [online] Available from:
661 <http://www.ncbi.nlm.nih.gov/pubmed/11669277>, 2001.
- 662 Weishaar, J. L., Aiken, G. R., Bergamaschi, B. A., Fram, M. S., Fujii, R. and Mopper, K.: Evaluation of specific
663 ultraviolet absorbance as an indicator of the chemical composition and reactivity of dissolved organic carbon.,
664 *Environ. Sci. Technol.*, 37(20), 4702–8 [online] Available from:
665 <http://www.ncbi.nlm.nih.gov/pubmed/14594381>, 2003.
- 666 Weltzin, J. F., Bridgham, S. D., Pastor, J., Chen, J. and Harth, C.: Potential effects of warming and drying on
667 peatland plant community composition, *Glob. Chang. Biol.*, 9(2), 141–151, doi:10.1046/j.1365-
668 2486.2003.00571.x, 2003.
- 669 Worrall, F. and Burt, T.: Time series analysis of long-term river dissolved organic carbon records, *Hydrol.*
670 *Process.*, 18(5), 893–911, doi:10.1002/hyp.1321, 2004.
- 671 Worrall, F., Burt, T. and Shedden, R.: Long term records of riverine dissolved organic matter, *Biogeochemistry*,
672 64, 165–178 [online] Available from: <http://link.springer.com/article/10.1023/A:1024924216148> (Accessed 17
673 February 2014), 2003.
- 674 Worrall, F., Harriman, R., Evans, C. D., Watts, C. D., Adamson, J., Neal, C., Tipping, E., Burt, T., Grieve, I.,
675 Monteith, D., Naden, P. S., Nisbet, T., Reynolds, B. and Stevens, P.: Trends in Dissolved Organic Carbon in UK
676 Rivers and Lakes, *Biogeochemistry*, 70(3), 369–402, doi:10.1007/s10533-004-8131-7, 2004.
- 677