

# *A deterministic evaluation of heat stress mitigation and feed cost under climate change within the smallholder dairy sector*

Article

Published Version

York, L., Heffernan, C., Rymer, C. ORCID: <https://orcid.org/0000-0002-3535-4330> and Panda, N. (2017) A deterministic evaluation of heat stress mitigation and feed cost under climate change within the smallholder dairy sector. *Animal*, 11 (5). pp. 900-909. ISSN 1751-7311 doi: 10.1017/S1751731116002706 Available at <https://centaur.reading.ac.uk/68761/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

To link to this article DOI: <http://dx.doi.org/10.1017/S1751731116002706>

Publisher: Cambridge University Press

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

[www.reading.ac.uk/centaur](http://www.reading.ac.uk/centaur)

**CentAUR**

Central Archive at the University of Reading

Reading's research outputs online

# A deterministic evaluation of heat stress mitigation and feed cost under climate change within the smallholder dairy sector

L. York<sup>1†</sup>, C. Heffernan<sup>1a</sup>, C. Rymer<sup>2</sup> and N. Panda<sup>3</sup>

<sup>1</sup>Livestock Development Group (LDG), Faculty of Life Sciences, University of Reading, Reading RG6 6AR, UK; <sup>2</sup>Food Production and Quality Division, Faculty of Life Sciences, University of Reading, Reading RG6 6AR, UK; <sup>3</sup>Department of Animal Nutrition, Faculty of Veterinary Science and Animal Husbandry, Orissa University of Agriculture and Technology, Bhubaneswar 751003, India

(Received 2 March 2016; Accepted 20 October 2016)

*In the global South, dairying is often promoted as a means of poverty alleviation. Yet, under conditions of climate warming, little is known regarding the ability of small-scale dairy producers to maintain production and/or the robustness of possible adaptation options in meeting the challenges presented, particularly heat stress. The authors created a simple, deterministic model to explore the influence of breed and heat stress relief options on smallholder dairy farmers in Odisha, India. Breeds included indigenous Indian (non-descript), low-grade Jersey crossbreed and high-grade Jersey crossbreed. Relief strategies included providing shade, fanning and bathing. The impact of predicted critical global climate parameters, a 2°C and 4°C temperature rise were explored. A feed price scenario was modelled to illustrate the importance of feed in impact estimation. Feed costs were increased by 10% to 30%. Across the simulations, high-grade Jersey crossbreeds maintained higher milk yields, despite being the most sensitive to the negative effects of temperature. Low-capital relief strategies were the most effective at reducing heat stress impacts on household income. However, as feed costs increased the lower-grade Jersey crossbreed became the most profitable breed. The high-grade Jersey crossbreed was only marginally (4.64%) more profitable than the indigenous breed. The results demonstrate the importance of understanding the factors and practical trade-offs that underpin adaptation. The model also highlights the need for hot-climate dairying projects and programmes to consider animal genetic resources alongside environmentally sustainable adaptation measures for greatest poverty impact.*

**Keywords:** India, climate change, cow, animal genetic resources, temperature–humidity index

## Implications

Climate change and adapting to climate change (via heat stress management) will have a profound impact on household income and food security. Relief strategies that required low capital investment were least effective at preventing losses in milk yield. However, due to their low cost of implementation such strategies provided farmers with more income. Feed cost was a significant determinant of profitability. This study questions national breeding policy as advocating high producing exotic breeds will limit the long-term sustainability of the dairy sector.

## Introduction

Since the commencement of Operation Flood in the 1970s, India's bovine milk production has increased by 84%

(111.2 million tonnes) (Government of India, 2014). Between 1999 and 2013 the average yield of exotic crossbreeds and indigenous breeds has increased by 9.18% (0.59 kg) and 24.87% (0.47 kg), respectively (Government of India, 2006 and 2014). Exotic crossbreeds yield 7.02 kg/cow per day whilst indigenous breeds yield 2.36 kg/cow per day (Government of India, 2014).

Operation Flood focussed on smallholders (owning <2 ha) who are responsible for 70% of India's bovine population (Datta *et al.*, 2015) and produce 80% of the milk (Cunningham, 2009). A feature of Operation Flood was the advocacy and introduction of exotic breeds for crossbreeding with indigenous breeds. The crossbreeding of indigenous breeds with higher-yielding exotic breeds became national breeding policy. By 2012–13, crossbred cattle represented 28.4% of the milking cow population (Government of India, 2014). However, as climate change become more severe, the pro-production (see Heffernan *et al.*, 2012) policies of Operation Flood may have placed dairy producers in a precarious position.

The temperatures expected under climate change will precipitate heat stress and cause declines in productivity.

<sup>a</sup> Present address: School of Veterinary Sciences, University of Bristol, Langford House, Langford, Bristol BS40 5DU, UK.

<sup>†</sup> E-mail: luke\_york@live.com.au

By 2020, heat stress will decrease milk production from India's dairy cows by 0.73 million litres (Upadhyay *et al.*, 2007). Heat stress is measured against a temperature–humidity index (THI), a combined measure of temperature and humidity to denote thermal comfort (Berman, 2011). The THI threshold above which decreased milk yield is observed in Holstein Friesian cows is accepted to be 72 (Igono *et al.*, 1992; Ravagnolo *et al.*, 2000). No widely accepted THI threshold values exist for other dairy cattle types. However, the Jersey (Bryant *et al.*, 2007; Smith *et al.*, 2013) and dairying *Bos indicus* breeds (such as Sahiwal and Gir) are considered less susceptible to heat stress (Sirohi and Michaelowa, 2007; Upadhyay *et al.*, 2007).

The heat tolerance of *B. indicus* breeds is derived from a raft of physiological adaptations (such as propensity to sweat, size of sweat glands and coat type) that help the animal dissipate metabolic heat (Thornton *et al.*, 2009). Some commentators argue that the heat tolerance of *B. indicus* breeds is a function of productivity; the lower the level of milk production, the lower the quantity of metabolic heat produced (Berman, 2011). The low productivity of Indian cows would afford some level of heat tolerance. However, heat tolerance as a function of productivity does not adequately explain the role of physiological adaptations associated with *B. indicus* breeds.

Indeed, selection of Gir cows for increased milk yield reduces heat tolerance (Santana *et al.*, 2015). Over 30 years, the Brazilian Gir breed improvement programme increased population productivity by 0.22 kg/year, whereas heat-stress-induced yield declines of 0.019 kg/year (Santana *et al.*, 2015). Thus, as productivity improves metabolic heat appears to overwhelm physiological adaptations. However, the literature is not clear as to whether *B. indicus* breeds are more (or less) heat tolerant than *Bos taurus* breeds at similar production levels. Despite this gap in knowledge, yield improvement via exotic genetics poses a significant challenge to maintaining heat tolerance characteristics within India's cattle.

The Indian state of Odisha is characterized by a high incidence of poverty (35.69% of rural households) (Government of India, 2014) and heavy investment in live-stock production (Datta *et al.*, 2015). Odisha is ranked amongst those states considered most susceptible to the effects of climate change (Upadhyay *et al.*, 2007). For example, O'Brien *et al.* (2004) found significant portions of Odisha's agricultural lands to be vulnerable. Annual heat stress in Odisha is expected to cost crossbreed producers Rs. 1775.56/head, whereas producers reliant on indigenous cows will experience declines of Rs. 33.91/head (in 2007 prices) (Upadhyay *et al.*, 2007).

Climate change is also expected to indirectly affect live-stock production via complex effects on plant growth. However, within India the situation is simplified by chronic feed shortages (Government of India, 2013). Deficiencies of green fodder, dry fodder and concentrates are in the realm of 62%, 23% and 63%, respectively (Sharma *et al.*, 2011). Indeed, climate change (via CO<sub>2</sub> fertilization and altered

precipitation patterns) coupled with technological advances and improved plant breeding may increase feed availability. However, it is unlikely that improvements will be significant enough to reduce deficits and keep pace with demand. As such, the feed deficit is expected to increase (Pathak and Devakumar, 2011).

A simple deterministic model was developed to evaluate the influence of breed and heat stress relief options on smallholder dairy farmers in Odisha, India. A feed price scenario was modelled to illustrate the importance of feed in climate change impact estimation.

## Material and methods

### Model framework


A deterministic model was constructed to examine the impact of heat stress on milk yield and household income at critical climate change temperature thresholds (+2°C and +4°C). The influence of breed (indigenous, low-grade Jersey crossbreed and high-grade Jersey crossbreed) and the effectiveness of microclimate heat stress management strategies (shading, bathing and fanning) in protecting household income were also examined. Feed availability was considered via a pricing scenario assuming that climate change will increase the price of feeds. The model framework is shown in Figure 1.

### Household-level sampling and data collection

Villages were randomly selected within the high potential dairying zone surrounding the state capital Bhubaneswar. Participating villages were within a 40-km area crossing two districts (Puri and Khurda) and characterized by sufficient water, market access and relatively reliable animal health infrastructure. In all, 115 cattle-owning households were purposively sampled from Puri ( $n = 31$ ) and Khurda ( $n = 84$ ) districts. Cattle ownership was the only selection criterion. Cattle-owning households were identified during key informant interviews with local community leaders. Sampling was not stratified. Households were smallholders maintaining a herd of less than five. Average herd size was 1.95 cattle aged 1 year or older. The survey was conducted in November 2012 in the local language (Oriya). All prices were in 2012 values. Responses were translated into English at the time of the interview. A voice recorder ensured all interviews were recorded verbatim. Interviews were transcribed into Microsoft Access 2010.

### The interview

Farmers were asked a range of questions regarding the scope and impact of their dairying operation. Breed definitions (Jersey, 1/4 Jersey and Local) were farmer derived based on the phenotypic appearance of the animals. Farmers categorized cows based on phenotypic appearance. The Jersey crossbreed cows were categorized at two levels: a low-grade Jersey crossbreed and a high-grade Jersey crossbreed. The Jersey was considered as the high-grade Jersey



**Figure 1** The simple deterministic model design outlining the various adaptations and feed availability scenarios under investigation for smallholder dairy producers in Odisha, India.

crossbreed, the 1/4 Jersey as the low-grade Jersey crossbreed and the Local as the indigenous (non-descript) breed. *B. indicus* traits were required for a crossbred cow to be categorized by farmers as 1/4 Jersey. The genetic background of each breed could not be determined due to breeding practices. No attempt was made to determine breed beyond farmer definitions. The local indigenous breed is not a breed *per se*. Rather, it is convention to group *B. indicus* cattle not belonging to a described Indian breed together as non-descript. Buffalo were not considered as they only form 6.76% of Odisha's dairy population (Government of India, 2014).

Farmers were also asked milk price and the average daily milk yield for each month of lactation. Responses were corroborated where possible with individual farm-level records supplied by local milk collection agents. However, as records contained sales information, the amount of milk kept for household consumption required farmer recall. Each value was multiplied by the number of days in the month to determine total milk production per lactation (Table 1). The resulting figure was divided by 300 (Moran, 2005) to determine the average daily production over a standardized lactation length. Average lactation length could also be used to indicate daily milk production (Table 1). However, as a 300-day lactation is the international standard it was selected as the unit of analysis. Milk yields and price were converted into kg/day via a milk density factor of 1.033 (International Farm Comparison Network, 2015). The THI load and quantity of milk lost to heat stress using actual lactation length is provided in Supplementary Table S1.

#### Temperature and temperature–humidity index

District-level meteorological data were procured for the period 2011–12 (Government of Odisha, 2013). Data included the mean daily minimum and maximum temperatures and relative humidity (RH%) at 0830 and 1730 h for each

month. Although temperature and RH% for the district of Khurda was not specifically included in the data set, temperature and RH% data for Bhubaneswar were utilized instead. Data from Bhubaneswar and Puri district were used to create an average THI estimate. Daily or weekly meteorological data were desired. However, no such data could be located. Similarly, no long term (i.e. >10 years) district-level data inclusive of temperature and RH% could be identified to allow Climate Normal calculation.

A standard THI formula (Bryant *et al.*, 2007) was used to combine temperature and RH%. Minimum temperature was used as it is the greatest determinant of heat stress in cattle (Igono *et al.*, 1992; Keister *et al.*, 2002; Bryant *et al.*, 2007). Humidity recorded at 0830 h was matched with minimum temperature (Mohapatra *et al.*, 2015). The mean minimum temperature for each month (N) was increased by 2°C (N+2) and 4°C (N+4) to simulate climate change.

#### Model parameters

**Temperature–humidity index threshold and milk loss.** The THI thresholds and rate of milk yield loss assumed are shown in Table 1. A total of 30 relevant documents were reviewed to determine values. Although a focus on the global south (India in particular) was desired, poorly described methodologies limited the usefulness of such documents. Thus, the sample was primarily composed ( $n = 15$ ) of peer-reviewed literature from the southern United States. Values that had relevance to the breeds under investigation were selected (i.e. a focus on Jersey cows (Keister *et al.*, 2002; Smith *et al.*, 2013) and/or Indian *B. indicus* cows (Upadhyay *et al.*, 2007)). However, limited robust research required additional inclusion criteria. Studies investigating lower-yielding cattle (e.g. Bouraoui *et al.*, 2002) or containing similar breeding themes such as crossbreeding (e.g. Bryant *et al.*, 2007) were

**Table 1** The model parameter values for milk yield, the temperature–humidity index (THI) thresholds and rate of decline in milk yield

Breeds	Total yield (kg/cow per lactation)	Lactation length (days)	Average milk yield (kg/cow per day)	300-day lactation milk yield (kg/cow per day)	THI threshold	Rate of decline (%/THI unit)	Quantity lost (kg/THI unit)	Method of calculation
Jersey	1374	255	5.39	4.58	>74	2.19	0.1	Milk yield: field data (Puri, <i>n</i> = 9; Khurda, <i>n</i> = 51) Lactation length: field data (Puri, <i>n</i> = 9; Khurda, <i>n</i> = 51) Threshold: average values offered by Keister <i>et al.</i> (2002) and Smith <i>et al.</i> (2013). Converted to minimum THI by reduction of 3 THI units (Zimbleman <i>et al.</i> , 2009)
1/4 Jersey	963	241	4	3.21	>76	0.76	0.02	Rate of decline: directly from Bouraoui <i>et al.</i> (2002) Milk yield: field data (Puri, <i>n</i> = 4; Khurda, <i>n</i> = 14) Lactation length: field data (Puri, <i>n</i> = 4; Khurda, <i>n</i> = 14) Threshold: adapted from Bryant <i>et al.</i> (2007), which indicated that crossbreeding improves heat tolerance by 2.4 THI units.
Local	444	159	2.8	1.48	>77	0.48	0.007	Rate of decline: directly from Ravagnolo <i>et al.</i> (2000) Milk yield: field data (Puri, <i>n</i> = 6; Khurda, <i>n</i> = 7) Lactation length: field data (Puri, <i>n</i> = 6; Khurda, <i>n</i> = 7) Threshold: directly from Upadhyay <i>et al.</i> (2007) Rate of decline: directly from Igono <i>et al.</i> (1992)

Jersey = high-grade Jersey crossbreed; 1/4 Jersey = low-grade Jersey crossbreed; Local = indigenous (non-descript) breed.  
The THI threshold is the point at which declines in milk yield would be expected during periods of heat stress.

included. Milk price was Rs. 18.14/kg. The severity of heat stress was not expected to result in differing rates of decline (Key and Sneeringer, 2014).

The transposal of values from higher-yielding cows risked inaccuracy. However, there is an absence of high-quality local data. Furthermore, the investigation is speculative, assuming the dairy sector will continue to implement northern production methods and breeds. Indian cows will become increasingly comparable with higher-yielding northern cows.

*Calculating the temperature–humidity index load and heat stress impact on milk yield.* The THI load refers to the total number of units above the threshold THI experienced throughout the year. In months which exceed the threshold, each day of that month was assumed above the threshold. An annual estimate of THI load was calculated for each breed by adding together the THI units above the threshold in each month. The THI load was multiplied by milk yield and rate of loss to determine total quantity of milk lost to heat stress.

Calving commonly occurs in the late winter to pre-monsoon (February to May) and post-monsoon (October to November) period. Cows calving in the October to November period avoid lactation during highest temperature periods. As breeding is not organized to avoid lactation during this time it was assumed cows would be lactating throughout the hottest period (i.e. May to September). Relief strategy cost was also considered in terms of productivity (Supplementary Table S2).

#### *Heat stress relief strategy: shading*

Two types of shade were considered: trees and a man-made structure (shed). To ensure maximum cooling capacity and animal welfare, each cow was provided with 4.65 m<sup>2</sup> of shade. The model assumed tree shade increased the THI threshold by 1 unit, the modern shed increased the threshold by 2 units and the traditional shed increased the threshold by 3 units. Assumptions were derived from the studies by Reena *et al.* (2014) and Mohapatra *et al.* (2015). The traditional shed is more effective at moderating microclimate temperature due to the rice straw thatched roof (Reena *et al.*, 2014; Mohapatra *et al.*, 2015).

*Trees.* Field data were used to determine the purchase price of trees. As the trees will not reach maturity for a number of years, only trees planted more than 10 years ago were included. On average, households spent Rs. 9.75 to 58.52/tree (depending on species). The model assumed trees have now reached maturity. There was no labour cost as seedlings were planted.

The number of trees required to create 4.65 m<sup>2</sup> of shade differed between species. The most expensive was papaya (*Carica papaya*) as nine trees were required (Rs. 490.68). Mango (*Mangifera indica*) was the cheapest as only one tree was required (Rs. 19.51).

*Shed.* The modern shed was made from modern materials such as concrete, bricks, iron roofing, whereas the traditional



**Table 2** The cost components associated with bathing different breeds of dairy cows (Jersey, 1/4 Jersey and Local) using different water sources (naturally occurring, wells and motorized pumps) in sampled locations of Odisha, India

Water sources	Breed category	Time (min/bath per cow)	Quantity of water required (l/bath)	Labour (Rs/bath)	Installation cost (Rs/installation)	Lifetime (years)	Electricity (Rs/bath)
Naturally occurring	Jersey ( <i>n</i> = 35)	27	–	8	–	–	–
	1/4 Jersey ( <i>n</i> = 7)	40	–	12			
	Local ( <i>n</i> = 6)	24	–	8			
Well	Jersey ( <i>n</i> = 84)	28	44	9	6770 ( <i>n</i> = 46)	5.25	–
	1/4 Jersey ( <i>n</i> = 14)	45	57	14			
	Local ( <i>n</i> = 10)	29	52	9			
Motorized pump	Jersey ( <i>n</i> = 18)	31	100	10	1947 ( <i>n</i> = 2)	2.625	0.44 ( <i>n</i> = 2)
	1/4 Jersey ( <i>n</i> = 1)	18	No data	6			
	Local ( <i>n</i> = 3)	60	No data	19			

Jersey = high-grade Jersey crossbreed; 1/4 Jersey = low-grade Jersey crossbreed; Local = indigenous (non-descript) breed.

shed was made from locally available items such as bamboo, rice straw thatch.

Modern material sheds were found to cost Rs. 8700/cow, including all materials and land rental. The modern shed depreciated over 10 years, which is the length of time households had leased the land. Land rentals for the 10-year period were paid before construction. Construction labour costs were estimated from interviews. On average, 3-day labour was required. The total cost of labour for construction was Rs. 529.20 (Rs. 176.40/day).

The traditional shed construction cost was Rs. 3563.16 (including all materials). Land rentals were not included as traditional sheds were constructed on the households' land. Construction took 1.5 days at a total labour cost of Rs. 264.60. The expected lifetime was 5 years due to the poor durability of construction materials. To improve comparability it was necessary to build a second traditional shed after 5 years. Field data indicated that the rice straw thatch would be replaced yearly at a cost of Rs. 730.63 for 365 bundles of straw (Rs. 2/bundle) requiring 1 day of labour (Rs. 176.40).

#### Heat stress relief strategy: bathing

The model considered the effects of altering current bathing practices by increasing wetting events from 1/day to 2/day. There is no data available to indicate the effectiveness of traditional bathing practices as a heat stress management strategy.

Three water sources were used: naturally occurring sources (such as ponds, rivers), wells and motorized pumps. Currently, 63% (*n* = 47) of sampled cows are bathed in the morning. A second bath can be undertaken at 1400 h to prevent the peak in cow temperature at 1600 h (Kendall *et al.*, 2007). A second bath will keep THI thresholds 2.5 units higher until 1800 h, as wetting will maintain body temperature within a thermoneutral zone for 4 h (Kendall *et al.*, 2007). Cows will be bathed twice daily during those months in which their THI threshold exceeded. Bathing water requirements (Table 2) were informed by field data. The entire body (except the face, head and ears) was wet to the skin. However, cows bathed in

natural sources are immersed in water up to the stomach. This additional cooling was not considered. Water temperature was assumed to be 22°C (Kendall *et al.*, 2007).

Women are primarily responsible for bathing the cows. Bathing is currently conducted for hygiene reasons. Those cows considered more valuable (i.e. Jersey) are bathed daily (data not shown). As such, they are less dirty than those bathed at longer intervals (i.e. Local). Therefore, the variation in bath duration is likely a function of time since bathing. Indeed, the motorized pump would be expected to be a labour-saving device. However, as bathing is conducted for hygiene purposes, farmers are likely to be more thorough when using the pump as the drudgery of taking water to the animal (or vice versa) is removed. The time required to bath (including time required to take the cow to water or vice versa) and installation costs are shown in Table 2. Time required did not vary by herd size and was considered additive. The price of women's labour was Rs. 18.56/h assuming 8 h work/day (Government of Odisha, 2012).

Naturally occurring water sources could be freely accessed. Labour was the only operating cost associated with naturally occurring sources. Installation costs of the well and motorized pumps (installed 2006–11) were adjusted (against a base of 2004–05) to account for inflation (Reserve Bank of India, 2013). A marginal installation cost was not included as costing considers the total relief strategy (i.e. 2 baths/day). Indeed, a marginal cost could account for baths conducted at other times of the year. However, such bathing is not related to heat stress management. A marginal installation cost risked underestimating the cost of the strategy. Field data indicated the lifetime of the pump to be half of that of the well.

#### Heat stress relief strategy: fanning

The average cost of ceiling fan installation was Rs. 1838.68. Households estimated fan electricity requirement (Rs. 55.5/month). Households fanning cows (*n* = 17) used fan for an average of 19.82 h/day for 6.72 months. The model assumed households would continue to use fan for an additional 2 months before thresholds are reached. Hourly usage was assumed to remain as currently practiced. Fan lifetime was

**Table 3** The cost of feeding a lactating dairy cow in Odisha, India under normal conditions (N) and warming scenarios of +2°C (N + 2) and +4°C (N + 4) as expected during climate change

Warming scenario	Effect on feed cost	Feed cost (Rs./cow per day)		
		Jersey (n = 87)	1/4 Jersey (n = 18)	Local (n = 16)
N	—	98.01	78.00	65.77
N + 2	+10%	107.81	85.80	72.35
	+15%	112.71	89.70	75.64
N + 4	+20%	117.61	93.60	78.92
	+30%	127.41	101.40	85.50

Jersey = high-grade Jersey crossbreed; 1/4 Jersey = low-grade Jersey crossbreed; Local = indigenous (non-descript) breed.

**Table 4** The number of temperature–humidity index units above the heat stress threshold during a 300-day lactation under normal conditions (N) and warming scenarios of +2°C (N + 2) and +4°C (N + 4) as expected during climate change under no adaptation and the adaptations of shading, bathing and fanning in Odisha, India

Warming scenario	Breed	No adaptation	Shading			Bathing (×2)	Fanning and shading	
			Tree	Modern shed	Traditional shed		Modern shed	Traditional shed
N	Jersey	551 (55.3)	398 (39.9)	245 (24.6)	92 (9.2)	169 (16.9)	169 (16.9)	46 (4.6)
	1/4 Jersey	245 (6.0)	92 (2.2)	0	0	0	0	0
	Local	92 (0.7)	0	0	0	0	0	0
N + 2	Jersey	1254 (125.8)	1009 (101.2)	795 (79.7)	612 (61.4)	704 (70.6)	704 (70.6)	536 (53.7)
	1/4 Jersey	795 (19.4)	612 (14.9)	459 (11.2)	306 (7.5)	383 (9.3)	383 (9.3)	230 (5.6)
	Local	612 (4.3)	459 (3.3)	306 (2.2)	153 (1.1)	230 (1.6)	230 (1.6)	92 (0.7)
N + 4	Jersey	2020 (202.6)	1775 (178.0)	1530 (153.5)	1285 (128.9)	1408 (141.2)	1408 (141.2)	1163 (116.6)
	1/4 Jersey	1530 (37.3)	1285 (31.4)	1040 (25.4)	826 (20.2)	933 (22.8)	933 (22.8)	735 (17.9)
	Local	1285 (9.1)	1040 (7.4)	826 (5.9)	643 (4.6)	735 (5.2)	735 (5.2)	566 (4.0)

Jersey = high-grade Jersey crossbreed; 1/4 Jersey = low-grade Jersey crossbreed; Local = indigenous (non-descript) breed.

The quantity of milk lost to heat stress (kg/cow per 300-day lactation) is provided in brackets.

assumed to be 5 years. Two fans are required to match the lifetime of the shed.

Fan wind speed was assumed to be >2 m/s (Little and Campbell, 2010). No studies could be identified investigating fanning as a single treatment against a THI. However, fanning will reduce the rate of rectal temperature increase by half compared with a shade-only group (Berman *et al.*, 1985). Thus, it was assumed that fanning would increase THI threshold by 0.5 units. The shading effect from the shed was assumed additive.

#### State-level impact

A state-level THI index was created (as outlined above) with the use of district-level data (Government of Odisha, 2013). Each district was weighted equally and the THI load experienced by different breeds was determined. Breed definitions (exotic *v.* indigenous) and productivity levels were derived from census data (Government of India, 2014). Milk production was 6.15 and 1.47 kg/cow per day for exotic and indigenous cows, respectively (Government of India, 2014). The 1/4 Jersey was not included.

The threshold values and rates of decline used for the Jersey and indigenous breed (Table 1) were used at a state level. Milk loss was calculated at the cow level (assuming 300-day

lactation) and extrapolated across the lactating population composed of 325 000 exotic and 1 397 000 indigenous cows (Government of India, 2014). Adaptation was not considered.

#### Feed analysis

Feed cost was determined from field data. Farmers provided the quantities and price of each ingredient throughout the year. The average daily cost of feeding during lactation informed the model (Table 3). Milk incomes (without adaptation) were reduced by feed cost. Expected price increases were based on estimates of future feed deficits (Indian Grassland and Fodder Research Institute, 2013).

## Results

#### Temperature–humidity index load and lost milk production

The THI load and quantity of milk lost to heat stress is provided in Table 4. The combined treatment of traditional shed and fan is most effective at preventing milk losses. None of the relief strategies completely prevent heat stress negatively affecting milk yields.

By modelling actual lactation length, the THI load experienced by the 1/4 Jersey and Local breeds was reduced due to



**Table 5** The amount of income (per 300-day lactation) the households received from milk sales without adaptation under normal conditions (N) and warming scenarios of +2°C (N + 2) and +4°C (N + 4) as expected during climate change in Odisha, India

Change in income (%)												
				Shading			Bathing			Fanning and shading		
		No adaptation (Rs./cow per 300-day lactation)		Papaya trees	Mango tree	Modern shed	Traditional shed	Naturally occurring	Well	Motorized pump	Modern shed	Traditional shed
Warming scenario	Breed											
	Jersey	23 922		-1.04	1.08	-1.53	-3.50	-7.93	-13.75	-13.38	-2.51	-4.71
	1/4 Jersey	17 360		-2.64	0.28	-4.69	-9.00	-21.14	-31.50	-14.12	-6.84	-11.16
N + 2	Local	8042		-6.40	-0.10	-11.33	-20.65	-17.24	-36.52	-53.23	-15.97	-25.29
	Jersey	22 643 (-5.35)		-0.36	1.88	-0.39	-2.23	-13.91	-20.34	-21.16	-2.10	-4.06
	1/4 Jersey	17 117 (-1.40)		-2.60	0.36	-4.52	-8.51	-25.34	-36.42	-15.97	-7.57	-11.55
N + 4	Local	7975 (-0.83)		-6.36	0.002	-11.08	-20.23	-28.54	-50.16	-82.73	-17.91	-27.09
	Jersey	21 249 (-11.17)		-0.38	2.01	-0.15	-1.58	-14.29	-21.14	-22.02	-1.71	-3.14
	1/4 Jersey	16 792 (-3.28)		-2.49	0.53	-4.20	-8.11	-34.47	-46.99	-20.18	-7.23	-11.17
	Local	7889 (-1.91)		-6.28	0.15	-10.95	-20.15	-46.31	-71.48	-128.39	-17.84	-27.06

Jersey = high-grade Jersey crossbreed; 1/4 Jersey = low-grade Jersey crossbreed; Local = indigenous (non-descript) breed. The percentage change in income following implementation of relief strategies compared with no adaptation is also provided. Brackets indicate the percentage of income lost without adaptation under warming scenarios N + 2 and N + 4 compared with income received under N conditions from milk sales.

their shorter lactation period (Supplementary Table S1). However, the quantity of milk lost is 15%, 20% and 47% larger for the Jersey, 1/4 Jersey and Local breeds, respectively (compared with the 300-day lactation) (Supplementary Table S3). This is due to the higher daily yields being susceptible to greater loss (albeit for a protracted period).

#### Heat stress relief strategy costings

Table 5 provides milk income under conditions of heat stress without adaptation. The percentage change in income after accounting for the cost of the relief strategies is also provided. The table indicates that bathing reduces income levels the most across all breeds. Even with relief the income received will decline under climate change.

#### State-level impact

Table 6 indicates that as climate change intensifies the indigenous cow population will lose significantly less milk than the exotic population. Heat stress will cost a total of Rs. 972.76 million. The majority of loss incurred is from the exotic cow population (89.87% or Rs. 874.23 million).

#### Feed analysis

Currently, the Jersey provides only marginal additional income compared with the 1/4 Jersey (Rs. 559 or 10.20%) (Figure 2). However, as climate change intensifies the 1/4 Jersey provides Rs. 33 347 (19.71%) more income than the Jersey. The profitability of the Jersey declines to such an extent that it only provides Rs. 787 (4.64%) more income than the indigenous breed.

## Discussion

#### The temperature–humidity index and temperature–humidity index load

Heat stress is a serious constraint as relief will be required for even the most tolerant breeds. The relief strategies demonstrate varying levels of effectiveness. At present, fanning and/or an additional bath is largely unnecessary for the less-susceptible breeds as no additional cooling benefit is provided that could not be achieved with shading. The indigenous (non-descript) cow is an extreme example as its apparent heat tolerance ensures that tree shade is currently sufficient to prevent heat stress. Thus, it is necessary to match relief strategies to the breed's heat stress susceptibility. Susceptible breeds require more drastic strategies to protect productivity.


#### Lost milk production

The coupled treatment of fanning and the traditional shed appear most effective at reducing losses in milk production. However, even with relief the quantity of milk lost never returns to zero. Although the quantities of milk lost appear relatively minor, the total yields are themselves small. Lost production represents a serious concern for households reliant on milk for income and food security. For example,

**Table 6** The temperature–humidity index (THI) load experienced by dairy cows under normal conditions (N) and warming scenarios of +2°C (N + 2) and +4°C (N + 4) as expected during climate change in Odisha, India

	Warming scenario	Exotic	Indigenous
THI load	N	61	0
	N + 2	520	61
	N + 4	1101	551
Milk losses (kg/cow per 300-day lactation)	N	8.22	0
	N + 2	70.04	0.43
	N + 4	148.29	3.89
State-level milk losses (tonnes/300-day lactation)	N	2670.13	0
	N + 2	22 761.77	601.29
	N + 4	48 193.66	5431.34

Exotic = *Bos taurus* crossbreed; Indigenous = *Bos indicus* (non-descript) breed.  
The quantity of milk lost at cow and state level is also provided.

**Figure 2** Income remaining after accounting for the effects of heat stress and increased feed costs under conditions of climate change.

without relief under the warming scenario N+4, Jersey cows will lose 14.8% of total yield. The most effective relief strategy (fanning and traditional shed) examined was able to reduce this loss to 8.5%.

This investigation is limited by the assumption that heat stress has a linear effect on milk yield. Indeed, the literature indicates a linear relationship (see Igono *et al.*, 1992; Ravagnolo *et al.*, 2000; Bouraoui *et al.*, 2002). However, care must be taken in interpreting the results as it is unlikely that the rate of decline experienced by cows under different warming scenarios will remain constant. Yet, yawning gaps in knowledge regarding the susceptibility of low producing cows to heat stress and the effectiveness of heat stress relief strategies in a smallholder context limit the extent to which this aspect can be assessed. This study is also underpinned by very small sample sizes which are taken as representative. Thus, this investigation is a proof of concept, indicative of an urgent need for further robust locally relevant field-based experimentation.

#### Relief strategy costings

The usefulness of relief strategies in supporting income varies based on the overall income derived from milk sales. For example, mango tree shade can increase total income beyond no relief simulations for the Jersey cow. Yet, for the

lower-yielding breeds the cost of relief often reduces income below what could be achieved if the farmer was to do nothing. The poor response to relief makes the capital-intensive strategies unprofitable. However, this model does not consider the effects of heat stress over time. Heat-stressed cows may be less productive in future lactations, exhibit poor reproductive performance and risk mortality (Thornton *et al.*, 2009).

In mitigating heat stress, households will face trade-offs. The primary trade-off is the additional labour to implement strategies (particularly bathing). Although this has been accounted for by the labour cost, additional trade-offs may affect uptake that need further investigation. For example, bathing may use water required by agriculture. Furthermore, the water footprint of milk is already greater than that of cereals with equivalent nutritional value (Hoekstra, 2012). Questions must be asked as to whether the use of water for heat stress relief will ensure the greatest benefit in terms of food security.

#### State-level impact

Heat stress is a very serious constraint for the development of Odisha's dairy sector if it is to be based on exotic breeds. The financial losses associated with the more numerous non-descript cows is relatively minor compared with the losses associated with the smaller crossbred population.

### Feed analysis

The coupling of impact pathways (i.e. heat stress and increased feed price) suggests that a holistic, production system-wide approach to climate change adaptation will be required. Thus, it is not appropriate to draw conclusions regarding the comparative importance of heat stress *v.* feed availability, particularly as intake is reduced during heat stress (Bouraoui *et al.*, 2002).

The heat-stress-induced yield declines identified are less severe than other Indian dairy sector heat stress models (namely, Upadhyay *et al.*, 2007) due to differences in the use of climate data. Upadhyay *et al.* (2007) calculated a THI based on weekly average temperature. Thus, the thresholds will be exceeded more often than was noted in the current investigation. Furthermore, the author assumed that indigenous and crossbred cows will experience the same rate of decline (0.77 l/cow per THI unit).

Heat stress modelling has also been conducted on dairy herds in the United States. St-Pierre *et al.* (2003) found high levels of abatement effective at limiting the cost of heat stress (St-Pierre *et al.*, 2003). However, the author does not consider any aspects of Animal Genetic Resources (AnGR). Furthermore, the yields examined were significantly higher than yields considered by this investigation. Thus, the response to abatement is much more pronounced. Key and Sneeringer (2014) modelled climate change impacts on US dairy farm efficiency. However, the author does not consider adaptation. Therefore, the current investigation is the first to illustrate the importance of AnGR in the selection of heat stress relief strategy.

This study illustrates the complexities of design and implementation of climate change adaptation. Although it is unlikely that the production levels of local breeds will be acceptable to farmers, the model shows that cows with high levels of exotic genetics will experience heat stress for the majority of the year. From an economic perspective, only those relief strategies that can be implemented at a low capital cost (e.g. tree shade) were found to be financially viable as other options reduced income to below what could be achieved without relief. However, to do nothing would compromise animal welfare and future productive capacity.

Therefore, the model questions the usefulness of national breeding policy in conditions of climate change. Heat stress will make milk production more challenging. However, to advocate the use of heat-stress-susceptible higher-yielding exotic breeds will limit sector sustainability as farmers will be required to invest in heat stress relief for little economic benefit. Yet, failure to do so will limit long-term productive capacity. Thus, hot-climate dairying projects and programmes need to take a more holistic approach to adaptation and consider AnGR as a means of improving dairy sector sustainability under conditions of climate change.

### Acknowledgements

The authors thank Dr Soumya Pani from the Orissa University of Agriculture and Technology, the secretary of the Women's Federation Sabita Sethy and the farmers.

### Supplementary material

To view supplementary material for this article, please visit <https://doi.org/10.1017/S1751731116002706>

### References

- Berman A 2011. Invited review: are adaptations present to support dairy cattle productivity in warm climates? *Journal of Dairy Science* 94, 2147–2158.
- Berman A, Folman Y, Kaim M, Mamen M, Herz Z, Wolfenson D, Arieli A and Graber Y 1985. Upper critical temperatures and forced ventilation effects for high-yielding dairy cows in a subtropical climate. *Journal of Dairy Science* 68, 1488–1495.
- Bouraoui R, Lahmar M, Abdesslem M, Djemali M and Belyea R 2002. The relationship of temperature-humidity index with milk production of dairy cows in a Mediterranean climate. *Animal Research* 51, 479–491.
- Bryant JR, Lopez-Villalobos N, Pryce JE, Holmes CW and Johnson DL 2007. Quantifying the effect of thermal environment on production traits in three breeds of dairy cattle in New Zealand. *New Zealand Journal of Agricultural Research* 50, 327–338.
- Cunningham K 2009. Connecting the milk grid: smallholder dairy in India. In *Millions fed: proven successes in agricultural development* (ed. DJ Spielman and R Pandya-Lorch), pp. 117–125. International Food Policy Research Institute, Washington, DC, USA.
- Datta TN, Shrestha and Chokkalingam G 2015. Agricultural land and bovine population in India: a critical review of agricultural census data. *Journal of Rural Development* 34, 167–186.
- Government of India 2006. Basic husbandry statistics. Government of India, New Delhi, India.
- Government of India 2013. National Livestock Policy 2013. Government of India, New Delhi, India.
- Government of India 2014. Basic animal husbandry and fisheries statistics. Government of India, New Delhi, India.
- Government of Odisha 2012. Labour and ESI Department: notification. Government of Orissa, Cuttack, India.
- Government of Odisha 2013. Odisha agricultural statistics 2011–2012. Directorate of Agriculture and Food Production, Bhubaneswar, India.
- Heffernan C, Salman M and York L 2012. Livestock infectious disease and climate change: a review of selected literature. *CAB Reviews* 7, 1–26.
- Hoekstra AY 2012. The hidden water resource use behind meat and dairy. *Animal Frontiers* 2, 3–8.
- Igono MO, Bjotvedt G and Sanford-Crane HT 1992. Environmental profile and critical temperature effects on milk production of Holstein cows in desert climate. *International Journal of Biometeorology* 36, 77–87.
- Indian Grassland and Fodder Research Institute 2013. Vision 2050. Indian Grassland and Fodder Research Institute – Indian Council of Agricultural Research, New Delhi, India.
- International Farm Comparison Network 2015. Calculator. Retrieved on 29 June 2015 from <http://www.ifcnetwork.org/en/calculator.php>.
- Keister ZO, Moss KD, Zhang HM, Teegerstrom T, Edling RA, Collier RJ and Ax RL 2002. Physiological responses in thermal stressed Jersey cows subjected to different management strategies. *Journal of Dairy Science* 85, 3217–3224.
- Kendall PE, Verkerk GA, Webster JR and Tucker CB 2007. Sprinklers and shade cool cows and reduce insect-avoidance behavior in pasture-based dairy systems. *Journal of Dairy Science* 90, 3671–3680.
- Key N and Sneeringer S 2014. Potential effects of climate change on the productivity of U.S. dairies. *American Journal of Agricultural Economics* 96, 1136–1156.
- Little S and Campbell J 2010. Cool cows: shade, sprinklers and fans on dairy farms. Dairy Australia, Southbank, VIC, Australia.
- Mohapatra AK, Swain SK, Dash AK and Behera D 2015. Microenvironment of a model bullock shed in different seasons compared to conventional shed and open shelter in tropical humid climatic condition of coastal Odisha. *Animal Science Reporter* 9, 22–27.
- Moran J 2005. Diet and milk production. In *Tropical dairy farming: feeding management for small holder dairy farmers in the humid tropics* (ed. J Moran), pp. 159–170. Landlinks Press, Kyabram, VIC, Australia.

- O'Brien K, Leichenko R, Kelkar U, Venema H, Aandahl G, Tompkins H, Javed A, Bhadwal S, Barg S, Nygaard L and West J 2004. Mapping vulnerability to multiple stressors: climate change and globalization in India. *Global Environmental Change* 14, 303–313.
- Pathak PS and Devakumar C 2011. Carrying capacity of Indian agriculture. National Academy of Agricultural Sciences, New Delhi, India.
- Ravagnolo O, Misztal I and Hoogenboom G 2000. Genetic component of heat stress in dairy cattle, development of heat index function. *Journal of Dairy Science* 83, 2120–2125.
- Reena K, Triveni D, Patel BHM, Amitava D, Chandran PC, Barari SK, Asit C and Bharat B 2014. Effect of shade materials on microclimate of crossbred calves during summer. *Veterinary World* 7, 776–783.
- Reserve Bank of India 2013. Database on Indian economy: monthly wholesale price index – inflation. Retrieved on 22 September 2013 from <http://dbie.rbi.org.in/DBIE/dbie.rbi?site=home>.
- Santana MLJ, Pereira RJ, Bignardi AB, Filho AEV, Menéndez-Buxadera A and El Faro L 2015. Detrimental effect of selection for milk yield on genetic tolerance to heat stress in purebred Zebu cattle: genetic parameters and trends. *Journal of Dairy Science* 98, 9035–9043.
- Sharma A, Tyagi S, Singh U and Mandal DK 2011. Vision 2030: Project Directorate on Cattle. Indian Council of Agricultural Research, Meerut Cantt, India.
- Sirohi S and Michaelowa A 2007. Sufferer and cause: Indian livestock and climate change. *Climatic Change* 85, 285–298.
- Smith DL, Smith T, Rude BJ and Ward SH 2013. Short communication: comparison of the effects of heat stress on milk and component yields and somatic cell score in Holstein and Jersey cows. *Journal of Dairy Science* 96, 3028–3033.
- St-Pierre NR, Cobanov B and Schnitkey G 2003. Economic losses from heat stress by US livestock industries. *Journal of Dairy Science* 86 (Suppl), E52–E77.
- Thornton PK, van de Steeg J, Notenbaert A and Herrero M 2009. The impacts of climate change on livestock and livestock systems in developing countries: a review of what we know and what we need to know. *Agricultural Systems* 101, 113–127.
- Upadhyay RC, Mohini M, Kansal VK, Singh SV, Ashutosh A, Sirohi SK and Kumar S 2007. Final report of the Network Project on Climate Change. National Dairy Research Institute, Karnal, India.
- Zimbleman RB, Rhoads RP, Rhoads ML, Duff GC, Baumgard LH and Collier RJ 2009. A re-valuation of the impact of temperature humidity index (THI) and black globe humidity index (BGHI) on milk production in high producing dairy cows. In the 24th Annual Southwest Nutrition and Management Conference, 9–11 March 2009, Reno, NV, USA, pp. 158–168.