

# *Pre-agricultural plant management in the uplands of the central Zagros: the archaeobotanical evidence from Sheikh-e Abad*

Article

Accepted Version

Whitlam, J., Bogaard, A., Matthews, R. ORCID: <https://orcid.org/0000-0002-8316-4312>, Matthews, W. ORCID: <https://orcid.org/0000-0002-7408-6885>, Mohammadifar, Y., Ilkhani, H. and Charles, M. (2018) Pre-agricultural plant management in the uplands of the central Zagros: the archaeobotanical evidence from Sheikh-e Abad. *Vegetation History and Archaeobotany*, 27 (6). pp. 817-831. ISSN 0939-6314 doi: <https://doi.org/10.1007/s00334-018-0675-x> Available at <https://centaur.reading.ac.uk/68880/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

To link to this article DOI: <http://dx.doi.org/10.1007/s00334-018-0675-x>

Publisher: Springer

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in

the [End User Agreement](#).

[www.reading.ac.uk/centaur](http://www.reading.ac.uk/centaur)

## **CentAUR**

Central Archive at the University of Reading

Reading's research outputs online

## Tables

**Table 1** Radiocarbon determinations from Sheikh-e Abad

<b>Trench</b>	<b>Deposit type</b>	<b>Material</b>	<b>Lab no.</b>	<b>Conventional date BP</b>	<b>Intercept with calibration curve</b>	<b>Calibrated date BC 2-sigma (95.4%)</b>	<b><math>\delta^{13}\text{C}</math> (PDB)</b>
1	Ash layer	Charred plant material (seed)	Beta-258647	10130±60	9810	10100–9450	n/a
1	Ash layer	Charred plant material (wood charcoal)	Beta-267508	9970±50	9400	9470–9730 and 9680–9300	-23.8‰
1	Ash layer	Charred plant material (seed)	Beta-267509	9730±60	9240	9280–9410	-22.7‰
2	Ash layer	Charred plant material (seed)	Beta-258646	8810±60	7960	8230–7730	-23.2‰
3	Building 2 (floor)	Animal bone: sheep/goat	Beta-258648	8600±40	7590	7640–7580	-18.9‰

**Table 2** Records of the frequency and abundance of all recorded taxa across Trenches 1, 2 and 3 at Sheikh-e Abad. ‘cf.’ denotes a lower certainty of identification and brackets the uncertainty of cultivated/domestic status. \* indicates taxa interpreted as possible “arable weeds” based on correspondence analysis as discussed in text

Taxa	Trench 1 (5 samples) 126 litres		Trench 2 (9 samples) 441 litres		Trench 3 (27 samples) 1326 litres	
	Sum	Ubiquity (%)	Sum	Ubiquity (%)	Sum	Ubiquity (%)
<b>Cereals</b>						
<i>Hordeum vulgare</i> hulled	.	.	14	78	2	7
<i>Hordeum vulgare</i>	.	.	12	56	1	4
<i>Hordeum vulgare/ spontaneum</i> , rachis	.	.	1	11	3	11
<i>Triticum monococcum/dicoccum</i>	.	.	3	11	.	.
<i>Triticum</i> 'new-type'/ <i>dicoccum</i> , glume base	.	.	.	.	2	4
Cereal indet.	.	.	3	22	1	4
<b>Pulses</b>						
<i>Lens</i> sp. Mill.	.	.	74	100	2	7
<i>Pisum</i> sp. L.	.	.	12	67	.	.
<i>Pisum</i> sp. L. (small-seeded)	.	.	7	33	.	.
<i>Trigonella</i> sp. L.	.	.	250	67	1	4
<i>Vicia ervilia</i> (L.) Willd.	.	.	57	100	1	4
<i>Vicia/Lathyrus</i> spp. L.	.	.	24	67	1	4
<i>Vicia</i> spp. L.	.	.	11	56	.	.
Large-seeded legume indet.	(1)	(20)	114	100	10	26
<b>Large-seeded grasses</b>						
<i>Hordeum spontaneum</i> C. Koch.	.	.	12	56	.	.
<b>Medium(large)-seeded grasses</b>						
<i>Aegilops</i> sp. L., spikelet base	5	20	22	11	.	.
<i>Piptatherum holciforme</i> (M.Bieb.) Roem. and Schult.	11	80	137	100	1	4
<i>Bromus</i> spp. Scop.	.	.	29	56	1	4
<b>Medium(medium)-seeded grasses</b>						
<i>Hordeum</i> (non-spontaneum) sp. L.	1	20	1	11	.	.
<i>Stipa</i> spp. L.	10	80	530	100	12	30
<i>Stipa</i> spp. L., awn	5	80	5	44	4	15
<i>Taeniatherum caput-medusae</i> (L.) Nevski.	5	40	245	100	2	7
<i>Taeniatherum caput-medusae</i> (L.) Nevski., rachis	1	20	230	89	2	4
<b>Medium(small)-seeded grasses</b>						
cf. <i>Eremopyrum</i> sp. (Ledeb.) Jaub. Et Spach.	.	.	3	11	.	.
<i>Lolium perenne/rigidum</i>	50	100	556	89	.	.
<b>Small-seeded grasses</b>						
<i>Crypsis</i> sp. Ait.	15	60	1064	67	3	11
<i>Poa</i> spp. L.	124	80	3296	67	1670	67
<i>Sporobolus</i> sp. R. Br.	.	.	72	33	1	4
<b>Medium-seeded Poaceae indet.</b>	<b>41</b>	<b>100</b>	<b>585</b>	<b>100</b>	<b>9</b>	<b>30</b>

<b>Small-seeded Poaceae indet.</b>	<b>5</b>	<b>20</b>	.	.	.	.
<b>Nut/fruit</b>						
<i>Pistacia</i> sp. L., nutshell	.	.	20	56	1	4
<i>Pistacia</i> indet., nutshell	1	20	13	56	6	19
<i>Prunus amygdalus</i> Batsch., nutshell	5	80	12	100	.	.
<i>Prunus</i> sp. L., nutshell	.	.	.	.	.	.
Nutshell spp./indet	2	40	13	89	7	26
<b>Wild/weed</b>						
<i>Coronilla</i> sp. L. *	2	20	4	22	.	.
Small-seeded legume indet.	66	100	319	100	10	11
<i>Chenopodium</i> spp. L.	41	100	13	56	.	.
<i>Bellevialia</i> sp. Lapeyr. *	2	40	30	89	1	4
Asteraceae spp.	.	.	10	33	.	.
<i>Heliotropium</i> sp.* L.	.	.	3	22	1	4
<i>Brassica/Sinapis</i> sp. L.	4	40	101	100	2	7
<i>Euclidium syriacum</i> (L.) R. Br.	.	.	44	78	2	7
<i>Neslia paniculata</i> (L.) Desv.	1	20	18	78	.	.
Brassicaceae spp./indet	4	60	17	89	2	7
Caryophyllaceae sp.	.	.	.	.	1	4
<i>Silene</i> sp. L.*	2	40	15	56	.	.
<i>Vaccaria pyramidata</i> Medik.*	2	20	12	56	.	.
Convolvulaceae sp.	.	.	2	22	.	.
<i>Bolboschoenous glaucus</i> (Lam.) S.G. Smith	49	100	362	100	9	30
<i>Carex</i> sp. L.	.	.	1	11	1	4
Cyperaceae indet.	11	80	89	56	8	22
Dipsacaceae/Apiaceae indet.	.	.	14	33	.	.
<i>Hypericum</i> sp. L.	.	.	253	44	2	7
<i>Ajuga</i> sp. L.	.	.	14	67	.	.
Lamiaceae spp.	11	60	5	44	1	4
<i>Fumaria</i> sp. L.*	1	20	3	11	.	.
<i>Glaucium</i> sp. Mill.*	2	40	208	78	.	.
<i>Papaver</i> spp. L.*	6	40	689	67	12	30
<i>Polygonum aviculare</i> L.	2	40	.	.	.	.
<i>Rumex</i> sp. L.	2	40	.	.	.	.
Polygonaceae indet	2	20	.	.	.	.
<i>Adonis</i> sp. L.*	.	.	15	67	.	.
<i>Galium</i> sp. L.*	1	20	10	33	.	.
<i>Hyoscyamus</i> sp. L.	.	.	2	22	2	4
cf. <i>Thymelea</i> sp. Mill.*	.	.	1	11	.	.
Pod types	2	40	30	78	6	22
Seed/fruit types	11	60	1007	100	10	30
<b>Culm</b>	<b>1</b>	<b>20</b>	<b>6</b>	<b>56</b>	.	.
<b>Animal Dung (ml)</b>	.	.	<b>1.2</b>	<b>33</b>	.	.
<b>Rodent Pellet</b>	.	.	<b>53</b>	<b>67</b>	.	.
<b>Average density (items per litre)</b>	<b>4.0</b>		<b>24.3</b>		<b>1.4</b>	
<b>Total sum</b>	<b>507</b>		<b>10724</b>		<b>1813</b>	

**Table 3** Classification of major wild plant taxa at Sheikh-e Abad based on their likelihood of surviving in dung (ranked from low to high) as discussed in text

Category	Taxa
Low	<i>Hordeum spontaneum</i> , <i>Bromus</i> spp., <i>Taeniatherum caput-medusae</i> , <i>Stipa</i> spp., medium-seeded grass indet., Brassicaceae indet.
Medium	<i>Lolium perenne/rigidum</i> , <i>Piptatherum holciforme</i> , <i>Bellevalia</i> sp., <i>Euclidium syriacum</i> , <i>Adonis</i> sp., <i>Ajuga</i> sp., <i>Neslia paniculata</i> , <i>Brassica/Sinapis</i> sp., Brassicaceae type 3, <i>Vaccaria pyramidata</i>
High	<i>Coronilla</i> sp., small-seeded legumes, <i>Bolboschoenus glaucus</i> , <i>Chenopodium</i> spp., <i>Galium</i> sp., Cyperaceae spp., <i>Heliotropium</i> sp., <i>Hyoscyomas</i> sp., Lamiaceae spp., <i>Hypericum</i> sp., <i>Poa</i> spp., <i>Crypsis</i> sp., <i>Sporobolus</i> sp., <i>Glaucium</i> sp., <i>Papaver</i> spp., <i>Silene</i> sp., small-seeded grass indet.