

Early hominins in north-west Europe: a punctuated long chronology?

Article

Supplemental Material

Creative Commons: Attribution-Noncommercial-No Derivative Works 4.0

Hosfield, R. ORCID: <https://orcid.org/0000-0001-6357-2805>
and Cole, J. (2018) Early hominins in north-west Europe: a
punctuated long chronology? *Quaternary Science Reviews*,
190. pp. 148-160. ISSN 0277-3791 doi:
10.1016/j.quascirev.2018.04.026 Available at
<https://centaur.reading.ac.uk/69498/>

It is advisable to refer to the publisher's version if you intend to cite from the
work. See [Guidance on citing](#).

To link to this article DOI: <http://dx.doi.org/10.1016/j.quascirev.2018.04.026>

Publisher: Elsevier

All outputs in CentAUR are protected by Intellectual Property Rights law,
including copyright law. Copyright and IPR is retained by the creators or other
copyright holders. Terms and conditions for use of this material are defined in
the [End User Agreement](#).

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online

Table 5: Winter and summer temperature estimates for Early and Middle Pleistocene British and German sites (with selected present day equivalents for reference). ¹Sensitivity tests on coleoptera-based MCR procedures suggest that winter temperature estimates are usually too warm (Pettitt and White, 2012: 35). ²The specific source of the palaeo-temperature estimates is not stated, but the fauna includes molluscs and ostracods; ³East Anglian data based on Met Office annual mean seasonal temperatures (1910–2016; http://www.metoffice.gov.uk/pub/data/weather/uk/climate/datasets/Tmean/date/East_Anglia.txt); ⁴Bilzingsleben data based on Deutscher Wetterdienst (German Weather Service) monthly mean January and July temperatures (1951–2017; Erfurt-Weimar station; <https://www.dwd.de/DE/leistungen/klimadatendeutschland/klarchivtagmonat.html>)

Site	T _{min} (°C)	T _{max} (°C)	Evidence ¹	Age (MIS)	Source
Early Pleistocene					
Happisburgh III (Bed E)	-3 – 0	+16 – +18	Coleoptera	Late 25 or late 21	(Ashton and Lewis, 2012; Parfitt et al., 2010)
Early Middle Pleistocene					
Pakefield (Bed Cii–Ciii)	-6 – +4	+17 – +23	Coleoptera	17 or later 19	(Ashton and Lewis, 2012; Coope, 2006b)
Boxgrove (Unit 4c & Freshwater Silt Bed ≈ Units 4b & 4c)	-4 – +4	+15 – +20	Ostracods (MOTR) & Herpetofauna (MCR)	13	(Ashton and Lewis, 2012; Holman, 1999; Holmes

					et al., 2010)
Happisburgh I (Organic Mud)	-11 – -3	+12 – +15	Coleoptera	13?	(Ashton and Lewis, 2012; Coope, 2006b)
High Lodge (Bed C1)	-4 – +1	+15 – +16	Coleoptera	13?	(Coope, 2006b)
Waverley Wood (Channel 2, Organic Mud)	-	+10 – +15	Coleoptera	13 or 15	(Coope, 2006b; Shotton et al., 1993)
Brooksby (Redland's Brooksby Channel)	-10 – +2	+15 – +16	Coleoptera	13 or 15	(Coope, 2006b)
Later Middle Pleistocene					
Barnham (Unit 5c; Holl)	-	+17 – +18	Herpetofauna	11c	(Holman, 1998)
Hoxne (Stratum D; Holla)	-10 – +6	+15 – +19	Coleoptera	11c	(Ashton et al., 2008a; Coope, 1993)
Bilzingsleben II	-0.5 – +3	+20 – +25	Mollusca & ostracods ²	11	(Mania, 1995; Mania and Mania, 2003)
Schöningen 13 II-4	-4 – -1	+16	Mollusca & ostracods	9	(Urban and Bigga, 2015)

Present Day Comparisons					
East Anglia (present day) ³	-0.7 – +6.9	+14.2 – + 18.0	-	-	-
Bilzingsleben (present day) ⁴	-3.3 – +2.0	+12.4 – +22.7	-	-	-