

Evaluation of 2-acetyl-1-pyrroline in foods, with an emphasis on rice flavour

Article

Accepted Version

Creative Commons: Attribution-Noncommercial-No Derivative Works 4.0

Wei, X., Handoko, D. D., Pather, L., Methven, L. and Elmore, J. S. ORCID: <https://orcid.org/0000-0002-2685-1773> (2017) Evaluation of 2-acetyl-1-pyrroline in foods, with an emphasis on rice flavour. *Food Chemistry*, 232. pp. 531-544. ISSN 0308-8146 doi: 10.1016/j.foodchem.2017.04.005 Available at <https://centaur.reading.ac.uk/69971/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

To link to this article DOI: <http://dx.doi.org/10.1016/j.foodchem.2017.04.005>

Publisher: Elsevier

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online

1 Evaluation of 2-acetyl-1-pyrroline in foods, with an emphasis on rice
2 flavour

3 Xuan Wei^a, Dody D. Handoko^b, Leela Pather^a, Lisa Methven^a, J. Stephen Elmore^{a*}

4 ^a Department of Food and Nutritional Sciences, University of Reading, Whiteknights,
5 Reading RG6 6AP, UK

6 ^b Indonesian Centre for Rice Research, Cikampek, Subang 41256, West Java, Indonesia

7

8 * Corresponding author. Tel.: +44 118 3787455; fax: +44 118 3787708.

9 E-mail address: j.s.elmore@reading.ac.uk (J.S. Elmore).

10

11 **ABSTRACT**

12 The popcorn-like aroma compound 2-acetyl-1-pyrroline (2-AP) is a key contributor to the
13 desirable aroma of fragrant rice and is also important in the aroma of other foods, such as
14 pandan leaf, popcorn and Mediterranean sausage. It can be formed enzymatically in the rice
15 grain as it grows and is also formed, as part of the Maillard reaction, when rice is heated. This
16 review examines the formation of 2-AP in rice and other foods, particularly its formation
17 during cooking, focusing on the importance of the Maillard reaction between reducing sugar
18 breakdown products and 1-pyrroline derived from the amino acids proline and ornithine. The
19 synthesis of 2-AP is discussed alongside the attempts that have been made to stabilise this
20 relatively unstable compound. The analysis of 2-AP by instrumental techniques, particularly
21 gas chromatography-mass spectrometry and gas chromatography-olfactometry, alongside the
22 use of sensory studies, is also discussed.

23

24 Keywords: 2-acetyl-1-pyrroline, 2-AP, flavour, rice, pandan, popcorn, Maillard reaction,
25 biosynthesis, analysis

26

27 **1. Introduction**

28 The IUPAC name of 2-acetyl-1-pyrroline (2-AP) is 1-(3,4-dihydro-2*H*-pyrrol-5-
29 yl)ethanone, its CAS number is 85213-22-5 and its FEMA (Flavor and Extract
30 Manufacturers Association) number is 4249. 2-AP was first identified in rice by
31 Buttery, Ling, and Juliano (1982), and is regarded as the most important aroma
32 compound in rice, especially fragrant rice (Buttery, Ling, Juliano, & Turnbaugh,
33 1983). In that study, 0.05 ppm 2-AP was described as popcorn-like and its odour
34 threshold in water was measured as 0.1 nL/L, while its odour threshold in air was
35 reported by Schieberle (1991) as 0.02 ng/L; this very low threshold makes it an
36 important contributor to a food's aroma when present. As well as rice, it is also a key
37 flavour compound in many cereal products, as well as some vegetable and animal
38 products (Adams & De Kimpe, 2006; Wakte, Zanan, Hinge, Khandagale, Nadaf, &
39 Henry, 2016).

40 Bioformation of 2-acetyl-1-pyrroline in both plants and microorganisms has been
41 studied and several types of bacteria are able to form this compound (see Part 3 of this
42 review). 2-Acetyl-1-pyrroline has also been shown to form in the Maillard reaction; it
43 can be formed from the reaction between proline and reducing sugars/sugar
44 degradation products upon heating (Schieberle, 1989).

45 Although there is a high commercial interest in 2-AP because of its desirable
46 sensory attributes, the instability of this compound is a significant problem for its
47 commercial application. Pure 2-AP will turn red and degrade within 10 minutes at

48 room temperature (Fang & Cadwallader, 2014), and there is significant short-term
49 reduction of 2-AP concentration in food products, such as popcorn (Schieberle, 1995)
50 and raw fragrant rice (Widjaja, Craske, & Wootton, 1996a).

51 The occurrence of 2-acetyl-1-pyrroline in food products, its bioformation and
52 thermal formation, synthesis, stabilisation, analysis and sensory evaluation will be
53 reviewed in this paper, with particular emphasis on the role of 2-AP in fragrant rice
54 aroma.

55

56 **2. Food sources of 2-acetyl-1-pyrroline**

57 *2.1. Rice*

58 Non-fragrant rice (long and medium grain *indicas* and short grain *japonicas*),
59 mainly grown in USA, Vietnam, Thailand and Australia, constitutes around 80% of
60 the world rice trade (Singh, Singh, & Khush, 2000). Major producers of fragrant rice
61 are India, Pakistan and Thailand. Most of the fragrant rice exported from India and
62 Pakistan is basmati, while fragrant jasmine rice is a major export of Thailand (Singh
63 et al., 2000). In 2010, Thailand was the biggest exporter of fragrant rice: 2.65 million
64 tonnes of jasmine rice were exported, followed by India (1.80 million tonnes basmati)
65 and Pakistan (1.05 million tonnes basmati) (Slayton & Muniroth, 2011).

66 The price of fragrant rice is much higher than that of non-fragrant rice. For
67 example, high-quality fragrant basmati rice has a three times higher price than high
68 quality non-fragrant rice. The commercial value of fragrant rice is higher than that of

69 non-fragrant rice, partly because fragrant rice varieties are relatively low yielding.
70 Fragrant rice is less resistant to disease and insect pests and is prone to high shedding,
71 leading to losses in yield (Berner & Hoff, 1986; Golam et al., 2011). It has been
72 shown that higher quality grains with stronger aromas are generated in crops grown in
73 drought and saline conditions (Yoshihashi, Nguyen, & Kabaki, 2004). These adverse
74 conditions do not favour high yields.

75 2-AP is the key discriminator between fragrant and non-fragrant rice and many
76 studies have focused on the concentration of 2-AP in different rice cultivars. 2-AP
77 concentrations in different fragrant cultivars vary substantially (Table 1). For example,
78 2-AP was present in milled Fowler Gourmet Aromatic rice (a US-grown aromatic
79 rice) at 999 $\mu\text{g}/\text{kg}$, while, in a set of five basmati samples, levels of 2-AP from 19
80 $\mu\text{g}/\text{kg}$ to 342 $\mu\text{g}/\text{kg}$ were measured (Bergman, Delgado, Bryant, Grimm, Cadwallader,
81 & Webb, 2000).

82 Milled rice (commonly referred to as white rice) is obtained from the milling of
83 brown rice to remove the outer bran layer. Whole rice grains are dehulled; then the
84 dehulled (brown) rice is milled twice. Generally, 20–22% of the rice grain is hull, and
85 another 8–10% is bran and embryo; therefore, the yield of milled rice is around 70%
86 (Singh et al., 2000). As can be seen in Table 1, in most cases more 2-AP is present in
87 brown rice compared to milled rice.

88 Caution should be applied when comparing data acquired by different authors. In
89 some cases 2-AP was measured in uncooked rice (e.g., Hopfer, Jodari, Negre-
90 Zakharov, Wylie, & Ebeler, 2016), while in other cases the rice was cooked before

91 analysis (e.g., Widjaja et al., 1996a; Widjaja, Craske, & Wootton, 1996b) and even
92 during analysis (e.g., Buttery et al., 1983). The effect of sample preparation on 2-AP
93 content in rice is covered in more detail in Part 7 of this review.

94 Soil and climate conditions during cropping can also influence 2-AP concentration
95 in rice cultivars. During cultivation, a dry climate or sandy soil with low moisture
96 retention can induce the fragrant rice cultivar Khao Dawk Mali 105 to produce more
97 2-AP (Yoshihashi et al., 2004). It appears that moisture during cultivation could be
98 one of the most important factors affecting 2-AP formation when rice grows.

99 Due to the instability of 2-AP, drying and storage of rice can also influence the 2-
100 AP content of the final product (Wongpornchai et al., 2004). The unstable nature of 2-
101 AP will be discussed in detail in Part 6 of this review.

102 *2.2. Pandan*

103 2-AP is an important component of pandan leaf; the aroma of 2-AP is often
104 described as pandan-like. Pandan plays an important role in south-east Asian cookery.
105 The leaf of this plant is often boiled with rice to enhance flavour. When boiled with
106 non-fragrant rice, it can provide the popcorn-like flavour associated with boiled
107 fragrant rice, allowing cheap non-fragrant cultivars to possess similar aroma to higher
108 value fragrant rice cultivars (Peter, 2006). The treatment of pandan leaf can affect 2-
109 AP content. The fresh or slightly withered leaf is normally torn into strips, tied in a
110 bunch and then boiled together with rice. The pandan leaves are removed from the
111 rice after cooking.

112 The concentration of 2-AP in pandan leaves ranges from 40 ± 10 to 450 ± 10
113 $\mu\text{g}/\text{kg}$ (Yahya, Lu, Santos, Fryer, & Bakalis, 2010). Dried and ground pandan leaves
114 were extracted in this study. However, those treatments disrupted the papillae
115 structure in epidermal cells on the surface of the pandan leaves. 2-AP is contained in
116 the papillae; therefore, a proportion of 2-AP is lost during drying and grinding.

117 *2.3. Cereal products*

118 2-Acetyl-1-pyrroline has also been detected in cooked cereal-based products.
119 Wheat bread crusts contain around $75 \mu\text{g}/\text{kg}$ 2-AP compared to $1\text{--}4 \mu\text{g}/\text{kg}$ in
120 sourdough processed rye bread (Schieberle & Grosch, 1987). Popcorn-like aroma
121 compound 2-AP is, unsurprisingly, present in popcorn. However, in popcorn 2-
122 acetyltetrahydropyridine and 2-propionyl-1-pyrroline also contribute roasty and
123 popcorn-like flavour. The alkyl side chains of those compounds are short; only one or
124 two carbon atoms length. In contrast, 2-butanoyl-1-pyrroline and 2-hexanoyl-1-
125 pyrroline, compounds with similar structure but with longer alkyl side-chains, do not
126 possess roasty or popcorn-like aroma (Schieberle, 1991).

127 2-AP was also identified in a cereal coffee brew at $8 \mu\text{g}/\text{L}$ and contributed intense
128 popcorn-like odour attributes when analysed by gas chromatography-olfactometry
129 (Majcher, Klensporf-Pawlik, Dziadas, & Jeleń, 2013). The cereal coffee was a roasted
130 mixture of 40% barley, 25% rye, 25% chicory, and 10% sugar beet.

131 *2.4. Other foods*

132 2-AP has also been detected in non-cereal-based food. A high concentration of 2-

133 AP of up to 750 µg/kg was found on the surface of Mediterranean dried sausages,
134 while values at the core were up to 100 µg/kg. *Penicillium nalgiovense*, the dominant
135 mould species present, was shown to synthesise 2-AP during sausage processing
136 (Stahnke, 2000). Using gas chromatography-olfactometry (GC-O), Blank, Devaud,
137 Fay, Cerny, Steiner, and Zurbriggen (2001) identified 2-AP as a key contributor to the
138 aromas of both Parma ham and Italian-type salami. They described the compound as
139 it eluted from the GC column as having a ‘roasty’ aroma in the Parma ham and a
140 ‘roasty, popcorn’ aroma in the salami.

141 2-AP was also isolated in Manuka honey at concentrations of 80–450 µg/kg. It
142 was formed from methylglyoxal, which is responsible for the antibacterial activity in
143 Manuka honey. Reaction of methylglyoxal with proline through the Strecker reaction
144 can form 2-AP (Ruckriemen, Schwarzenbolz, Adam, & Henle, 2015). In addition, 2-
145 AP was also isolated from two kinds of cooked edible fungus: huitlacoche and austern
146 pilzen (Lizarraga-Guerra, Guth, & Lopez, 1997), but the compound was mistakenly
147 identified as 2-acetyl-2-pyrroline (Adams & De Kimpe, 2006). The importance of 2-
148 AP in mushroom (*Agaricus bisporus*) aroma increased significantly as a result of pan-
149 frying (Grosshauser & Schieberle, 2013), its concentration rising from 0.4 to 5.3
150 µg/kg. Similarly, 2-AP was also detected in both raw and roasted hazelnuts; a
151 significant increase of 2-AP concentration was observed, from trace levels (< 3 µg/kg)
152 to 85 µg/kg, when hazelnuts were roasted (Kiefl, Pollner, & Schieberle, 2013).

153 2-AP may not always make a positive contribution to food aroma. An undesirable
154 ‘mousy’ flavour in wetted raw pearl millet grits was attributed to 2-AP. Although 2-

155 AP concentration was not quantified in this study, it was implied that there was a
156 higher concentration of 2-AP in millet than in rice, which was reflected in the
157 difference in their odour quality (Seitz, Wright, Waniska, & Rooney, 1993).

158 2-AP has been identified and quantified in many food products. Table 2 shows
159 those foods other than rice where 2-AP has been quantified. Even at a very low
160 concentration, such as 3 µg/kg in milk chocolate (Liu, et al., 2015), this compound
161 can still be considered a key odorant. In a recent review, a comprehensive list of food
162 sources of 2-AP was provided, which included fruit and vegetables, fungi, cooked
163 meat and fish, dairy and egg products (Wakte et al., 2016).

164 *2.5. 2-AP as a flavouring*

165 Several patents have suggested that 2-AP could be applied as a food flavouring. A
166 food coating with a content of at least 40 ppb 2-AP, made from fragrant rice, was
167 applied to increase popcorn odour in several products (Richard, 2001). In distilled
168 alcoholic beverages, 0.2 to 200 ppb 2-AP contributed to a fragrant rice flavour (Asano
169 et al., 2000). 2-AP was included in GRAS 22 (Smith et al., 2005) and the average and
170 maximum levels for its addition to various food products have been summarised
171 (Adams & De Kimpe, 2006).

172

173 **3. Biological formation of 2-acetyl-1-pyrroline**

174 *3.1. Fragrant rice*

175 It was originally thought that 2-AP was only produced during the cooking of rice

176 *via* the Maillard reaction (Buttery et al., 1982). However, further research has shown
177 that 2-AP is produced by the rice plant, and is detected in the majority of plant tissues
178 (Sakthivel, Sundaram, Rani, Balachandran, & Neeraja, 2009; Sood & Siddiq, 1978;
179 Yoshihashi, Huong, & Inatomi, 2002). It is now generally accepted that although
180 some 2-AP in rice is produced during cooking, 2-AP is predominantly biosynthesised
181 in rice.

182 Yoshihashi (2002) reported that 2-AP cannot be formed during the cooking of
183 fragrant rice (when heated with or without water at 90 °C for 8, 10, 12, 14 min, the
184 concentration of 2-AP showed a slight decrease), nor in postharvest processes like
185 drying and storage. It can only be formed in the aerial parts of plants during growing
186 in paddy fields. In a later paper by the same author, excised callus (cells covering a
187 plant wound) and seedlings were floated on labelled amino acid (200 ppm ¹⁵N-
188 glycine, ¹⁵N-L-proline or 1-¹³C-L-proline; pH 5.5) solutions. After incubation at 27 °C
189 in darkness for 8 hours, increasing 2-AP concentrations were detected. Results
190 showed clearly that the labelled derivative was only found in seeding and callus
191 incubated with ¹⁵N-L-proline. This result indicated that one of the precursors in 2-AP
192 biosynthesis could be proline but not glycine and that the nitrogen source of 2-AP is
193 proline. On the other hand, because no labelled derivative was found in the 1-¹³C-L-
194 proline sample, the acetyl group in 2-AP could not be provided by proline
195 (Yoshihashi, Huong, & Inatomi, 2002).

196 It appears that moisture during cultivation could be one of the most important
197 factors affecting 2-AP formation when rice grows. 2-AP concentrations in fragrant

198 rice Khao Dawk Mali 105 from the Tung Kula Rong Hai region in north-east
199 Thailand, where there is a drought-prone climate with sandy soil, were much higher
200 than in the same rice grown in other areas of Thailand. Rice samples planted in clay
201 soil can retain moisture during growth, resulting in lower 2-AP concentrations than
202 those grown in sandy soil (Yoshihashi et al., 2004). Numerous studies have shown
203 that proline accumulation occurs in higher plants due to different environmental
204 stresses, such as drought, high salinity, high light and UV irradiation, heavy metals,
205 oxidative stress and in response to biotic stresses (Szabados & Savouré, 2010). For
206 example, Rhodes, Handa, and Bressan (1986) showed that proline will accumulate in
207 water-deficient plant cells through the glutamate pathway. In this study, tomato cells
208 adapted to water stress induced with polyethylene glycol (PEG); a tenfold increase of
209 proline synthesis was observed in the water-stressed cells. This research and that of
210 Yoshihashi et al. (2004) suggest that more 2-AP will be synthesised when rice is
211 grown in a dry climate, due to increased accumulation of its precursor proline.

212 In addition, a cool climate and early harvest could increase 2-AP concentration in
213 fragrant rice varieties. Between 1992 and 1994, three brown fragrant rice cultivars
214 from Japan (Hieri, Miyakaori and Sari Queen) were harvested once a year and their 2-
215 AP concentration was analysed. It was found that 2-AP was higher in rice crops
216 exposed to low temperature (day 25 °C/ night 20 °C) than high (day 35 °C/night
217 30 °C) or moderate temperature (day 30 °C/night 25 °C). In addition, from results
218 across three years, 2-AP concentrations of early harvest samples were higher than
219 samples harvested at normal time (Itani, Tamaki, Hayata, Fushimi, & Hashizume,

220 2004).

221 In a recent study, a partial least squares model was built, based on planting and
222 harvesting conditions, that could predict 2-AP concentrations in Thai Jasmine
223 Pathumthani 1 rice (Funsueb, Krongchai, Mahatheeranont, & Kittiwachana, 2016).
224 The status of the rice plants was recorded during cultivation and after harvest, such as
225 number of tillers (grain-bearing branches), plant height, root length, number of grains
226 per plant and grain weight. Nitrogen and sodium concentrations, rice yield, shoot dry
227 weight and number of tillers per plant all had significant influences on 2-AP
228 concentration.

229 There are two mechanisms proposed for the accumulation of 2-AP in mature
230 grains. In the first 2-AP is synthesised in leaves and stem sheaths and transported to
231 mature grains, while in the second proline translocates from leaves into grains and 2-
232 AP synthesis occurs in grains. Hinge, Patil, and Nadaf (2016) showed maximum 2-AP
233 concentrations in mature grains, with less proline in the grain at that time than at other
234 developmental stages. These results suggested that the first mechanism was more
235 likely in the fragrant rice cultivars they were studying (Ambemohar-157 and Basmati-
236 370).

237 The gene *BADH2* encodes an enzyme, betaine aldehyde hydrogenase (BADH2)
238 (Bradbury, Fitzgerald, Henry, Jin, & Waters, 2005), which catalyses the oxidation of
239 4-aminobutanal to 4-aminobutanoic acid (GABA). 4-Aminobutanal is a high affinity
240 substrate for the BADH2 enzyme (Oishi & Ebina, 2005; Trossat, Rathinasabapathi, &
241 Hanson, 1997). In solution 4-aminobutanal exists in equilibrium with its cyclic form,

242 1-pyrroline (Struve & Christophersen 2003), a 2-AP precursor. Hence the oxidation of
243 4-aminobutyraldehyde reduces the potential for 2-AP synthesis (Kovach, Calingacion,
244 Fitzgerald, & McCouch, 2009). Bradbury et al. (2005) identified a mutated version of
245 the *BADH2* gene as being responsible for determining fragrance in rice, which has
246 since been confirmed (Arikrit et al., 2011; Fitzgerald, Waters, Brools, & Henry, 2010;
247 Kovach et al., 2009; Siddiq, Vemireddy, & Nagaraju, 2012). The mutated *BADH2*
248 gene incurs a deletion of eight base pairs in exon 7, leading to early gene termination
249 and production of a truncated non-functional BADH2 enzyme (Bradbury et al., 2005).
250 Non-fragrant rice cultivars contain the *BADH2* gene and hence a functional BADH2
251 enzyme; whereas fragrant cultivars have the mutated *BADH2* gene and so produce a
252 non-functional enzyme (Bradbury, Gillies, Brusheet, Waters, & Henry, 2008). This
253 non-functional enzyme will not be able to oxidise 4-aminobutyraldehyde, leading to a
254 build-up of 1-pyrroline and hence increased 2-AP synthesis. Recent studies have
255 shown that there are various other mutations in the *BADH2* gene that may also lead to
256 increased 2-AP production, such as a deletion of seven base pairs in exon 2
257 (Amarawathi, Singh, Singh, Singh, Mohaoatra, & Sharma, 2008; He & Park, 2015).
258 Similar biosynthetic pathways for the formation of 2-AP have been reported in soy
259 beans (Arikrit et al., 2011) and sorghum (Zanan, Khandagale, Hinge, Elangovan,
260 Henry, & Nadaf 2016).

261 Another biosynthetic pathway of 2-AP was proposed by Huang, Teng, Chang,
262 Chuang, Ho, and Wu (2008) that did not involve BADH2. Higher levels of pyrroline-
263 5-carboxylate synthase enzyme, and hence increased conversion of glutamate to 1-

264 pyrroline-5-carboxylate, occurred in fragrant cultivars, in comparison to non-fragrant
265 cultivars. It was suggested that the 1-pyrroline-5-carboxylate undergoes a reaction
266 with methylglyoxal, giving rise to 2-AP, either directly or *via* degradation to 1-
267 pyrroline.

268 Although very different (a comparison can be seen in Figure 1), both pathways
269 could require 1-pyrroline in order to produce 2-AP. 1-Pyrroline has been shown to be
270 a limiting substrate of the biosynthesis of 2-AP in a recent study, where both fragrant
271 and non-fragrant rice calli were incubated with 1-pyrroline. In both cases, a
272 significant increase in 2-AP production was observed, proving 1-pyrroline to be a key
273 intermediate of 2-AP biosynthesis (Poonlaphdecha et al., 2016).

274 *3.2. Formation of 2-AP by microorganisms*

275 Microorganisms could also play an important role in 2-AP formation. During
276 cocoa bean fermentation, yeasts, lactic acid, acetic acid, and various spore-forming
277 bacteria, such as *Bacillus cereus*, are involved in the flavour-forming reactions. Some
278 *Bacillus cereus* strains produce popcorn-like notes and 2-AP was produced by several
279 of these strains incubated on standard plate count agar at 35 °C; 30–75 µg/kg 2-AP
280 was produced after 2 days. A series of ¹³C and ¹⁵N experiments showed that 2-AP
281 could be formed from glucose as carbon source, and glutamic acid and proline as
282 nitrogen sources, through *Bacillus cereus* metabolism (Romanczyk, McClelland, Post,
283 & Aitken, 1995).

284 In Mediterranean dried sausages, which have a popcorn-like odour and are very

285 different from Northern European sausages, 2-AP is also regarded as the key aroma
286 compound. The main difference between Northern European sausages and
287 Mediterranean dried sausages is a coverage of mould on the latter. 2-AP concentration
288 on the surface of Mediterranean dried sausages is much higher than at the core.
289 Therefore, it was suggested that the mould on the surface of Mediterranean dried
290 sausages is able to produce 2-AP. *Penicillium nalgiovense* was isolated from the
291 sausage surface and it was the dominating mould species. When incubated with and
292 without various supplements, it was found that *Penicillium nalgiovense* could only
293 produce popcorn odour when the sausage was present (Stahnke, 2000).

294 2-AP, together with other *N*-heterocyclic compounds 2-ethyltetrahydropyridine
295 and 6-acetyl-1,2,3,4-tetrahydropyridine, could cause mousy off-flavour in wine
296 (Herderich, Costello, Grbin, & Henschke, 1995; Strauss & Heresztyn, 1984), through
297 the action of lactic acid bacteria (LAB). *Lactobacillus hilgardii* DSM 20176 was
298 incubated with a defined *N*-heterocycle assay medium, which included D-fructose,
299 ethanol, L-lysine, L-ornithine and mineral salts. It was found that L-ornithine
300 stimulated 2-AP formation and repressed 6-acetyl-1,2,3,4-tetrahydropyridine
301 formation, while L-lysine had the opposite effect (Costello & Henschke, 2002). It had
302 previously been suggested that D-fructose and ethanol could provide the acetyl side-
303 chain for 2-AP and 6-acetyl-1,2,3,4-tetrahydropyridine (Strauss & Heresztyn, 1984).
304 A possible mechanism of fermentable carbohydrate and amino acid, forming 2-AP
305 and 6-acetyl-1,2,3,4-tetrahydropyridine through LAB fermentation was proposed by
306 Costello and Henschke (2002). This pathway is shown in Figure 2. L-Lysine could

307 form the intermediate 1-piperideine *via* cadaverine pathways, with the enzymes L-
308 lysine decarboxylase and cadaverine aminotransferase involved (Fothergill & Guest,
309 1977). Pathways from putrescine to succinate *via* 1-pyrroline in *P. fluorescens* and *E.*
310 *coli* (Jacoby & Fredericks, 1959; Kim, 1964) have been reported. Putrescine is the
311 decarboxylation product of ornithine (Fothergill & Guest, 1977); hence, 1-pyrroline
312 could be formed through the putrescine pathway from L-ornithine. Due to the
313 presence of carbohydrates, such as ethanol and glucose/fructose, acetyl-CoA
314 accumulated through the heterolactic pathway and reacted with intermediates 1-
315 pyrroline and 1-piperideine to form 2-AP and 6-acetyl-1,2,3,4-tetrahydropyridine ,
316 respectively.

317 Adams and De Kimpe (2007) reproduced the work of Romancyk et al. (1995) and
318 suggested *B. cereus* formed 2-AP by enzymatic acetylation of 1-pyrroline. 1-Pyrroline
319 was formed from the degradation of ornithine and proline, as proposed by Costello
320 and Henschke (2002; see above and Figure 2).

321

322 **4. Formation of 2-acetyl-1-pyrroline through the Maillard reaction**

323 2-AP is not only present in raw food, like rice and pandan leaf, but is also formed
324 in many cooked products. Therefore, the Maillard reaction is also an important route
325 to 2-acetyl-1-pyrroline. Schieberle (1988) tested several model systems containing
326 different amino acids, and showed that, when heated with reducing sugars, only
327 proline, lysine and alanine could form 2-AP, with proline giving the highest yield.
328 Two ¹³C-labelling experiments were designed: in the first experiment, 1-¹³C-proline

329 reacted with unlabelled glucose, and in the second experiment, unlabelled proline
330 reacted with U-¹³C-glucose (all six carbon atoms are labelled with ¹³C). Both
331 experiments were carried out at 170 °C for 30 min. In both experiments labelled
332 carbon was only found in the acetyl group of 2-AP and much more ¹³C was detected
333 in the second experiment, which indicated that glucose could provide the acetyl group
334 in 2-AP formation (Schieberle, 1988).

335 Schieberle (1990) showed that both proline and ornithine could react with 2-
336 oxopropanal to form 1-pyrroline, the most important intermediate in 2-AP formation,
337 and there was a higher yield with ornithine than with proline. Figure 3a shows the
338 formation of 1-pyrroline *via* Strecker degradation of ornithine; both ornithine and
339 citrulline, another amino acid, can generate 4-aminobutanal. Schieberle (1995) also
340 hypothesised a mechanism of 1-pyrroline formation from proline and 1-deoxyosone
341 through Strecker degradation (Figure 3b). This reaction starts with the formation of an
342 iminium ion. After decarboxylation and water elimination, 1-pyrroline can be
343 generated from hydrolysis of the iminium ion.

344 Rewicki et al. (1993) reacted unlabelled proline with 1-¹³C-glucose, and noted the
345 formation of a 1:1 mixture of unlabelled and labelled 2-AP. A proposed mechanism is
346 shown in Figure 3c. Two isomers of 1-deoxy-2,3-glucosone form in a ratio of 1:1
347 from the labelled sugar. They are converted to the dihydro form of diacetylformoin,
348 which reacts with 1-pyrroline, to form 2-acetylpyrrolidine, which then oxidises to 1:1
349 labelled and non-labelled 2-AP (Rewicki et al., 1993). The 1:1 ¹³C label was due to
350 the 1:1 ratio of the reducing sugar fragments. The ¹³C from labelled proline did not

351 exist in the final 2-AP product; supporting the theory that 2-AP is formed by acylation
352 of 1-pyrroline by a two-carbon sugar fragment. Hofmann and Schieberle (1998a) also
353 reported that 2-acetylpyrrolidine could oxidise to 2-AP and proposed that 1-pyrroline
354 and 2-oxopropanal formed 2-acetylpyrrolidine *via* a number of steps. 2-
355 Acetylpyrrolidine was then readily oxidised to 2-AP.

356 Phosphate ion could significantly increase the yield of 2-AP, through increased
357 formation of 2-oxopropanal *via* 1,3-dihydroxyacetone phosphate (Schieberle, 1989).
358 If malonate buffer replaced phosphate buffer, there was a one-third reduction of 2-AP
359 formation (Schieberle, 1995). Blank, Devaud, Matthey-Doret, and Robert (2003)
360 examined the effect of pH and heating time on the formation of various Maillard-
361 derived compounds in two phosphate-buffered model systems: one an equimolar
362 mixture of proline and glucose, the other the Amadori compound fructose-proline. 2-
363 AP yield was similar in both systems across all treatments and was shown to increase
364 with increasing pH and heating time, when samples were refluxed for 1, 2 and 4 hours
365 at pH 6, 7 and 8.

366 From the above hypothesised mechanisms of 2-AP thermal formation, it is agreed
367 that 2-AP is formed through an acylation of 1-pyrroline. Certain amino acids, i.e.,
368 proline, ornithine and citrulline, reacting with 2-oxopropanal from reducing sugar
369 fragmentation, are the most important intermediates of 2-AP formation during the
370 Maillard reaction (Adams & De Kimpe, 2006).

371

372 **5. Synthesis of 2-acetyl-1-pyrroline**

373 The first synthesis of 2-AP was reported by Buttery et al. (1983) and is shown in
374 Figure 4a. This method is based on an earlier synthesis of a six-membered ring
375 compound 2-acetyl-1,4,5,6-tetrahydropyridine (Buchi & Wuest, 1971). However, the
376 yield of 2-AP from this reaction was only around 10%.

377 The first large-scale method for 2-AP synthesis was developed in 1993. Methyl
378 proline is oxidised to 2-(methoxycarbonyl)-1-pyrroline, which then reacts with
379 methylmagnesium iodide in a Grignard reaction. However, this is not a completed
380 reaction with a 45–83% yield and 8–39% starting material in the final product (De
381 Kimpe, Stevens, & Keppens, 1993). Methyl lithium in ether also converted 2-
382 (methoxycarbonyl)-1-pyrroline into a mixture of 2-acetyl-1-pyrroline (47%) and a
383 side-product, 2-(1-hydroxy-1-methylethyl)-1-pyrroline (32%) (Fig. 4b). To prevent
384 formation of this side-product, which was also formed in the methylmagnesium iodide
385 reaction, a cyanide functional group can replace the ester (Figure 4c). In this modified
386 method, the reaction started with oxidation of pyrrolidine to tripyrroline. The
387 tripyrroline was hydrocyanated into 2-cyanopyrrolidine, which can form 2-cyano-1-
388 pyrroline through the Grignard reaction. 2-Cyano-1-pyrroline can form 2-AP with a
389 yield of 60% when treated with methylmagnesium iodide (De Kimpe et al., 1993).

390 Over subsequent years, other synthesis methods focused on the stabilisation of 2-
391 AP during the reaction, by using protected carbonyl groups, e.g., Duby and Huynh-Ba
392 (1993), or amino groups, e.g., De Kimpe and Keppens (1996). De Kimpe and
393 Keppens (1996) used diacetyl as a starting material to generate an α -diimine, which
394 then reacted with a stabase derivative to form the 1-pyrroline ring structure (Fig. 4d).

395 Another synthesis method applied the high substrate selectivity of immobilised
396 penicillin G acylase (PGA) as the catalyst in the last reaction step (Favino, Fronza,
397 Fuganti, Fuganti, Grasselli, & Mele, 1996). 1-Aminohex-4-yne reacted with
398 phenylacetyl chloride, and then ozone oxidised the product to form 1-[*N*-
399 (phenylacetyl)amino]-4,5-dioxohexane, which when treated with PGA could form 2-
400 AP spontaneously, as shown in Figure 4e. An 80% yield of 2-AP could be achieved
401 using this method.

402 A four-step synthesis was reported by Hofmann and Schieberle (1998b) starting
403 from *N-tert*-butoxycarbonyl-protected proline, while 2-pyrrolidinone was selected as a
404 raw material for 2-AP synthesis by Harrison and Dake (2005). Another ‘popcorn’
405 compound, 2-acetyl-1,4,5,6-tetrahydropyridine, was a by-product of this latter
406 method. A three-step synthesis was reported by Fuganti, Gatti, and Serra (2007),
407 starting from the reaction of *N*-Boc-pyrrolidinone with ethylmagnesium bromide. The
408 yield of 2-AP was only 20–30% but with 98% purity. Maraval et al. (2010) formed
409 *N*,5-diacetylpyrrolidin-2-one from L-glutamic acid and acetic anhydride with 78%
410 yield. Sodium carbonate was used for deacetylation to form 5-acetylpyrrolidin-2-one;
411 then lithium aluminium hydride was used for reduction to form 2-(1-hydroxyethyl)-
412 pyrrolidine, which was oxidised by silver carbonate to 2-AP. The overall yield of 5-
413 acetylpyrrolidin-2-one formation was 37% but the yield of 2-AP was not reported.

414 A recent publication reported the synthesis of three major popcorn-like Maillard
415 aroma compounds, 2-acetyl-1-pyrroline, 2-acetyl-1,4,5,6-tetrahydropyridine and 2-
416 acetyl-5,6-dihydro-4*H*-1,3-thiazine (Deblander, Van Aeken, Adams, De Kimpe, &

417 Abbaspour Tehrani, 2015). The authors noted that existing synthetic procedures for
418 these compounds suffered a number of problems, including extensiveness of some
419 reaction pathways, and the use of costly and/or harmful reagents. The 2-AP synthesis
420 they proposed started from *N*-Boc-prolinate to give a final yield of 2 AP of 28%, *via* a
421 four-step reaction (Fig. 4f). The authors considered this method to be relatively
422 straightforward, as the starting materials were readily available, only one general
423 procedure was involved and the vinyl ether intermediate prepared was a stable
424 precursor, which could be readily converted to 2-AP.

425 Although numerous procedures have been published for the synthesis of 2-AP,
426 these methods all require an experienced organic chemist to make them work.
427 Synthesis of 2-AP is difficult, due to the unstable nature of this compound, which
428 degrades very rapidly upon standing (see Part 6). This is reflected in the high price of
429 commercial 2-AP and the small number of 2-AP suppliers.

430

431 **6. Stability and stabilisation of 2-acetyl-1-pyrroline**

432 Stability is very important for a flavour compound in food products.
433 Unfortunately, 2-acetyl-1-pyrroline has limited stability. Pure 2-AP will turn red and
434 degrade within 10 min at room temperature (Fang & Cadwallader, 2014). This
435 instability of 2-AP was noticed when it was first identified by Buttery et al. (1982),
436 and this instability was assumed to be due to polymerisation. Loss of 2-AP in stored
437 foods could be due to complexation, decomposition, diffusion to the environment and

438 generation of other compounds (Adams & De Kimpe, 2006).

439 An experiment was designed to investigate the effect of storage on 2-AP levels in
440 rice (Widjaja et al., 1996a). In this experiment, using the fragrant cultivar YRF9,
441 paddy (rice with husk and rice bran), brown (rice without husk but with rice bran) and
442 white (rice without husk and rice bran) rice samples were stored under two conditions:
443 atmospheric pressure and reduced pressure, at 84% RH and 30 °C. After three months'
444 storage, 2-AP level was reduced by 40–50% in all cases. Another study aimed to
445 compare the effect of different drying methods and storage time on 2-AP reduction in
446 fragrant rice (Wongpornchai et al., 2004). Six different drying methods (sun drying,
447 30 °C modified air, 40 °C modified air, 40 °C air, 50 °C air and 70 °C air) were
448 applied to fresh paddy rice, to reduce moisture content from 28% to below 14%, and
449 then the rice was stored in gunnysacks at 20–35 °C. 2-AP concentration in 10 months
450 stored rice was only 25% of freshly dried rice and it was shown in a concentration–
451 storage time curve that a significant decrease occurred at the beginning of storage.
452 The sun-dried sample retained less 2-AP than the other drying methods; this could be
453 due to the longer drying time. Sun drying took 54 hours in this study, while the
454 average time for the other drying methods was 10 hours. Although the authors did not
455 provide details of the modified air used, drying with this kind of air maintained 2-AP
456 in rice better than normal hot air drying, while lower air temperatures also resulted in
457 less 2-AP loss (Wongpornchai et al., 2004).

458 In addition to rice, 2-AP decreases in other food products during storage. Hot air
459 popped popcorn was sealed in commercial polyethylene food bags and stored in the

460 dark at room temperature. After two days, the 2-AP level reduced by 20% and after
461 seven days storage, it reduced by 75% (Schieberle, 1995).

462 Therefore, it is important to develop a stabilisation method to defer 2-AP
463 breakdown. Encapsulation is a popular technique to protect unstable volatile
464 compounds for commercial processing, and several studies have applied this
465 technique. Encapsulation of 2-AP by β -cyclodextrin (Duby & Huynh-Ba, 1996)
466 showed some success. When stored at room temperature (20 °C), 99% 2-AP
467 decomposed after 110 days' storage, when the 2-AP load of the β -cyclodextrin was
468 1%. However, if the storage temperature decreased, encapsulation performed better,
469 with 10% losses at 4°C and no losses at -20 °C. If the loading of cyclodextrin was
470 increased to 10%, the stability of the 2-AP was reduced.

471 Apintanapong and Noomhorm (2003) extracted 2-AP from pandan leaves and
472 examined its stability at 30 ppm in acidic and basic solution at room temperature.
473 They also microencapsulated 2-AP in various maltodextrin and gum acacia mixtures.
474 In basic solution, 2-AP was reduced by 63% after 7 days, and in acidic solution by
475 30% after 35 days. When 2-AP was microencapsulated with 70:30 gum
476 acacia:maltodextrin, only 28% of the encapsulated 2-AP was lost after 72 days at
477 room temperature. Gum acacia and/or starch mixed materials were used in a patented
478 form by Srinivas, Sulochanamma, Raghavan, and Gurudutt (2006), to form a stable 2-
479 AP powder using spray drying, but the stability of this powder was not reported.

480 Fang and Cadwallader (2014) recently reported a novel stabilisation method, using
481 zinc ions to solve 2-AP powder storage problems. Anhydrous 2-AP and ZnI_2 were

482 added into diethyl ether to form a yellowish precipitate, which was the desired 2-AP-
483 ZnI₂ complex. Excess ZnI₂ and other impurities were removed through dissolving the
484 complex in anhydrous diethyl ether. The complex compound was obtained as a
485 powder after drying through nitrogen evaporation. Other 2-AP-zinc halide complexes
486 could be obtained in the same way. When stored at 25 °C, there was only 6% loss of
487 2-AP from a 2-AP-ZnI₂ complex (2-AP content = 14.4%) after 3 months, and 3%
488 reduction of 2-AP after 3 months when a 12.5% 2-AP content complex was stored at –
489 20 °C. It was found that compared with ZnI₂(2-acetyl-1-pyrroline)_n, which had a yield
490 of 62%, complexes of ZnBr₂(2-acetyl-1-pyrroline)_n and ZnCl₂(2-acetyl-1-pyrroline)_n
491 had better yields of 96% and 86%, respectively. A ZnCl₂-2-AP complex would be the
492 preferred food agent because ZnCl₂ has been approved for food use (CFR – Code of
493 Federal Regulations Title 21; April 1st, 2016). This method can also applied to similar
494 volatile compounds, such as 2-propionyl-1-pyrroline, 2-acetyl-1,4,5,6-
495 tetrahydropyridine, 2-acetyl-2-thiazoline, 2-acetylthiazole, 2-acetylpyrazine and 2-
496 acetylpyridine. Although this is an effective technique for 2-AP stabilisation compared
497 with others, this high yield was only confirmed in the dry powdered complex. It may
498 be reduced by moisture, temperature and other conditions when applied in food.
499 Therefore, it may be necessary to combine this technique with an encapsulation
500 technique to protect the 2-AP-zinc halide complex in a changeable food environment
501 (Fang & Cadwallader, 2014).

502

503 **7. Extraction and instrumental analysis of 2-acetyl-1-pyrroline**

504 *7.1 Solvent-based extraction techniques*

505 Simultaneous distillation extraction (SDE) was used as the extraction method
506 when 2-AP was first discovered by Buttery et al. (1982). SDE was widely used in the
507 1980s and 1990s for volatile compound extraction (Chi, Yeung, & MacLeod, 1981).
508 The sample is heated in water to produce steam and the steam transfers volatile
509 material to a boiling non-polar solvent, which condenses to give an aroma extract
510 (Likens & Nickerson, 1964). However, this vigorous heating process may cause
511 volatile compound formation or breakdown. Buttery et al. (1983) reported that the
512 boiling conditions used in this extraction may decompose 2-AP in rice and cause a
513 lower concentration than in raw samples.

514 In Buttery's study, 500 g rice were extracted with 6 L water in a Likens-Nickerson
515 type extraction equipment (Likens & Nickerson, 1964); diethyl ether was used as
516 solvent and the isolation process was carried out at atmospheric pressure for 2 hours.
517 After concentration, the solvent extract was dissolved in hexane and then extracted
518 with 3 N hydrochloric acid and then ether. The ether extract was then concentrated to
519 a small amount for analysis. For subsequent quantitative measurements (Buttery,
520 Ling, & Mon, 1986), an internal standard (5 mL of 30 ppm 2,4,6-trimethylpyridine
521 (collidine) solution) was added to the rice before extraction. Buttery's group published
522 a number of papers on 2-AP in fragrant rice. They detected 2-AP in 10 different
523 varieties of cooked rice (both milled and brown) using SDE and found that the
524 concentrations of 2-AP in brown rice were much higher than in white rice (Buttery et
525 al., 1983).

526 Likens-Nickerson extraction was continuously developed and used in the
527 following decade for 2-AP extraction in rice, pandan leaf and other food samples.
528 Addition of magnesium sulfate ($MgSO_4$) during rice SDE inhibited starch
529 gelatinisation, water absorption, swelling of rice and foaming of the mixture during
530 distillation (Widjaja et al., 1996a & b). Dichloromethane (DCM) has also been used as
531 the extraction solvent in SDE of 2-AP (Nadaf, Krishnan, & Wakte, 2006).

532 Rice was boiled before extraction in some studies and in others the rice was boiled
533 during SDE. For example, when fresh and stored brown and white fragrant YRF9 rice
534 were compared, samples were boiled during the SDE process (Widjaja et al., 1996a &
535 b). When white Italian Line B5-3 and basmati, two fragrant rice species, were
536 compared, they were also boiled during SDE. A 4-fold higher 2-AP concentration was
537 found in the Italian variety than in basmati (Tava & Bocchi, 1999). Several different
538 varieties of brown fragrant rice (Malagkit Sungsong, 370 basmati, Khashkani and
539 Indica) were boiled for 25 min in tap water before SDE analysis. 2-AP was found in
540 all four species but the concentration in Indica was much lower than in the others
541 (Jezussek, Juliano, & Schieberle, 2002). 2-AP was also detected in five boiled fragrant
542 rice cultivars; four of them were white rice (Hyangmiby eo 1, Hyangmiby eo 2, Royal,
543 Golden Elephant) and one a Korean black rice called Goemjeongssal. Boiled non-
544 fragrant rice Jeongilpum also contained 2-AP. Those six cultivars were boiled 30 min
545 with distilled water (Yang, Shewfelt, Lee, & Kays, 2008). Three white fragrant
546 cultivars (Aychade, Fidji, and Giano) and one white non-fragrant cultivar (Ruille)
547 were boiled for 20 min. 2-AP was found in all four cultivars, but the concentration in

548 Ruille was too low to quantify; it was lower than 2 µg/kg while the concentrations in
549 the other fragrant cultivars were 150–300 µg/kg (Maraval et al., 2008).

550 When using SDE at atmospheric pressure, sample and water mixture are boiled.
551 Therefore, this kind of extraction technique cannot be used to study uncooked foods.
552 Buttery et al., when first identifying 2-AP, used simultaneous distillation/extraction
553 under vacuum (V-SDE) to study rice aroma. However, compared to SDE at
554 atmospheric pressure (A-SDE), V-SDE showed a low efficiency of extraction
555 (Buttery et al., 1983). Levels of 2-AP in rice extracted by A-SDE were 10 times
556 higher than in rice that was cooked and then extracted with V-SDE. The authors
557 suggested that most of the 2-AP may be lost during cooking. Therefore, compared
558 with A-SDE, where rice is cooked during the isolation process, less 2-AP is present in
559 the already cooked sample in V-SDE. In addition, 2-AP may be generated during
560 cooking, which can also cause the significant difference in concentrations obtained
561 between V-SDE and A-SDE.

562 Another solvent-based extraction technique, solvent-assisted flavour evaporation
563 (SAFE), first introduced in 1999 (Engel, Bahr, & Schieberle, 1999), is a useful
564 technique for 2-AP extraction. The volatile compounds in a solvent extract, usually in
565 diethyl ether or dichloromethane, are removed from non-volatile material using high-
566 vacuum distillation. The procedure takes place at around 30 °C, keeping sample
567 decomposition to a minimum. When using 1:1 diethyl ether:dichloromethane as the
568 solvent, 2-AP was isolated from cereal coffee brew (Majcher et al., 2013) and this
569 compound was also isolated from hazelnut when using diethyl ether as solvent (Kiefl

570 et al., 2013).

571 Solid-phase extraction (SPE) can also be applied for 2-AP extraction. Several
572 commercial SPE cartridges (Strata™ X from Phenomenex, LiChrolut® EN from
573 Merck Millipore and Isolute® ENV+ from Biotage) have been successfully used for
574 volatile compound extraction (Du & Qian, 2008; Metafa & Economou, 2013),
575 particularly for isolation of relatively polar aroma compounds, such as 2-AP. An
576 advantage of SPE is that no heating is applied when using this technique, which is the
577 same as SAFE, but SPE is much easier to perform than SAFE. 2-AP was generated
578 during high-temperature cooking of fragrant rice (180 °C for 20 min in an open
579 system), using SPE, followed by GC-MS (Handoko, 2014). This result suggests that
580 there may be a component of fragrant rice that is formed enzymatically, which can be
581 converted to 2-AP by the application of higher temperatures. Although there was no 2-
582 AP detected in Ciherang rice (a non-fragrant rice) heated under the same conditions, a
583 sensory panel perceived popcorn-like odour (Handoko, 2014), suggesting that
584 compounds besides 2-AP that could cause popcorn-like odour in rice heated at
585 180 °C.

586 *7.2 Headspace techniques*

587 Dynamic headspace extraction using an adsorbent polymer such as Tenax can also
588 be used in 2-AP extraction, Buttery, Turnbaugh, and Ling (1988) used this technique
589 to analyse the volatile compounds in cooked rice. Seventeen volatile compounds
590 include 2-AP were identified through this method. Around 0.6 µg/kg 2-AP were found
591 in white Californian long-grain rice (a kind of non-fragrant rice) boiled in water for 20

592 min before Tenax trapping (Buttery et al., 1988). Around 30 odorants were identified
593 in cooked rice using the same technique (Yang et al., 2008).

594 Headspace solid-phase microextraction (HS-SPME) is now the most widely used
595 extraction method for 2-AP. As with all extraction techniques, increased extraction
596 time or higher temperatures could result in better release of 2-AP from the sample.
597 However, the reported instability of 2-AP may cause its loss during isolation at higher
598 temperatures, while at lower temperatures enzymatic changes may occur during this
599 extraction. These conflicting reactions make 2-AP quantification difficult.

600 One large study used manual SPME to analyse 91 different uncooked cultivars,
601 including 77 non-basmati fragrant cultivars, 9 basmati fragrant cultivars and 5 non-
602 fragrant cultivars. A 1-cm divinylbenzene/Carboxen/polydimethylsiloxane
603 (DVB/CAR/PDMS) fibre was used in this experiment. Samples were extracted for 15
604 min at 80 °C after a 30-min equilibration period. 2-AP was detected in some non-
605 fragrant cultivars but its average concentration was around 10-fold higher in basmati
606 fragrant cultivars and around 20-fold higher in selected non-basmati fragrant cultivars
607 (Mathure, Wakte, Jawali, & Nadaf, 2011; Mathure, Jawali, Thengane, & Nadaf,
608 2014). Bryant and McClung (2011) used an automated SPME system to compare
609 seven uncooked fragrant and two uncooked non-fragrant rice samples extracted with a
610 1-cm DVB/CAR/PDMS SPME fibre at 80 °C for 18 min after a 5-min equilibration
611 period. 2-AP was only found in the fragrant rice samples.

612 A recent study measured 2-AP in 48 fragrant rice samples using manual SPME,
613 followed by GC-MS/MS. The ion transition from m/z 111 to m/z 82 was used to

614 quantify 2-AP. Optimised conditions were 10 minutes extraction at 40 °C after a 5-
615 min thermostating period. The technique was sensitive enough both to quantify 2-AP
616 below its odour threshold concentration (Buttery et al., 1983) and to obtain successful
617 2-AP measurements using a single grain of rice (Hopfer et al., 2016). Although
618 DVB/CAR/PDMS fibres performed better than DVB/PDMS fibres, the latter were
619 preferred, because carry-over (the presence of volatile material from the fibre in a
620 subsequent blank analysis) observed when using the DVB/CAR/PDMS fibre was not
621 observed with the DVB/PDMS fibre. A limit of quantification of 103 ng per kg was
622 reported for 2-AP in this work.

623 *7.3 Gas chromatography-mass spectrometry*

624 Gas chromatography-mass spectrometry (GC-MS) is the most common technique
625 used for volatile compound analysis. Column choice in GC-MS analysis of 2-AP is
626 very important. A polar phase (e.g., Carbowax) is normally chosen; 2-AP is a
627 relatively polar compound and its peak shape is more symmetrical and sharper on a
628 polar column. The use of base-deactivated phases is recommended for basic
629 compounds that may possess poor peak shape under normal GC conditions (De
630 Zeeuw, Stricek, & Stidsen, 2011). A base-deactivated column may also be useful for
631 quantifying 1-pyrroline (Poonlaphdecha et al., 2016). Because of the instability of 2-
632 AP, Buttery et al. (1986) recommended a relatively low injector temperature of 150-
633 170 °C to minimise its decomposition. Base-deactivated injection port liners may also
634 have a protective role (De Zeeuw et al., 2011).

635 When quantifying 2-AP using GC-MS, separation of 2-AP from other compounds

636 is a common challenge. This interference problem was first reported by Paule and
637 Powers (1989) when using a packed column coated with 10% Carbowax 20M on
638 Chromosorb® W; 1-hexanol eluted very close to 2-AP and could interfere in its
639 quantification. The mass spectrum of 6-methyl-5-hepten-2-one contains all the major
640 ions present in the mass spectrum of 2-AP (m/z 43, 41, 111, 83, 68, 69); these two
641 compounds often co-elute in fragrant rice extracts run on polar columns, affecting the
642 quantification of 2-AP, especially when the concentration of 2-AP is similar to or
643 lower than that of 6-methyl-5-hepten-2-one. A long isothermal stage of 65 °C for 70
644 min at the start of the GC run was reported by Tanchotikul and Hsieh (1991) when
645 performing sample analysis by GC-MS using a 60-m length Supelcowax® 10
646 (Supelco, Bellefonte, PA) column. A similar method was reported by Seitz et al.
647 (1993), to obtain better separation of 2-AP and 6-methyl-5-hepten-2-one. They used a
648 shorter, 30 m Supelcowax 10 column in this analysis at an initial temperature of 60 °C
649 for 15 min.

650 Although electron ionisation (EI) is the usual ionisation mode used for GC-MS,
651 chemical ionisation (CI) is an option. CI is a softer ionisation technique, which can
652 reduce interference during MS analysis compared with EI, and hence could result in
653 increased signal-to-noise ratio for compounds of interest. In a study on bread flavour
654 by Schieberle and Grosch (1987), 2-AP was analysed in CI mode using isobutane as
655 reagent gas. Compared with EI mode in GC-MS analysis, Maraval et al. (2010)
656 reported that positive ion CI mode could be better for 2-AP quantification in rice,
657 especially when MS-MS was applied for analysis. The EI mass spectrum of 2-AP

658 possesses few defining peaks: a characteristic ion at m/z 83 and a less intense
659 molecular ion at m/z 111. However, in PCI mode, using acetonitrile as the reagent gas,
660 only an intense pseudomolecular ion at m/z 112 was observed. Under MS-MS
661 conditions, m/z 112 ion yielded a fragment ion at m/z 70 and this transformation was
662 used for 2-AP quantification, with a low limit of quantification of 0.4 $\mu\text{g}/\text{kg}$.

663 *7.4 Quantification of 2-acetyl-1-pyrroline*

664 When Buttery et al. (1983) first quantified 2-AP in rice they measured peak areas
665 by flame ionisation detector and performed an approximate quantification, in order to
666 determine the relative amounts of 2-AP in the 16 types of rice that they analysed. In
667 subsequent work they used collidine (2,4,6-trimethylpyridine) as an internal standard,
668 adding it in solution to the rice/water mixture prior to extraction. Collidine was
669 chosen because it has similar physicochemical properties to 2-AP (basic, similar water
670 solubility, similar volatility), is stable, has a GC retention time similar to 2-AP on a
671 wax column and is commercially available (Buttery et al., 1986). Known amounts of
672 2-AP were added to the rice prior to extraction alongside a fixed amount of collidine,
673 in order to provide a calibration curve for quantification. Collidine was subsequently
674 used as the internal standard in a number of papers where 2-AP was quantified in rice
675 (Tanchotikul & Hsieh, 1991; Widjaja, et al. 1996a & b; Tava & Bocchi, 1999;
676 Bergman et al., 2000).

677 Stable isotope dilution assays (SIDA) are now widely used in flavour science. An
678 isotopomer of the compound of interest is added to the sample under study, in order to
679 permit accurate quantification of the compound of interest. The quantification of 2-AP

680 by SIDA was carried out for the first time by Schieberle and Grosch (1987). They
681 prepared a 2-AP analogue, which was partially deuterated in the heterocyclic ring,
682 giving a product with a range of molecular masses from 113 to 116. They then used
683 the deuterated isotopomer to quantify 2-AP in wheat and rye bread.

684 SIDA was used to measure 2-AP in rice for the first time by Yoshihashi et al.
685 (2004). Instead of deuteration, a ^{13}C atom was introduced in the methyl position of the
686 acetyl side-chain giving an isotopomer with a mass of 112. Naturally-occurring 2-AP
687 has an M+1 ion with a mass of 112, which has 7% of the intensity of its molecular
688 ion. It is not clear if this was taken into account by the authors in their calculations.
689 This issue was highlighted by Maraval et al. (2010), who used SPME with deuterated
690 2-AP to quantify 2-AP in rice. Unlike Schieberle and Grosch (1987), the deuteration
691 was defined. Deuterium-hydrogen exchange can occur in aqueous solution and to
692 reduce the chances of this happening the authors replaced both hydrogen atoms at the
693 5-position of the heterocyclic ring with deuterium. In order to provide a calibration
694 curve for quantification, ground leaves from a non-scented rice cultivar were spiked
695 with nine different amounts of 2-AP in solution.

696 The key reason for performing SIDA is that several steps of enrichment of the
697 compounds can be performed without losses in accuracy, provided that the initial ratio
698 between the compound and its labelled analogue remains unchanged during the entire
699 procedure (Schieberle & Grosch, 1987). As the compound of interest and its
700 isotopomer should have the same physicochemical properties, SIDA provides a degree
701 of confidence that is lacking when other internal standards are used. The incompletely

702 deuterated 2-AP synthesised by Schieberle and Grosch is now available commercially
703 from AromaLab AG (Planegg, Germany) and has been used to quantify 2-AP in rice
704 (Hopfer et al., 2016).

705

706 **8. Sensory evaluation of 2-acetyl-1-pyrroline**

707 2-AP is described as a popcorn-like odour compound and it has a very low odour
708 threshold. When Buttery et al. (1983) first identified this compound, they ranked the
709 amount of popcorn-like odour in different rice varieties. Malagkit Sungsong, a kind of
710 Philippine fragrant rice, had the greatest popcorn aroma and Texas Long Grain was
711 determined as the rice with the least popcorn aroma. The most famous fragrant rice,
712 basmati, was ranked in the middle of this list. When the Malagkit Sungsong was
713 compared with Calrose (a non-fragrant rice), it was easy to distinguish them.
714 However, when a 2-AP solution was added to the Calrose rice, they became much
715 more difficult to tell apart. It was clear that the popcorn aroma of 2-AP is a key
716 component of rice flavour.

717 Lexicons of aroma and flavour of rice are being continuously developed by
718 researchers (Goodwin et al., 1996; Piggott, Morrison, & Clyne, 1991; Yau & Liu,
719 1999). When comparing these studies, some descriptors are similar, but some are
720 different; it is difficult to estimate which research has an intact lexicon and which
721 needs more development. The choice of descriptors depends on the culture and
722 familiarity with the sample of the panellists in each study (Paule & Powers, 1989). A
723 study aiming to build an intact lexicon tested 36 different varieties of rice, which were

724 mainly jasmine and basmati rice samples from different regions, but also included
725 many other fragrant and non-fragrant rice species (Limpawattana & Shewfelt, 2010).
726 Twenty-four attributes were listed by 8 trained panellists, of which 6 did not vary
727 across the 36 varieties. The 18 attributes finally used in this study were ‘popcorn’,
728 ‘starchy’, ‘woody’, ‘cooked-grain’, ‘grain’, ‘sulfury’, ‘corn’, ‘nutty’, ‘floral’, ‘dairy’,
729 ‘hay-like’, ‘barny’, ‘buttery’, ‘green’, ‘rancid’, ‘waxy’, and ‘earthy’. A standard and
730 intensity of standard for each attribute was also defined. Of the 18 significant
731 attributes, ‘popcorn’, which was mainly attributed to 2-AP, was positively correlated
732 with ‘buttery’ and ‘corn’ and negatively correlated with ‘earthy’ and ‘smoky’.

733 In an earlier study from Limpawattana’s group, sensory profiling was conducted
734 on 13 varieties of rice by using trained panels, and aroma-active compounds were
735 analysed by GC-olfactometry (GC-O) and GC-MS (Limpawattana, Yang, Kays, &
736 Shewfelt, 2008). In this study, a predictive model was built for correlation analysis of
737 attributes and volatile compounds. Unexpectedly, ‘popcorn’ in this model was
738 negatively correlated with guaiacol and (*E,E*)-2,4-decadienal, while 2-AP was not
739 present in this model as a ‘popcorn’ descriptor. Guaiacol and (*E,E*)-2,4-decadienal
740 contributed smoky and fatty attributes, respectively. The authors suggested that the
741 thermal process of reference standard preparation may have influenced the ‘popcorn’
742 descriptor analysis. In addition, these authors suggested that the contribution of 2-AP
743 to popcorn-like odour was always overemphasised relative to many other compounds
744 which also contribute to this aroma in fragrant rice.

745 Three types of fragrant rice (jasmine, basmati, and Jasmati) were studied in a

746 recent paper, to compare their main aroma active compounds using GC-O and GC-
747 MS (Mahattanatawee & Rouseff, 2014). Hexanal, octanal, 2-AP, (*E,E*)-2,4-
748 nonadienal, (*E*)-2-nonenal, 4-vinyl-2-methoxyphenol and indole were identified as the
749 aroma-active compounds common to all three species. Across all three types of rice,
750 30 compounds were identified as aroma-active compounds and were described by 8
751 attributes. Jasmati contained 35% less ‘roasty/nutty’ total aroma intensity than jasmine
752 and basmati, while ‘medicine’ flavour was not detected in jasmine rice. Jasmine
753 contained 35% more ‘sweet fruity/floral’ total intensity than basmati and 79% more
754 than Jasmati.

755

756 **9. Conclusions**

757 2-AP contributes important aroma in many foods, like pandan leaf, mushroom and
758 especially fragrant rice. Amino acids, in particular proline and ornithine, have been
759 identified, alongside reducing sugars, as precursors of 2-AP in both biosynthesis and
760 Maillard reaction. The presence of a non-functional betaine aldehyde dehydrogenase
761 (non-functional BADH2) allows the formation of 2-AP in fragrant rice and several
762 bacteria like *Bacillus cereus* and *Penicillium nalgiovence* may also form 2-AP.

763 It appears that 1-pyrroline is a key intermediate in both biosynthesis and thermal
764 formation of 2-AP, and this intermediate could form 2-AP through an acylation
765 reaction. 2-Oxopropanal and 2-acetylpyrrolidine are other intermediates hypothesised
766 to form 2-AP during the Maillard reaction and the presence of phosphate ion could
767 increase yields of 2-AP. 2-AP formation mechanisms, particularly in rice, still need to

768 be researched. The work of Poonlaphdecha et al. (2016) showed the importance of 1-
769 pyrroline in 2-AP formation, and future work with this intermediate may provide
770 useful information.

771 Synthesis of 2-AP is still difficult but its stabilisation in a zinc halide complex has
772 increased its applicability. New synthesis strategies and stabilisation techniques could
773 reduce the cost of 2-AP, which may increase its use in the food industry, adding
774 desirable popcorn-like aroma to rice products such as rice cakes. Another possibility
775 is the addition of 2-AP intermediates, such as 1-pyrroline, to rice, which can then
776 readily form 2-AP during processing, providing a desirable fragrance to rice products.

777

778 **References:**

- 779 Adams, A., & De Kimpe, N. (2006). Chemistry of 2-acetyl-1-pyrroline, 6-acetyl-
780 1,2,3,4-tetrahydropyridine, 2-acetyl-2-thiazoline, and 5-acetyl-2,3-dihydro-
781 4H-thiazine: Extraordinary Maillard flavor compounds. *Chemical Reviews*,
782 106(6), 2299-2319.
- 783 Adams, A., & De Kimpe, N. (2007). Formation of pyrazines and 2-acetyl-1-pyrroline
784 by *Bacillus cereus*. *Food Chemistry*, 101(3), 1230-1238.
- 785 Amarawathi, Y., Singh, R., Singh, A. K., Singh, V. P., Mohaoatra, T., & Sharma, T. R.
786 (2008). Mapping of quantitative trait loci for basmati quality traits in rice
787 (*Oryza sativa L.*). *Molecular Breeding*, 21(1), 49-65.
- 788 Apintanapong, M., & Noomhorm, A. (2003). The use of spray drying to
789 microencapsulate 2-acetyl-1-pyrroline, a major flavour component of aromatic
790 rice. *International Journal of Food Science and Technology*, 38(2), 95-102.
- 791 Arikait, S., Yoshihashi, T., Wanchana, S., Uyen, T. T., Huong, N. T. T., Wongpornchai,
792 S., & Vanavichit, A. (2011). Deficiency in the amino aldehyde dehydrogenase
793 encoded by GmAMADH2, the homologue of rice Os2AP, enhances 2-acetyl-
794 1-pyrroline biosynthesis in soybeans (*Glycine max L.*). *Plant Biotechnology*
795 *Journal*, 9(1), 75-87.
- 796 Asano, K., Hirai, N., Kawakita, S., Kurose, N., Nagatomo, M., Ogawa, K., Sakai, K.,
797 Takahashi, K., & Tarumi, S. (2000). Distilled liquor product and method for
798 producing the same. Japanese Patent No. JP2002034549A.
- 799 Bergman, C. J., Delgado, J. T., Bryant, R., Grimm, C., Cadwallader, K. R., & Webb,

800 B. D. (2000). Rapid gas chromatographic technique for quantifying 2-acetyl-1-
801 pyrroline and hexanal in rice (*Oryza sativa*, L.). *Cereal Chemistry*, 77(4), 454-
802 458.

803 Berner, D. K., & Hoff, B. J. (1986). Inheritance of scent in American long grain rice.
804 *Crop Science*, 26(5), 876-878.

805 Blank, I., Devaud, S., Fay, L. B., Cerny, C., Steiner, M., & Zurbriggen, B. (2001).
806 Odor-active compounds of dry-cured meat: Italian-type salami and Parma
807 ham. In G. R. Takeoka, M. Güntert & K.-H. Engel (Eds.), *Aroma Active*
808 *Compounds in Foods*, (pp. 9-20). Washington, DC: American Chemical
809 Society.

810 Blank, I., Devaud, S., Matthey-Doret, W., & Robert, F. (2003). Formation of odorants
811 in Maillard model systems based on L-proline as affected by pH. *Journal of*
812 *Agricultural and Food Chemistry*, 51(12), 3643-3650.

813 Bradbury, L., Fitzgerald, T. L., Henry, R. J., Jin, Q., & Waters, D. (2005). The gene for
814 fragrance in rice. *Plant Biotechnology Journal*, 3, 363-370.

815 Bradbury, L., Gillies, S. A., Brusheet, D. J., Waters, D., & Henry, R. J. (2008).
816 Inactivation of an aminoaldehyde dehydrogenase is responsible for fragrance
817 in rice. *Plant Molecular Biology*, 68(4-5), 439-449.

818 Bryant, R. J., & McClung, A. M. (2011). Volatile profiles of aromatic and non-
819 aromatic rice cultivars using SPME/GC-MS. *Food Chemistry*, 124, 501-513.

820 Buchi, G., & Wuest, H. (1971). Synthesis of 2-acetyl-1,4,5,6-tetrahydropyridine, a
821 constituent of bread aroma. *Journal of Organic Chemistry*, 36, 609-610.

822 Buttery, R. G., Ling, L. C., & Juliano, B. O. (1982). 2-Acetyl-1-pyrroline: an
823 important aroma component of cooked rice. *Chemistry and Industry*, 12, 958-
824 959.

825 Buttery, R. G., Ling, L. C., Juliano, B. O., & Turnbaugh, J. G. (1983). Cooked Rice
826 Aroma and 2-Acetyl-1-Pyrroline. *Journal of Agricultural and Food Chemistry*,
827 31, 823-826.

828 Buttery, R. G., Ling, L. C., & Mon, T. R. (1986). Quantitative-analysis of 2-acetyl-1-
829 pyrroline in rice. *Journal of Agricultural and Food Chemistry*, 34(1), 112-114.

830 Buttery, R. G., Turnbaugh, J. G., & Ling, L. C. (1988). Contribution of volatiles to
831 rice aroma. *Journal of Agricultural and Food Chemistry*, 36(5), 1006-1009.

832 Carrascon, V., Escudero, A., Ferreira, V., & Lopez, R. (2014). Characterisation of the
833 key odorants in a squid broth (*Illex argentinus*). *LWT-Food Science and*
834 *Technology*, 57(2), 656-662.

835 Chi, Y., Yeung, A., & MacLeod, A. J. (1981). A comparison of the efficiency of the
836 Linkens and Nickerson extractor for aqueous, lipid/aqueous, and lipid
837 samples. *Journal of Agricultural and Food Chemistry*, 29, 502-505.

838 Costello, P. J., & Henschke, P. A. (2002). Mousy off-flavor of wine: Precursors and
839 biosynthesis of the causative *N*-heterocycles 2-ethyltetrahydropyridine, 2-
840 acetyltetrahydropyridine, and 2-acetyl-1-pyrroline by *Lactobacillus hilgardii*
841 DSM 20176. *Journal of Agricultural and Food Chemistry*, 50(24), 7079-7087.

842 Deblander, J., Van Aeken, S., Adams, A., De Kimpe, N., & Abbaspour Tehrani, K.
843 (2015). New short and general synthesis of three key Maillard flavour

844 compounds: 2-Acetyl-1-pyrroline, 6-acetyl-1,2,3,4-tetrahydropyridine and 5-
845 acetyl-2,3-dihydro-4H-1,4-thiazine. *Food Chemistry*, 168, 327-331.

846 De Kimpe, N. G., Stevens, C. V., & Keppens, M. A. (1993). Synthesis of 2-acetyl-1-
847 pyrroline, the principal rice flavor component. *Journal of Agricultural and*
848 *Food Chemistry*, 41(9), 1458-1461.

849 De Kimpe, N. G., & Keppens, M. A.. (1996). Novel syntheses of the major flavor
850 components of bread and cooked rice. *Journal of Agricultural and Food*
851 *Chemistry*, 44(6), 1515-1519.

852 De Zeeuw, J., Stricek, R., & Stidsen, G. (2011). An advanced base-deactivated
853 capillary column for analysis of volatile amines, ammonia, and alcohols.
854 *American Laboratory*, 43(2), 37-40.

855 Du, X., & Qian, M. (2008). Quantification of 2,5-dimethyl-4-hydroxy-3(2H)-furanone
856 using solid-phase extraction and direct microvial insert thermal desorption gas
857 chromatography–mass spectrometry. *Journal of Chromatography A*, 1208,
858 197-201.

859 Duby, P., & Huynh-Ba, T. (1993). Process for the preparation of 2-acetyl-1-pyrroline
860 in encapsulated form, the intermediate compound and end product obtained.
861 Canada Patent No. CA2084256 (A1).

862 Duby, P., & Huynh-Ba, T. (1996). Compositions containing 2-acetyl-1-pyrroline.
863 United States Patent No. US5512290(A).

864 Engel, W., Bahr, W., & Schieberle, P. (1999). Solvent assisted flavour evaporation - a
865 new and versatile technique for the careful and direct isolation of aroma

866 compounds from complex food matrices. *European Food Research and*
867 *Technology*, 209, 237-241.

868 Erten, E. S., & Cadwallader, K. (2017). Identification of predominant aroma
869 components of raw, dry roasted and oil roasted almonds. *Food Chemistry*, 217,
870 244-253.

871 Fang, M.-C., & Cadwallader, K. R. (2014). Stabilization of the potent odorant 2-
872 acetyl-1-pyrroline and structural analogues by complexation with zinc halides.
873 *Journal of Agricultural and Food Chemistry*, 62(35), 8808-8813.

874 Favino, T. F., Fronza, G., Fuganti, C., Fuganti, D., Grasselli, P., & Mele, A. (1996).
875 Penicillin acylase-mediated synthesis of 2-acetyl-1-pyrroline and of 2-
876 propionyl-1-pyrroline, key roast-smelling odorants in food. Inclusion
877 complexes with beta-cyclodextrin and their NMR and MS characterization.
878 *Journal of Organic Chemistry*, 61(25), 8975-8979.

879 Fitzgerald, T. L., Waters, D., Brools, L. O., & Henry, R. J. (2010). Fragrance in rice
880 (*Oryza sativa*) is associated with reduced yield under salt treatment.
881 *Environmental and Experimental Botany*, 68(3), 292-300.

882 Fothergill, J. C., & Guest, J. R. (1977). Catabolism of L-lysine by *Pseudomonas*
883 *aeruginosa*. *Journal of General Microbiology*, 99(1), 139-155.

884 Frauendorfer, F., & Schieberle, P. (2008). Changes in key aroma compounds of criollo
885 cocoa beans during roasting. *Journal of Agricultural and Food Chemistry*,
886 56(21), 10244-10251.

887 Fuganti, C., Gatti, F. G., & Serra, S. (2007). A general method for the synthesis of the

888 most powerful naturally occurring Maillard flavours. *Tetrahedron*, 63(22),
889 4762-4767.

890 Funsueb, S., Krongchai, C., Mahatheeranont, S., & Kittiwachana, S. (2016).
891 Prediction of 2-acetyl-1-pyrroline content in grains of Thai Jasmine rice based
892 on planting condition, plant growth and yield component data using
893 chemometrics. *Chemometrics and Intelligent Laboratory Systems*, 156, 203-
894 210.

895 Golam, F., Yin, Y. H., Masitah, A., Afnierna, N., Majid, N. A., Khalid, N., & Osman,
896 M. (2011). Analysis of aroma and yield components of aromatic rice in
897 Malaysian tropical environment. *Australian Journal of Crop Science*, 5(11),
898 1318-1325.

899 Goodwin, J., Koop, L. A., Rister, M. E., Miller, R. K., Maca, J. V., Chambers, E.,
900 Hollingsworth, M., Bett, K., Webb, B. D., & Mcclung, A. M. (1996).
901 Developing a common language for the U. S. rice industry: linkages among
902 breeders, producers, processors and consumers. *TAMRC Consumer Product*
903 *Market Research CP2-96*. Texas A&M: College Station.

904 Grosshauser, S., & Schieberle, P. (2013). Characterization of the key odorants in pan-
905 fried white mushrooms (*Agaricus bisporus* L.) by means of molecular sensory
906 science: comparison with the raw mushroom tissue. *Journal of Agricultural*
907 *and Food Chemistry*, 61(16), 3804-3813.

908 Handoko, D. D. (2014). *Ph.D. Thesis: An investigation of Flavour Formation in*
909 *Cooked Rice under Boiling and High Temperature Heating Condition.*

910 University of Reading.

911 Harrison, T. J., & Dake, G. R. (2005). Expeditious, high-yielding construction of the
912 food aroma compounds 6-acetyl-1,2,3,4-tetrahydropyridine and 2-acetyl-1-
913 pyrroline. *Journal of Organic Chemistry*, 70(26), 10872-10874.

914 He, Q., & Park, Y. J. (2015). Discovery of a novel fragrant allele and development of
915 functional markers for fragrance in rice. *Molecular Breeding* 35(11), article
916 217.

917 Herderich, M., Costello, P. J., Grbin, P. R., & Henschke, P. A. (1995). Occurrence of
918 2-acetyl-1-pyrroline in mousy wines. *Natural Product Letters*, 7(2), 129-132.

919 Hinge, V. R., Patil, H. B., & Nadaf, A. B. (2016). Aroma volatile analyses and 2AP
920 characterization at various developmental stages in Basmati and Non-Basmati
921 scented rice (*Oryza sativa* L.) cultivars. *Rice*, 9, 22.

922 Hofmann, T., & Schieberle, P. (1998a). New and convenient syntheses of the
923 important roasty, popcorn-like smelling food aroma compounds 2-acetyl-1-
924 pyrroline and 2-acetyltetrahydropyridine from their corresponding cyclic
925 alpha-amino acids. *Journal of Agricultural and Food Chemistry*, 46(2), 616-
926 619.

927 Hofmann, T., & Schieberle, P. (1998b). 2-Oxopropanal, hydroxy-2-propanone, and 1-
928 pyrroline: Important intermediates in the generation of the roast-smelling food
929 flavor compounds 2-acetyl-1-pyrroline and 2-acetyltetrahydropyridine.
930 *Journal of Agricultural and Food Chemistry*, 46(6), 2270-2277.

931 Hopfer, H., Jodari, F., Negre-Zakharov, F., Wylie, P. L., & Ebeler, S. E. (2016). HS-

932 SPME-GC-MS/MS Method for the Rapid and Sensitive Quantitation of 2-
933 Acetyl-1-pyrroline in Single Rice Kernels. *Journal of Agricultural and Food*
934 *Chemistry* 64(20), 4114-4120.

935 Huang, T. C., Teng, C. S., Chang, J. L., Chuang, H. S., Ho, C. T., & Wu, M. L. (2008).
936 Biosynthetic mechanism of 2-acetyl-1-pyrroline and its relationship with 1-
937 pyrroline-5-carboxylic acid and methylglyoxal in aromatic rice (*Oryza sativa*
938 L.) callus. *Journal of Agricultural and Food Chemistry*, 56(16), 7399-7404.

939 Itani, T., Tamaki, M., Hayata, Y., Fushimi, T., & Hashizume, K. (2004). Variation of 2-
940 acetyl-1-pyrroline concentration in aromatic rice grains collected in the same
941 region in Japan and factors affecting its concentration. *Plant Production*
942 *Science*, 7(2), 178-183.

943 Jacoby, W. B., & Fredericks, J. (1959). Pyrrolidine and putrescine metabolism: γ -
944 aminobutyraldehyde dehydrogenase. *Journal of Biological Chemistry*, 234(8),
945 2145-2150.

946 Jezussek, M., Juliano, B. O., & Schieberle, P. (2002). Comparison of key aroma
947 compounds in cooked brown rice varieties based on aroma extract dilution
948 analyses. *Journal of Agricultural and Food Chemistry*, 50(5), 1101-1105.

949 Kaneko, S., Sakai, R., Kumazawa, K., Usuki, M., & Hishimura, O. (2013). Key aroma
950 compounds in roasted in-shell peanuts. *Bioscience, Biotechnology, and*
951 *Biochemistry*, 77(7), 1467-1473.

952 Kiefl, J., Pollner, G., & Schieberle, P. (2013). Sensomics analysis of key hazelnut
953 odorants (*Corylus avellana* L. 'Tonda gentile') using comprehensive two-

954 dimensional gas chromatography in combination with time-of-flight mass
955 spectrometry (GC×GC-TOF-MS). *Journal of Agricultural and Food*
956 *Chemistry*, 61, 5226-5235.

957 Kim, K. H. (1964). Purification and properties of a diamine α -ketoglutarate
958 transaminase from *Escherichia coli*. *Journal of Biological Chemistry*, 239,
959 783-786.

960 Kovach, M. J., Calingacion, M. N., Fitzgerald, M. A., & McCouch, S. R. (2009). The
961 origin and evolution of fragrance in rice (*Oryza sativa* L.). *PNAS*, 106(34),
962 14444-14449.

963 Lasekan, O., Buettner, A., & Christlbauer, M. (2007). Investigation of important
964 odorants of palm wine (*Elaeis guineensis*). *Food Chemistry*, 105(1), 15-23.

965 Likens, S. T., & Nickerson, G. B. (1964). Determination of certain hop oil
966 constituents in brewing products. *American Society of Brewing Chemistry*
967 *Proceedings*, 5-13.

968 Limpawattana, M., & Shewfelt, R. L. (2010). Flavor lecion for sensory descriptive
969 profiling of different rice types. *Journal of Food Science*, 75, 199-205.

970 Limpawattana, M., Yang, D. S., Kays, S. J., & Shewfelt, R. L. (2008). Relating
971 sensory descriptors to volatile components in flavor for specialty rice types.
972 *Journal of Food Science*, 73(9), 456-461.

973 Liu, J., Liu, M., He, C., Song, H., Guo, J., Wang, Y., Yang, H., & Su, X. (2015). A
974 comparative study of aroma-active compounds between dark and milk
975 chocolate: relationship to sensory perception. *Journal of the Science of Food*

976 *and Agriculture, 95, 1362-1372.*

977 Lizarraga-Guerra, R., Guth, H., & Lopez, M. G. (1997). Identification of the most
978 potent odorants in huitlacoche (*Ustilago maydis*) and austern pilzen (*Pleurotus*
979 *sp.*) by aroma extract dilution analysis and static head-space samples. *Journal*
980 *of Agricultural and Food Chemistry, 45, 1329-1332.*

981 Mahattanatawee, K., & Rouseff, R. L. (2014). Comparison of aroma active and sulfur
982 volatiles in three fragrant rice cultivars using GC-Olfactometry and GC-PFPD.
983 *Food Chemistry, 154, 1-6.*

984 Majcher, M. A., Klensporf-Pawlik, D., Dziadas, M., & Jeleń, H. H. (2013).
985 Identification of aroma active compounds of cereal coffee brew and its roasted
986 ingredients. *Journal of Agricultural and Food Chemistry, 61, 2648-2654.*

987 Maraval, I., Mestres, C., Pernin, K., Ribeyre, F., Boulanger, R., Guichard, E., &
988 Gunata, Z. (2008). Odor-active compounds in cooked rice cultivars from
989 Camargue (France) analyzed by GC-O and GC-MS. *Journal of Agricultural*
990 *and Food Chemistry, 56(13), 5291-5298.*

991 Maraval, I., Sen, K., Agrebi, A., Menut, C., Morere, A., Boulanger, R., Gay, F.,
992 Mestres, C., & Gunata, Z. (2010). Quantification of 2-acetyl-1-pyrroline in
993 rice by stable isotope dilution assay through headspace solid-phase
994 microextraction coupled to gas chromatography-tandem mass spectrometry.
995 *Analytica Chimica Acta, 675, 148-155.*

996 Mathure, S. V., Wakte, K. V., Jawali, N., & Nadaf, A. B. (2011). Quantification of 2-
997 acetyl-1-pyrroline and other rice aroma volatiles among Indian scented rice

998 cultivars by HS-SPME/GC-FID. *Food Analytical Methods*, 4(3), 326-333.

999 Mathure, S. V., Jawali, N., Thengane, R. J., & Nadaf, A. B. (2014). Comparative
1000 quantitative analysis of headspace volatiles and their association with BADH2
1001 marker in non-basmati scented, basmati and non-scented rice (*Oryza sativa* L.)
1002 cultivars of India. *Food Chemistry*, 142, 383-391.

1003 Metafa, M., & Economou, A. (2013). Comparison of solid-phase extraction sorbents
1004 for the fractionation and determination of important free and glycosidically-
1005 bound variental aroma compounds in wines by gas chromatography-mass
1006 spectrometry. *Central European Journal of Chemistry*, 11(2), 228-247.

1007 Nadaf, A. B., Krishnan, S., & Wakte, K. V. (2006). Histochemical and biochemical
1008 analysis of major aroma compound (2-acetyl-1-pyrroline) in basmati and other
1009 scented rice (*Oryza sativa* L.). *Current Science*, 91(11), 1533-1536.

1010 Oishi, H., & Ebina, M. (2005). Isolation of cDNA and enzymatic properties of betaine
1011 aldehyde dehydrogenase from *Zoysia tenuifolia*. *Journal of Plant Physiology*,
1012 162(10), 1077-1086.

1013 Paule, C. M., & Powers, J. J. (1989). Sensory and chemical examination of aromatic
1014 and nonaromatic rice. *Journal of Food Science*, 54, 1515-1520.

1015 Peter, K. V. (2006). *Handbook of Herbs and Spices* (1 ed. Vol. 3). Cambridge:
1016 Woodhead Publishing Limited.

1017 Piggott, J. R., Morrison, W. R., & Clyne, J. (1991). Changes in lipids and in sensory
1018 attributes on storage of rice amilled to different degrees. *International Journal*
1019 *of Food Science and Technology*, 26, 615-628.

1020 Poonlaphdecha, J., Gantet, P., Maraval, I., Sauvage, F., Menut, C., Morere, A.,
1021 Boulanger, R., Wust, M., & Gunata, Z. (2016). Biosynthesis of 2-acetyl-1-
1022 pyrroline in rice calli cultures: demonstration of 1-pyrroline as a limiting
1023 substrate. *Food Chemistry*, *197*, 965-971.

1024 Rewicki, D., Tressl, R., Ellerbeck, U., Kersten, E., Burgert, W., Gorzynski, M.,
1025 Hauck, R. S., & Helak, B. (1993). Formation and synthesis of some Maillard
1026 generated aroma compounds. In: Peter Schreier & Peter Winterhalter (eds.)
1027 *Progress in Flavor Precursor Studies* (pp. 301-314). Carol Stream: Allured
1028 Publishing Corporation, .

1029 Rhodes, D., Handa, S., & Bressan, R. A. (1986). Metabolic changes associated with
1030 adaptation of plant cells to water stress. *Plant Physiology*, *82*, 890-903.

1031 Richard, T. (2001). Rice composition for coating foods. United States Patent No.
1032 US6274183 B1.

1033 Romanczyk, L. J., McClelland, C. A., Post, L. S., & Aitken, W. M. (1995). Formation
1034 of 2-acetyl-1-pyrroline by several *Bacillus cereus* strains isolated from cocoa
1035 fermentation boxes. *Journal of Agricultural and Food Chemistry*, *43*(2), 469-
1036 475.

1037 Ruckriemen, J., Schwarzenbolz, U., Adam, S., & Henle, T. (2015). Identification and
1038 quantitation of 2-acetyl-1-pyrroline in Manuka honey (*Leptospermum*
1039 *scoparium*). *Journal of Agricultural and Food Chemistry*, *63*, 8488-8492.

1040 Sakthivel, K., Sundaram, R. M., Rani, N. S., Balachandran, S. M., & Neeraja, C. N.
1041 (2009). Genetic and molecular basis of fragrance in rice. *Biotechnology*

1042 *Advances*, 27(4), 468-473.

1043 Schieberle, P. (1988). Effects of additions of proline on formation of the aroma
1044 compound 2-acetyl-1-pyrroline in white bread crust. *Getreide, Mehl und Brot*,
1045 42(11), 334-335.

1046 Schieberle, P. (1989). Formation of 2-acetyl-1-pyrroline and other important flavor
1047 compounds in wheat bread crust. In T. H. Parliment, R. J. McGorin & C. T.
1048 Ho (Eds.), *Thermal Generation of Aromas* (pp. 268-275). Washington, D.C.:
1049 American Chemical Society.

1050 Schieberle, P. (1990). The role of free amino-acids present in yeast as precursors of
1051 the odorants 2-acetyl-1-pyrroline and 2-acetyltetrahydropyridine in wheat
1052 bread crust. *Zeitschrift für Lebensmittel-Untersuchung und -Forschung*,
1053 191(3), 206-209.

1054 Schieberle, P. (1991). Primary odorants in popcorn. *Journal of Agricultural and Food*
1055 *Chemistry*, 39(6), 1141-1144.

1056 Schieberle, P. (1995). Quantitation of important roast-smelling odorants in popcorn by
1057 stable-isotope dilution assays and model studies on flavor formation during
1058 popping. *Journal of Agricultural and Food Chemistry*, 43(9), 2442-2448.

1059 Schieberle, P., & Grosch, W. (1987). Quantitative-analysis of aroma compounds in
1060 wheat and rye bread crusts using a stable isotope-dilution assay. *Journal of*
1061 *Agricultural and Food Chemistry*, 35(2), 252-257.

1062 Seitz, L. M., Wright, R. L., Waniska, R. D., & Rooney, L. W. (1993). Contribution of
1063 2-acetyl-1-pyrroline to odors from wetted ground pearl-millet. *Journal of*

1064 *Agricultural and Food Chemistry*, 41(6), 955-958.

1065 Siddiq, E. A., Vemireddy, L. R., & Nagaraju, J. (2012). Basmati rices: genetics,
1066 breeding and trade. *Agricultural Research*, 1(1), 25-36.

1067 Singh, R. K., Singh, U. S., & Khush, G. S. (2000). *Aromatic Rices*. New Delhi:
1068 Oxford & IBH Publishing.

1069 Slayton, T., & Muniroth, S. (2015). Rice sector review: a more detailed road map for
1070 Cambodian rice exports. *World Bank Working Paper*, 98834.

1071 Smith, R. L., Cohen, S. M., Doull, J., Feron, V. J., Goodman, J. I., Marnett, L. J.,
1072 Portoghese, P. S., Waddell, W. J., & Wanger, B. M. (2005). GRAS Flavouring
1073 Substances 22. *Food Technology* 27, 24-62.

1074 Sood, B. C., & Siddiq, E. A. (1978). A rapid technique for scent determination in rice.
1075 *Indian Journal of Genetics and Plant Breeding*, 38(2), 268-275.

1076 Srinivas, P., Sulochanamma, G., Raghavan, B., & Gurudutt, K. N. (2006). Process for
1077 the stabilization of 2-acetyl-1-pyrroline, the basmati rice flavorant. United
1078 States Patent No. US 20060147596 A1.

1079 Stahnke, L. H. (2000). 2-acetyl-1-pyrroline - Key aroma compound in Mediterranean
1080 dried sausages. In P. Schieberle & K. H. Engel (Eds.), *Frontiers of Flavour*
1081 *Science* (pp. 361-365). Garching, Germany: Deutsche Forschungsanstalt für
1082 Lebensmittelchemie.

1083 Strauss, C. R., & Heresztyn, T. (1984). 2-Acetyltetrahydropyridines - a cause of the
1084 mousy taint in wine. *Chemistry & Industry*, 3, 109-110.

1085 Struve, C., & Christophersen, C. (2003). Structural equilibrium and ring-chain

1086 tautomerism of aqueous solutions of 4-aminobutyraldehyde. *Heterocycles*,
1087 60(8), 1907-1914.

1088 Szabados, L., & Savouré, A. (2010). Proline: a multifunctional amino acid. *Trends in*
1089 *Plant Science*, 15(2), 89-97.

1090 Tanchotikul, U., & Hsieh, T. C. Y. (1991). An improved method for quantification of
1091 2- acetyl-1-pyrroline a popcorn-like aroma, in aromatic rice by high-resolution
1092 gas chromatography/mass spectrometry/selected ion monitoring. *Journal of*
1093 *Agricultural and Food Chemistry*, 39, 944-947.

1094 Tava, A., & Bocchi, S. (1999). Aroma of cooked rice (*Oryza sativa*): Comparison
1095 between commercial Basmati and Italina Line B5-3. *Cereal Chemistry*, 76(4),
1096 526-529.

1097 Trossat, C., Rathinasabapathi, B., & Hanson, A. D. (1997). Transgenically expressed
1098 betaine aldehyde dehydrogenase efficiently catalyzes oxidation of
1099 dimethylsulfoniopropionaldehyde and [ω]-aminoaldehydes. *Plant*
1100 *Physiology*, 113(4), 1457-1461.

1101 Wagber, J., Granvogl, M., & Schieberle, P. (2016). Characterization of the key aroma
1102 compounds in raw licorice (*Glycyrrhiza glabra* L.) by means of molecular
1103 sensory science. *Journal of Agricultural and Food Chemistry*, 64(44), 8388-
1104 8396.

1105 Wakte, K., Zanan, R., Hinge, V., Khandagale, K., Nadaf, A., & Henry, R. (2017).
1106 Thirty-three years of 2-acetyl-1-pyrroline, a principal basmati aroma
1107 compound in scented rice (*Oryza sativa* L.): a status review. *Journal of the*

- 1108 *Science of Food and Agriculture*, 97(2), 384-395..
- 1109 Widjaja, R., Craske, J. D., & Wootton, M. (1996a). Changes in volatile components of
1110 paddy, brown and white fragrant rice during storage. *Journal of the Science of*
1111 *Food and Agriculture*, 71, 218-224.
- 1112 Widjaja, R., Craske, J., & Wootton, M. (1996b). Comparative studies on volatile
1113 components of non-fragrant and fragrant rices. *Journal of the Science of Food*
1114 *and Agriculture*, 70, 151-161.
- 1115 Wongpornchai, S., Dumri, K., Jongkaewattana, S., & Sirri, B. (2004). Effects of
1116 drying methods and storage time on the aroma and milling quality of rice
1117 (*Oryza sativa* L.) cv. Khao Dawk Mali 105. *Food Chemistry*, 87(3), 407-414.
- 1118 Wongpornchai, S., Sriseadka, T., & Choonvisase, S. (2003). Identification and
1119 quantitation of the rice aroma compound, 2-acetyl-1-pyrroline, in bread
1120 flowers (*Vallisneria spiralis* L.). *Journal of Agricultural and Food Chemistry*,
1121 51(2), 457-462.
- 1122 Yahya, F., Lu, T., Santos, R. C. D., Fryer, P. J., & Bakalis, S. (2010). Supercritical
1123 carbon dioxide and solvent extraction of 2-acetyl-1-pyrroline from pandan
1124 leaf: The effect of pre-treatment. *Journal of Supercritical Fluids*, 55(1), 200-
1125 207.
- 1126 Yang, D. S., Shewfelt, R. L., Lee, K. S., & Kays, S. J. (2008). Comparison of odor-
1127 active compounds from six distinctly different rice flavor types. *Journal of*
1128 *Agricultural and Food Chemistry*, 56(8), 2780-2787.
- 1129 Yau, N. J. N., & Liu, T. T. (1999). Instrumental and sensory analysis of volatile aroma

1130 of cooked rice. *Journal of Sensory Studies*, 14(2), 209-233.

1131 Yoshihashi, T. (2002). Quantitative analysis on 2-acetyl-1-pyrroline of an aromatic
1132 rice by stable isotope dilution method and model studies on its formation
1133 during cooking. *Journal of Food Science*, 67(2), 619-622.

1134 Yoshihashi, T., Huong, N. T. T., & Inatomi, H. (2002). Precursors of 2-acetyl-1-
1135 pyrroline, a potent flavor compound of an aromatic rice variety. *Journal of*
1136 *Agricultural and Food Chemistry*, 50(7), 2001-2004.

1137 Yoshihashi, T., Nguyen, T. T. H., & Kabaki, N. (2004). Area dependency of 2-acetyl-
1138 1-pyrroline content in an aromatic rice variety, Khao Dawk Mali 105. *Japan*
1139 *Agricultural Research Quarterly*, 38(2), 105-109.

1140 Zanan, R., Khandagale, K., Hinge, V., Elangovan, M., Henry, R. J., & Nadaf, A.
1141 (2016). Characterization of fragrance in sorghum (*Sorghum bicolor* (L.)
1142 Moench) grain and development of a gene-based marker for selection in
1143 breeding. *Molecular Breeding*, 36(11), 12.

1144

1145 **FIGURE CAPTIONS**

1146 **Figure 1.** A comparison of the (a) BADH2-dependent 2AP biosynthetic pathway
1147 (Bradbury, Gillies, Brusheet, Waters, & Henry, 2008) and the (b) BADH2-
1148 independent 2AP biosynthetic pathway (Sakthivel et al., 2009; Huang et al.,
1149 2008).

1150 **Figure 2.** Mechanism of 2-AP formation through the heterolactic pathway (Costello
1151 & Henschke, 2002)


1152 **Figure 3.** 2-AP Maillard Reaction formation pathways: a) formation of 1-pyrroline
1153 from ornithine (Schieberle, 1990); b) formation of 1-pyrroline from proline
1154 and 1-deoxyosone (Schieberle, 1995); c) ¹³C-labelled and unlabelled 2-AP
1155 formation from 1-pyrroline and ¹³C-glucose (Rewicki et al., 1993).

1156 **Figure 4.** 2-AP synthesis strategies: a) Buttery, Ling, Juliano, and Turnbaugh, 1983;
1157 b) De Kimpe, Stevens, and Keppens, 1993; c. De Kimpe, Stevens, and
1158 Keppens, 1993; d) De Kimpe, and Keppens, 1996; e) Favino, Fronza, Fuganti,
1159 Fuganti, Grasselli, and Mele, 1996; f) Deblander, Van Aeken, Adams, De
1160 Kimpe, & Abbaspour Tehrani (2015).


1161 LDA: lithium diisopropylamide; THF: tetrahydrofuran; PGA: immobilised penicillin
1162 G acylase; DMT: dimethyltitanocene; TFA: trifluoroacetic acid

1163

1164




1165 Figure 1




1166

1167 Figure 2



1168

1169 Figure 3



1170

1171 Figure 4

1172 **Table 1:** 2-AP concentrations in fragrant and non-fragrant rice
 1173

rice variety	2-AP concentration ($\mu\text{g}/\text{kg}$)	
	milled	brown
<i>fragrant rice</i>		
Basmati	60 ^a	170 ^a
	87 ^d	610 ^b
	588 ^g	119 ^h
	19–342 ^h	
	434 ^k	
Khao Dawk Mali 105	70 ^a	200 ^a
	87–532 ⁱ	
Malagkit Sungsong	90 ^a	200 ^a
		760 ^b
Milagross	70 ^a	
Seratus Malam	60 ^a	
Azucena	40 ^a	160 ^a
Hieri	40 ^a	100 ^a
Ir841-76-1	70 ^a	200 ^a
		560 ^b
Jasmine	156 ^d	550 ^h
	810 ^h	
Della	76 ^d	
Goolarah	691 ^e	
Yrf9	670 ^e	344 ^f
B5-3	2746 ^g	
Amber Aromatic (Lundberg)		345 ^h
Aromatic (Fowler Gourmet)	999 ^h	
Black Thai (Bulk)		259 ^h
Jasmati (Rice Tec)	526 ^h	
Kasmati (Rice Tec)	496 ^h	
Texmati (Rice Tec)	266 ^h	
Aychade	575–638 ^j	
Fidji	45–475 ^j	
Giano	28–336 ^j	
Kala Bhat	920 ^k	
Kali Kumud	732 ^k	
Amritbhog	787 ^k	
<i>non-fragrant rice</i>		
Calrose	<6 ^a	
California Long-Grain	0.6 ^c	
Pelde	15 ^e	

Texas Long Grain	<8 ^a
	6 ^b
Ariette	10.6 ^j
Ruille	24.7 ^j
Sonsali	72 ^k
Kolamb	125 ^k

1175

1176 Data are from the following references: ^aButtery et al., 1983; ^bButtery et al., 1986;

1177 ^cButtery et al., 1988; ^dTanchotikul and Hsieh, 1991; ^eWidjaja et al., 1996a; ^fWidjaja et

1178 al., 1996b; ^gTava and Bocchi, 1999; ^hBergman et al., 2000; ⁱYoshihashi et al., 2004;

1179 ^jMaraval et al., 2010; ^kMathure et al., 2014.

1180

1181 **Table 2:** 2-AP concentrations in foods other than rice
 1182

food sample	2-AP concentration ($\mu\text{g}/\text{kg}$)
wheat bread crusts	75 ^a
Mediterranean dried sausages	750 ^b
bread flowers (<i>Vallaris glabra</i> ktze)	3.36 (fresh) 26.1 (dry) ^c
palm wine	11.4 ^d
roasted Criollo cocoa beans	4.2 ^e
pandan leaves	40–450 ^f
pan-fried mushrooms	4.2–7.0 ^g
roasted in-shell peanuts	1920 ^h
roasted hazelnuts	85 ⁱ
cereal coffee brew	8 ^j
squid broth	97.3 ^k
dark chocolate	21 ^l
milk chocolate	3 ^l
cocoa liquor	11 ^l
Manuka honey	80–450 ^m
raw licorice	9.41 ⁿ
roasted almonds	12 (dry roasted) ^o 30 (oil roasted) ^o

1183
 1184 Data are from the following references: ^aSchieberle and Grosch, 1987; ^bStahnke,
 1185 2000; ^cWongpornchai et al., 2003; ^dLasekan et al., 2007; ^eFrauendorfer and
 1186 Schieberle, 2008; ^fYahya et al., 2010; ^gGrosshauser & Schieberle, 2013; ^hKaneko et
 1187 al., 2013; ⁱKiefl et al., 2013; ^jMajcher et al., 2013; ^kCarrascon et al., 2014; ^lLiu et al.,
 1188 2015; ^mRuckriemen et al., 2015; ⁿWagber et al., 2016; ^oErten et al., 2017.

1189
 1190
 1191