

The effect of caloric restriction on the forelimb skeletal muscle fibers of 1 the hypertrophic myostatin null mice

Article

Accepted Version

Creative Commons: Attribution-Noncommercial-No Derivative Works 4.0

Elashry, M. I., Matsakas, A., Wenisch, S., Arnhold, S. and Patel, K. (2017) The effect of caloric restriction on the forelimb skeletal muscle fibers of 1 the hypertrophic myostatin null mice. *Acta Histochemica*. ISSN 0065-1281 doi: 10.1016/j.acthis.2017.06.002 Available at <https://centaur.reading.ac.uk/70674/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

Published version at: <https://www.journals.elsevier.com/acta-histochemica>

To link to this article DOI: <http://dx.doi.org/10.1016/j.acthis.2017.06.002>

Publisher: Elsevier

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online

The effect of caloric restriction on the forelimb skeletal muscle fibers of the hypertrophic myostatin null mice.

Mohamed I. Elashry^{3, 4}; Antonios Matsakas²; Sabine Wenisch⁵; Stefan Arnhold⁴ and Ketan Patel¹

¹School of Biological Sciences, Hopkins building, Whiteknights, University of Reading, RG6 6UB, United Kingdom

² Hull York Medical School, University of Hull

³Anatomy and Embryology Department, Faculty of Veterinary Medicine, University of Mansoura 35516, Egypt

⁴Institute of Veterinary Anatomy, Histology and Embryology, Justus Liebig University of Giessen 35392, Germany

⁵Clinic of Small Animals, c/o Institute of Veterinary Anatomy, Histology and Embryology, Justus Liebig University of Giessen 35392, Germany

Corresponding author:

Mohamed Ismail Elashry

Institute of Veterinary Anatomy, Histology and Embryology

Justus Liebig University of Gießen

Frankfurter Str. 98,

35392 Gießen

Email: mohammed.elashry@vetmed.uni-giessen.de

Abstract

Skeletal muscle mass loss has a broad impact on body performance and physical activity. Muscle wasting occurs due to genetic mutation as in muscular dystrophy, age-related muscle loss (sarcopenia) as well as in chronic wasting disorders as in cancer cachexia. Food restriction reduces muscle mass underpinned by increased muscle protein break down. However the influence of dietary restriction on the morphometry and phenotype of forelimb muscles in a genetically modified *myostatin* null mice are not fully characterized. The effect of a five week dietary limitation on five anatomically and structurally different forelimb muscles was examined. C57/BL6 wild type (*Mstn*^{+/+}) and *myostatin* null (*Mstn*^{-/-}) mice were either given a standard rodent normal daily diet *ad libitum* (ND) or 60% food restriction (FR) for a 5 week period. *M. triceps brachii Caput laterale* (T.lateral), *M. triceps brachii Caput longum* (T.long), *M. triceps brachii Caput mediale* (T.medial), *M. extensor carpi ulnaris* (ECU) and *M. flexor carpi ulnaris* (FCU) were dissected, weighted and processed for immunohistochemistry. Muscle mass, fibers cross sectional areas (CSA) and myosin heavy chain types IIB, IIX, IIA and type I were analyzed. We provide evidence that caloric restriction results in muscle specific weight reduction with the fast myofibers being more prone to atrophy. We show that slow fibers are less liable to dietary restriction induced muscle atrophy. The effect of dietary restriction was more pronounced in *Mstn*^{-/-} muscles to implicate the oxidative fibers compared to *Mstn*^{+/+}. Furthermore, peripherally located myofibers are more susceptible to dietary induced reduction compared to deep fibers. We additionally report that dietary restriction alters the glycolytic phenotype of the *Mstn*^{-/-} into the oxidative form in a muscle dependent manner. In summary our study shows that calorie restriction alters muscle fiber profile of forelimb muscles of Myostatin null mice.

Keywords: Skeletal muscle, Myostatin, food restriction, myosin heavy chain

1. Introduction

Skeletal muscle the main protein reservoir in the body, is a highly adaptable tissue that changes its physical as well as composition based on physiological demands. Mechanical and nutritional stimuli cause an increase in muscle mass. In contrast undernutrition, aging and diseases as cancer cachexia reduces the muscle mass. Unbalanced nutrient intake with increased energy requirements enhances the muscle catabolism resulting in fiber atrophy and muscle mass loss (Koskela et al., 1990). Muscle loss also occurring in progressive wasting diseases e.g. cancer cachexia (Bruggeman et al., 2016) and age-related sarcopenia leads to a decrease in muscle mass and strength (Thomas, 2007). Additionally the role of physiological and metabolic stimuli e.g. physical activity, disuse and immobilization, food restriction, drugs and diseases cause molecular and cellular dysregulation which results in loss of muscle mass (Carmeli and Reznick, 1994). It has been reported that ageing induces muscle mass loss of 16%, 18%, 37% and 38% for soleus, extensor digitorum longus, plantaris and gastrocnemius muscles respectively with greater fiber area loss compared to the decrease in body mass (Brown and Hasser, 1996). Similarly it was reported that dietary restriction causes a decrease in muscle fiber size but not fiber number in chicken and rabbits (Tanaka et al., 1992 and Timson et al., 1983). In addition the glycolytic fibers were selectively decreased in the cross sectional area lead to a proportional increase in the area of oxidative fibers in bovine muscle (Greenwood et al., 2009). Muscle phenotype also changed according to the dietary challenges, It has been reported that dietary deprivation for 48 hours upregulates the expression of fast myosin heavy chain 2b mRNA with no change in fiber type composition for EDL and soleus muscles in rat (Mizunoya et al., 2013b) However 4 weeks fatty diet administration induced reduction in the fast MHC2b and improved the oxidative metabolism in the EDL of rat (Matsakas et al., 2013; Mizunoya et al., 2013a). Previously, it was shown that fasting induces muscle proteolysis via glucocorticoid activation (Wing and Goldberg, 1993) as well as causing an increase in the ATP-dependent proteolysis and upregulation of ubiquitin conjugates and polyubiquitins in rat skeletal muscle (Medina et al., 1991).

Myostatin is a TGF- β family member of secreted proteins that negatively regulates skeletal muscle proliferation (Sharma et al., 2001). *Myostatin* protein interference via knock out results in significant myofiber hyperplasia and hypertrophy (McPherron et al., 1997). Furthermore, systemic administration of *Myostatin* induced muscle atrophy in mice (Zimmers et al., 2002). Increased myostatin expression was also associated with muscle wasting in cancer cachexia, ageing, after HIV infection and in the course of chronic obstructive pulmonary disease (COPD) (Costelli et al., 2008; Gonzalez-Cadavid et al., 1998; Plant et al., 2010; Yarasheski et al., 2002). *Myostatin* knock out mice showed a greater atrophy in response to unloading (McMahon et al., 2003). Our previous study demonstrated a massive reduction in forelimb muscles mass of aged *myostatin* null compared to wild type mice (Elashry et al., 2009). These studies suggest that hypertrophic muscle has decreased adaptability compared to WT. In this study we hypothesized that five weeks of 60% dietary restriction would impact to a greater degree on the muscle of *Mstn*^{-/-} compared to WT. We provide evidence that caloric restriction results in muscle specific weight reduction with the fast myofibers being more prone to atrophy. We show that slow fibers are less liable to dietary restriction induced muscle atrophy. Furthermore, peripherally located myofibers are more susceptible to dietary induced reduction. Additionally, we report that dietary restriction alters the glycolytic phenotype of the myostatin null into more oxidative form in a muscle dependent manner.

2. Materials and Methods

2. 1. Animals

Three months old male transgenic myostatin knockout (*Mstn*^{-/-}) and C57/BL6 wild type littermates (*Mstn*^{+/+}) mice were bred in the biological resource unit, Reading University, UK. *Mstn*^{-/-} mice were generously provided from Dr Lee (McPherron et al., 1997). The experiments were performed under a project license from the United Kingdom Home Office in agreement with the Animals (Scientific Procedures) Act 1986 (license number 7516). The University of Reading Animal Care and Ethical Review Committee (AWERB) approved all procedures. The mice were divided into 4 subgroups: a normal diet group (N=8)

Mstn^{+/+} and *Mstn*^{-/-} kept with a normal diet with food and water provided *ad libitum* and a food restriction group (N=8) with an up to 60% reduction of the daily diet compared to normal mice for a 5 week period. Animals were humanely sacrificed via Schedule 1 killing.

2. 2. Tissue isolation and processing

Aged matched mice were sacrificed through schedule-1 killing following the approval of the ethics committee. Five forelimb muscles were carefully chosen to represent the arm and the forearm region *M. triceps brachii Caput longum*, T.long; *M. triceps brachii Caput laterale*, T.lateral; *M. triceps brachii Caput mediale*, T.medial; *M. extensor carpi ulnaris*, ECU; and *M. flexor carpi ulnaris*, FCU. The muscles were dissected, snap frozen using isopentane pre-cooled with liquid nitrogen and were embedded in tissue tech OCT (Sakura, VWR) using dry ice cooled ethanol. 10 µm transverse mid-belly cryosections for each muscle were obtained on a poly-L-lysine coated slides (VWR) and left to dry at room temperature for 1 hour and stored at -80°C.

2. 3. Immunohistochemical staining

Frozen muscle sections were washed in PBS for 10 minutes, permeabilized using buffer solution containing 20 mM Hepes, 300 mM sucrose, 50 mM NaCl, 3 mM MgCl₂ and 0.5% Triton-X100 (pH7) at for 15 minutes. Non-specific binding was blocked using wash buffer (5% fetal calf serum (v/v) and 0.05% Triton X-100 (v/v) in PBS) for 30 minutes at room temperature before the addition of antibodies. Myosin heavy chain (MHC) type I, IIA and IIB expressing myofibers were identified by incubating the slides with A4.840 mouse IgM (1:1), A.474 mouse IgG (1:4) and BFF3 mouse IgM (1:1) monoclonal primary antibodies (DSHB) respectively as previously described (Matsakas et al., 2009). The primary antibodies were tested separately against the corresponding MHC type on a muscle section and the following section with no primary antibody was served as a negative control (Fig. 1g-l). Additionally, MHC type IIA⁺ and IIB⁺ myofibers were stained together and MHC type I⁺ myofibers were stained on consecutive section. All primary antibodies were pre-blocked in washing buffer 30 minutes prior to use and incubated on samples overnight at 4°C. Primary antibodies were highlighted via incubation with Alexa Fluor 633

goat anti-mouse IgM (Molecular Probes A21046, 1:200) for MHC I and MHC IIB and Alexa Fluor 488 Goat-anti-mouse IgG (Molecular probes A11029, 1:200) for MHC IIA secondary antibodies in dark at room temperature for 45 minutes. Slides were mounted using a fluorescent mounting medium (Mowiol 4-88, Calbiochem) containing 2.5 µg/ml DAPI.

2. 4. Imaging and quantifications

Muscle sections were examined and photographed using a Zeiss Axioscop2 fluorescence microscope loaded with a digital camera and connected to a computer equipped with the axiovision computer software (Zeiss). The whole muscle cross sections was reconstructed, total number, number of IIB, IIA and type I fibers quantifications were performed by manual counting using the Photoshop CS6 extended software. Type IIX fibers were counted via subtraction of all other fibers from the total fiber number. Cross sectional area (CSA) for each fiber type for all muscles were measured using the axiovision software (Zeiss).

2. 5. Statistical analysis

The effect of genotype (Mstn^{+/+} vs. Mstn^{-/-}) and diet regime (ND vs. FR) on total muscle fiber numbers, CSA and the percentage of type IIB, IIX, IIA and I fibers were analyzed by two-way ANOVA for T.lateral, T.long, T.medial, ECU and FCU muscles. The effect of genotype, diet regime and region of myofiber (peripheral vs central) on CSA for each muscle were analyzed via three-way ANOVA. Multiple comparisons and the interactions were evaluated using Tukey's Post Hoc test. Chi-square test was performed to examine frequency distribution of fiber types CSA following FR. Statistical analysis was conducted by using Graph Pad prism 6 software. All values are presented as mean ± SEM, *P* values less than 0.05 were considered significant.

3. Results

3. 1. Dietary restriction induces body weight and muscle mass specific reduction

In order to evaluate the effect of dietary restriction on skeletal muscle of *Mstn*^{+/-} and *Mstn*^{-/-} mice for a five week period, average body weight measurements (g) revealed that *Mstn*^{-/-}ND mice were significantly heavier compared to *Mstn*^{+/-}ND mice ($P<0.001$). However both genotypes showed approximately 33% reduction in the body mass following dietary restriction (Fig. 1a). Individual muscle weight measurements showed that *Mstn*^{-/-}ND exhibited increases in muscle mass of T.lateral ($P<0.0001$), T.long ($P<0.0001$), T.medial ($P<0.05$), ECU ($P<0.01$) and FCU muscles ($P<0.0001$) compared to *Mstn*^{+/-}ND. To evaluate whether dietary restriction impact on different muscles mass. Our data showed that dietary restriction induced muscle specific weight loss e.g. T.lateral showed large reduction in muscle mass for *Mstn*^{-/-}FR and *Mstn*^{+/-}FR ($P<0.0001$ and $P<0.001$) respectively compared to genotype-matched ND as well as significant interaction ($P<0.001$) between diet and genotype (Fig.1b). Similarly, T.long muscle exhibited more pronounced reduction in the *Mstn*^{-/-}FR and *Mstn*^{+/-}FR ($P<0.0001$ and $P<0.001$) respectively compared to ND matched control also with significant interaction ($P<0.0001$) between the effect of diet and the genotype (Fig. 1c). Furthermore, ECU muscle showed more weight reduction following FR in *Mstn*^{-/-} than *Mstn*^{+/-} ($P<0.001$ and $P<0.05$) compared to genotype matched control (Fig.1e). On the other hand, T.medial and FCU muscles displayed a diet induced decreases in the muscle mass only in *Mstn*^{-/-} mice ($P<0.01$ and 0.0001) compared to genotype-matched control (Fig.1d and f). Representative immunofluorescent staining for T.long muscle of *Mstn*^{+/-}ND show MHC types IIA, IIB and I expressing myofibers compared to the negative control (Fig. 1g-l).

3. 2. Caloric restriction does not alter muscle fiber number

Based on previous experiments (Elashry et al., 2009) we examined whether a FR related muscle mass decrease was due to either loss in fiber number or loss of fiber size. Quantification of the average fibers number revealed that myostatin deletion increases muscle fiber number for T.lateral and T.long ($P<0.0001$), ECU and FCU ($P<0.01$) muscles on normal diet could be detected. In contrast, T.medial muscle displayed no fiber number change in *Mstn*^{-/-}. However dietary restriction for five weeks caused no change in the fibers

number for all muscles of both genotypes (Fig. 2a, b, c, d and e). Representatives whole muscle reconstruction showed clearly the effect of *myostatin* deletion and dietary restriction on the cross section of the FCU muscle of *Mstn*^{+/+} and *Mstn*^{-/-} compared to normal diet control (Fig. 2f-i).

3. 3. Fast fibers are more prone to caloric restriction-induced muscle mass loss

Next we evaluated the impact of dietary restriction on individual muscle fiber size via quantification of the CSA (μm^2) of IIB, IIX, IIA and type I fibers per muscle. A common trend that dietary restriction reduced the CSA of glycolytic type IIB fibers in *Mstn*^{-/-} and *Mstn*^{+/+} ($P < 0.0001$) compared to controls was observed (Fig. 3). Type IIB fibers CSA measurement of T.lateral muscle displayed pronounced reduction (2074 ± 7 to $1232 \pm 3 \mu\text{m}^2$) in *Mstn*^{-/-} ($P < 0.0001$) compared to (1605 ± 4 to $1246 \pm 3 \mu\text{m}^2$) in *Mstn*^{+/+} ($P < 0.0001$) following FR. These results were supported with significant interaction ($P < 0.0001$) indicating the effect of FR on *Mstn*^{-/-} compared to *Mstn*^{+/+} (Fig. 3a). Furthermore, T.long muscle showed similar reduction in type IIB fibers CSA (3596 ± 52 to $1304 \pm 3 \mu\text{m}^2$) and (3118 ± 8 to $1085 \pm 3 \mu\text{m}^2$) for *Mstn*^{-/-} and *Mstn*^{+/+} respectively ($P < 0.0001$) following dietary restriction and compared to genotype-matched control (Fig. 3b). On the other hand, measurements of type IIB fibers CSA showed more loss in the T.medial and ECU muscles of *Mstn*^{-/-} ($P < 0.0001$) compared to *Mstn*^{+/+} following FR. E.g. T.medial muscle of *Mstn*^{-/-} demonstrated 1647 ± 42 to $1090.6 \pm 3 \mu\text{m}^2$ CSA reduction ($P < 0.0001$) compared to 1479.5 ± 4 to $1257 \pm 3 \mu\text{m}^2$, ($P < 0.0001$) reduction in *Mstn*^{+/+} provided with significant interaction ($P < 0.0001$) between the effect of diet and genotype (Fig. 3c and d). Next, type IIX fibers CSA measurement for T.lateral, ECU and FCU muscles displayed significant decreases in the CSA for both *Mstn*^{-/-} ($P < 0.05$, 0.01 and 0.0001) and *Mstn*^{+/+} ($P < 0.01$ and 0.0001) following dietary restriction compared to control (Fig. 3a, d and e) however IIX fibers measurements showed more loss in T.long and T.medial muscles in the *Mstn*^{-/-} ($P < 0.0001$ and $P < 0.01$) compared to *Mstn*^{+/+} following FR (Fig. 3b and c). Additionally, we examined whether the effect of dietary limitation involves the oxidative fibers, analysis of type IIA fibers CSA demonstrated reduction in the T.long and T.medial muscles

($P < 0.0001$ and 0.05) respectively only in the *Mstn*^{-/-} compared to *Mstn*^{+/+} under the FR regime (Fig. 3b and c). Surprisingly, IIA fibers CSA for ECU and FCU muscles showed significant increase in *Mstn*^{-/-}FR ($P < 0.01$ and 0.0001) compared to *Mstn*^{-/-}ND (Fig. 3d and e). Furthermore, we tested the oxidative capacity of the muscles following dietary restriction by analyzing type I (slow oxidative) fibers CSA, our results displayed large reduction in T.long muscle of *Mstn*^{-/-} ($P < 0.0001$) compared to *Mstn*^{+/+} (Fig. 3b). These data demonstrated that glycolytic fibers were more liable to dietary restriction followed by IIX, IIA and I oxidative fibers. Furthermore the impact of dietary restriction was more pronounced in some of the *Mstn*^{-/-} muscles to implicate the oxidative fibers (IIA and I) compared to *Mstn*^{+/+}. In order to confirm our data, T.long fibers CSA measurements were categorized into small ($0-1000$ and $1000-2000 \mu\text{m}^2$), medium ($2000-4000 \mu\text{m}^2$) and large ($4000-6000 \mu\text{m}^2$) for type IIB, small ($0-1000 \mu\text{m}^2$), medium ($1000-2000 \mu\text{m}^2$) and large fibers ($2000-4000 \mu\text{m}^2$) for IIX, IIA and type I fibers. Frequency distribution assessment of type IIB, IIX, IIA and I fibers of T.long muscle showed significant CSA shift from large size into the small size population ($P < 0.0001$) following dietary restriction compared to genotype-matched control (Fig. 3f, g, h and i).

3. 4. Dietary restriction causes peripheral myofiber atrophy

Our results show that dietary restriction primarily target fast MHCIIB fibers. To address whether the anatomical location of the fiber had an effect on the myofibers CSA following caloric restriction, we analyzed the CSA of the peripheral (PF) and central (CT) fibers within the mid-belly of the corresponding muscle. Our results revealed that in T.lateral, T.long and ECU muscles but not T.medial muscle showed a common observation that type IIB myofibers CSA were larger in the PF ($P > 0.0001$) compared to CT region for *Mstn*^{-/-} and *Mstn*^{+/+} on normal diet (Fig. 4a, c, f and i). Furthermore, FCU muscles demonstrated larger CSA of PF IIB compared to CT fibers ($P < 0.0001$) only in the *Mstn*^{-/-} on a normal diet (Fig. 4l). Furthermore, FR induced significant reduction in the CSA of PF IIB fibers ($P < 0.0001$) compared to CT fibers e.g. T.lateral muscle showed (850 ± 87 vs. $1247 \pm 67 \mu\text{m}^2$) and (609 ± 1 vs. $1348 \pm 2 \mu\text{m}^2$) for *Mstn*^{-/-} and *Mstn*^{+/+} respectively also significant interaction was detected ($P < 0.0001$) indicating the

interference of diet and genotype (Fig. 4a). Next we analyzed type IIX fibers, larger IIX fibers CSA of both regions in T.lateral muscle of *Mstn*^{+/+} (P<0.0001) and T.long muscle of *Mstn*^{-/-} (P<0.0001) on normal diet were observed. Also, PF IIX fibers of *Mstn*^{-/-}FR showed more reduction in the CSA in T.lateral and T.long muscles (P<0.05 and P<0.0001) compared to *Mstn*^{-/-}ND respectively (Fig. 4b and d). However, reduction in the PF IIX fibers of both genotypes were noticed in the T.medial (P<0.01), ECU and FCU muscles (P<0.0001) following FR (Fig. 4g, j and m). Moreover, CT type IIX fibers displayed reduction in T.long and T.medial muscles of *Mstn*^{-/-} (P<0.0001) as well as T.lateral muscle of *Mstn*^{+/+} (P<0.0001) following FR. Additionally, PF type IIX fiber CSA demonstrated further reduction in T.medial (P<0.001 and P<0.0001) and ECU muscle (P<0.0001) of both genotypes compared to CT fibers (Fig. 4 g and j) following FR. On the other hand, more reduction in the PF IIX fibers was detected in T.lateral and FCU muscles of only *Mstn*^{-/-} (P<0.0001) compared to CT fibers following FR (Fig. 4b and m), also significant interaction (P<0.01) indicating the effect of diet on *Mstn*^{-/-}. In contrast, analysis of type IIA fibers showed some variations, PF IIA fibers of T.medial muscle displayed more reduction in the CSA (P<0.01) compared to CT fibers only in *Mstn*^{+/+} (Fig. 4h) however PF fibers of ECU muscle showed more CSA reduction compared to CT fibers (P<0.0001 and P<0.001) for *Mstn*^{-/-} and *Mstn*^{+/+} respectively following dietary restriction (Fig. 4k). However, FCU muscle displayed larger CSA reduction in PF IIA fibers only in the *Mstn*^{-/-} compared to CT fibers after dietary restriction (Fig. 4n). Type I fibers was only localized in the central part of the muscle therefore were excluded from this comparison.

3. 5. Caloric restriction induces a slow phenotype conversion in a muscle-specific manner

Our previous study demonstrated that *myostatin* deletion induced various degrees of slow to fast fiber type shift in the forelimb muscles (Elashry et al., 2009). Therefore, we examined whether dietary restriction alters the muscle phenotype of the *Mstn*^{-/-} compared to *Mstn*^{+/+}. Quantification of the average percentage of each fiber type population per muscle demonstrated percentage increases of type IIB fibers in the T.lateral, T.long, T.medial, ECU and FCU

muscles *Mstn*^{-/-}ND (P<0.0001) compared to *Mstn*^{+/+}ND. However FR altered the fiber type composition for both genotypes (P<0.001) e.g. T.lateral muscle demonstrated a reduction in the type IIB fibers (P<0.0001 and P<0.05) and increases of type IIX fibers (P<0.0001 and P<0.05) for *Mstn*^{-/-}FR and *Mstn*^{+/+}FR respectively compared to genotype matched control (Fig. 5a and b). Also FR induced an increase in the type IIA fibers of *Mstn*^{-/-}FR compared to ND, however *Mstn*^{+/+}FR showed a reduction in type IIA fibers compared to ND. These data were confirmed by significant interaction between both genotypes (P<0.01, Fig. 5c). Furthermore, T.long muscle showed a reduction in type IIB fibers (P<0.0001) with an increase in type IIX fibers (P<0.001 and P<0.0001) for *Mstn*^{-/-}FR and *Mstn*^{+/+}FR respectively compared to genotype-matched ND (Fig. 5d and e). Although there were no significant changes, *Mstn*^{-/-}FR demonstrated marked increases in type IIA fibers compared to *Mstn*^{-/-}ND. However *Mstn*^{+/+}FR showed significant increase in type IIA fibers (P<0.01) compared to *Mstn*^{+/+}ND (Fig. 5f) as well as FR resulted in marked increase in type I fibers in both genotypes (P<0.01) compared to ND control (Fig. 5g). On the other hand, T.medial muscle analysis revealed fiber type population of 33%, 34%, 26% and 7% for IIB, IIX, IIA and I fibers respectively in the *Mstn*^{+/+}ND and showed almost no fiber type shift following dietary restriction in the *Mstn*^{-/-} however *Mstn*^{+/+} FR displayed fiber shift into type I fibers (IIB→IIX→IIA→I) following the same treatment (P<0.05) compared to ND. These data was supported by significant interaction (P<0.05) between diet and genotype effect (Fig. 5h, i and j and k). On the other hand, ECU muscle displayed a different fiber shift pattern with dietary restriction whereas IIB to IIX fibers transformation in the *Mstn*^{-/-} (P<0.0001 and P<0.01) respectively (Fig. 5l and m) and exhibited type IIA to type I fiber shift in *Mstn*^{+/+} (P<0.001) compared to ND (Fig. 5n and o). Similarly, FCU muscle FR induced type IIB to IIX fiber shift for *Mstn*^{-/-} (P<0.0001) compared to ND (Fig. 5p and q) however demonstrated fibers shift from IIB directly into IIA fibers but not type IIX fibers in *Mstn*^{+/+} (P<0.0001 and P<0.05) compared to *Mstn*^{+/+}ND (Fig. 5r and s). Representative double immunofluorescent images of T.long muscle for *Mstn*^{+/+} and *Mstn*^{-/-} showed the transformation of the glycolytic fibers type IIB and IIX into the oxidative type IIA fibers following dietary restriction. A negative control without addition of primary

antibodies was performed on a consecutive muscle section (Fig. 5t, u, v, w and x).

4. Discussion

Dietary intake plays a critical role to maintain the physiological composition of the skeletal muscle. Previous studies revealed that *myostatin* deletion results in an increase in muscle mass due to an increase in fiber number and fiber size (McPherron et al., 1997). However, the impact of dietary restriction on anatomically and structurally different forelimb muscles of the *myostatin* null are not fully characterized. Our results suggest that the impact of dietary restriction was greater in the *Mstn*^{-/-} than in wild-type muscle. In agreement with our results, previous work showed that dietary limitation caused more muscle loss in the hind limb muscles of the *Mstn*^{-/-} compared to *Mstn*^{+/+} suggesting higher catabolic activities (Matsakas et al., 2013). These results suggest that the hypertrophic muscle of *Mstn*^{-/-} is more prone to protein breakdown than that of wild-types. In support of this notion is the finding that *Mstn*^{-/-} mice exhibited more muscle mass loss during hind limb unloading (McMahon et al., 2003). Our results indicate that muscle size (enlarged) and composition (fast) might play a role in the degree of muscle loss following dietary limitation. This in line with previous reports indicating that fast twitch IIB fibers of rodents and monkeys are more sensitive to starvation compared to slow twitch fiber (Goldspink and Ward, 1979; Li and Goldberg, 1976; Matsakas et al., 2013; McKiernan et al., 2012). However we show that there is also a regional specificity of MHC types in terms of atrophy.

We propose that the dietary restriction induced muscle loss is regulated via inhibition of IGF-1 signaling which results in upregulation of atrogen-1 and MuRF-1 expression. A similar study showed that undernutrition for 28 days causes reduced growth rate via regulation of IGF-1 and myostatin expressions (Jeanplong et al., 2015). We shows that the oxidative fibers are resistant to dietary restriction. In agreement with our results, it has been shown that the atrophy resistance of the oxidative fibers was due to a protection afforded by peroxisome proliferator-activated receptor gamma coactivator-1 alpha (PGC1α) (Wang and Pessin, 2013). Unexpectedly, dietary restriction causes

CSA increases in type IIA fibers for ECU and FCU of *Mstn*^{-/-} which might be an adaptive mechanism to compensate the decrease in glycolytic fibers number.

Analysis of the muscles phenotypes revealed a general conversion toward slow oxidative phenotype following dietary restriction. However, we observed only the degree and the mechanism of fiber shift was dependent on the original fiber type population e.g. T.long and T.lateral muscles displayed IIB→IIX→IIA in the *Mstn*^{-/-} however only showed a IIB→IIX fiber type conversion in *Mstn*^{+/+}. Moreover, ECU and FCU muscles showed a IIB→IIX fiber shift in *Mstn*^{-/-} however demonstrated a IIA→I and IIB→IIA shift respectively in *Mstn*^{+/+}. These data indicate that myostatin deletion and dietary limitation interactions alter muscle adaptation and plasticity. Muscle oxidative conversion was previously reported with ageing, although there was trophic activity in type II fibers, additionally there was a relative increase in type I fibers (Kallman et al., 1990). Furthermore, a recent study in muscular dystrophy showed that muscle atrophy was accompanied with a myofiber oxidative shift perhaps due to an increase in the expression of PGC1α (Rocchi et al., 2016). On the other hand, soleus muscle immobilization, hind limb unloading and spinal cord transection induced muscle atrophy plus type I into type II fibers transformation see review by (Cho et al., 2016). However in our model the physical performance and muscle function were preserved compared to previous models which resulted in improved phenotype particularly in *Mstn*^{-/-}. It could be explain that muscle adaptability is dependent on the extend of the external stimuli therefore the high demand for protein supplement in the *Mstn*^{-/-} induced fiber atrophy and fiber type transformation. This is far greater compared to *Mstn*^{+/+} such a change in muscle metabolism energy saving mechanisms are required.

Our data discussed the impact of caloric restriction in combination with *myostatin* deletion on five muscles from different anatomical locations. We provide evidence that 60% of dietary limitation results in body weight and muscle specific mass reduction with fast myofibers more prone to atrophy compared to slow myofibers. The effect of dietary restriction was more pronounced in some of the *Mstn*^{-/-} muscles to implicate the oxidative fibers compared to *Mstn*^{+/+} mice. Furthermore, peripheral myofibers are more

susceptible to dietary induced reduction compared to deep fibers. We provide evidence that dietary restriction alters the fast phenotype of the *Mstn*^{-/-} into the slower form in a muscle dependent manner. These results allow the conclusion that caloric reduction alters the muscle fiber composition and the oxidative pattern to compensate muscle mass loss which might impact on muscle function particularly in the hypermuscular *myostatin* null animals.

Competing interest

All the authors have declared no conflict of interest regarding the publication of this paper.

Acknowledgement

We gratefully thank University of Reading, Wellcome Trust (078649), UK, University of Mansoura, Egypt, Engemann research foundation and Faculty of Veterinary medicine, University of Gießen, Germany for generous funding of this work.

5. References

- Brown M, Hasser EM. Complexity of age-related change in skeletal muscle. The journals of gerontology. Series A, Biological sciences and medical sciences 1996; 51(2):B117-23.
- Bruggeman AR, Kamal AH, LeBlanc TW, Ma JD, Baracos VE, Roeland EJ. Cancer Cachexia: Beyond Weight Loss. Journal of oncology practice 2016; 12(11):1163–71.
- Carmeli E, Reznick AZ. The physiology and biochemistry of skeletal muscle atrophy as a function of age. Proceedings of the Society for Experimental Biology and Medicine. Society for Experimental Biology and Medicine (New York, N.Y.) 1994; 206(2):103–13.
- Cho SH, Kim JH, Song W. In Vivo Rodent Models of Skeletal Muscle Adaptation to Decreased Use. Endocrinology and metabolism (Seoul, Korea) 2016; 31(1):31–7.

- Costelli P, Muscaritoli M, Bonetto A, Penna F, Reffo P, Bossola M, et al. Muscle myostatin signalling is enhanced in experimental cancer cachexia. *European journal of clinical investigation* 2008; 38(7):531–8.
- Elashry MI, Otto A, Matsakas A, El-Morsy SE, Patel K. Morphology and myofiber composition of skeletal musculature of the forelimb in young and aged wild type and myostatin null mice. *Rejuvenation research* 2009; 12(4):269–81.
- Goldspink G, Ward PS. Changes in rodent muscle fibre types during post-natal growth, undernutrition and exercise. *The Journal of physiology* 1979; 296:453–69.
- Gonzalez-Cadavid NF, Taylor WE, Yarasheski K, Sinha-Hikim I, Ma K, Ezzat S, et al. Organization of the human myostatin gene and expression in healthy men and HIV-infected men with muscle wasting. *Proceedings of the National Academy of Sciences of the United States of America* 1998; 95(25):14938–43.
- Greenwood PL, Tomkins NW, Hunter RA, Allingham PG, Harden S, Harper GS. Bovine myofiber characteristics are influenced by postweaning nutrition. *Journal of animal science* 2009; 87(10):3114–23.
- Jeanplong F, Osepchook CC, Falconer SJ, Smith HK, Bass JJ, McMahon CD, et al. Undernutrition regulates the expression of a novel splice variant of myostatin and insulin-like growth factor 1 in ovine skeletal muscle. *Domestic animal endocrinology* 2015; 52:17–24.
- Kallman DA, Plato CC, Tobin JD. The role of muscle loss in the age-related decline of grip strength: cross-sectional and longitudinal perspectives. *Journal of gerontology* 1990; 45(3):M82-8.
- Koskelo EK, Saarinen UM, Siimes MA. Skeletal muscle wasting and protein-energy malnutrition in children with a newly diagnosed acute leukemia. *Cancer* 1990; 66(2):373–6.
- Li JB, Goldberg AL. Effects of food deprivation on protein synthesis and degradation in rat skeletal muscles. *The American journal of physiology* 1976; 231(2):441–8.
- Matsakas A, Foster K, Otto A, Macharia R, Elashry MI, Feist S, et al. Molecular, cellular and physiological investigation of myostatin propeptide-mediated

muscle growth in adult mice. *Neuromuscular disorders* NMD 2009; 19(7):489–99.

Matsakas A, Romanello V, Sartori R, Masiero E, Macharia R, Otto A, et al. Food restriction reverses the hyper-muscular phenotype and force generation capacity deficit of the myostatin null mouse. *International journal of sports medicine* 2013; 34(3):223–31.

McKiernan SH, Colman RJ, Aiken E, Evans TD, Beasley TM, Aiken JM, et al. Cellular adaptation contributes to calorie restriction-induced preservation of skeletal muscle in aged rhesus monkeys. *Experimental gerontology* 2012; 47(3):229–36.

McMahon CD, Popovic L, Oldham JM, Jeanplong F, Smith HK, Kambadur R, et al. Myostatin-deficient mice lose more skeletal muscle mass than wild-type controls during hindlimb suspension. *American journal of physiology. Endocrinology and metabolism* 2003; 285(1):E82-7.

McPherron AC, Lawler AM, Lee SJ. Regulation of skeletal muscle mass in mice by a new TGF-beta superfamily member. *Nature* 1997; 387(6628):83–90.

Medina R, Wing SS, Haas A, Goldberg AL. Activation of the ubiquitin-ATP-dependent proteolytic system in skeletal muscle during fasting and denervation atrophy. *Biomedica biochimica acta* 1991; 50(4-6):347–56.

Mendias CL, Kayupov E, Bradley JR, Brooks SV, Claflin DR. Decreased specific force and power production of muscle fibers from myostatin-deficient mice are associated with a suppression of protein degradation. *Journal of applied physiology (Bethesda, Md. 1985)* 2011; 111(1):185–91.

Mizunoya W, Iwamoto Y, Shirouchi B, Sato M, Komiya Y, Razin FR, et al. Dietary fat influences the expression of contractile and metabolic genes in rat skeletal muscle. *PloS one* 2013a; 8(11):e80152.

Mizunoya W, Sawano S, Iwamoto Y, Sato Y, Tatsumi R, Ikeuchi Y. Effect of 48-h food deprivation on the expressions of myosin heavy-chain isoforms and fiber type-related factors in rats. *Journal of nutritional science and vitaminology* 2013b; 59(4):289–98.

Plant PJ, Brooks D, Faughnan M, Bayley T, Bain J, Singer L, et al. Cellular markers of muscle atrophy in chronic obstructive pulmonary disease.

American journal of respiratory cell and molecular biology 2010; 42(4):461–71.

Rocchi A, Milioto C, Parodi S, Armirotti A, Borgia D, Pellegrini M, et al. Glycolytic-to-oxidative fiber-type switch and mTOR signaling activation are early-onset features of SBMA muscle modified by high-fat diet. *Acta neuropathologica* 2016; 132(1):127–44.

Sharma M, Langley B, Bass J, Kambadur R. Myostatin in muscle growth and repair. *Exercise and sport sciences reviews* 2001; 29(4):155–8.

Tanaka N, Hayakawa T, Zyo K, Hori S. The morphological and biochemical changes in skeletal muscle fibers by dietary protein restriction. *Journal of nutritional science and vitaminology* 1992; 38(5):523–8.

Thomas DR. Loss of skeletal muscle mass in aging: examining the relationship of starvation, sarcopenia and cachexia. *Clinical nutrition (Edinburgh, Scotland)* 2007; 26(4):389–99.

Timson BF, Chi MS, Bowlin BK. The effect of reduced dietary protein on the anterior latissimus dorsi muscle fibers in the Single Comb White Leghorn pullet. *Poultry science* 1983; 62(11):2230–3.

Wang Y, Pessin JE. Mechanisms for fiber-type specificity of skeletal muscle atrophy. *Current opinion in clinical nutrition and metabolic care* 2013; 16(3):243–50.

Wing SS, Goldberg AL. Glucocorticoids activate the ATP-ubiquitin-dependent proteolytic system in skeletal muscle during fasting. *The American journal of physiology* 1993; 264(4 Pt 1):E668-76.

Yarasheski KE, Bhasin S, Sinha-Hikim I, Pak-Loduca J, Gonzalez-Cadavid NF. Serum myostatin-immunoreactive protein is increased in 60-92 year old women and men with muscle wasting. *The journal of nutrition, health & aging* 2002; 6(5):343–8.

Zimmers TA, Davies MV, Koniaris LG, Haynes P, Esquela AF, Tomkinson KN, et al. Induction of cachexia in mice by systemically administered myostatin. *Science (New York, N.Y.)* 2002; 296(5572):1486–8.

Figure legends

Figure 1 Dietary restriction induces body weight and muscle mass specific reduction.

a Average body weight (g) for normal diet (ND) and food restricted (FR) *Mstn*^{+/+} and *Mstn*^{-/-} mice (N=4 for all groups). Significant increase in the *Mstn*^{-/-} ($P<0.001$) compared to the *Mstn*^{+/+} on normal diet. FR decreases the body weight for *Mstn*^{+/+} and *Mstn*^{-/-} compared to genotype-matched ND mice. **b-f** Average muscle weight (g) for **b** T.lateral, **c** T.long, **d** T.medial, **e** ECU, and **f** FCU muscles. Specific muscle mass increases in *Mstn*^{-/-} ND compared to *Mstn*^{+/+} ND. FR induces more muscle mass reduction in T.lateral and T.long muscles of *Mstn*^{-/-} (diet and genotype interaction $P<0.001$ and 0.0001) respectively compared to genotype matched control. *Mstn*^{-/-} display more muscle loss in T.medial, ECU and FCU muscles compared to *Mstn*^{+/+} following FR. **g-l** Immunofluorescent images for T.long muscle of *Mstn*^{+/+}ND against A.474, BFF3 and A.4840 specific MHC antibodies show **g** type IIA⁺ (green), **h** type IIB⁺ (red) and **i** type I⁺ fibers (green) compared to **j**, **k** and **l** the corresponding negative control (NC) respectively. DAPI was used to visualize the myonuclei. All values displayed as mean \pm SEM. * = $P<0.05$, ** = $P<0.01$, *** = $P<0.001$ and ****= $P<0.0001$). Scale bar in **g-l** =20 μ m.

Figure 2 Caloric restriction does not alter muscle fiber number.

a-e Average total myofiber number for **a** T.lateral, **b** T.long, **c** T.medial, **d** ECU, and **e** FCU muscles of ND and FR *Mstn*^{+/+} and *Mstn*^{-/-} mice. The muscles of *Mstn*^{-/-} show increases in the myofiber number compared to *Mstn*^{+/+}. T.medial muscle display no fiber number change in both genotypes. FR reveals no fiber number loss compared to ND mice. **f-i** Reconstructive immunolabelled images for FCU muscle of *Mstn*^{+/+} and *Mstn*^{-/-} ND and FR against MHCIIIB (red), MHCIIA (green), MHCIIIX (unstained). *Myostatin* deletion increases the total CSA of the muscle compare to wild type (**f** vs **g**). FR reduces the CSA of *Mstn*^{+/+} (**f** vs. **h**) and *Mstn*^{-/-} (**g** vs. **i**). All values displayed as mean \pm SEM. *= $P<0.05$, **= $P<0.01$ and ****= $P<0.0001$).

Figure 3 Fast fibers are more prone to caloric restriction-induced muscle mass loss.

a-e Average CSA measurements (μm^2) for type IIB, IIX, IIA and type I fibers of **a** T.lateral, **b** T.long, **c** T.medial, **d** ECU and **e** FCU muscles of *Mstn*^{+/+} and *Mstn*^{-/-} ND and FR mice. FR causes significant reduction in the CSA of type IIB fibers in T.lateral, T.medial and ECU muscles of *Mstn*^{-/-} compared to *Mstn*^{+/+}. FR induces significant type IIX reduction in T.medial muscle of only *Mstn*^{-/-}, T.lateral muscle of only *Mstn*^{+/+} and in T.long, ECU and FCU muscles for both genotypes compared to ND. FR causes reduction in CSA of IIA fibers of T.long and T.medial muscles of *Mstn*^{-/-} however increases the CSA of ECU and FCU muscles. FR reveals no change in type IIA fibers for all *Mstn*^{+/+} muscles. FR induces only significant reduction in type I fibers of T.long muscle but not significant (NS) in T.medial muscle. **f-i** T.long muscle frequency distribution of fiber CSA measurements (μm^2) for **f** type IIB, **g** type IIX, **h** type IIA and **i** type I fibers for of *Mstn*^{+/+} and *Mstn*^{-/-} ND and FR mice. FR induces fibers shift into small size population (arrow marked) compared to ND for both genotypes. All values displayed as mean \pm SEM. *=*P*< 0.05, ** = *P*< 0.01 and ****=*P*<0.0001.

Figure 4 Dietary restriction causes peripheral myofiber atrophy.

a-n Average CSA (μm^2) measurements for peripheral (PF) and central fibers (CT) type IIB, IIX and IIA fibers for the muscles T.lateral **a, b**; T.long **c, d, e**; T.medial **f, g, h**; ECU **i, j, k** and FCU **l, m, n** of ND and FR *Mstn*^{+/+} and *Mstn*^{-/-} mice. For all muscle except T.medial, PF IIB fibers show CSA increases in *Mstn*^{-/-} and *Mstn*^{+/+} ND compared to CT fibers. PF IIB fibers of T.lateral, ECU and FCU muscles display smaller CSA compared to CT fibers following dietary restriction (**a, c, f, i** and **l**). FR reduces the CSA of PF IIX fibers compared to CT fibers for *Mstn*^{-/-} and *Mstn*^{+/+} (**b, d, g, j** and **m**). FR show larger CSA in the CT IIA fibers of ECU and FCU muscles of *Mstn*^{-/-} (**k** and **n**). All values displayed as mean \pm SEM. *=*P*< 0.05, ** = *P*< 0.01, *** = *P*< 0.001 and ****=*P*<0.0001.

Figure 5 Caloric restriction induces a slow phenotype conversion in a muscle-specific manner.

Average percentages of type IIB, IIX, IIA, and I fibers for T.lateral **a, b, c**; T.long **d, e, f, g**; T.medial **h, i, j, k**; ECU **l, m, n, o**; and FCU muscles **p, q, r, s** of ND and FR *Mstn*^{+/+} and *Mstn*^{-/-} mice. *Mstn*^{-/-} ND show increases in type IIB and IIX fibers compared to *Mstn*^{+/+}ND for all muscles. FR induces an increase in type IIX and type IIA fibers of T.lateral muscle in *Mstn*^{-/-} and only type IIX increases in *Mstn*^{+/+}. T.long muscle exhibits type IIB to IIA fiber shift and IIB to IIX fiber shift for *Mstn*^{+/+} and *Mstn*^{-/-} following FR respectively. FR causes an increase in type I fibers for T.medial muscle of *Mstn*^{+/+}. ECU muscles show IIB to IIX fiber type shift for *Mstn*^{-/-} compared to IIA to I fibers for *Mstn*^{+/+} following FR. FCU muscle displays IIB to IIA fiber shift for *Mstn*^{+/+} following FR. **t-x** Representative double immunofluorescent images of mid-belly section for T.long muscle in ND and FR *Mstn*^{+/+} and *Mstn*^{-/-} mice. FR induces myofiber transformation from glycolytic type IIB fibers (red) into oxidative type IIA. Type IIX fibers remained unstained (black). A consecutive muscle section with only secondary antibodies and DAPI was used as a negative control (NC). Scale bar =100 μm. All values displayed as mean± SEM.




Figure 1 Dietary restriction induces body weight and muscle mass specific reduction.




Figure 2 Caloric restriction does not alter muscle fiber number.




Figure 3 Fast fibers are more prone to caloric restriction-induced muscle mass loss.





Figure 4 Dietary restriction causes peripheral myofiber atrophy.



637

638 **Figure 5 Caloric restriction induces a slow phenotype conversion in a**
 639 **muscle-specific manner.**

640

641