

Trace ideal criteria for embeddings and composition operators on model spaces

Article

Accepted Version

Creative Commons: Attribution-Noncommercial-No Derivative Works 4.0

Aleman, A., Lyubarskii, Y., Malinnikova, E. and Perfekt, K.-M. (2016) Trace ideal criteria for embeddings and composition operators on model spaces. *Journal of Functional Analysis*, 270 (3). pp. 861-883. ISSN 0022-1236 doi: 10.1016/j.jfa.2015.11.006 Available at <https://centaur.reading.ac.uk/71469/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

To link to this article DOI: <http://dx.doi.org/10.1016/j.jfa.2015.11.006>

Publisher: Elsevier

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online

TRACE IDEAL CRITERIA FOR EMBEDDINGS AND COMPOSITION OPERATORS ON MODEL SPACES

ALEXANDRU ALEMAN, YURII LYUBARSKII, EUGENIA MALINNIKOVA,
KARL-MIKAEL PERFEKT

ABSTRACT. Let K_ϑ be a model space generated by an inner function ϑ . We study the Schatten class membership of embeddings $I : K_\vartheta \hookrightarrow L^2(\mu)$, μ a positive measure, and of composition operators $C_\varphi : K_\vartheta \rightarrow H^2(\mathbb{D})$ with a holomorphic function $\varphi : \mathbb{D} \rightarrow \mathbb{D}$. In the case of one-component inner functions ϑ we show that the problem can be reduced to the study of natural extensions of I and C_φ to the Hardy-Smirnov space $E^2(D)$ in some domain $D \supset \mathbb{D}$. In particular, we obtain a characterization of Schatten membership of C_φ in terms of Nevanlinna counting function. By example this characterization does not hold true for general ϑ .

1. INTRODUCTION

Let $\mathbb{D} = \{z : |z| < 1\}$ be the unit disk and $\mathbb{T} = \{z : |z| = 1\}$ be the unit circle. A bounded analytic function ϑ in \mathbb{D} is said to be *inner* if its non-tangential boundary values satisfy $|\vartheta| = 1$ almost everywhere on \mathbb{T} . We denote by $H^2(\mathbb{D})$ the Hardy space on \mathbb{D} and by $K_\vartheta = H^2(\mathbb{D}) \ominus \vartheta H^2(\mathbb{D})$ the corresponding model space.

In this article two classes of operators are considered: embeddings $I_\mu : K_\vartheta \rightarrow L^2(\mu)$, where μ is a finite positive measure supported on $\overline{\mathbb{D}}$, and composition operators $C_\varphi : f \mapsto f \circ \varphi$ acting from K_ϑ into $H^2(\mathbb{D})$, where $\varphi : \mathbb{D} \rightarrow \mathbb{D}$ is a holomorphic function. In fact, it is well-known that the latter type of operator may be considered as a special case of the former for a certain pullback measure μ_φ . We mention that embeddings of model spaces have been studied by a number of authors [4–7, 28]; composition operators on Hardy (and more general) spaces is by now a classical subject – we refer the reader to [24] for a description of the current state of the art and a history survey. In this article we study the composition operator acting from the model space K_ϑ into $H^2(\mathbb{D})$ thus emphasis interaction between the boundary behavior of φ and the spectrum of the inner function ϑ . In

2010 *Mathematics Subject Classification.* Primary 47B33; Secondary 30H10, 30J05, 47A45.

Key words and phrases. Embedding, composition operator, model space, Nevanlinna counting function, one-component inner function, Schatten classes.

This work was carried out at the Center for Advanced Study, Norwegian Academy of Science and Letters. Yu.L. and E.M. are partially supported by project 213638 of the Norwegian Research Council.

such setting the problem has been considered in [20]. Our main goal is to understand when such embedding and composition operators belong to the Schatten trace ideal \mathcal{S}_p , $0 < p < \infty$.

The embedding operators on K_ϑ have proved easier to analyze when ϑ is a *one-component* inner function, see [2, 4, 5, 28]. In particular, the Schatten ideal membership of I_μ has been characterized by Baranov [4] for one-component ϑ . In Section 3 we suggest a different approach to the problem. We return to the original ideas of Cohn [6] and extend embedding operators on K_ϑ to operators acting on the Hardy-Smirnov space $E^2(D)$ of a certain domain $D \supset \mathbb{D}$. This allows us to obtain a geometrical criterion for the inclusion of I_μ in \mathcal{S}_p . In particular we recover the aforementioned result in [4].

For composition operators C_φ we further refine our result to give trace ideal criteria in terms of the Nevanlinna counting function N_φ ,

$$N_\varphi(z) = \sum_{\varphi(\zeta)=z} \log \frac{1}{|\zeta|}.$$

We combine the geometric approach with recent results [16, 17] that clarify the connection between the Nevanlinna counting function N_φ and the measure μ_φ , in order to obtain the following characterization.

Theorem 4.2. *Let ϑ be a one-component inner function. The operator $C_\varphi : K_\vartheta \rightarrow H^2$ is in \mathcal{S}_p , $0 < p < \infty$, if and only if*

$$\int_{\mathbb{D}} \left(\frac{N_\varphi(z)(1 - |\vartheta(z)|^2)}{1 - |z|^2} \right)^{p/2} \left(\frac{1 - |\vartheta(z)|^2}{1 - |z|^2} \right)^2 dA < \infty.$$

The article is organized as follows. The next section contains preliminary information about one-component inner functions and the corresponding model spaces. In Section 3 we reduce the trace ideal problem of embedding operators on K_ϑ to a corresponding problem in the Hardy-Smirnov space in a larger domain, leading to a geometric characterization. In Section 4 we use these results in order to describe when $C_\varphi \in \mathcal{S}_p$ in terms of the Nevanlinna counting function of φ . We also give some geometric examples, illustrating the Schatten class behavior of composition operators on the Paley-Wiener space. General model spaces are treated in Section 5, giving an example that the one-component requirement of Theorem 4.2 can not be dropped, and providing sufficient conditions for C_φ to belong to \mathcal{S}_p , $p \geq 2$.

2. PRELIMINARIES

2.1. One-component inner functions. We recall that the class of *one-component* inner functions, introduced in [6], consists of those inner functions ϑ such that, for some $q_0 \in (0, 1)$, the set

$$D_\epsilon = \{z \in \mathbb{D} : |\vartheta(z)| < \epsilon\}$$

is connected for all $\epsilon \in (q_0, 1)$. We refer the reader to [6] and [7] for the basic facts about one-component inner functions. For the remainder of this section we fix a one-component function ϑ and a corresponding number q_0 .

Consider the canonical factorization of ϑ ,

$$\vartheta(z) = B_\Lambda(z) \exp \left(- \int_{\mathbb{T}} \frac{\xi + z}{\xi - z} d\omega(\xi) \right),$$

where Λ is the zero set of ϑ , B_Λ is the corresponding Blaschke product, and ω is a singular measure on \mathbb{T} . Functions in K_ϑ admit analytic continuation through $\mathbb{T} \setminus \Sigma(\vartheta)$, where

$$\Sigma(\vartheta) = (\mathbb{T} \cap \text{Clos}(\Lambda)) \cup \text{supp}(\omega)$$

is the *spectrum* of ϑ (see [22], Lecture 3). The function ϑ itself can be reflected over $\mathbb{T} \setminus \Sigma(\vartheta)$ and thus definition of the domain D_ϵ makes sense for any $\epsilon \in (q_0, q_0^{-1})$.

Our construction is based on the following results from [6].

Proposition 2.1. *Let $\delta \in (1, q_0^{-1})$. Then*

- *The set D_δ is simply connected, its boundary ∂D_δ is a rectifiable Jordan curve, and the linear measure on ∂D_δ is a Carleson measure with respect to $\mathbb{D}^- = \{z : |z| > 1\}$.*
- *Any function $f \in K_\vartheta$ extends analytically to a function in $E^2(D_\delta)$ and*

$$\|f\|_{K_\vartheta} \simeq \|f\|_{E^2(D_\delta)}, \quad f \in K_\vartheta.$$

Here and in what follows $E^2(D_\delta)$ and $E_-^2(D_\delta)$ denote the Hardy-Smirnov spaces in the interior and exterior of D_δ (see e.g. [10]). We mention that functions $g \in E_-^2(D_\delta)$ are required to assume the value 0 at ∞ .

Recall that a rectifiable curve Γ is called *Ahlfors regular* if for each $z \in \mathbb{C}$ and $r > 0$ we have $H^1(\Gamma \cap B(z, r)) \leq Cr$, where $H^1(\cdot)$ denotes arc length, $B(z, r) = \{\zeta : |\zeta - z| < r\}$, and $C = C(\Gamma) > 0$ is some constant.

Lemma 1. *Let $\delta \in (1, q_0^{-1})$. Then the boundary of D_δ satisfies the Ahlfors regularity condition.*

Proof. Denote $q = \delta^{-1}$. The boundary of D_δ is a rectifiable Jordan curve that is the reflection of the curve $\partial D_q = \vartheta^{-1}(|z| = q) \cup \Sigma(\vartheta)$ with respect to the unit circle, see [6].

First, if $B(z, r)$ is a ball centered at a point $z \in \Sigma(\vartheta)$, $r > 0$ sufficiently small, there exists a constant $C > 0$ such that the Carleson box $S(\tilde{z})$ centered at $\tilde{z} = (1 - Cr)z$ contains $B(z, r) \cap \partial D_q$. Since the arc length on ∂D_q is a Carleson measure, we obtain

$$H^1(B(z, r) \cap \partial D_q) \leq H^1(S(\tilde{z}) \cap \partial D_q) \lesssim 1 - |\tilde{z}| \leq Cr,$$

which is precisely the Ahlfors regularity estimate for points $z \in \Sigma(\vartheta)$.

For points in the level set $\Gamma_q = \vartheta^{-1}(|z| = q)$, we again want to show the existence of constant C , independent of z , such that $H^1(\Gamma_q \cap B(z, r)) \leq Cr$,

for all $z \in \Gamma_q$. By the same argument as in the previous paragraph, for a fixed $a > 0$ we have desired estimate for all balls with radius $r > a(1 - |z|)$. For comparatively smaller balls, we note that Theorem 1.1 and Lemma 2.1 of [6] imply that there exists $\gamma = \gamma(q)$ such that for any $z \in \Gamma_q$ the restriction $\vartheta|B(z, \gamma(1 - |z|))$ is univalent. The proof is then completed by the Hayman-Wu theorem [14], since it implies that $H^1(\vartheta^{-1}(|z| = q) \cap B(z, r)) \lesssim r$ whenever $r < \gamma(1 - |z|)$. \square

Corollary 2.2. *The space $L^2(\partial D_\delta)$ admits the direct sum decomposition*

$$L^2(\partial D_\delta) = E^2(D_\delta) \dot{+} E_-^2(D_\delta).$$

The corresponding projectors P_\pm are bounded and have the form

$$(1) \quad P_\pm f(z) = \pm \frac{1}{2} f(z) + \frac{1}{2i\pi} \int_{\partial D_\delta} \frac{f(\zeta)}{\zeta - z} d\zeta, \quad z \in \partial D_\delta.$$

The proof is straightforward; it suffices to mention that the Ahlfors regularity yields the boundedness of P_\pm in $L^2(\partial D_\delta)$ by David's theorem [9].

2.2. Model spaces. Each function $f \in K_\vartheta = H^2(\mathbb{D}) \cap \vartheta H_-^2(\mathbb{D})$ admits an extension to a function in $E^2(D_\delta)$. Denote by J the operator of analytic prolongation from \mathbb{D} to D_δ , and let $\mathcal{K}_\vartheta = E^2(D_\delta) \cap \vartheta E_-^2(D_\delta)$.

Proposition 2.3. $J(K_\vartheta) = \mathcal{K}_\vartheta$.

Proof. The inclusion $J(K_\vartheta) \subset \mathcal{K}_\vartheta$ follows from Cohn's extension construction [6] which shows that $J(K_\vartheta) \subset E^2(D_\delta)$, and the observation that if $f \in K_\vartheta$ then $\vartheta^{-1}f$ has an analytic continuation in $\mathbb{C} \setminus \mathbb{D}$.

In order to prove the opposite inclusion we first observe that the linear measure $ds = |dz|$ on \mathbb{T} is a Carleson measure for $E^2(D_\delta)$. In other words

$$\int_{\mathbb{T}} |f(z)|^2 ds \lesssim \|f\|_{E^2(D_\delta)}^2, \quad f \in E^2(D_\delta).$$

Indeed, let $f \in E^2(D_\delta)$. It is sufficient to bound $\int_{\mathbb{T}} f h ds$ for $h \in L^2(\mathbb{T})$ that are compactly supported in $\mathbb{T} \setminus \Sigma(\vartheta)$. For such h we have, by Cauchy's formula

$$\int_{\mathbb{T}} f(z) h(z) ds = \int_{\partial D_\delta} f(\zeta) \int_{\mathbb{T}} \frac{h(z)}{\zeta - z} ds(z) d\zeta.$$

The function $H(\zeta) = \int_{\mathbb{T}} \frac{h(z)}{\zeta - z} ds(z)$ belongs to $H_-^2(\mathbb{D})$. Since the arc length measure on ∂D_δ is a Carleson measure for $H_-^2(\mathbb{D})$ we have

$$\|H\|_{L^2(\partial D_\delta)} \lesssim \|H\|_{H_-^2(\mathbb{D})} \lesssim \|h\|_{L^2(\mathbb{T})}$$

and therefore

$$\left| \int_{\mathbb{T}} f(z) h(z) ds \right| \lesssim \|f\|_{E^2(\partial D_\delta)} \|h\|_{L^2(\mathbb{T})},$$

which is the required estimate.

The proposition now follows readily. Indeed, let $f \in \mathcal{K}_\vartheta = E^2(D_\delta) \cap \vartheta E_-^2(D_\delta)$, so that $f|_{\mathbb{D}} \in H^2(\mathbb{D})$ and $f = \vartheta g$, where $g \in E_-^2(D_\delta)$. Since f

and ϑ are holomorphic and $1 \leq |\vartheta(z)| \leq \delta$ in $D_\delta \setminus \mathbb{D}$, we see that g admits prolongation to a function in $H_-^2(\mathbb{D})$. Hence $f|_{\mathbb{D}} \in H^2(\mathbb{D}) \cap \vartheta H_-^2(\mathbb{D}) = K_\vartheta$. \square

3. OPERATOR EXTENSION

3.1. Embeddings of $E^2(D_\delta)$. We have seen that K_ϑ can be considered as a subspace of $E^2(D_\delta)$. The following theorem reduces trace ideal problems for embeddings of K_ϑ to trace ideal problems for embeddings of the whole space $E^2(D_\delta)$. For a positive measure μ , we denote by I_μ the embedding operator (inclusion map) from a space into $L^2(\mu)$.

Theorem 3.1. *Let ϑ be a one-component inner function, μ be a positive measure on $\overline{\mathbb{D}} \setminus \Sigma(\vartheta)$, and $0 < p < \infty$. Then the embedding $I_\mu : K_\vartheta \rightarrow L^2(\mu)$ is bounded, compact, or belongs to \mathcal{S}_p if and only if $I_\mu : E^2(D_\delta) \rightarrow L^2(\mu)$ is bounded, compact, or belongs to \mathcal{S}_p , respectively.*

Proof. We focus on showing that $I_\mu : K_\vartheta \rightarrow L^2(\mu)$ belongs to \mathcal{S}_p if and only if $I_\mu : E^2(D_\delta) \rightarrow L^2(\mu)$ does. The statements concerning boundedness and compactness have very similar, but slightly easier proofs. We may further assume that $\text{dist}(\Sigma(\vartheta), \text{supp } \mu) > 0$, as one can easily see that all estimates are uniform with respect to this quantity.

By Proposition 2.3, $I_\mu : K_\vartheta \rightarrow L^2(\mu)$ belongs to \mathcal{S}_p if and only if $I_\mu : \mathcal{K}_\vartheta \rightarrow L^2(\mu)$ does. From Corollary 2.2 follows the decomposition

$$E^2(D_\delta) = \mathcal{K}_\vartheta \dot{+} \vartheta E^2(D_\delta),$$

with bounded projectors $P_1 = M_\vartheta P_- M_{\vartheta^{-1}}$ and $Q_1 = M_\vartheta P_+ M_{\vartheta^{-1}}$ onto \mathcal{K}_ϑ and $\vartheta E^2(D_\delta)$, respectively. Here P_\pm are defined in (1), and M_h is the multiplication operator with symbol h . We identify the functions in $E^2(D_\delta)$ with their boundary values and consider by extension P_1 and Q_1 as operators on $L^2(\partial D_\delta)$. Let $C = \max(\|P_+\|, \|P_-\|)$. Since $|\vartheta| = \delta$ a.e. on ∂D_δ , we have $\|P_1\|, \|Q_1\| \leq C$ and also $\|P_+ M_{\vartheta^{-1}}\| \leq a := C\delta^{-1}$. We estimate the singular value of the second summand in the decomposition

$$(2) \quad I_\mu = I_\mu P_1 + I_\mu Q_1$$

Introducing $\tilde{\vartheta}(z) = \vartheta(z)\mathbf{1}_{|\text{supp } \mu}$, we have $I_\mu Q_1 = M_{\tilde{\vartheta}} I_\mu P_+ M_{\vartheta^{-1}}$. We obtain the following estimate of the j 'th singular value of $I_\mu Q_1$:

$$s_j(I_\mu Q_1) \leq \|M_{\tilde{\vartheta}}\| s_j(I_\mu) \|P_+ M_{\vartheta^{-1}}\| \leq a s_j(I_\mu).$$

Here we used the fact that $|\tilde{\vartheta}(z)| \leq 1$, $z \in \text{supp } \mu$.

Combining (2) with known (see e.g. [12]) inequalities for singular numbers we obtain

$$\sum_1^n s_j(I_\mu) \leq \sum_1^n s_j(I_\mu P_1) + \sum_1^n s_j(I_\mu Q_1) \leq \sum_1^n s_j(I_\mu P_1) + a \sum_1^n s_j(I_\mu).$$

If $a < 1$ and $p \geq 1$ this yields that

$$\|I_\mu : E^2(D_\delta) \rightarrow L^2(\mu)\|_{\mathcal{S}_p} \sim \|I_\mu : \mathcal{K}_\vartheta \rightarrow L^2(\mu)\|_{\mathcal{S}_p}.$$

More generally, such an equivalence of ideal norms holds for any *symmetrically normed ideal* of compact operators [12]. For $p < 1$ note that $s_{2j-1}(I_\mu) \leq s_j(I_\mu P_1) + a s_j(I_\mu)$. If $a < a_0$ is small enough, where $2^{p+1}a_0^p = 1$, we see that

$$\sum_j s_j(I_\mu P_1)^p \sim \sum_j s_j(I_\mu)^p,$$

finishing the proof also in this case.

To deal with general values of a , we note that $I_\mu : K_\vartheta \rightarrow L^2(\mu)$ is in \mathcal{S}_p if and only if $I_\mu : K_{\vartheta^2} \rightarrow L^2(\mu)$ is in \mathcal{S}_p , since $K_\vartheta \subset K_{\vartheta^2} = K_\vartheta \oplus \vartheta K_\vartheta$. Replacing ϑ by a sufficiently large power ϑ^n we will obtain a new value $a_1 = C\delta^{-n}$ such that $\|P_+ M_{\vartheta^{-n}}\| \leq a_1 < a_0 < 1$; note that in moving from the study of K_ϑ to that of K_{ϑ^n} we do not change the domain D_δ , so that the projections P_\pm stay the same for all values of n . \square

3.2. Whitney decomposition. To pass from the domain D_δ to the unit disk, let σ be a conformal mapping of \mathbb{D} onto D_δ , and ψ be its inverse. For $f \in E^2(D_\delta)$, let $h_f(w) = f(\sigma(w))(\sigma'(w))^{1/2}$, $w \in \mathbb{D}$. Let further ν denote the measure on \mathbb{D} given by

$$\nu(E) = \int_{\sigma(E)} |\psi'| d\mu, \quad E \subset \mathbb{D}.$$

Then for $f, g \in E^2(D_\delta)$ we have

$$(I_\mu^* I_\mu f, g) = \int_{\mathbb{D} \setminus \Sigma(\vartheta)} f(z) \overline{g}(z) d\mu(z) = \int_{\mathbb{D}} h_f(w) \overline{h_g}(w) d\nu(w).$$

Therefore, by Theorem 3.1, the embedding $I_\mu : K_\vartheta \rightarrow L^2(\mu)$ is in \mathcal{S}_p if and only if the embedding $I_\nu : H^2(\mathbb{D}) \rightarrow L^2(\nu)$ is in \mathcal{S}_p . We can now apply the results from Luecking [18] to conclude that $I_\mu : K_\vartheta \rightarrow L^2(\mu)$ is in the Schatten ideal \mathcal{S}_p , $0 < p < \infty$, if and only if

$$(3) \quad \sum_j \left(\frac{\nu(R_j)}{d(R_j)} \right)^{p/2} < \infty,$$

where $\{R_j\}$ is the standard dyadic decomposition of the unit disk, and $d(R_j) = \text{diam}(R_j)$.

For a domain Ω in the plane, we say that a family of Borel sets $\{G_i\}_i$ is a *Whitney-type decomposition* of Ω if $\Omega = \cup_i G_i$, the covering is of finite multiplicity, and there exist constants a, b, c such that: (i) if $z_1, z_2 \in G_i$ then $\text{dist}(z_1, \partial\Omega) \leq c \text{dist}(z_2, \partial\Omega)$, (ii) for each i there exists $z \in G_i$ such that $B(z, ad) \subset G_i \subset B(z, bd)$, where $d = \text{dist}(z, \partial\Omega)$.

We need the following simple observation:

Given two Whitney-type decompositions $\{G_i\}_i$ and $\{F_j\}_j$ of a domain Ω , let $J(i) = \{j, G_i \cap F_j \neq \emptyset\}$. Then $M := \sup_i |J(i)| < \infty$.

Indeed, just note that all F_j that intersect G_i have diameter proportional to $d_i = \text{dist}(G_i, \partial\Omega)$ and the area of each such F_j is proportional to d_i^2 .

Furthermore, if $\phi : \Omega \rightarrow \mathbb{C}$ is univalent, then $\{\phi(G_i)\}_i$ is a Whitney-type decomposition of $\phi(\Omega)$. This follows from standard estimates for univalent functions, see e.g. [23]. Together with Luecking's condition (3) this yields the following corollary.

Corollary 3.2. *Let ϑ be a one-component inner function, D_δ be a corresponding level set and μ be a positive measure on $\overline{\mathbb{D}} \setminus \Sigma(\vartheta)$. Further, let $\{G_i\}$ be any Whitney-type decomposition of D_δ . Then the embedding $K_\vartheta \rightarrow L^2(\mu)$ belongs to the Schatten ideal \mathcal{S}_p , $0 < p < \infty$, if and only if*

$$(4) \quad \sum_i \left(\frac{\mu(G_i)}{d(G_i)} \right)^{p/2} < \infty,$$

where $d(G_i) = \text{diam}(G_i)$.

Proof. First let $Q_j = \sigma(R_j)$, where $\sigma : \mathbb{D} \rightarrow D_\delta$ is a conformal mapping, as above. Then $\{Q_j\}_j$ is a Whitney-type decomposition of D_δ and since $|\sigma'| \sim \text{dist}(Q_j, \partial D_\delta) / \text{dist}(R_j, \partial \mathbb{D}) \sim \text{diam}(Q_j) / \text{diam } R_j$ (see [23]), (3) is equivalent to

$$\sum_i \left(\frac{\mu(Q_j)}{d(Q_j)} \right)^{p/2} < \infty.$$

But for any $\alpha > 0$ we have

$$\sum_i \left(\frac{\mu(G_i)}{d(G_i)} \right)^\alpha \sim \sum_{i,j} \frac{\mu(G_i \cap Q_j)^\alpha}{d(G_i)^\alpha} \sim \sum_{i,j} \frac{\mu(G_i \cap Q_j)^\alpha}{d(Q_j)^\alpha} \sim \sum_i \left(\frac{\mu(Q_j)}{d(Q_j)} \right)^\alpha,$$

proving the corollary. \square

We remark that Corollary 3.2, which will be the main ingredient in the proof of Theorem 4.2 below, can be deduced from a result of Baranov [4]. We think however that Theorem 3.1 may be of independent interest and it has some applications which do not appear to us to be immediate consequences of Corollary 3.2, see Section 4.3.

The proof of Corollary 3.2 also gives a simple and natural criterion for the Schatten class memberships of embeddings of the Hardy-Smirnov space $E^2(D)$ into $L^2(\mu)$ for measures μ in D ; one has to check the Luecking condition for an arbitrary Whitney-type decomposition of D .

4. COMPOSITION OPERATORS

4.1. Preliminaries. For a holomorphic function $\varphi : \mathbb{D} \rightarrow \mathbb{D}$, we denote by $C_\varphi : f \mapsto f \circ \varphi$ the composition operator acting on holomorphic functions f in \mathbb{D} . This operator is bounded on the Hardy space $H^2(\mathbb{D})$ (see e.g. [26]). We study the operator $C_\varphi : K_\vartheta \rightarrow H^2(\mathbb{D})$, where ϑ is an inner function in \mathbb{D} . The compactness of C_φ in terms of the Nevanlinna counting function

$$N_\varphi(z) = \sum_{\varphi(\zeta)=z} \log \frac{1}{|\zeta|}$$

was characterized in [20]; $C_\varphi : K_\vartheta \rightarrow H^2$ is compact if and only if

$$(5) \quad \limsup_{|z| \rightarrow 1} \frac{N_\varphi(z)(1 - |\vartheta(z)|^2)}{1 - |z|^2} = 0.$$

The basic tools in the argument are the Stanton formula

$$(6) \quad \|C_\varphi f\|^2 = 2 \int_{\mathbb{D}} |f'(z)|^2 N_\varphi(z) dA(z) + |f(\varphi(0))|^2,$$

where A is the normalized area measure, and also the norm inequality due to Axler, Chang and Sarason [3]

$$(7) \quad \int_{\mathbb{D}} |f'(z)|^2 \frac{1 - |z|^2}{(1 - |\vartheta(z)|^2)^b} \leq C \|f\|_2^2, \quad f \in K_\vartheta, \quad b \in (0, 1/2).$$

In this section we discuss when C_φ belongs to the Schatten ideals \mathcal{S}_p in the one-component case, aiming to capture the interaction between the symbol φ and the inner function ϑ that defines the model space. We recall the known description of the Schatten ideals for composition operators on the whole of H^2 , due to Luecking and Zhu [19]. The operator C_φ belongs to $\mathcal{S}_p(H^2)$ if and only if

$$(8) \quad \int_{\mathbb{D}} \left(\frac{N_\varphi(z)}{1 - |z|^2} \right)^{p/2} \frac{dA(z)}{(1 - |z|^2)^2} < \infty.$$

It is well understood that the composition operators can be considered as a special case of the embedding operators, see e.g. [8]. We shall now clarify this connection in our context, so that we may apply Theorem 3.1 and Corollary 3.2. We associate $\varphi : \mathbb{D} \rightarrow \mathbb{D}$ with its pullback measure μ_φ on $\overline{\mathbb{D}}$;

$$\mu_\varphi(E) = s(\varphi^{-1}(E) \cap \mathbb{T}), \quad E \subset \overline{\mathbb{D}},$$

where s denotes the Lebesgue measure on \mathbb{T} .

It is clear that $C_\varphi : K_\vartheta \rightarrow H^2$ is unitarily equivalent to the embedding operator $I_{\mu_\varphi} : K_\vartheta \rightarrow L^2(\mu_\varphi)$, and similarly that $C_\varphi : E^2(D_\delta) \rightarrow H^2$ is equivalent to $I_{\mu_\varphi} : E^2(D_\delta) \rightarrow L^2(\mu_\varphi)$. Before applying Theorem 3.1 we need to verify that $\mu_\varphi(\Sigma(\vartheta)) = 0$. This is true in view of the following lemma and the fact that $\Sigma(\vartheta)$ has zero linear Lebesgue measure when ϑ is one-component, see [2].

Lemma 2. $\mu_\varphi|_{\mathbb{T}}$ is absolutely continuous.

Proof. It is sufficient to verify that $\mu_\varphi(E) = 0$ for every closed measure zero set $E \subset \mathbb{T}$. We follow the approach of the original proof of the F. and M. Riesz theorem. Namely, there exists a continuous function $G : \overline{\mathbb{D}} \rightarrow \overline{\mathbb{D}}$, holomorphic in \mathbb{D} , such that $G(z) = 1$ for $z \in E$ and $|G(z)| < 1$ for $z \in \overline{\mathbb{D}} \setminus E$. Then $\lim_k \int_{\mathbb{D}} G^k d\mu_\varphi = \mu_\varphi(E)$. On the other hand, the sequence $G^k \circ \varphi$ converges pointwise to zero in \mathbb{D} and is uniformly bounded. Therefore,

$$\mu_\varphi(E) = \lim_k \int_{\mathbb{D}} G^k d\mu_\varphi = \lim_k \int_{\mathbb{T}} G^k \circ \varphi ds = \lim_k (G(\varphi(0)))^k = 0.$$

□

Theorem 3.1 now yields

Corollary 4.1. *Let ϑ be a one-component inner function and $\varphi : \mathbb{D} \rightarrow \mathbb{D}$ be a holomorphic function. Given any Whitney-type decomposition $\{G_j\}$ of D_δ , the operator $C_\varphi : K_\vartheta \rightarrow H^2$ belongs to \mathcal{S}_p , $0 < p < \infty$, if and only if*

$$(9) \quad \sum_i \left(\frac{\mu_\varphi(G_i)}{d(G_i)} \right)^{p/2} < \infty.$$

4.2. Nevanlinna counting function. In this subsection we implement the approach of [16, 17] in our more general setting, with the goal of showing that (9) is equivalent to

$$\int_{D_\delta} \left(\frac{N_\varphi(z)}{\text{dist}(z, \partial D_\delta)} \right)^{p/2} \frac{dA(z)}{\text{dist}(z, \partial D_\delta)^2} < \infty.$$

Theorem 4.2 then immediately follows from the relation

$$(10) \quad \text{dist}(z, \partial D_\delta) \sim \frac{1 - |z|^2}{1 - |\vartheta(z)|^2}, \quad z \in \overline{\mathbb{D}} \setminus \Sigma(\vartheta),$$

which holds for one-component inner functions, by Theorems 1.1 and 1.2 in Aleksandrov [2].

We begin by fixing a convenient Whitney-type decomposition of D_δ . Clearly we are interested only in domains that intersect $\overline{\mathbb{D}}$. We construct a decomposition of $A = \{1/2 < |z| \leq 1\} \setminus \Sigma(\vartheta)$ as follows. First we divide A into four equal parts, one for each quadrant. Each part is roughly a Carleson square. Fix some $\gamma > 0$. We say that a Carleson square S is *good* if $\text{dist}(S, \partial D_\delta) > \gamma d(S)$. If a square is good we include it in our family of sets $\{G_i\}$. Otherwise we include its upper half into the family $\{G_i\}$ and divide the lower half into two new Carleson squares. We repeat the procedure inductively, obtaining a countable family of sets $\{G_i\}$ that covers A . In particular, every point $z \in \mathbb{T} \setminus \Sigma(\vartheta)$ is included in a good square, since $\Sigma(\vartheta)$ is a closed set. We claim that $\text{dist}(G_i, \partial D_\delta) \simeq d(G_i)$ for each G_i . For if G_i is the upper half of a bad square S , it automatically satisfies $d(G_i) \lesssim \text{dist}(G_i, \partial D_\delta)$. Since S was bad we obtain the reverse inequality,

$$\text{dist}(G_i, \partial D_\delta) \lesssim d(G_i) + \text{dist}(S, \partial D_\delta) \leq d(G_i) + \gamma d(S) \lesssim d(G_i).$$

A similar argument works for the good squares G_i . Further, $\{G_i\}$ can be extended to a Whitney-type decomposition of the whole D_δ , since $\text{supp } \mu \subset \overline{\mathbb{D}}$. We omit the corresponding terms.

For each G_i we let W_i be the corresponding Carleson square (i.e. either $W_i = G_i$ or G_i is the upper half of W_i). Given a Carleson square W supported by the arc $I \subset \mathbb{T}$ and $a > 0$ we denote by aW the Carleson square supported by the arc $aI \subset \mathbb{T}$. aI has the same center as I and

$|aI| = a|I|$. According to [16] and [17] there exists $a > 1$ such that for some constant $c > 0$ the following estimates hold,

$$(11) \quad \mu(W_i)^{p/2} \leq \frac{c}{A(W_i)} \int_{\tilde{W}_i} N_\varphi^{p/2} dA,$$

$$(12) \quad \sup_{W_i} N_\varphi \leq c\mu(\tilde{W}_i),$$

where $\tilde{W}_i = aW_i$.

To obtain the integral condition, we follow the argument of [15, Proposition 3.3] and prove that (9) for the above decomposition $\{G_i\}_i$ is equivalent to

$$(13) \quad \sum_i \left(\frac{\mu_\varphi(\tilde{W}_i)}{d(W_i)} \right)^{p/2} < \infty.$$

Clearly (13) implies (9). Now we prove the converse statement.

For each j and n let $I(j) = \{i : \tilde{W}_i \cap G_j \neq \emptyset\}$ and

$$S_{n,j} = \{i \in I(j) : 2^n d(G_j) \leq d(W_i) < 2^{n+1} d(G_j)\}.$$

We note that $d(G_j) \sim \text{dist}(G_j, \partial D_\delta) \lesssim d(W_i)$ for any $i \in I(j)$, yielding that $S_{n,j} = \emptyset$ when $n < n_0$ for some fixed (negative) n_0 . We also have $\text{dist}(G_j, W_i) \lesssim d(W_i)$ which yields $C := \sup_{j,n} |S_{n,j}| < \infty$.

For $\alpha := p/2 \leq 1$ we now obtain

$$\begin{aligned} \sum_i \left(\frac{\mu_\varphi(\tilde{W}_i)}{d(W_i)} \right)^\alpha &\leq \sum_i \sum_{j:i \in I(j)} \frac{\mu_\varphi(G_j)^\alpha}{d(W_i)^\alpha} = \\ &= \sum_j \mu_\varphi(G_j)^\alpha \sum_{i \in I(j)} d(W_i)^{-\alpha} \lesssim \sum_j \left(\frac{\mu_\varphi(G_j)}{d(G_j)} \right)^\alpha, \end{aligned}$$

as desired.

For $\alpha > 1$ we also follow the argument of [15]: for each index i , consider the sets

$$s_{n,i} = \{j : i \in I(j), 2^{-n-1} d(W_i) \leq d(G_j) < 2^{-n} d(W_i)\}.$$

We have $|s_{n,i}| \leq C2^n$ for some constant C . Let β be the conjugate exponent of α and choose $\gamma \in (1 - 1/\alpha, 1)$ such that $1 - \gamma\beta < 0$. By the Hölder inequality

$$\begin{aligned} \mu_\varphi(\tilde{W}_i) &\leq \left(\sum_{j:i \in I(j)} d(G_j)^{\gamma\beta} \right)^{1/\beta} \left(\sum_{j:i \in I(j)} d(G_j)^{-\gamma\alpha} \mu_\varphi(G_j)^\alpha \right)^{1/\alpha} \lesssim \\ &\quad d(W_i)^\gamma \left(\sum_{j:i \in I(j)} d(G_j)^{-\gamma\alpha} \mu_\varphi(G_j)^\alpha \right)^{1/\alpha}, \end{aligned}$$

so that finally,

$$\sum_i \left(\frac{\mu_\varphi(\tilde{W}_i)}{d(W_i)} \right)^\alpha \leq \sum_j \mu_\varphi(G_j)^\alpha d(G_j)^{-\gamma\alpha} \sum_{i \in I(j)} d(W_i)^{\gamma\alpha-\alpha} \lesssim \sum_j \left(\frac{\mu_\varphi(G_j)}{d(G_j)} \right)^\alpha.$$

The sought after criterion in terms of the Nevanlinna counting function now follows readily.

Theorem 4.2. *Let ϑ be a one-component inner function. The operator $C_\varphi : K_\vartheta \rightarrow H^2$ is in \mathcal{S}_p , $0 < p < \infty$, if and only if*

$$\int_{\mathbb{D}} \left(\frac{N_\varphi(z)(1 - |\vartheta(z)|^2)}{1 - |z|^2} \right)^{p/2} \left(\frac{1 - |\vartheta(z)|^2}{1 - |z|^2} \right)^2 dA < \infty.$$

Proof. We follow the proof for the Hardy space given in [17, Theorem 6.1]. Inequality (12) for the Nevanlinna counting function implies

$$\begin{aligned} \int_{\mathbb{D}} \left(\frac{N_\varphi(z)}{\text{dist}(z, \partial D_\delta)} \right)^{p/2} \frac{dA(z)}{\text{dist}(z, \partial D_\delta)^2} &\lesssim \\ \sum_i d(G_i)^{-2-p/2} \int_{G_i} N_\varphi(z)^{p/2} dA(z) &\lesssim \sum_i d(G_i)^{-p/2} \mu_\varphi(\tilde{W}_i)^{p/2}. \end{aligned}$$

The converse follows from inequality (11). Specifically,

$$\begin{aligned} \sum_i d(G_i)^{-p/2} \mu_\varphi(G_i)^{p/2} &\leq \sum_i d(G_i)^{-p/2-2} \int_{\tilde{W}_i} N_\varphi(z)^{p/2} dA(z) \\ &\lesssim \sum_j \sum_{i \in I(j)} d(G_i)^{-p/2-2} \int_{G_j} N_\varphi(z)^{p/2} dA(z) \\ &\lesssim \int_{\mathbb{D}} \left(\frac{N_\varphi(z)}{\text{dist}(z, \partial D_\delta)} \right)^{p/2} \frac{dA(z)}{\text{dist}(z, \partial D_\delta)^2}, \end{aligned}$$

where the last inequality follows as in the discussion preceding the statement of the theorem. \square

4.3. Weighted composition operators on H^2 . We complete the study of composition operators on model spaces generated by one-component inner functions, by establishing the connection given by Theorem 3.1 between composition operators on K_ϑ and weighted composition operators on $H^2(\mathbb{D})$. As previously noted, $C_\varphi : K_\vartheta \rightarrow H^2$ is of \mathcal{S}_p -class if and only if $C_\varphi : E^2(D_\delta) \rightarrow H^2$ has the same property. That is, if and only if the weighted composition operator $C : H^2 \rightarrow H^2$ is of \mathcal{S}_p -class, where

$$Ch = (\psi' \circ \varphi)^{1/2} h \circ \psi \circ \varphi, \quad h \in H^2.$$

We mention [11, 21], where composition operators on Hardy-Smirnov spaces $E^2(D)$ have been studied as weighted composition operators on $H^2(\mathbb{D})$. Note that $Ch = h(M_{\psi \circ \varphi})(\psi' \circ \varphi)^{1/2}$, at least for polynomials h . Here $M_{\psi \circ \varphi}$ denotes a multiplication operator on H^2 , and we hence understand Ch as the action

of the operator $h(M_{\psi \circ \varphi})$ on $(\psi' \circ \varphi)^{1/2} \in H^2$. For $p \geq 1$, Harper and Smith [13] have utilized the theory of contractive semigroups to characterize the Schatten membership of such operators in terms of Berezin transform-type conditions. Note that in their notation, $C = \Lambda_{M_{\psi \circ \varphi}, (\psi' \circ \varphi)^{1/2}}$.

Theorem 4.3. *Denote by $G_z(w) = \frac{(1-|z|^2)^{1/2}}{1-\bar{z}w}$ the normalized reproducing kernel of H^2 at z and by $H_z(w) = \frac{(1-|z|^2)^{3/2}w}{(1-\bar{z}w)^2}$ the normalized derivative of the reproducing kernel at z . Let ϑ be a one-component inner function and let C be defined as above.*

(1) *If $1 \leq p \leq 2$, then $C_\varphi : K_\vartheta \rightarrow H^2$ belongs to \mathcal{S}_p if and only if*

$$\int_{\mathbb{D}} \|CH_z\|_{H^2}^p \frac{dA(z)}{(1-|z|^2)^2} < \infty.$$

(2) *If $2 < p < \infty$, then $C_\varphi : K_\vartheta \rightarrow H^2$ belongs to \mathcal{S}_p if and only if*

$$\int_{\mathbb{D}} \|CG_z\|_{H^2}^p \frac{dA(z)}{(1-|z|^2)^2} < \infty.$$

Remark. For $\vartheta = 0$, $p \geq 2$, we obtain the characterization of the Schatten classes in terms of the Berezin transforms found in [29]. For $p = 2$, note that

$$\begin{aligned} \int_{\mathbb{D}} \|CH_z\|_{H^2}^2 \frac{dA(z)}{(1-|z|^2)^2} &= \int_{\mathbb{D}} |\psi'(w)| |\psi(w)|^2 \int_{\mathbb{D}} \frac{1-|z|^2}{|1-\bar{z}\psi(w)|^4} dA(z) d\mu_\varphi(w) \\ &\sim \int_{\mathbb{D}} \frac{|\psi'(w)| |\psi(w)|^2}{1-|\psi(w)|^2} d\mu_\varphi(w). \end{aligned}$$

Note that $1-|\psi(w)|^2 \sim \text{dist}(w, \partial D_\delta) |\psi'(w)|$ by standard estimates [23], and that we may reexpress (10) as

$$\text{dist}(w, \partial D_\delta) \sim \frac{1-|w|^2}{1-|\vartheta(w)|^2} = \|k_w\|^{-2}, \quad w \in \overline{\mathbb{D}} \setminus \Sigma(\vartheta),$$

where k_w is the reproducing kernel of K_ϑ at w . In summary,

$$\int_{\mathbb{D}} \|CH_z\|_{H^2}^2 \frac{dA(z)}{(1-|z|^2)^2} \sim \int_{\mathbb{D}} \|k_w\|^2 d\mu_\varphi(w).$$

Theorem 4.3 could hence be viewed as a generalization of the simple fact that $C_\varphi : K_\vartheta \rightarrow H^2$ is Hilbert-Schmidt if and only if $\int_{\mathbb{D}} \|k_w\|^2 d\mu_\varphi < \infty$.

4.4. Examples on the Paley–Wiener space. In this section we consider the special case $\vartheta(z) = \exp\left(-\frac{1+z}{1-z}\right)$. The space K_ϑ can then be naturally identified with the classical Paley–Wiener space of entire functions.

For $0 < \alpha < 1$ we construct domains $U_\alpha \subset \mathbb{D}$ satisfying the following proposition. The domain U_α will be chosen so that the boundary ∂U_α intersects \mathbb{T} only at 1 and has a corner of angle $\pi\alpha$ there.

Proposition 4.4. *For each α , $0 < \alpha < 1$, there exists a domain U_α such that any corresponding Riemann map $\varphi_\alpha : \mathbb{D} \rightarrow U_\alpha$ satisfies*

- (1) $C_{\varphi_\alpha} : H^2 \rightarrow H^2$ is compact but not in any Schatten class \mathcal{S}_p ,
- (2) $C_{\varphi_\alpha} : K_\vartheta \rightarrow H^2$ is in \mathcal{S}_p for $p > 2\alpha/(\alpha - 1)$.

Note that $C_{\varphi_\alpha} : K_\vartheta \rightarrow H^2$ is in \mathcal{S}_p for one Riemann map $\varphi_\alpha : \mathbb{D} \rightarrow U_\alpha$ if and only if it is in \mathcal{S}_p for all such Riemann maps. The same statement is obviously true for $C_{\varphi_\alpha} : H^2 \rightarrow H^2$.

Let $V_\alpha \subset \mathbb{D}$ be the simply connected domain whose boundary consists of the upper half-circle $\{z : |z| = 1, \Im z \geq 0\}$ and the circular arc with terminal points -1 and 1 , intersecting the upper half-circle at an interior angle $\pi\alpha$ at those points. Note that $V_\alpha = \kappa(A_\alpha)$, where

$$A_\alpha = \{re^{i\theta} : 0 < r < \infty, 0 < \theta < \pi\alpha\},$$

and κ is the Möbius map

$$\kappa(w) = \frac{1 + iw}{1 - iw}.$$

Let $\psi_\alpha : \mathbb{D} \rightarrow V_\alpha$ be a Riemann map. The next lemma says that $C_{\psi_\alpha} : K_\vartheta \rightarrow H^2$ is in \mathcal{S}_p if and only if $p > 2\alpha/(\alpha - 1)$. This will immediately imply item (2) of Proposition 4.4, as we will choose U_α as a subset of V_α , so that $C_{\varphi_\alpha} = C_\tau C_{\psi_\alpha}$ for a composition operator $C_\tau : H^2 \rightarrow H^2$ with τ a Riemann map from \mathbb{D} to $\psi_\alpha^{-1}(U_\alpha)$.

Lemma 3. $C_{\psi_\alpha} : K_\vartheta \rightarrow H^2$ is p -Schatten if and only if $p > 2\alpha/(\alpha - 1)$.

Proof. Y. Zhu [30] obtains the corresponding result on H^2 , for domains similar to V_α , rescaled and translated to only touch (tangentially from one side) the unit circle at $z = 1$. In view of the fact that $C_{\psi_\alpha} : K_\vartheta \rightarrow H^2$ belongs to \mathcal{S}_p if and only if $C_{\psi_\alpha} : E^2(D_\delta) \rightarrow H^2$ belongs to \mathcal{S}_p and the observation that ∂D_δ in this case is a circle such that \mathbb{T} is internally tangent to ∂D_δ at 1 , the lemma follows. A direct computational proof based on our integral criterion can be also given. \square

We now define U_α . Let γ_α be the circular arc of ∂V_α which is not the upper half-circle, $\gamma_\alpha = \{z \in \partial V_\alpha : |z| \neq 1\} \cup \{-1, 1\}$. Close to 1 , we choose ∂U_α to coincide with the union of γ_α and

$$\tau = \{re^{i\theta} : 0 < \theta \leq \pi/2, r = 1 - e^{-1/\theta}\}.$$

That is, let U_α be a simply connected domain contained in V_α , such that $\partial U_\alpha \cap \mathbb{T} = \{1\}$ and for sufficiently small $\epsilon > 0$ we have

$$\partial U_\alpha \cap B_\epsilon(1) = (\gamma_\alpha \cup \tau) \cap B_\epsilon(1),$$

where $B_\epsilon(1)$ is a ball centered at 1 of radius ϵ .

Proof of Proposition 4.4. We have already proven the validity of item (2). It remains to prove that $C_{\varphi_\alpha} : H^2 \rightarrow H^2$ is compact but fails to lie in any Schatten class \mathcal{S}_p . For simplicity in presentation we will assume that $\alpha < 1/2$. The proof for $\alpha \geq 1/2$ is very similar and follows exactly the same ideas. Without loss of generality we further assume that $\varphi_\alpha(1) = 1$.

Let $\eta_\alpha = \varphi_\alpha^{-1}$. For $z \in U_\alpha$ close to 1 we have that $N_{\varphi_\alpha}(z) \sim 1 - |\eta_\alpha(z)|$. We will again make use of the inequality

$$1 - |\eta_\alpha(z)| \sim \text{dist}(z, \partial U_\alpha) |\eta'_\alpha(z)|.$$

Since η_α maps U_α , a domain with a corner of angle $\pi\alpha$ at 1, conformally onto the unit disc, it follows that $|\eta'_\alpha(z)| \sim |z - 1|^{1/\alpha - 1}$ for z close to 1 (see e.g. [23]). Therefore, for $z \in U_\alpha$ close to 1 we have

$$N_{\varphi_\alpha}(z) \sim \text{dist}(z, \partial U_\alpha) |z - 1|^{1/\alpha - 1}.$$

This estimate obviously implies that $\lim_{|z| \rightarrow 1} \frac{N_{\varphi_\alpha}(z)}{1 - |z|} = 0$ so that C_{φ_α} is compact on H^2 [26], and we will now use it to show that

$$\int_{U_\alpha \cap B_\epsilon(1)} \frac{N_{\psi_\alpha}(z)^{p/2}}{(1 - |z|^2)^{p/2+2}} dA(z) = \infty$$

for every $p > 0$, proving in view of (8) the desired statement.

We will restrict ourselves to considering points in the set

$$U'_\alpha = \{z = re^{i\theta} \in B_\epsilon(1) : r < 1 - 2e^{-1/\theta}, \theta > k(1 - r)\},$$

for a fixed large constant k . k should be chosen so large and then ϵ so small that there exists a constant $C > 1/\tan(\pi\alpha)$ such that every point $re^{i\theta} \in U'_\alpha$ satisfies $r \sin \theta > C(1 - r \cos \theta)$. In other words, U'_α should lie above the line $\Im z = C(1 - \Re z)$, which intersects the unit circle in the point 1 at an angle smaller than $\pi\alpha$. If necessary, we decrease ϵ further, attaining that $U'_\alpha \subset U_\alpha$ and

$$\text{dist}(z, \partial U_\alpha) \sim 1 - e^{-1/\theta} - r \sim 1 - |z|, \quad z \in U'_\alpha.$$

The proof is now finished by the following chain of inequalities.

$$\begin{aligned} \int_{U'_\alpha} \frac{N_{\psi_\alpha}(z)^{p/2}}{(1 - |z|^2)^{p/2+2}} dA(z) &\sim \int_{U'_\alpha} \frac{|1 - z|^{\frac{p(1-\alpha)}{2\alpha}}}{(1 - |z|^2)^2} dA(z) \\ &\gtrsim \int_{1-\epsilon'}^1 \int_{k(1-r)}^{\frac{1}{\log \frac{2}{1-r}}} \frac{(1 - 2r \cos \theta + r^2)^{\frac{p(1-\alpha)}{4\alpha}}}{(1 - r)^2} d\theta dr \\ &\gtrsim \int_{1-\epsilon'}^1 \int_{k(1-r)}^{\frac{1}{\log \frac{2}{1-r}}} \frac{\theta^{\frac{p(1-\alpha)}{2\alpha}}}{(1 - r)^2} d\theta dr \gtrsim \int_{1-\epsilon'}^1 \frac{\left(\frac{1}{\log \frac{2}{1-r}}\right)^{\frac{p(1-\alpha)}{2\alpha}+1}}{(1 - r)^2} d\theta dr = \infty, \end{aligned}$$

where $0 < \epsilon' < \epsilon$ is sufficiently small. \square

5. REMARKS ON COMPOSITION OPERATORS ON GENERAL MODEL SPACES

In this section we do not assume that the inner function ϑ is one-component.

5.1. The Hilbert-Schmidt norm. Denote the reproducing kernel of K_ϑ at w by $k(w, \cdot)$,

$$k(w, z) = \frac{1 - \overline{\vartheta(w)}\vartheta(z)}{1 - \bar{w}z}, \quad z \in \mathbb{D}.$$

Proposition 5.1. *The Hilbert-Schmidt norm of the composition operator $C_\varphi : K_\vartheta \rightarrow H^2$ is given by the expression*

$$\|C_\varphi\|_{HS}^2 = 1 - |\vartheta(0)|^2 + \frac{1}{2} \int_{\mathbb{D}} \Delta k(z, z) N_\varphi(z) dA(z).$$

Proof. Let $\{e_j(z)\}$ be any orthonormal basis in K_ϑ . Then $k(z, z) = \sum_j |e_j(z)|^2$ and the Stanton formula (6) yields

$$\|C_\varphi\|_{HS}^2 = \sum_j (C_\varphi e_j, C_\varphi e_j) = \sum_j \left(|e_j(0)|^2 + 2 \int_{\mathbb{D}} |e'_j|^2 N_\varphi dA \right).$$

Since $\sum_j |e'_j(z)|^2 = \frac{1}{4} \Delta \sum_j |e_j(z)|^2$, we obtain

$$\|C_\varphi\|_{HS}^2 = k(0, 0) + \frac{1}{2} \int_{\mathbb{D}} \Delta k(z, z) N_\varphi(z) dA(z).$$

□

Corollary 5.2. (i) *If*

$$\int_{\mathbb{D}} \frac{1 - |\vartheta(z)|^2}{(1 - |z|^2)^3} N_\varphi(z) dA(z) < \infty$$

then $C_\varphi : K_\vartheta \rightarrow H^2$ is a Hilbert-Schmidt operator;

(ii) *If $C_\varphi : K_\vartheta \rightarrow H^2$ is a Hilbert-Schmidt operator then*

$$\int_{\mathbb{D}} \frac{(1 - |\vartheta(z)|^2)^3}{(1 - |z|^2)^3} N_\varphi(z) dA(z) < \infty.$$

Proof. A direct calculation shows that

$$\frac{1}{4} \Delta k(z, z) = \frac{(1 + |z|^2)(1 - |\vartheta(z)|^2)}{(1 - |z|^2)^3} - 2 \frac{\Re(z \overline{\vartheta(z)} \vartheta'(z))}{(1 - |z|^2)^2} - \frac{|\vartheta'(z)|^2}{1 - |z|^2}.$$

Applying the standard inequality $|\vartheta'(z)| \leq (1 - |\vartheta(z)|^2)(1 - |z|^2)^{-1}$,

$$(14) \quad 4 \frac{(|z| - |\vartheta(z)|)^2 (1 - |\vartheta(z)|^2)}{(1 - |z|^2)^3} \leq \Delta k(z, z) \leq 4 \frac{(1 + |z|)^2 (1 - |\vartheta(z)|^2)}{(1 - |z|^2)^3}.$$

Statement (i) is an immediate consequence of the right hand side inequality in (14).

In order to prove statement (ii) we first observe that if $C_\varphi : K_\vartheta \rightarrow H^2$ is a Hilbert-Schmidt operator then so is $C_\varphi : K_{z\vartheta} \rightarrow H^2$, since $K_{z\vartheta} = K_z \oplus zK_\vartheta$

and $\dim K_z = 1$. Denote by $\tilde{k}(w, z)$ the reproducing kernel of $K_{z\vartheta}$ at w . We then have

$$\begin{aligned} \int_{\mathbb{D} \setminus \frac{1}{2}\mathbb{D}} \frac{(1 - |\vartheta(z)|^2)^3}{(1 - |z|^2)^3} N_\varphi(z) &\leq \\ 4 \int_{\mathbb{D}} \frac{(1 - |z\vartheta(z)|^2)(|z| - |z\vartheta(z)|)^2}{(1 - |z|^2)^3} N_\varphi(z) dA(z) &\leq \\ \int_{\mathbb{D}} \Delta \tilde{k}(z, z) N_\varphi(z) dA(z) &\leq 2 \|C_\varphi : K_{z\vartheta} \rightarrow H^2\|_{HS}^2. \end{aligned}$$

□

Remark. In the previous section we proved that the inequality

$$(15) \quad \int_{\mathbb{D}} \frac{(1 - |\vartheta(z)|^2)^3}{(1 - |z|^2)^3} N_\varphi(z) dA(z) < \infty$$

gives a complete description of the Hilbert-Schmidt composition operators $C_\varphi : K_\vartheta \rightarrow H^2$ in the case of one-component ϑ . This condition is not sufficient in general, see Section 5.3 below.

5.2. A sufficient condition for larger Schatten ideals. The Hilbert-Schmidt norm characterization, discussed for composition operators so far, remains true for the differentiation operator acting between K_ϑ and $L^2(\mathbb{D}, \mu)$, where μ is a positive, finite measure on \mathbb{D} . Using complex interpolation we can obtain a sufficient condition for a composition operator to be in the Schatten ideal \mathcal{S}_p when $2 \leq p < \infty$.

First we assume that $\Phi(z)$ is a positive function on \mathbb{D} such that the differentiation $D : K_\vartheta \rightarrow L^2(\Phi(z) dA)$ is a bounded operator. It follows from (7) that we can take

$$\Phi(z) = \frac{1 - |z|^2}{(1 - |\vartheta(z)|^2)^b},$$

with $b \in (0, 1/2)$. For the special case when ϑ is a one-component inner function we may choose

$$\Phi(z) = \frac{1 - |z|^2}{1 - |\vartheta(z)|^2},$$

see [7, Theorem 1].

Proposition 5.3. *Let $\Phi(z)$ be as above and $p \geq 2$. If*

$$\int_{\mathbb{D}} \left(\frac{N_\varphi(z)}{\Phi(z)} \right)^{p/2} \Delta k(z, z) \Phi(z) dA(z) < \infty$$

then $C_\varphi : K_\vartheta \rightarrow H^2$ belongs to \mathcal{S}_p .

Proof. The proof follows an idea of Luecking [18]; see also [4]. By the Stanton formula (6), we want to prove that the differentiation operator $D : K_\vartheta \rightarrow L^2(N_\varphi dA)$ belongs to the Schatten ideal \mathcal{S}_p . It is enough to prove the corresponding statement where N_φ has been replaced by $N = \chi_r N_\varphi$, χ_r

being the characteristic function of the disk $r\mathbb{D}$, as long as the corresponding Schatten norm is independent of r .

Consider the following holomorphic family of compact operators

$$(T_\zeta f)(z) = N(z)^{p(1-\zeta)/4} \Phi(z)^{1/2-p(1-\zeta)/4} f'(z),$$

where $\zeta \in \mathbb{C}$, $0 \leq \Re \zeta \leq 1$. We note that $T_{1+iy} : K_\vartheta \rightarrow L^2(dA)$ is bounded by a constant C_1 that depends on Φ only. Further, $T_{iy} : K_\vartheta \rightarrow L^2(dA)$ is a Hilbert-Schmidt operator with norm (see the proof of Proposition 5.1)

$$\|T_{iy}\|_{HS} \leq 1 + \int_{\mathbb{D}} \Delta K(z, z) N(z)^{p/2} \Phi(z)^{1-p/2} dA(z) \leq C_2$$

by the assumption of the theorem. Then $T_{1-2/p} \in \mathcal{S}_p$ and $\|T_{1-2/p}\|_{\mathcal{S}_p} \leq C_1^{2/p} C_2^{1-2/p}$ by Theorem 13.1 in [12]. But $T_{1-2/p} f(z) = N(z)^{1/2} f'(z)$. \square

Note that for $\vartheta = 0$, Proposition 5.3 gives us the known condition of Luecking and Zhu [19].

5.3. Integral condition insufficient when ϑ is not one-component.

Let

$$\alpha_n = 2^{-n}, \quad r_n = 1 - \frac{1}{n} 2^{-2n}, \quad z_n = r_n e^{i\alpha_n}, \quad n = 1, 2, \dots,$$

and let ϑ be the Blaschke product

$$\vartheta(z) = \prod_{n=1}^{\infty} \frac{-\bar{z}_n}{|z_n|} \frac{z - z_n}{1 - \bar{z}_n z}, \quad z \in \mathbb{D}.$$

The sequence $\{z_n\}$ is interpolating (for H^∞), since it is hyperbolically separated in \mathbb{D} and $\sum_n (1 - |z_n|) \delta_{z_n}$ is a Carleson measure for H^2 . Therefore the sequence of normalized reproducing kernels

$$\{(1 - |z_n|)^{1/2} k_{z_n}\}_n, \quad k_{z_n} = \frac{1}{1 - \bar{z}_n z},$$

is a Riesz basis for K_ϑ . In particular any Hilbert-Schmidt operator $C : K_\vartheta \rightarrow H^2$ satisfies

$$(16) \quad \sum_n (1 - |z_n|) \langle C k_{z_n}, C k_{z_n} \rangle < \infty.$$

Let $\Delta = \{z : |z - \frac{1}{2}| < \frac{1}{2}\}$ and let $\varphi(z) = (1+z)/2$, a conformal mapping of \mathbb{D} onto Δ . We will show that (16) with $C = C_\varphi$ fails, while

$$(17) \quad \int_{\mathbb{D}} \left(\frac{1 - |\vartheta(z)|^2}{1 - |z|^2} \right)^3 N_\varphi(z) dA(z) < \infty.$$

Indeed, since $N_\varphi(z) = -\log |2z - 1| \gtrsim \text{dist}(z, \Delta)$ for $z \in \Delta$, we have

$$\begin{aligned} (1 - |z_n|) \langle C_\varphi k_{z_n}, C_\varphi k_{z_n} \rangle_{H^2} &\gtrsim (1 - |z_n|) \int_{\Delta} \left| z - \frac{1}{\bar{z}_n} \right|^{-4} N_\varphi(z) dA(z) \\ &\gtrsim \frac{1 - |z_n|}{\text{dist}(z_n, \Delta)} \sim \frac{1}{n}. \end{aligned}$$

This yields divergence of the series on the right hand side of (16). Therefore C_φ is not a Hilbert-Schmidt operator.

To prove (17) we need an auxiliary statement.

Lemma 4. *For any $a > 0$, the sum*

$$\sum_{n=1}^{\infty} \left(\frac{1 - |z_n|}{|1 - \bar{z}_n z|} \right)^a$$

is uniformly convergent in $\overline{\mathbb{D}}$. In particular, it is uniformly bounded.

Proof. For $z \in \overline{\mathbb{D}}$ and $0 < t < 1$, let $M(t, z) = \{n \geq 1 : \frac{1 - |z_n|}{|1 - \bar{z}_n z|} > t\}$ and let $N(t, z) = |M(t, z)|$ be the number of points in $M(t, z)$. Pick n and k such that $n > k \geq N(t, z) - 1$ and $z_n, z_k \in M(t, z)$. Then

$$2^{-k}|z| \lesssim |\overline{(z_n - z_k)}z| \lesssim \frac{1 - |z_k|}{t} \leq \frac{2^{-2k}}{t},$$

from which $|z|t \lesssim 2^{-k} \lesssim 2^{-N(t, z)}$. That is, there exists a constant $C > 0$ such that

$$N(t, z) \leq \log \frac{C}{|z|t}.$$

In combination with the identity

$$\sum_{n=1}^{\infty} \left(\frac{1 - |z_n|}{|1 - \bar{z}_n z|} \right)^\alpha = \alpha \int_0^1 t^{\alpha-1} N(t, z) dt,$$

the statement of the lemma clearly follows. \square

In order to prove (17) we first observe that for $0 < \epsilon < 1$

$$\begin{aligned} \frac{1 - |\vartheta(z)|^2}{1 - |z|^2} &\leq \sum_n \frac{1 - |z_n|^2}{|1 - \bar{z}_n z|^2} \\ &\leq \left(\sum_n \left(\frac{1 - |z_n|^2}{|1 - \bar{z}_n z|} \right)^{1-\epsilon} \right)^{2/3} \left(\sum_n \frac{(1 - |z_n|^2)^{1+2\epsilon}}{|1 - \bar{z}_n z|^{4+2\epsilon}} \right)^{1/3}. \end{aligned}$$

Let $\Delta' = \{z : |z - 1/2| < 1/4\}$. Using Lemma 4 and the fact that $N_\varphi(z) \sim \text{dist}(z, \Delta)$ for $z \in \Delta \setminus \Delta'$, we obtain

$$\begin{aligned} \int_{\mathbb{D} \setminus \Delta'} \left(\frac{1 - |\vartheta(z)|^2}{1 - |z|^2} \right)^3 N_\varphi(z) dA(z) &\lesssim \\ \sum_n (1 - |z_n|^2)^{1+2\epsilon} \int_{\Delta} \left| z - \frac{1}{\bar{z}_n} \right|^{-4-2\epsilon} \text{dist}(z, \Delta) dA(z) &\sim \\ \sum_n \left(\frac{1 - |z_n|}{\text{dist}(z_n, \Delta)} \right)^{1+2\epsilon} &\sim \sum_n \frac{1}{n^{1+2\epsilon}} < \infty, \end{aligned}$$

which of course implies (17).

REFERENCES

- [1] A. B. Aleksandrov, A simple proof of the Vol'berg-Treil' theorem on the embedding of covariant subspaces of the shift operator, *Zap. Nauch. Sem. POMI*, **217** (1994), 26–35.
- [2] A. B. Aleksandrov, On embedding theorems for coinvariant subspaces of the shift operator, II, *Zap. Nauch. sem. POMI*, **262** (1999), 5–48.
- [3] S. Axler, S. Y. Chang, D. Sarason, Products of Toeplitz operators, *Integral Equations Operator Theory*, **1** (1978), 285–309.
- [4] A. D. Baranov, Embeddings of model subspaces of the Hardy space: compactness and Schatten - von Neumann ideals, *Izv. Math.*, **73** (2009), no. 6, 1077–1100.
- [5] J. A. Cima, A. L. Matheson, On Carleson embeddings of star-invariant subspaces, *Quaest. Math.*, **26** (2003), no. 3, 279–288.
- [6] W. S. Cohn, Carleson measures for functions orthogonal to invariant subspaces, *Pacific J. Math.* **103** (1982), no. 2, 347–364.
- [7] W. S. Cohn, Carleson measures and operators on star-invariant subspaces, *J. Operator Theory* **15** (1986), 181–202.
- [8] C. C. Cowen, B. MacCluer, Composition operators on spaces of analytic functions, *Studies in Adv. math.*, CRC Press, Boca Raton, FL, 1995.
- [9] G. David, Opérateurs intégraux singuliers sur certaines courbes du plan complexe, *Ann. Sci. cole Norm. Sup.* (4) **17** (1984), no. 1, 157–189.
- [10] P. Duren, Theory of H^p spaces, Pure and applied Math., vol. 38, Academic Press, NY, 1970.
- [11] E.A. Gallardo-Gutiérrez, M. González, A. Nicolau, Composition operators on Hardy spaces on Lavrentiev domains, *Trans. AMS* **360** (2008), no. 1, 395–410.
- [12] I. C. Gohberg, M. G. Krein, Introduction to the Theory of Non-selfadjoint Linear Operators, Nauka, Moscow, 1965; English translation, Transl. Math. monographs, 18, AMS, Providence, RI, 1969.
- [13] Z. Harper, M. Smith, Testing Schatten class Hankel operators, Carleson embeddings and weighted composition operators on reproducing kernels, *J. Operator Theory* **55** (2006), no. 2, 349–371.
- [14] W. K. Hayman, J. M. G. Wu, Level sets of univalent functions, *Comment. Math. Helv.* **56** (1981), no. 3, 366–403.
- [15] P. Lefèvre, D. Li, H. Queffélec, L. Rodriguez-Piazza, Some examples of compact composition operators on H^2 , *J. Funct. Anal.*, **255** (2008), 3098–3124.
- [16] P. Lefèvre, D. Li, H. Queffélec, L. Rodriguez-Piazza, Nevanlinna counting function and Carleson function of analytic maps, *Math. Ann.*, **351** (2011), 305–326.
- [17] P. Lefèvre, D. Li, H. Queffélec, L. Rodriguez-Piazza, Some revisited results about composition operators on Hardy spaces, *Rev. Math. Iberoam.*, **28** (2012), 57–76.
- [18] D. Luecking, Trace ideal criteria for Toeplitz operators, *J. Funct. Anal.* **73** (1987), 345–368.
- [19] D. Luecking, K. Zhu, Composition operators belonging to Schatten Ideals, *American J. of Math.*, **114** (1992), no. 5, 1127–1145.
- [20] Yu. Lyubarskii, E. Malinnikova, Composition operators on model spaces, *Recent trends in Analysis: Proceedings of the conference in honor of Nikolai Nikolski*, Bordeaux, 2011. Theta foundation, Bucharest, 149–157, 2013.
- [21] V. Matache, Weighted composition operators on H^2 and applications, *Complex Anal. Oper. Theory* **2** (2008), no. 1, 169–197.
- [22] N. K. Nikol'skii, *Treatise on the shift operator. Spectral function theory*. Grundlehren der Mathematischen Wissenschaften, 273. Springer-Verlag, Berlin, 1986.
- [23] Ch. Pommerenke, Boundary behaviour of conformal mappings, *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*, 299. Springer-Verlag, Berlin, 1992.

- [24] E. Saksman, An elementary introduction to Clark measures, *Topics in Complex Analysis and Operator Theory*, Proceedings of the Winter School held in Antequera, Malaga, Spain (February 5–9, 2006), 85–136.
- [25] B. Simon, Trace ideals and their applications, *London Mathematical Society Lecture Note Series*, 35. Cambridge University Press, Cambridge-New York, 1979.
- [26] J. H. Shapiro, The essential norm of the composition operator, *Ann. of Math.*, 125 (1987), no. 2, 375–404.
- [27] J. H. Shapiro, Composition operators and classical function theory, *Universitext: Tracts in Mathematics*, Springer-Verlag, NY, 1993.
- [28] A. L. Vol'berg, S. R. Treil', Embedding theorems for invariant subspaces of the inverse shift operator, *Zap. Nauchn. Sem. LOMI*, **149** (1986), 38–51.
- [29] K. H. Zhu, Positive Toeplitz operators on weighted Bergman spaces of bounded symmetric domains, *J. Operator Theory* **20** (1988), no. 2, 329–357.
- [30] Y. Zhu, Geometric properties of composition operators belonging to Schatten classes, *Int. J. Math. Math. Sci.* **26** (2001), 239–248.