

# *Lake surface temperature [in “State of the Climate in 2016”]*

Article

Accepted Version

Woolway, R. I. ORCID: <https://orcid.org/0000-0003-0498-7968>,  
Carrea, L. ORCID: <https://orcid.org/0000-0002-3280-2767>,  
Merchant, C. J. ORCID: <https://orcid.org/0000-0003-4687-9850>, Dokulil, M., de Eyto, E., DeGasperi, C., Korhonen, J.,  
Marszelewski, W., May, L., Paterson, A., Rimmer, A., Rusak, J.,  
Schladow, G., Schmid, M., Shimaraeva, S., Silow, E., Timofeev, M.,  
Verburg, P., Watanabe, S. and Weyhenmeyer, G. (2017) Lake surface temperature [in “State of the Climate in 2016”]. *Bulletin of the American Meteorological Society*, 98 (8). pp. 13-14. ISSN 1520-0477 doi: <https://doi.org/10.1175/2017BAMSStateoftheClimate.1>  
Available at <https://centaur.reading.ac.uk/72001/>

It is advisable to refer to the publisher’s version if you intend to cite from the work. See [Guidance on citing](#).

To link to this article DOI:

<http://dx.doi.org/10.1175/2017BAMSStateoftheClimate.1>

Publisher: American Meteorological Society

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other

copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

[www.reading.ac.uk/centaur](http://www.reading.ac.uk/centaur)

## **CentAUR**

Central Archive at the University of Reading

Reading's research outputs online

1 **2.b.2 Lake surface temperature**—*R. I. Woolway, L. Carrea, C. J. Merchant, M. T. Dokulil,*  
2 *E. de Eyto, C. L. DeGasperi, J. Korhonen, W. Marszelewski, L. May, A. M. Paterson, A.*  
3 *Rimmer, J. A. Rusak, S. G. Schladow, M. Schmid, S. V. Shimaraeva, E. Silow, M. A.*  
4 *Timofeev, P. Verburg, S. Watanabe, and G. A. Weyhenmeyer*

5 Observed lake surface water temperature (LSWT) thermal anomalies in 2016 are placed in  
6 the context of the recent warming observed in global surface air temperature (section 2b1) by  
7 collating long-term in situ LSWT observations from some of the world’s best-studied lakes  
8 and a satellite-derived global LSWT dataset. The period 1996–2015, 20 years for which  
9 satellite-derived LSWTs are available, is used as the base period for all LSWT anomaly  
10 calculations. Warm-season averages (July–September in the Northern Hemisphere and  
11 January–March in the Southern Hemisphere) are analyzed to avoid ice cover, in line with  
12 previous LSWT analyses (Schneider and Hook 2010; Hook et al. 2012; O’Reilly et al. 2015;  
13 Torbick et al. 2016; Woolway et al. 2016).

14

15 In situ observations from 48 lakes show an average warm-season LSWT anomaly of 1.0°C in  
16 2016 (Fig. 2.b.2.1). The LSWT anomaly in Lake Baikal (Russia), the largest (by volume) and  
17 deepest of the world’s freshwater lakes, was more than 2.3°C warmer in 2016. Comparable  
18 anomalies were observed in the North American Great Lakes, with an average anomaly of  
19 +2°C in 2016. Warming is not restricted to the largest lakes. For example, Harp Lake in  
20 Dorset, Ontario (Canada; surface area ~1 km<sup>2</sup>) was 1.1°C warmer in 2016, compared to its  
21 20-year average. High LSWT anomalies were also observed in central Europe, with LSWT  
22 anomalies >+0.5°C, and in Scandinavia, with the second largest lake in Sweden, Lake  
23 Vättern, having a LSWT anomaly of +1.3°C. Higher-than-average LSWTs were also evident  
24 in the Southern Hemisphere, with Lakes Rotorua and Taupo (New Zealand) showing an  
25 average LSWT anomaly exceeding +1°C, and the smaller lakes in the Bay of Plenty region  
26 (New Zealand) experiencing an average anomaly of +1°C in 2016.

27

28 Satellite-derived warm-season LSWTs generated within the Globolakes project  
29 ([www.globolakes.ac.uk/](http://www.globolakes.ac.uk/)) for 681 lakes are used in this analysis to investigate global  
30 variations in LSWT. LSWTs were retrieved during the day using the retrieval methods of  
31 MacCallum and Merchant (2012) on image pixels filled with water according to both the  
32 inland water dataset of Carrea et al. (2015) and a reflectance-based water detection scheme  
33 (Xu 2006). The satellite temperatures represent mid-morning observations throughout the

34 record. The observations were generated using data from the ATSR (Along Track Scanning  
35 Radiometer) series including ATSR-2 (1995–2003) and the Advanced ATSR (AATSR)  
36 (2002–12), extended with MetOp-A AVHRR (Advanced Very High Resolution Radiometer)  
37 (2007–16).

38

39 Globally and regionally averaged warming rates calculated from the satellite data show  
40 widespread warming tendencies in recent years (Figs 2.b.2.2), being most evident in the  
41 extratropical Northern Hemisphere ( $>30^{\circ}\text{N}$ ), with a hemispheric average LSWT trend of  
42  $+0.31^{\circ}\text{C decade}^{-1}$  ( $p = 0.06$ ). Warming ( $+0.21^{\circ}\text{C decade}^{-1}$ ,  $p = 0.07$ ) is also found for the  
43 Southern Hemisphere ( $<30^{\circ}\text{S}$ ), but not in the tropics ( $30^{\circ}\text{N}–30^{\circ}\text{S}$ ;  $p = 0.4$ ). Using all  
44 available data, and weighting equally the northern, southern, and tropical regions, we obtain a  
45 global LSWT trend of  $+0.24^{\circ}\text{C decade}^{-1}$  ( $p = 0.01$ ). Europe is the region showing the largest  
46 and most consistent LSWT warming trend (Fig. 2.b.2.2b), inline with previous studies (Hook  
47 et al. 2012), with a regional average LSWT trend of  $+0.55^{\circ}\text{C decade}^{-1}$ . Other regions such as  
48 northeastern North America ( $+0.43^{\circ}\text{C decade}^{-1}$ ) and southern South America (notably those  
49 in southern Chile and Argentina;  $+0.3^{\circ}\text{C decade}^{-1}$ ) also experience significant regionally  
50 averaged warming.

51

52 In the year 2016, lakes were particularly warm with a global and equally weighted LSWT  
53 anomaly of  $+0.65^{\circ}\text{C}$ . LSWT anomalies in the Northern Hemisphere ( $+0.72^{\circ}\text{C}$ ), Southern  
54 Hemisphere ( $+0.70^{\circ}\text{C}$ ), and the tropics ( $+0.52^{\circ}\text{C}$ ) were all anomalously high (Figs 2.b.2.2a;  
55 Plate 2.1) in 2016. About 83% of satellite-observed LSWT anomalies in 2016 were warmer  
56 than their 20-year average.

57

58 Global in situ and satellite measurements both point to LSWTs in 2016 being anomalously  
59 high, the warmest year in the 21-year record, reflecting the observed warming in global  
60 surface air temperature. Rising LSWTs have major implications for lake ecosystems  
61 (O'Reilly et al. 2003; Smol et al. 2005; Smol and Douglas 2007) and can, among other  
62 things, increase the occurrence of toxic cyanobacterial blooms (Kosten et al. 2012) and  
63 subsequently threaten water quality (Huisman et al. 2005). Warming of LSWT has been  
64 observed since 1996 and was particularly striking in 2016. If this trend continues, local  
65 economies dependent on lakes for drinking water, agricultural irrigation, recreation, and  
66 tourism are likely to be increasingly affected.

67

68 **References:**

- 69 Carrea, L., O. Embury, and C. J. Merchant, 2015: Datasets related to in-land water for  
70 limnology and remote sensing applications: Distance-to-land, distance-to-water,  
71 water-body identifier and lake-centre co-ordinates. *Geosci. Data J.*, **2** (2), 83–97,  
72 doi:10.1002/gdj3.32.
- 73 Hook, S. R., R. C. Wilson, S. MacCallum, and C. J. Merchant, 2012: Lake surface  
74 temperature [in "State of the Climate in 2011"]. *Bull. Amer. Meteor. Soc.*, **93** (7), S18–  
75 S19, doi:10.1175/2012BAMSSStateoftheClimate.1.
- 76 Huisman, J., H. C. P. Matthijs, and P. M. Visser, Eds., 2005: *Harmful Cyanobacteria*.  
77 Springer, 243 pp.
- 78 Kosten, S., and Coauthors, 2012: Warmer climates boost cyanobacterial dominance in  
79 shallow lakes. *Global Change Biol.*, **18**, 118–126, doi:10.1111/j.1365-  
80 2486.2011.02488.x.
- 81 MacCallum, S. N., and C. J. Merchant, 2012: Surface water temperature observations of large  
82 lakes by optimal estimation. *Canadian J. Remote Sens.*, **38**, 25–45, doi:10.5589/m12-  
83 010.
- 84 O'Reilly, C. M., and Coauthors, 2015: Rapid and highly variable warming of lake surface  
85 waters around the globe. *Geophys. Res. Lett.*, **42**, 10,773–710,781,  
86 doi:10.1002/2015GL066235.
- 87 O'Reilly, C. M., S. R. Alin, P.-D. Plisnier, A. S. Cohen, and B. A. McKee, 2003: Climate  
88 change decreases aquatic ecosystem productivity of Lake Tanganyika, Africa. *Nature*,  
89 **424**, 766–768, doi:10.1038/nature01833.
- 90 Schneider, P., and S. J. Hook, 2010: Space observations of inland water bodies show rapid  
91 surface warming since 1985. *Geophys. Res. Lett.*, **37**, L22405,  
92 doi:10.1029/2010GL045059.

93 Smol, J. P., and Coauthors, 2005: Climate-driven regime shifts in the biological communities  
94 of Arctic lakes. *Proc. Natl. Acad. Sci. USA*, **102**, 4397–4402,  
95 doi:10.1073/pnas.0500245102.


96 Smol, J. P., and M. S. V. Douglas, 2007: Crossing the final ecological threshold in high  
97 Arctic ponds. *Proc. Natl. Acad. Sci. USA*, **104**, 12395–12397,  
98 doi:10.1073/pnas.0702777104.

99 Torbick, N., B. Ziniti, S. Wu, and E. Linder, 2016: Spatiotemporal lake skin summer  
100 temperature trends in the northeast United States. *Earth Interact.*, **20** (25), 1–21,  
101 doi:10.1175/ei-d-16-0015.1.


102 Woolway, R. I., and Coauthors, 2016: Lake surface temperatures [in "State of the Climate in  
103 2015"]. *Bull. Amer. Meteor. Soc.*, **97** (8), S17–S18,  
104 doi:10.1175/2016BAMSSStateoftheClimate.1.

105 Xu, H., 2006: Modification of normalised difference water index (NDWI) to enhance open  
106 water features in remotely sensed imagery. *Int. J. Remote Sens.*, **27**, 3025–3033,  
107 doi:10.1080/01431160600589179.

108



112 **Figure 2.b.2.1.** In situ LSWT observations from 48 globally distributed lakes, showing the annually  
113 averaged warm season (Jul–Sep in NH; Jan–Mar in SH) anomalies (°C; relative to 1996–2015).




114

115 **Figure 2.b.2.2.** Satellite-derived LSWT measurements from 681 lakes showing (a) global and  
 116 regional annual average anomalies (°C), and (b) 1996–2016 LSWT trend (°C decade<sup>-1</sup>). Annual  
 117 LSWTs are calculated for the warm season (Jul–Sep in NH; Jan–Mar in SH) and LSWT trends are  
 118 calculated on these anomalies.

119





120

121 **Plate 2.1.** Satellite-derived LSWT anomalies in 2016. Annual LSWTs are calculated for the warm  
122 season (Jul–Sep in NH; Jan–Mar in SH).

123