

Spatial dependency and technical efficiency: an application of a Bayesian stochastic frontier model to irrigated and rainfed rice farmers in Bohol, Philippines

Article

Accepted Version

Pede, V. O., Areal, F. J., Singbo, A., McKinley, J. and Kajisa, K. (2018) Spatial dependency and technical efficiency: an application of a Bayesian stochastic frontier model to irrigated and rainfed rice farmers in Bohol, Philippines. *Agricultural Economics*, 49 (3). pp. 301-312. ISSN 0169-5150 doi: 10.1111/agec.12417 Available at <https://centaur.reading.ac.uk/73582/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

To link to this article DOI: <http://dx.doi.org/10.1111/agec.12417>

Publisher: Wiley

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online

Spatial Dependency and Technical Efficiency: An Application of a Bayesian Stochastic Frontier Model to Irrigated and Rainfed Rice Farmers in Bohol, Philippines

Valerien O. Pede¹, Francisco Areal², Alphonse Singbo³, Justin McKinley^{1,4}, and Kei Kajisa^{1,5}

¹International Rice Research Institute (IRRI), Los Banos, Philippines

²University of Reading, UK

³International Crops Research Institute for the Semi-Arid Tropics (ICRISAT), Bamako, Mali

⁴Monash University, Australia

⁵Aoyama Gakuin University, Tokyo, Japan

Corresponding author:

Valerien O. Pede, PhD

Senior Agricultural Economist

Agri-food Policy Platform

International Rice Research Institute

DAPO Box 7777

Metro Manila, Philippines

Ph: +63 2 580 5600 (ext. 2721)

Email: v.pede@irri.org

Abstract

We investigated the role of spatial dependency in the technical efficiency estimates of rice farmers using panel data from the Central Visayan island of Bohol in the Philippines. Household-level data were collected from irrigated and rainfed agroecosystems. In each ecosystem, the geographical information on residential and farm-plot neighborhood structures was recorded to compare household-level spatial dependency among four types of neighborhoods. A Bayesian stochastic frontier approach that integrates spatial dependency was used to address the effects of neighborhood structures on farmers' performance. Incorporating the spatial dimension into the neighborhood structures allowed for identification of the relationships between spatial dependency and technical efficiency through comparison with non-spatial models. The neighborhood structure at the residence and plot levels were defined with a spatial weight matrix where cut-off distances ranged from 100m to 1,000m. We found that spatial dependency exists at the residential and plot levels and is stronger for irrigated farms than rainfed farms. We also found that technical inefficiency levels decrease as spatial effects are more taken into account. Because the spatial effects increase with a shorter network distance, the decreasing technical inefficiency implies that the unobserved inefficiencies can be explained better by considering small networks of relatively close farmers over large networks of distant farmers.

Keywords: Rice farming, Spatial dependency, Bayesian approach, Efficiency

JEL codes:

1. Introduction

Numerous attempts have been made to measure technical efficiency (TE) and other efficiency estimates in farming (Idiong, 2007; Michler and Shively, 2014; Karagiannis and Tzouvelekas, 2009; Alvarez, 2004; Balde et al., 2014; Hossain and Rahman, 2012; Coelli and Battese, 1996); to this end, one of the common econometric approaches is the stochastic frontier analysis (Aigner et al., 1977; Meeusen and van Den Broeck, 1977). Previous studies have contributed to the understanding of how large TE is; how different TE levels are among individual farmers; and what are the factors that underlie the differences. These studies generated useful policy implications for efficient farming, especially in developing countries where wide productivity variations have been observed. Despite the aforementioned research, spatial dependency among farmers has yet to be adequately analyzed. Farrell (1957) expressed concerns about spatial factors such as how climate and location influence efficiency. Although the concerns existed at the time, the econometric techniques required to complete such an analysis were not available during the time of Farrell's research. The importance of making use of spatial information in agricultural economics, and in particular the little attention paid to spatial autocorrelation in land use data has still been highlighted in more recent times (Bockstael, 1996).

Recent developments in spatial econometrics have made it possible to observe the spatial effects in the stochastic frontier analysis (Areal et al., 2012; Glass et al., 2014; Glass et al., 2015; Tsionas and Michaelides, 2015). Furthermore, Druska and Horrace (2004) extended the estimator presented by Kelejian and Prucha (1999) and applied it to a stochastic frontier model for the panel data of 171 Indonesian rice farmers. Another innovation in this area was the adoption of the Bayesian paradigm in the estimation procedure (Schmidt et al., 2008). With this approach, Koop and Steel (2001) and Kumbhakar and Tsionas (2005) investigated geographical variations of outputs and

farm productivity for 370 municipalities in Brazil. Similarly, Areal et al. (2012) also investigated the spatial dependence of 215 dairy farms in England at a 10 km grid-square level using the Bayesian paradigm. All these studies have used a meso-level data to measure the spatial distribution of farmers.

Although these meso-level studies are valuable in recognizing the importance of spatial dependency in agriculture, important questions surrounding this topic remain unanswered. To illustrate, one unanswered question is how and through what kinds of networks the spatial dependency of TE shows up at the farm level. There are several reasons why spatial dependency may differ by type of farm-level networks. First, despite the presence of agricultural extension agents, it is still expected that farmers rely on their social networks for information on input allocation, management practices, etc. (Ward and Pede 2014). Additionally, the intensity of information sharing may differ between residential and plot neighbors. Second, farmers who belong to the same agro-ecological system and share a common resource pool depend on the regulations set within the resource pool for their farming management. For farmers in irrigated areas who share water within a water users group, this kind of dependency may emerge more strongly among them as compared to the rainfed farmers who farm more independently. Lastly, farmers who belong to the same water users group in irrigated areas may face similar institutional shocks and regulations. Notably, they may have formed similar social preferences through collective irrigation management (Tsusaka et al. 2015). In this manner, farm-level spatial dependency may arise through socio-economic, agro-climatic, or institutional similarities.

The purpose of this paper is to investigate the role of spatial dependency in TE, using unique micro-level farm panel data from individual rice farmers in Bohol, Philippines. We aim to identify the types of networks in which spatial dependency arises in TE. Our empirical study is unique in two

ways. First, we use farm-specific micro-level data, whereas previous studies were aggregated to sub-municipalities (Koop and Steel 2001; Kumbhakar and Tsionas 2005). In this way, micro-level studies are highly relevant in investigating spatial effects for smallholder farmers that occur on a small scale in meters. To the best of our knowledge, this has not been investigated in previous efficiency work. Furthermore, to the extent that previous studies have used spatial information at aggregated levels, this was not possible to be investigated. At aggregated levels, this type of small-scale spatial dependence may be unfeasible to detect, i.e., the area studied is larger than the area at which spatial dependence occurs, and consequently may be overlooked. This oversight could lead to either accepting that spatial dependency in smaller areas may be present, but cannot be observed, or that there are no small-scale spatial processes present. The advantage of focusing on farm-specific micro-level data in small areas for TE analysis is that some interesting spatial characteristics can be incorporated that may include those from nature, e.g., climatic condition, but also techno-sociological, e.g., cooperation, information sharing, and resource sharing. Since the area of study is relatively small we are confident that those aspects are controlled for and we are left with variation on the techno-sociological aspects. Analysis at micro level also presents the opportunity of better targeting agricultural development policies towards local conditions. The second way in which this study is unique is that we compare spatial dependency among two separate neighborhood structures (residential neighborhood and farm plot neighborhood) in two different agro-ecosystems (irrigated and rainfed ecosystems), and explore the panel nature of our dataset which covers four growing seasons (two years).

The data were collected for four consecutive rice growing seasons from 2009 to 2011, coupled with detailed geographical information to capture different kinds of networks among sample farmers. This data set allowed us to compare spatial dependency among two separate neighborhood structures

(residential neighborhood and farm plot neighborhood) in two different agro-ecosystem (irrigated and rainfed ecosystems). Taking advantage of the panel data structure, analyses were performed following a one-step procedure as described in Areal et al. (2012), which integrates spatial dependency into the stochastic frontier analysis with a Bayesian estimation approach. The rest of the article is organized as follows. The next section provides some background information about the major characteristics of the two rice farming systems in Bohol. Section 3 presents the empirical model used to estimate the technical efficiency and the endogenous spatial effect of rice farming technical efficiency. Section 4 describes the data-set used in this study. Section 5 presents the estimation results and discussions. Section 6 concludes and derives policy implications for rice farming productivity in Philippines.

Rice Farming in Bohol

Rice production in Bohol consists of two agroecosystems: irrigated and rainfed farming. The Bohol Irrigation System (BIS) started its operation in 2009, currently spans 14 villages in three municipalities, and is expected to service as many as 4,104 hectares in the future (JICA, 2012). The BIS works through a gravity irrigation system composed of a reservoir dam, a main canal, secondary canals and laterals, turnouts, and farm ditches. Most of the farmers in the project site converted their rainfed plots to irrigated plots as long as their plots were accessible to the irrigation facilities. Our sample famers were randomly taken from these irrigated famers. The rainfed sample farmers were randomly taken from adjacent villages that have similar cultural and climatic background (Figure 1). Rainfed rice farming is conducted in a traditonal manner with moderate use of modern inputs and little use of machineries. The same scenario applied to the irrigated area until the start of irrigation in 2009.




Fig. 1. Location of study sites designated by ecosystem.

Farmers in the irrigated area must form a water users group. A group consisting of 20 individual farmers on average and its members rely on the same intake gate on a canal and thus share irrigation water with each other. Since the location and the water supply capacity of each intake gate is determined by the capacity of the canal and the topography of the area, the size and composition of the water users group is basically determined exogenously. In addition, our field observation tells us that no farmer exchanged their plots in order to move to a particular water user group. This means that there is no self-selection behavior in the formation of the water user group.

Member farmers are expected to pay an irrigation service fee equivalent to 150 kg of paddy per hectare per season to the National Irrigation Administration (NIA)¹. The members of the water users group are expected to manage local irrigation facilities collectively. Since they share irrigation water, the synchronization of farming practices is needed among them. Meanwhile, rice farming under rainfed conditions is conducted more independently. In this regard, the opportunities for networking are more frequent, and the demand for strict coordination is higher among irrigated farmers than rainfed farmers.

In the study site, rice is the dominant crop and is cultivated twice a year. The Bohol Island belongs to a climatic area characterized by even rainfall distribution throughout the year. During our survey period of four agricultural seasons in two years (2009–2011), our study site experienced two weather shocks: severe drought in the second season and flood in the fourth season. Furthermore, rainfed areas suffered directly from these variations. Meanwhile, the water supply condition among irrigated farmers was mitigated by the irrigation system, to some extent. Hence, the irrigated farmers suffered fewer water shortages in the second season than the rainfed farmers. Since the BIS has no drainage system, all the farmers suffered flood in the fourth season.

A notable feature of the study site, which is important in network analyses, is that the places of residence are relatively scattered over a wide geographical area; although we can still find the center of a village where residences and small businesses are concentrated. Hence, the data presented in this study has wide geographical variations in residential networks, which is different from another type of common residential pattern in which residents are highly concentrated in a particular place.

¹ With a market price of Php 14-20 per kg, the 150 kg of paddy is equivalent to about Php 2,500-3,000. As of February 2017 1USD=49.7 Php.

2. Modeling

Since the seminal works of Aigner et al. (1977) and Meeusen and van Den Broeck (1977), the stochastic frontier approach (SFA) has become the most commonly used method of modeling the production and measure efficiency of farm-level data. The SFA approach estimates the parametric form of a production function and recognizes the presence of two random error terms in the data. One component of the error term reflects the inefficiency in production while the other component represents the random effects outside of the producer's control. The production frontier itself is stochastic since it varies randomly across farms due to the presence of the random error component. Following the model proposed by Areal et al. (2012), the stochastic frontier production function for a balanced panel data assuming efficiency is constant over time² is defined as³:

$$y_{it} = x_{it}\beta + z_{it}\theta + p_{it}\psi + v_{it} - u_i \quad (1)$$

² This is not an uncommon assumption to make especially when the time series is relatively short as in this case (2 years).

³ A specification including time and its interactions was estimated but no significant time effects were found. A referee has noted that the model does not include heteroscedasticity terms. This is an area that has not been explored within this context. We have run the non-spatial model and extracted the errors and the inefficiency terms. However, there is not a prior reason to believe that allowing for heteroscedasticity would make it necessarily a better model. We have conducted a Levene's test on the errors to test whether the variance changes through the periods. We found that the variance for year 2 is actually different (p-value<0.05), but the standard deviations for the years are not very different in absolute values: Year 1: 0.246; year 2: 0.306; year 3: 0.230; year 4: 0.239

where y_{it} denotes the production of farm i ($i = 1, 2, \dots, N$) at season t ($t = 1, \dots, T$) with $T = 4$; x_{it} represents a $(N \times T) \times k$ matrix of inputs of production; z_{it} is a $N \times m$ matrix of non-stochastic environmental variables (farmer's level of education, household size, household head, being a female, remittance), associated with the i th farm at the t th observation (farm-specific variables); p_{it} is a $N \times (T - 1)$ matrix of dummy variables for periods 2 to 4; β , θ and ψ are a $k \times 1$ and a $m \times 1$ vectors of unknown parameters to be estimated; v_{it} is the random error, and u_i represents the inefficiency of the i th farm. Stacking all variables into matrices we obtain:

$$y = x\beta + z\theta + p\psi + v - (u \otimes 1_T) \quad (2)$$

where the inefficiency term in the standard efficiency analysis usually assume u to follow an exponential of half-Normal distribution. However, u can be made spatially dependent by defining it as:

$$u = \rho W u + \tilde{u} \quad (3)$$

where W is a weight matrix; ρ is the spatial coefficient, which is assumed to be between 0 and 1; and u and \tilde{u} are latent variables whose distributional form is unknown. In the context of farming,

ρWu captures the effects of shocks spreading among neighboring farmers through similarity in socio-economic, agro-ecological, and institutional backgrounds of the group defined by W .

Estimation of spatial models requires specification for the spatial structure of observation units considered in the study. As such, a distance-based weight matrix, W , of a Boolean type with elements w_{ij} was defined as follows⁴:

$$w_{ij} = \exp\left(\frac{-d_{ij}^2}{s^2}\right) \quad (4)$$

where d_{ij} is the distance in km between the residence/farm location i and the residence/farm location j ; s is the distance from the residence/farm where spatial dependence may be relevant, i.e., the cut-off point of spatial dependence. Finding the appropriate cut-off distance is an empirical issue (Roe et al., 2002) that is commonly dealt with by estimating the spatial model using different cutoff distances (Bell and Bockstael, 2000; Roe et al., 2002; Kim et al., 2003; Areal et al., 2012; Areal and Riesgo, 2014).

Therefore, we use two types of Bayesian models, one standard SFA (non-spatial) and spatial models (with cut-off distance ranging from 100m to 1,000m), which allow for an investigation of the relationship between spatial dependency and efficiency under different farm environments. Results from the non-spatial model and the spatial model with the highest spatial dependency are compared as follows. Once the farm efficiency estimates from both models are obtained, the efficiency percentage change between the spatial and non-spatial model is calculated per household and farm

⁴ The weight matrix W is of dimension $N \times N$ and has 0 as diagonal elements.

environment (residential or farm plot). This allows us to explore how much accounting for spatial dependency can help in explaining efficiency. If the area used to determine the neighborhood is relatively large, we may find spatial dependence; however this may not help in explaining efficiency. The same would occur if certain spatial effects that were accounted for are not relevant in explaining inefficiency, i.e., the spatial models in this case would be underperforming compared with the non-spatial models. Having found spatial dependency, a farm with a positive percentage change in their efficiency level would indicate that such farm's efficiency level would have been underestimated under the non-spatial approach (i.e. positive aspects such as sharing information that make farms more efficient were not taken into account). We would expect this to be the case of farms that work closely and share knowledge under similar environment. On the other hand, we may expect farms that work more independently to show lower levels of spatial dependence; and no or small percentage changes in cases where the spatial model matches the performance of the non-spatial model or even negative percentage change in cases where the non-spatial model outperforms the spatial models.

A translog functional form was chosen for the stochastic frontier production analysis. To explain, the translog is a flexible functional form that can be viewed as a second-order Taylor expansion in logarithms of any function of unknown form. Unlike the Cobb-Douglas function, it imposes no restriction a priori on the elasticities of substitution between inputs and outputs. As mentioned above, some non-stochastic environmental variables were incorporated directly into the non-stochastic component of the production frontier accounting for changes in the production level.⁵

⁵ There are two general approaches to incorporate non-stochastic environmental variables into technical efficiency analysis (Coelli et al., 2005). The first, the one used here, is to incorporate them in the non-stochastic part of the frontier model whereas the second approach incorporates them into the stochastic component of the production frontier

Thus, the variable, *Education*, was included and consists of the years of formal schooling of the primary decision-maker of the household; the variable *Size*, which is the total number of people living in the household; *Gender*, a binary variable taking a value of one when the household head is female; and *Remittance* consisting of the ratio of remittance as it relates to total household income. The first three variables capture the human capital endowment of the sample farmer: education for quality, size for amount, and the gender for advantage or disadvantage of female head. Educated farmers are generally assumed to have better farming capacity and access to information; therefore, they are more productive (Battese and Coelli, 1995). The amount of remittance indicates that farmers have alternative income sources other than rice farming. Hence, we hypothesize that remittance, which captures an unimportance of rice farming, has a negative effect on production levels.

The data for all inputs and outputs are normalized by their respective geometric means prior to estimation. This makes the model's parameter estimates directly interpretable as elasticities that are evaluated at the geometric mean of the data. To cope with the great number of zero observations for fertilizer inputs, the procedure proposed by Battese (1997) was followed. The original variable for fertilizer was replaced with $x_{it}^k = \max(x_{it}^k, D_{it}^k)$, where D_{it}^k is a dummy variable defined by

(Kumbhakar, Ghosh and McGukin, 1991). We decided to use the first approach to distinguish between observed information, which is included in the production side and non-observed information (spatial aspects) into the stochastic part of the frontier. However, following the suggestion of a reviewer we also conducted the second approach in which *Education*, *Size*, *Gender* and *Remittance* are removed from the non-stochastic part and are used in a second stage as explanatory variables for the estimated efficiency. The coefficient estimates of the non-stochastic part are similar. Regarding the explanatory variables for efficiency, *Education* was found to be associated with higher levels of efficiency in all cases whereas *Remittance* was found to be associated with lower levels of efficiency (i.e. farmers recipients of relatively larger amounts of remittances were found to be less efficient than those receiving lower amounts of remittances). Results are available in Appendix.

$D_{it}^k = 1$ if $x_{it}^k = 0$ and $D_{it}^k = 0$ if $x_{it}^k > 0$. Thus, the final estimable form of the translog stochastic production function becomes:

$$\begin{aligned} \ln y_{it} = & \alpha_0 + \sum_k \alpha_k \ln(x_{it}^k) + \frac{1}{2} \sum_k \sum_j \alpha_{kj} \ln(x_{it}^k) \ln(x_{it}^j) + \beta_k D_{it}^k + \theta_1 Education_{it} \\ & + \theta_2 HHsize_{it} + \theta_3 Gender_{it} + \theta_4 Remittance_{it} + \sum_{l=2}^4 \psi_l p_l + V_{it} - U_i \end{aligned} \quad (5)$$

Where y is the output, t is a time index ($t = 1, \dots, T$), k and j are the inputs, and $\alpha_0, \alpha_k, \alpha_{kj}, \theta_1, \theta_2, \theta_3, \theta_4, \psi_l, \beta_k$ are the parameters to be estimated. The symmetry property was imposed by restricting $\alpha_{kj} = \alpha_{jk}$. The U_i are farm specific inefficiency terms as defined above. The estimation was conducted using a Bayesian approach that integrates the latent distributions of u and \tilde{u} into the estimation process as defined in Equation 3 (Areal et al. 2012). Thus, a standard form for the conditional likelihood function was assumed in efficiency analysis with a spatial component added to:

$$p(y|\beta, h, \rho, \mu_u^{-1}, \tilde{u}) = \prod_{i=1}^N \frac{h^{\frac{T}{2}}}{(2\pi)^{\frac{T}{2}}} \exp\left(-h \frac{\varepsilon' \varepsilon}{2}\right) \quad (6)$$

By re-parameterizing $\tilde{y} = [y + (I - \rho W)^{-1} \tilde{u} \otimes 1_T]$, $x = x + z + p$ the expression for the conditional likelihood function was obtained:

$$p(y|\beta, h, \rho, \mu_u^{-1}, \tilde{u}) \propto h^{\frac{TN}{2}} \exp\left(-\frac{h}{2} (\tilde{y} - x\beta)' (\tilde{y} - x\beta)\right) \quad (7)$$

The prior distribution for the parameters $\beta, h, \mu_z^{-1}, \tilde{u}, \rho$ are an independent Normal-Gamma prior for β and h ; the prior for μ_u^{-1} is assumed to be Gamma with parameters 2 and $-\ln(r^*)$, where r^* is the median of the prior distribution, and the conditional distribution for \tilde{u} is:

$$p(\tilde{u}_i | \alpha, \mu_u^{-1}) = \frac{\tilde{u}_i^{\alpha-1}}{\mu_u^j \Gamma(\alpha)} \exp(-\mu_u^{-1} \tilde{u}_i) \quad (8)$$

where $\Gamma(\alpha)$ is the Gamma function with parameter $\alpha = 1$, which is commonly used in the literature.

The prior for ρ is assumed to have a positive impact on the efficiency and is defined as an indicator function $I(\cdot) = 1$ if $\rho \in [0,1]$, or otherwise $I(\cdot) = 0$.

The following conditional posteriors are obtained from the joint posterior distribution, $p(\beta, h, \rho, \mu_u^{-1}, u | y)$: the conditional posterior for β and h are a Normal distribution and Gamma distribution as in Koop (2003). The conditional posterior distribution for μ_u^{-1} is $p(\mu_u^{-1} | \beta, h, \rho, \tilde{u}, y) \sim G(m, \eta)$ where $m = \frac{N+1}{\sum_{i=1}^N \tilde{u}_i - \ln(r^*)}$ and $\eta = 2N + 2$. Furthermore, the conditional posterior distribution for \tilde{u}_i is

$$p(\tilde{u}_i | \beta, h, \rho, \mu_u^{-1}, y) \propto \exp \left[-\frac{hT}{2} \left[z_i - \left(\bar{x}_i \beta - \bar{y}_i + \frac{\mu_u^{-1}}{Th} \right) + (\tilde{u}_i - u_i) \mu_u^{-1} \right] \right] \quad (9)$$

where $\bar{x}_i = \sum_{t=1}^T \frac{x_{it}}{T}$ and $\bar{y}_i = \sum_{t=1}^T \frac{y_{it}}{T}$ and the conditional posterior for the spatial dependence parameter ρ is $p(\rho|\beta, h, \mu_u^{-1}, \tilde{u}, y) \propto \exp\left(-h \frac{\varepsilon' \varepsilon}{2}\right) \times I(\rho \in (0,1))$. Finally, the conditional posterior distributions for \tilde{u}_i and ρ each requires a posterior Metropolis-Hastings algorithm step (Metropolis et al. 1953; Hastings 1970)⁶.

3. Data

The data for this study was collected by the International Rice Research Institute (IRRI) from 2009 to 2011 to conduct an impact assessment of the Bohol Irrigation Development Project in the Philippines. There were 496 observations per season from two different ecosystems; 205 and 291 observations from rainfed and irrigated ecosystems, respectively. Therefore, the panel used for the stochastic frontier analysis has a size of 820 and 1,164 for rainfed and irrigated, respectively. Data on household characteristics, inputs, and output for rice farming were collected with a structured questionnaire. Additionally, the data set also contains geographical coordinates at both the farm plot and farmer residences.

Descriptive statistics for the variables used in the efficiency analysis are available in Table 1. Capital is defined as the sum of the current values of agricultural machineries such as tractors,

⁶ We use a random walk chain Metropolis-Hastings algorithm, which takes draws proportionately in different regions of the posterior making sure that the chain moves in the appropriate direction (Koop, 2003), where a new set of \tilde{u}_i is proposed using a Metropolis based on $(\tilde{u}_i|\beta, h, \rho, \mu_u^{-1}, y) \propto \exp\left[-\frac{hT}{2}\left[z_i - \left(\bar{x}_i\beta - \bar{y}_i + \frac{\mu_u^{-1}}{Tn}\right) + (\tilde{u}_i - u_i)\mu_u^{-1}\right]\right]$. On the other hand, to draw ρ the Metropolis is based on $p(\rho|\beta, h, \mu_u^{-1}, \tilde{u}, y) \propto \exp\left(-h \frac{\varepsilon' \varepsilon}{2}\right) \times I(\rho \in (0,1))$.

sprayers, and other farming devices. Since the level of mechanization in the area is low, the capital value is not very large in either area. Notably, it is apparent from Table 1 that farmers in the irrigated areas perform more intensified rice farming (high inputs and high output), particularly with regard to the level of fertilizer and labor. For comparison's sake, we report the yield at the bottom of Table 1, which supports the notion of higher productivity in the irrigated areas. Additionally, education attainment is nearly the same for farmers from the two ecosystems. Moreover, socio-economic characteristics such as household size, female-led households, and remittances as a percent of total income were not found to be significantly different between rainfed and irrigated systems.

Table 1: Summary statistics of production inputs and socio-economic characteristics.

	Rainfed (n=820)	Irrigated (n=1164)	Difference
Output (kg)	724.430 (636.956)	1364.897 (1107.032)	640.467***
Seed (kg)	30.996 (23.123)	37.773 (26.179)	6.777***
Fertilizer (kg)	27.674 (35.886)	43.023 (34.791)	15.349***
Labor (Mandays)	32.868 (18.363)	43.685 (25.570)	10.817***
Plot Size (Ha)	0.573 (0.409)	0.619 (0.412)	0.045***
Capital (PHP)	1028.623 (860.518)	1151.318 (923.425)	122.695***
Education (Yrs.)	6.080 (3.468)	5.728 (3.011)	0.352
Household size	5.606 (2.322)	5.601 (2.582)	0.005
Female Household Head (%)	7.44%	4.90%	2.54%
Remittance [†] (%)	7.35%	4.69%	2.66%
Yield (Ton/ha)	1.436 (0.949)	2.352 (1.172)	0.916***

Note: '***', '**', and '*' mean the difference is statistically significant at the 1, 5, and 10 percent levels, respectively

[†]: Calculated as remittance as a portion of total income

4. Results

We estimated both spatial and non-spatial models for each ecosystem (rainfed and irrigated) by considering residential and plot neighborhood structures. We also estimated the spatial models by considering various definitions of the weight matrix based on 10 cut-off distances from 100m to 1,000m by 100m.⁷ We found no significant differences in the estimated coefficients of the non-spatial models in comparison to the spatial counterparts. Coefficient estimates associated with production inputs were consistent with what we would expect, which was that inputs have a positive relationship with outputs.

Table 2 shows the summary results obtained for the spatial dependence parameter ρ with cut-off distance (100m to 1,000m). The spatial dependence parameter rapidly decreases as the cut-off distance increases, reaching its highest average value at a 100m cut-off distance. This is an expected finding that means non-observables explain efficiency at distances equal or below 100m. Moreover, this finding is also in accordance with Tobler's First Law of Geography which says that near things tend to be more related than distant things. Another interesting result is that spatial dependence was stronger for irrigated farms than for rainfed farms (Table 2). Thus, for the 100m

⁷ We find no significant differences among 10 different cut-off distance models in the coefficients associated to the production inputs and the environmental factors. Estimated parameters for spatial (all distance cut-off) and non-spatial models are available in Online appendix Tables A1 and A2.

Table 2: Spatial dependence at different cut-off distances.

Distance (metres)	Spatial parameter rho							
	Residence				Plot			
	Irrigated		Rainfed		Irrigated		Rainfed	
100	0.218	(0.014,0.565)	0.153	(0.010,0.375)	0.195	(0.015,0.478)	0.086	(0.006,0.209)
200	0.095	(0.006,0.237)	0.067	(0.004,0.160)	0.076	(0.004,0.185)	0.055	(0.003,0.130)
300	0.057	(0.003,0.134)	0.041	(0.002,0.099)	0.045	(0.002,0.107)	0.041	(0.002,0.097)
400	0.042	(0.003,0.096)	0.026	(0.001,0.065)	0.032	(0.001,0.076)	0.029	(0.006,0.209)
500	0.033	(0.002,0.071)	0.019	(0.009,0.381)	0.026	(0.001,0.060)	0.023	(0.001,0.057)
600	0.028	(0.002,0.058)	0.014	(0.001,0.036)	0.022	(0.015,0.048)	0.018	(0.001,0.045)
700	0.022	(0.002,0.047)	0.012	(0.001,0.032)	0.018	(0.001,0.040)	0.015	(0.001,0.037)
800	0.020	(0.002,0.039)	0.010	(0.001,0.027)	0.016	(0.001,0.035)	0.012	(0.001,0.031)
900	0.016	(0.001,0.033)	0.009	(0.001,0.022)	0.014	(0.001,0.030)	0.011	(4E-4,0.027)
1,000	0.015	(0.002,0.028)	0.008	(5E-4,0.019)	0.012	(0.001,0.026)	0.009	(4E-4,0.022)

Note: The interpretation of the Bayesian 95% coverage posterior (a,b) is that according to our data and model the parameter is between a and b with a 0.95 probability

model, the probability that the spatial dependence parameter is greater in irrigated farms than rainfed farms is 63% and 78%, respectively, for the residence and plot neighborhoods⁸. For irrigated farms, the probability that the spatial effect is greater under the residential neighborhood structure than under the plot neighborhood structure is 54%, whereas for the rainfed farms this probability is 71%.

⁸ These were obtained comparing the conditional posterior distributions obtained for ρ for rainfed farms for residence and plot neighborhoods after 25,000 draws from the conditional distributions with 5,000 draws discarded and 20,000 retained. The comparison was done between each of the 20,000 values of the conditional posterior distributions for ρ . When the spatial dependence of on type of farm (1) is greater than another (2) a value of 1 is given, and 0 otherwise.

Lastly, as the spatial dependence increases with shorter distances, the mean efficiency also increases, suggesting that the more unobservable aspects (e.g. cooperation, information sharing) are explained with the spatial models the more “inefficiency” from the non-spatial models is controlled for. Thus, the estimated mean efficiency for the irrigated farms models with residence spatial dependence at 100m, 400m, 700m, and 1,000m are 0.91, 0.90, 0.88, and 0.87, respectively. Additionally, the estimated mean efficiency for the rainfed farms models with residence spatial dependence at 100m, 400m, 700m, and 1,000m are 0.88, 0.87, 0.86, and 0.86, respectively. As for the plot spatial dependence, the results are consistent with what we found in the residence spatial models. The estimated mean efficiency for the irrigated farms models with plot spatial dependence at 100m, 400m, 700m, and 1,000m are 0.91, 0.90, 0.89, and 0.88, respectively. The estimated mean efficiency for the rainfed farms models with residence spatial dependence at 100m, 400m, 700m, and 1,000m are 0.88, 0.87, 0.86, 0.86, respectively. However, although spatial dependency increases with shorter distances, this does not mean that spatial models always explain efficiency better than a non-spatial model. The use of non-spatial models may be able to explain efficiency as well as or even better than spatial models when cut-off distances are relatively large. Notably, the average efficiency levels of non-spatial models for irrigated and rainfed farms is 0.89 and 0.85 respectively for both residence and plot coordinates models, which suggests that for irrigated farms, only spatial models with cut-off distances at 100m and 400m explain efficiency better than the non-spatial model. For the case of rainfed farms all spatial models outperform the non-spatial model. This result suggests that spatial effects at relatively small distances (< 400 m) (e.g. sharing information, use of common resources) are important determinants for irrigated rice production. For datasets that cover relatively large areas, accounting for spatial dependency in this way helps control for some of the unobserved heterogeneity in the sample, e.g., climatic and topographical conditions. However, the

source and processes behind the spatial dependence cannot be explained due to the variety of heterogeneous possible reasons. In this study, the fact that the sample is relatively homogeneous works as an advantage in explaining such spatial dependence. Since the observed spatial dependence exists at such small cut-off distances (100m), it cannot be a result of any climatic condition.

Figures 2a shows the distribution of efficiency for irrigated and rainfed farms using non-spatial model and spatial models (at 100m, 400m, 700m and 1,000m) in the case of residential neighborhood. Interestingly, in all four scenarios, the distribution is skewed towards the right and has a relatively long left tale. Very few farmers have efficiency levels less than 0.5. The distribution of efficiency varies not only by ecosystem, but also by type of neighborhood. In every case, the distribution of the non-spatial model is very distinct from the spatial models. This finding exposes the biases in efficiency levels that arise when spatial considerations are ignored, i.e., cases where the efficiency distribution from spatial models is located to the right of the non-spatial efficiency distribution. More specifically, for the case using residential neighborhood, models for irrigated farms where spatial effects were found to be relatively high (100m and 400m) have a narrow distribution to the right of the non-spatial efficiency distribution. This suggests that part of the farm inefficiency not captured under the non-spatial model can be explained by these spatial models. Also, the fact that the distribution shape is narrower indicates that differences between farm efficiency levels have been reduced once spatial effects have been taken into account.

Hence, although we find that the shorter the distance, the greater the spatial dependence in both cases of irrigated and rainfed, we can see in Figure 2a (bottom) that for rainfed farms the effect of such increase in spatial dependence with distance is relatively small in explaining efficiency (efficiency distributions are closer to each other than for Figure 2a (top)). Additionally, considering the non-spatial efficiency distribution as a reference, we found that the efficiency distribution using

the spatial model (100m, 400m) for irrigated farms is situated to the right of the non-spatial case (Figure 2a). This means that for irrigated farms, spatial dependence may help explaining inefficiency, i.e., irrigated farmers working more closely, but for rainfed farms, which work more independently, spatial dependence does not contribute as much in explaining efficiency as the case for irrigated farms.

This information is not surprising since spatial effects related to rainfed farmer's residence should be relatively less important in explaining efficiency, as compared to irrigated farms. For instance, for irrigated farms, information sharing about technology among residential neighbors may determine production levels. This may not be the case for rainfed farms whose practices may be determined more independently, and the level of production may be more dependent on the plot's location, i.e., specific agronomic conditions rather than sharing knowledge. The same argument can be used when comparing spatial models that use large neighborhood areas and their contributions to explaining inefficiency, e.g., comparing efficiency distributions using the spatial dependence model (cut-off distance 1,000m) vs. non-spatial dependence model (cut-off distance 100m) for irrigated farms. We found small spatial dependence in our longer distance spatial model.




Figure 2a: Distribution of efficiency in irrigated farms (top) and rainfed farms (bottom) using residence coordinates.

Figures 2b shows the distribution of efficiency for irrigated (top) and rainfed (bottom) farms, using a non-spatial model and spatial models (at 100m, 400m, 700m and 1,000m) in the case of plot neighborhood structure. For the models on irrigated farms, the same findings were produced as in the case of residence neighborhood, suggesting that both natural conditions of the spatial area and communication between farmers with neighbor plots plays a role in explaining part of the

inefficiency detected by the non-spatial models. As in the residence coordinates case, the efficiency distribution for rainfed farms when spatial effects are taken into account are different from the efficiency distribution obtained by the non-spatial model.

This aforementioned finding suggests that natural conditions are likely playing a role in explaining the estimated inefficiency levels gathered by the non-spatial model.




Figure 2b: Distribution of efficiency in irrigated farms (top) and rainfed farms (bottom) using plot coordinates.

Spatial dependence can explain why the level of connectedness, i.e., working together and sharing information, is important in explaining efficiency levels. We found that different neighborhood (residential and plot) explain different spatial processes, and there are two types of processes that are captured by residential neighborhood and plot neighborhood. While social conditions are captured for farmers' residence location, both social and environmental conditions are captured by the geographical location of the plot. Using residential neighborhood, Figure 3a shows the map of percentage change in farm efficiency levels for both irrigated and rainfed farms in this study area. Rainfed farms tend to have greater increases in efficiency levels once the spatial dependency is incorporated into the analysis. Irrigated farms have relatively less increase in efficiency levels once spatial dependency is incorporated.

The finding above needs some clarification because we found stronger spatial dependency in irrigated area. Irrigated farms being more spatially dependent means that the efficiency levels of neighboring irrigated farms are more similar between them than the efficiency levels of neighboring rainfed farms. This is possibly due to conditions and practices under irrigation being more similar between neighboring irrigated farms than the conditions and practices under rainfed between neighboring rainfed farms. The fact that we can capture this with the spatial models helps us identify better farm efficiency levels (i.e. avoiding attributing unobservable environmental conditions to inefficiency). The nature of the spatial dependency is what determines its effect of spatial dependency on the farm efficiency estimation. Thus, for irrigated farms the nature of spatial dependency may come from similar environmental conditions and practices (e.g. through sharing information) whereas for rainfed farms it may come from more variable conditions (e.g. climatic and topographical conditions)". Being able to capture unobservable variable conditions was found to be relatively more important in explaining efficiency levels for rainfed farms than capturing

unobservable environmental conditions and practices for explaining farm efficiency levels for irrigated farms (i.e. accounting for more variable conditions such as weather conditions are more determinant than more “controlled” conditions in explaining efficiency levels).

Figure 3b shows the map of percentage change in farm efficiency levels for both irrigated and rainfed farms in the study area using plot neighborhood. In this case, we found similar results as in the residential neighborhood. We found a higher efficiency increase on the rainfed area than on the irrigated area. Again, we expected this result since natural conditions are expected to be more important in explaining efficiency for rainfed farms than for irrigated farms. Still, we find increase in efficiency levels for irrigated farms. This finding may be a result of the plot neighborhood, or it may be a result of partially capturing the social aspect.

The percentage average increase in efficiency is, on average, higher for rainfed farms (3.4% and 3.3% for residential and plot neighborhood) than for irrigated farms (2.9% and 2.6% for residential and plot neighborhood), in light of the average efficiency levels mentioned above for the spatial model using the 100m cut-off distance for irrigated and rainfed farm (0.91 and 0.88), and the efficiency levels obtained from the equivalent non-spatial models (0.89 and 0.85). Hence, we found that although the spatial dependence parameter (ρ) tells us the strength of the spatial dependence, which is generally greater for irrigated than for rainfed farms, such strength, i.e., incorporating spatial dependency into the analysis, follows a non-linear relationship with how well the spatial model performs compared with the non-spatial model in terms of percentage change in efficiency between spatial and non-spatial models. To explain, using the plot neighborhood structure, the spatial dependence parameter at 100m for irrigated and rainfed farms is 0.195 and 0.086, respectively, and the percent efficiency increase is 2.6% and 3.3% for irrigated and rainfed farms, respectively.




Figure 3a: Percentage change in efficiency score between spatial model (100m) and non spatial (residence coordinates).




Figure 3b: Percentage change in efficiency score between spatial model (100m) and non spatial (plot coordinates).

The estimated models also show noteworthy results. *Education* has a positive and significant effect in irrigated as well as rainfed environments. Even though the rainfed farmers are more educated by about 0.4 years than irrigated farmers, there was no significant difference in means between the two ecosystems (see Table 1). In the irrigated area, the rice farming is more modernized in the sense that farmers use new and improved varieties and chemical inputs, as well as following standardized agronomic practices under controlled irrigation. Formal education for literacy as well as basic scientific knowledge is important to understand these types of practices. The fact that education significantly contributed to improve output in the rainfed environment also makes sense because even though farming is less intensified in the rainfed environment; formal education is still useful to the rainfed farmers. In fact, the results in Appendix A1 & A2 show the largest educational impact in the rainfed farmers' plot neighbor model (0.013). Additionally, *Household size* was found to be insignificant in all estimated models. This finding was expected because more farmers reach out to hired labor for farm operations. Finally, regarding the dummy variables for the studied periods we found that results corroborate the expected effects where rice production in periods 2 and 4 levels was lower than the first period (i.e. the benchmark period) due to the severe drought in the second season and flood in the fourth season mentioned above.

Conclusions and Policy Implications

This paper investigates the role of spatial dependency in technical efficiency for different ecosystems and neighborhood structures focusing on rice farmers in Bohol, Philippines. A spatial econometrics Bayesian approach was used to estimate the stochastic production parameters, as well as the spatial dependency parameters. The results were compared with non-spatial Bayesian

stochastic frontier approach. We found that spatial dependency exists at the residential and plot levels, maintaining more strength for irrigated than rainfed farms. We also found that technical inefficiency levels decrease as spatial effects are more taken into account. Since the spatial effects increases with a shorter network distance, the decreasing technical inefficiency means that the unobserved inefficiencies can be explained better by considering small networks of relatively close farmers over large networks of distant farmers, reflecting the location-specific nature of farming.

Two policy implications can be drawn from this study. First, a stronger spatial dependency in the irrigated area indicates the existence of stronger externalities; a positive shock on one farmer's TE improves the TE of the nearby farmers. The existence of externalities may justify public interventions. However, it is important to note that we also found that the size of the spatially dependent network is small. Hence, such an externality may be easily internalized through collective actions within the small group. In irrigated area, the water users group may serve as an appropriate unit for this purpose. Although this is an important practical issue, it is beyond the scope of this paper and requires future study. Additionally, since the rainfed farming is more individualistic, policies which are targeted to individual farmers are relatively more important, in comparison to the case of irrigated area. Having observed a strong impact of schooling years, educational support or extension may work effectively in improving rainfed farmers' technical efficiency, which is currently lower than the irrigated farmers.

Although our analysis focuses on technical efficiency for rice production technology change and scale effect are also relevant aspects to be considered in long-term studies. Our data covers only two years which did not allow for TFP growth and technological progress estimation as done in studies like Singbo and Larue (2016), Umetsu et al. (2003), Nin et al. (2003) and Coelli et al. (2005). In addition, evidence of spatial dependency in technological progress has been largely demonstrated

in the economic literature. The role of spatial dependence in technological progress has mainly been stressed in the context of regional productivity (see Nelson and Phelps, 1966; Benhabib and Spiegel, 1994). Similar to the endogenous growth theory in economic literature, technological progress varies across farmers and it depends on farmers' ability to innovate or use the improved technologies. As we highlighted above, farmers use the technology differently with human capital or the level of education being commonly cited as drivers of technological progress. In addition, farmers located below the frontier require sufficient social capabilities to allow them to successfully exploiting the technologies employed by the most efficient farmers.

Acknowledgment: We would like to thank the Japan International Cooperation Agency and Japan International Research Center for Agricultural Science for their financial support of the survey; Shigeki Yokoyama for co-managing the project; Takuji W. Tsusaka for data analysis; Modesto Membreve, Franklyn Fusingan, Cesar Niluag, Baby Descallar and Felipa Danoso of the National Irrigation Administration for arranging the interviews with farmers; and Pie Moya, Lolit Garcia, Shiela Valencia, Elmer Suñaz, Edmund Mendez, Evangeline Austria, Ma. Indira Jose, Neale Paguirigan, Arnel Rala and Cornelia Garcia for data collection. Authors would like to thank the editor- in-charge and two anonymous reviewers for their helpful comments. Pede's time on this research was supported by the RICE CGIAR Research Program.

References

- Aigner, D., Lovell, C.A.K., and Schmidt, P., 1977. Formulation and Estimations of Stochastic Frontier Production Function Models. *J. Econ.* 6(1). 21–37.
- Alvarez, A., 2004. Technical efficiency and farm size: A conditional analysis. *Agric. Econ.* 30(3). 241–250.
- Areal, F.J., Balcombe, K., and Tiffin, R., 2012. Integrating spatial dependence into Stochastic Frontier Analysis. *Aust. J. Agr. Res. Econ.* 56(4). 21–541.
- Areal, F.J., and Riesgo, L., 2014. Farmers' views on the future of olive farming in Andalusia, Spain. *Land. Use. Pol.* 36, 543–553.
- Balde, B.S., Kobayashi, H., Nohmi, M., Ishida, A., Esham, M., and Tolno, E., 2014. An analysis of technical efficiency of mangrove rice production in the Guinean Coastal Area. *J. Agr. Sci.* 6(8).179–196.
- Battese, G.E., 1997. A note on the estimation of Cobb-Douglas production functions when some explanatory variables have zero values. *J. Agr. Econ.* 48(1-3), 250–252.
- Battese, G.E., and Coelli, T.J., 1995. A model for technical inefficiency effects in a Stochastic frontier production function for panel data. *Emp. Econ.* 20(2), 325–332.
- Bell, K.P., and Bocksteal, N.E., 2000. Applying the generalized-moments estimation approach to spatial problems involving microlevels data. *Rev. Econ. Stat.* 82(1), 72–82.
- Benhabib, J. and Spiegel, M., 1994. The Role of Human Capital in Economic Development: Evidence from Aggregate Cross-Country Data. *J. Mon. Econ.* 34, 143–173.
- Bockstael, N.E., 1996. Modeling economics and ecology: the importance of a spatial perspective. *Am. J. Agr. Econ.* 78(5),1168–1180.
- Coelli, T.J., and Battese, G.E., 1996. Identification of factors which influence the technical inefficiency of Indian farmers. *Aust. J. Agr. Econ.* 40(2), 103–128.

- Coelli, T.J., Rao, D.S.P., O'Donnell, C.J. and Battese, G.E., 2005. An introduction to efficiency and productivity analysis. 2nd ed. Springer, New York.
- Druska, V., and Horrace, W.C., 2004. Generalized moments estimation for spatial panel data: Indonesian rice farming. *American J. Agr. Econ.* 86(1), 185–198.
- Farrell, M.J., 1957. The measurement of productive efficiency. *J. Roy. Stat. Soc.* 120(3), 253–290.
- Glass, A.J., Kenjegalieva, K. and Sickles, R.C., 2014. Estimating efficiency spillovers with state level evidence for manufacturing in the US. *Econ. Let.* 123, 154–159.
- Glass, A., Kenjegalieva, K. and Sickles, R.C., 2015. A Spatial autoregressive stochastic frontier model for panel data with asymmetric efficiency spillovers. RISE working paper.
- Hastings, W.K. 1970. Monte Carlo sampling methods using Markov chains and their applications, *Biometrika* 57, 97–109.
- Hossain, E., and Rahman, Z., 2012. Technical efficiency analysis of rice farmers in Naogaon district: an application of the stochastic frontier approach. *J. Econ. Dev. Stud.* 1(1), 2–22.
- Idiong, I.C., 2007. Estimation of farm level technical efficiency in smallscale swamp rice production in Cross River State of Nigeria: a stochastic frontier approach. *Worl. J. Agr. Sci.* 3(5), 653–658.
- JICA., 2012. Impact evaluation of Bohol irrigation project (Phase2) in the Republic of the Philippines. JICA, Tokyo.
- Karagiannis, G., and Tzouvelekas, V., 2009. Measuring technical efficiency in the stochastic varying coefficient frontier model. *Agric. Econ.* 40(4), 389–396.
- Kelejian, H.H., and Prucha, I.R., 1999. A generalized moments estimator for the autoregressive parameter in a spatial model. *Int. Econ. Rev.* 40(2), 509–533.
- Kim, C.W., Phippa, T.T., and Anselin, L., 2003. Measuring the benefits of air quality improvement: a spatial hedonic approach. *J. Env. Econ. Man.* 45(1), 24–39.

- Koop, G., and Steel, M.F.J., 2001. Bayesian analysis of stochastic frontier models, in B. H. Baltagi, ed., *A companion to theoretical econometrics*, Blackwell, Malden, MA , USA.
- Koop, G., 2003. *Bayesian econometrics*. John Wiley & Sons Inc., Chicester, West Sussex, UK.
- Kumbhakar, S.C., and Tsionas, E.G., 2005. Measuring technical and allocative inefficiency in the translog cost system: a Bayesian approach. *J. Econ.*126(2), 355–384.
- Kumbhakar, S.C., Ghosh, S., and McGukin, J.T., 1991. A generalized production frontier approach for estimating determinants of inefficiency in U.S. dairy farms. *J. Bus. Econ. Stat.* 9(3), 279-286.
- Meeusen, W., and van Den Broeck, J., 1977. Efficiency estimation from Cobb-Douglas production functions with composed error. *Int. Econ. Rev.* 18(2), 435-444.
- Metropolis, N., Rosenbluth, A.W., Rosenbluth, M.N., Teller, A. and Teller, E. 1953. Equations of state calculations by fast computing machines, *J.Chem. Phys.* 21, 1087-1092.
- Michler, J.D., and Shively, G.E., 2014. Land tenure, tenure security and farm efficiency: panel evidence from the Philippines. *J. Agr. Econ.* 66(1), 155-169.
- Nelson, R. and Phelps, E., 1966. “Investment in Human, Technological Diffusion, and Economic Growth” *Am. Econ. Rev.* 56:65–75.
- Nin, A., C. Arndt and Preckel, P., 2003. Is agricultural productivity in developing countries really shrinking? New evidence using a modified nonparametric approach. *J. Dev. Econ.* 71, 395–415.
- Roe, B., Irwin, E.G., and Sharp, J.S., 2002. Pigs in space: modeling the spatial structure of hog production in traditional and nontraditional production regions. *Am. J. Agr. Econ.* 84(2), 259–278.
- Schmidt, A.M., Moreira, A.R.B., Helfand, S.M., and Fonseca, T.C.O., 2008. Spatial stochastic frontier models: accounting for unobserved local determinants of inefficiency. *J. Prod. Anal.* 31(2), 101–112.
- Singbo, A. and Larue, B., 2016. Scale economies and the sources of TFP growth of Quebec Dairy farms. *Can. J. Agric. Econ.* 64(2), 339-363.

- Tsionas, E.G. and Michaelides, P.G., 2015. 'A spatial stochastic frontier model with spillovers: Evidence for Italian regions', *Scot. J. Pol. Econ.* 1–14, doi: 10.1111/sjpe.12081.
- Tsusaka, T.W., Kajisa, K. , Pedde, V.O., and Aoyagi, K., 2015. Neighborhood effects and social behavior: the case of irrigated and rainfed farmers in Bohol, the Philippines. *J. Econ. Beh. Org.* 118, 227–246.
- Umetsu, C., Lekprichakul, T. and Charavorty, U., 2003: Efficiency and Technical Change in the Philippine Rice Sector: A Malmquist Total Factor Productivity Analysis. *Am. J. Agric. Econ.* 85(4), 943-963.
- Ward, P.S., and Pedde, V.O., 2014. Capturing social network effects in technology adoption: The spatial diffusion of hybrid rice in Bangladesh. *Aust. J. Agr. Res. Econ.* 58, 1–17.