

Effect of dietary vitamin D3 and 25-hydroxyvitamin D3 supplementation on plasma and milk 25-hydroxyvitamin D3 concentration in dairy cows

Article

Accepted Version

Letter to Editor

Guo, J., Jones, A. K., Givens, D. I. ORCID: <https://orcid.org/0000-0002-6754-6935>, Lovegrove, J. A. ORCID: <https://orcid.org/0000-0001-7633-9455> and Kliem, K. E. ORCID: <https://orcid.org/0000-0002-0058-8225> (2018) Effect of dietary vitamin D3 and 25-hydroxyvitamin D3 supplementation on plasma and milk 25-hydroxyvitamin D3 concentration in dairy cows. *Journal of Dairy Science*, 101 (4). pp. 3545-3553. ISSN 0022-0302 doi: 10.3168/jds.2017-13824 Available at <https://centaur.reading.ac.uk/74751/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

To link to this article DOI: <http://dx.doi.org/10.3168/jds.2017-13824>

Publisher: American Dairy Science Association

copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online

Reply to the Letter to the Editor for “A 25-hydroxycholecalciferol-fortified dairy drink is more effective at raising a marker of postprandial vitamin D status than cholecalciferol in men with suboptimal vitamin D status.” (Manuscript doi: 10.3945/jn.117.254789) by Jing Guo, Kim G Jackson, Che Suhaili binti Che Taha, Yue Li, David I Givens, and Julie A Lovegrove

We thanks Dr Thomas R Hill and Dr Ilias Kyriazakis for their comments and feedback on our article.

We agreed to derive vitamin D enriched foods by ‘biofortification’ via adding vitamin D supplements to the diet of animal is practice and possible. Our recently review article (1) has summarised previous vitamin D biofortified studies on eggs and milk and found most of vitamin D biofortified studies have higher vitamin D dose than EU diet limit (1, 3). Furthermore, there are few human randomised controlled trial on the effect of those vitamin D enriched foods on the human vitamin D status. In addition, our lab work (4) by adding vitamin D₃ maximum dose within the EU diet limit (2) into dairy cows’ diet and showed negligible increased of vitamin D (vitamin D₃ and 25(OH) D₃) in the milk produciton. Therefore, vitamin D fortified milk by adding vitamin D supplement directly into milk are needed rather than biofortified milk.

Current finding of ‘dairy drink fortified with 25(OH) D₃ was more effective at raising plasma 25(OH) D₃ concentrations than dairy drink fortified with vitamin D₃ in men with suboptimal vitamin D status’ agreed with previous studies (5, 6), which may indicates the potential application of enriched foods with 25(OH) D₃, although long-term intervention period is needed to confirm the effect of 25(OH) D₃ fortified dairy drinks on human vitamin D status. Furthermore, different vitamin D fortified food are needed to accommodate dietary

diversity (7), thus, future studies are needed to investigate the effect of other foods fortified with 25(OH) D₃ compared with vitamin D₃ on human vitamin D status.

References

1. Guo J, Lovegrove JA, Givens DI. 25(OH) D₃ enriched or fortified foods are more efficient at tackling inadequate vitamin D status than vitamin D₃. *Proc Nutr Soc.* 2017 [Accepted and in press].
2. EFSA FEEDAP Panel (EFSA Panel on Additives and Products or Substances used in Animal Feed) (2012). Scientific Opinion on the safety and efficacy of vitamin D₃ (cholecalciferol) as a feed additive for chickens for fattening, turkeys, other poultry, pigs, piglets (suckling), calves for rearing, calves for fattening, bovines, ovines, equines, fish and other animal species or categories, based on a dossier submitted by DSM. *EFSA J* 10, 2968, 26 pp.
3. Commission Implementing Regulation (EU) No 2017/1492 of 21 August 2017 concerning the authorisation of cholecalciferol as a feed additive for all animal species. *Official Journal of the European Union* L216/19.
4. Guo J, Jones AK, Givens DI, Lovegrove JA, Kliem KE. Effect of dietary vitamin D₃ and 25(OH) D₃ supplementation on plasma and milk 25(OH) D₃ concentration in dairy cows. *J Dairy Sci* [under review].
5. Cashman KD, Seamans KM, Lucey AJ, Stocklin E, Weber P, Kiely M, Hill TR. Relative effectiveness of oral 25-hydroxyvitamin D₃ and vitamin D₃ in raising wintertime serum 25-hydroxyvitamin D in older adults. *Am J Clin Nutr* 2012;95:1350–6.
6. Jetter A, Egli A, Dawson-Hughes B, Staehelin HB, Stoecklin E, Goessl R, Henschkowski J, Bischoff-Ferrari HA. Pharmacokinetics of oral vitamin D-3 and calcifediol. *Bone* 2014;59:14–9.

7. Cashman KD, Kiely M. Tackling inadequate vitamin D intakes within the population: fortification of dairy products with vitamin D may not be enough. *Endocrine* 2016;51:38-46.