

Apportioning bacterial carbon source utilization in soil using ^{14}C isotope analysis of FISH-targeted bacterial populations sorted by Fluorescence Activated Cell Sorting (FACS): ^{14}C -FISH-FACS

Article

Accepted Version

Gougoulias, C., Meade, A. and Shaw, L. (2018) Apportioning bacterial carbon source utilization in soil using ^{14}C isotope analysis of FISH-targeted bacterial populations sorted by Fluorescence Activated Cell Sorting (FACS): ^{14}C -FISH-FACS. *Environmental Microbiology Reports*, 10 (3). pp. 245-254. ISSN 1758-2229 doi: <https://doi.org/10.1111/1758-2229.12631> Available at <https://centaur.reading.ac.uk/75599/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

To link to this article DOI: <http://dx.doi.org/10.1111/1758-2229.12631>

Publisher: Wiley-Blackwell

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other

copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online

Apportioning bacterial carbon source utilization in soil using ^{14}C isotope analysis of FISH-targeted bacterial populations sorted by Fluorescence Activated Cell Sorting (FACS): ^{14}C -FISH-FACS

Christos Gougoulias^{1†}, Andrew Meade² and Liz J. Shaw^{1*} 

¹Soil Research Centre, Department of Geography and Environmental Science, School of Human and Environmental Sciences, University of Reading, Whiteknights, Reading, RG6 6DW U.K.

²School of Biological Sciences, University of Reading, Whiteknights, Reading, RG6 6 6BX U.K.

*Correspondence: e-mail: e.j.shaw@reading.ac.uk; Phone: +44 (0)118 378 6971; Fax: +44 (0)118 378 6666

†Present address: APHA Scientific, APHA Weybridge, New Haw, Addlestone, KT15 3NB U.K.

Running title: Apportioning bacterial carbon source utilization in soil

Originality-significance statement: We present a novel combination of methodology (^{14}C -FISH-FACS) to address an unresolved need in microbial ecology: the ability to quantify *in situ* microbial substrate carbon use at the population level within a whole system carbon mass balance. Using ^{14}C -FISH-FACS, we demonstrate assimilation of salicylic acid by targeted *Pseudomonas* spp. and *Burkholderia* spp. populations. In conjunction with analysis of the

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process which may lead to differences between this version and the Version of Record. Please cite this article as an 'Accepted Article', doi: 10.1111/1758-2229.12631

taxonomic distribution of the salicylic acid biodegradation trait, we evidence that the ability of the targeted populations to capture a large proportion (~50%) of the added C was not due to conservation of this trait within the targeted group but instead due to competitiveness of this population for the added carbon. The sensitivity of ^{14}C -FISH-FACS and its compatibility with obtaining a full-system mass balance allows a quantitative dissection of C flow within the microbial biomass black box that has hitherto not been achieved to the same detail. ^{14}C -FISH-FACS will enable advances in understanding of population-specific C competitiveness, C use traits and their role in controlling overall C dynamics and microbial community composition in soil and rhizosphere.

Summary

An unresolved need in microbial ecology is methodology to enable quantitative analysis of *in situ* microbial substrate carbon use at the population level. Here, we evaluated if a novel combination of radiocarbon- labelled substrate tracing, Fluorescence *in situ* Hybridisation (FISH) and Fluorescence-Activated Cell Sorting (FACS) to sort the FISH-targeted population for quantification of incorporated radioactivity (^{14}C -FISH-FACS) can address this need. Our test scenario used FISH probe PSE1284 targeting *Pseudomonas* spp. (and some *Burkholderia* spp.) and salicylic acid added to rhizosphere soil. We examined salicylic acid- ^{14}C fate (mineralized, cell-incorporated, extractable and non-extractable) and mass balance (0-24 h) and show that the PSE1284 population captured ~50% of the Nycodenz extracted biomass ^{14}C . Analysis of the taxonomic distribution of the salicylic acid biodegradation trait suggested that PSE1284 population success was not due to conservation of this trait but due to competitiveness for the added carbon. Adding 50KBq of ^{14}C sample $^{-1}$ enabled detection of ^{14}C in the sorted population at ~60-600 times background; a sensitivity which demonstrates potential extension to analysis of rarer/ less active populations. Given its sensitivity and compatibility with obtaining a C mass balance, ^{14}C -FISH-FACS allows quantitative dissection of C flow within the microbial biomass that has hitherto not been achieved.

Introduction

Soil microorganisms play a critical role in driving the cycling of soil organic carbon (Nielsen et al., 2011; Gougoulias et al., 2014; Wieder et al., 2015) and in recognition of this importance there has been considerable discussion about the representation of microbial-scale processes and

the composition of the microbial biomass in carbon cycle models (McGuire and Treseder, 2010; Trivedi et al., 2013; Wieder et al., 2015). Such representation of microbial communities requires knowledge of the relationship between microbial community composition and the rate of carbon cycling processes (Trivedi et al., 2013). In a synthesis of findings from studies that have investigated the relationship between soil biodiversity and ecosystem functions related to C cycling, Nielsen et al. (2011) have concluded that whilst the overall richness of soil communities may have minimal effects on soil C cycling, changes in community composition that result in a changed abundance and activity of particularly influential species may be much more important. Therefore, it might be species traits rather than species richness per se that primarily control C cycling processes (Nielsen et al., 2011). For example, the efficiency with which microorganisms convert available C to biosynthesized products versus respired CO₂ (i.e. carbon use efficiency) is a trait with the potential to vary at the microbial population level (Geyer et al., 2016) that ultimately influences the proportion of substrate C that is incorporated in to stable soil organic matter (Miltner et al., 2012; Geyer et al., 2016). However, how the use efficiency of carbon and its destination within the metabolome (i.e. biomass versus exudate production) at the microbial population level scales to influence community and ecosystem level C cycling is underexplored, partly due to technical difficulties in empirically examining the *quantitative* role of individual microbial species in processing C *in situ* (Geyer et al., 2016).

The population specific microbial use of C in the rhizosphere has particular significance for global biogeochemical cycles given the magnitude of the rhizodeposition C flux, its role in priming the decomposition of existing soil organic matter and in driving heterotrophic pathways responsible for the cycling and dissimilation of other elements. We know that through

rhizodeposition, plants shape their microbiome with dynamic C availabilities selecting for fast-growing, competitive genotypes (Lundberg et al., 2012; Peiffer et al., 2013; Chen et al., 2016).

This selection has functional consequences additional to those related to nutrient cycling in terms of the recruitment of microorganisms that are beneficial or deleterious to plant health (Berendsen et al., 2012). Whilst there have been many studies that have examined the impact of plant genotype or plant growth stage on the rhizosphere microbiome, we know less about the competitiveness of specific microbial populations for components of rhizosphere C flow. Such understanding might be indispensable for engineering the rhizosphere for benefit but, again, is partly hampered by technical difficulties in empirically examining the quantitative role of individual microbial species in processing C *in situ*.

A central goal in microbial ecology has long been to link carbon use to specific microbial taxa in the environment and a suite of methods involving the tracing of ^{13}C - or ^{14}C -labelled substrates (reviewed by Gutierrez-Zamora and Manfield, 2010; Musat et al., 2012; Wang et al., 2016) have been developed in order to achieve this goal without the need for cultivation and characterization of isolates. These methods have been classified into two groups (Gutierrez-Zamora and Manfield, 2010): (i) isotope probing methods (e.g. DNA-, RNA-, PLFA- and protein-stable isotope probing (SIP)) which allow the identification of functional members of a community without prior knowledge of their identity; and, (ii) probe-based methods with either *in situ* (e.g. microautoradiography–fluorescence *in situ* hybridisation (MAR-FISH), FISH-Raman microspectroscopy, Nano secondary ion mass spectrometry *in situ* hybridization (NanoSIMS-ISH)) or *ex situ* (e.g. CHIP-SIP) hybridisation which require prior knowledge of the identity of the population of focus. The development of these methods has revolutionised microbial

ecology over the last two decades, however, the methods vary in the extent to which they can *quantitatively* apportion the utilisation of C to specific microbial populations *in situ*. With nucleic acid- based isotope probing methods, it is not generally possible to determine the extent of isotopic incorporation per phylogenetic group due to reliance on PCR for analysis of microbial identity which dilutes the isotopic content of the initial DNA or RNA (Gutierrez-Zamora and Manefield, 2010). In contrast, quantification of isotopic incorporation is possible with Protein-SIP and PLFA-SIP (Gutierrez-Zamora and Manefield, 2010; Lunsmann et al., 2016). Probe-based methods also generally allow, with appropriate calibration, the microbial incorporation of C from C isotope-labelled substrates to be quantified. In “Chip-SIP” isotope array analysis, *ex situ* hybridisation to a high-density microarray is used to phylogenetically sort extracted rRNA into individual rRNAs, and then their isotopic enrichment is quantified with NanoSIMS (Mayali et al., 2012). In *in situ* methods, isotope incorporation at the single cell level can be quantified by MAR, NanoSIMS or Raman microspectroscopy giving indispensable insight in to microbial processes at the scale approaching that at which they occur (Gutierrez-Zamora and Manefield, 2010; Eichorst et al., 2015).

The above quantitative identity-function methods add a great deal to the methodological tool box in microbial ecology, however, they also have certain limitations. Compared to nucleic-acid based methods, protein-SIP and PLFA-SIP are limited in phylogenetic resolution; labelled peptides or labelled PLFAs can often only be assigned to bacterial families or orders (Jehmlich et al., 2016) or are biomarkers for broad groups of microorganisms (Gutierrez-Zamora and Manefield, 2010), respectively. For probe-based methods, even including Chip-SIP where multiple probes are deployed in parallel, a concern is that taxa coverage is dependent on

selection of probes and therefore analysis might miss activity in unexpected taxa (Mayali et al., 2012). In addition, the application of NanoSIMS to soil samples is not straightforward due to the large background of soil particles and sensitivity of the technique to sample topographical unevenness and label dilution due to redeposition of sputtered material (Eichorst et al., 2015). Similarly, Raman microspectroscopy also requires cells that are dispersed from soil constituents to retrieve reliable cell spectra (Eichorst et al., 2015). NanoSIMS-ISH, FISH-Raman, Protein-, PLFA- and Chip-SIP also rely on access to costly and specialised analytical instrumentation (LC-MS/MS; GC-c-IRMS; NanoSIMS; Fluorescence-equipped Raman Microscope; Array Synthesis) meaning that access might be an issue for many researchers.

When the interest is quantifying flows of total substrate carbon, however, a large constraint is that methods which quantify C incorporation into specific classes of biomolecules (protein-, PLFA- and Chip-SIP) do not provide a whole-cell view of C incorporation by a microbial group of interest. Extrapolation of C incorporation from the biomolecule to the whole cell level that is prerequisite to apportioning C use to a population within the context of a system mass balance is problematic because, at the cell level, microorganisms will not allocate substrate derived carbon equally within their metabolome and for biomolecule synthesis. Out of the methods reviewed, only the *in situ* methods detecting C isotopic incorporation at the whole cell level (by MAR, NanoSIMS) negate the need to extrapolate from biomolecules to whole cells if trying to understand and quantify the flows of total substrate carbon. However, these methods are limited in the number of cells per sample that can be analysed (Clode et al., 2009; Wang et al., 2016). Consequently, it might also be difficult to quantify C apportionment to a given population within

the total C mass balance of added substrate due to the challenges of scaling-up results obtained from a few cells.

Therefore, as a response to some of these issues, here we develop a ^{14}C labelling and probe-based approach in which FISH-hybridised whole cells are isolated by fluorescence activated cell sorting (FACS) prior to quantification of ^{14}C incorporation (^{14}C -FISH-FACS). This approach has methodological steps in common with the established technique of MAR-FISH but employs FACS and scintillation counting instead of MAR for high throughput quantification of ^{14}C incorporation by hybridised cells. In previous work (Gougoulis and Shaw, 2012), we have evaluated the ability of a Nycodenz -based density gradient centrifugation protocol to obtain representative cell extracts suitable for FACS from soil and also validated the environmental specificity of FISH probe PSE1284 primarily targeting *Pseudomonas* spp. (but also some *Burkholderia* spp.) through 16S rDNA sequencing of FACS-sorted hybridized populations. Accordingly, in this research, we utilised probe PSE1284 and, as a test scenario for the ^{14}C -FISH-FACS method, chose salicylic acid as the applied carbon source. Salicylic acid is a plant hormone and has been identified as a major root exudate in several plant species (Schmidt et al., 2000; Hao et al., 2010; Khorassani et al., 2011). Members of the *Pseudomonas* genera, the main group targeted here, are known degraders of salicylic acid and also important players in the rhizosphere in consequence of their roles in promotion of plant health and nutrition (Santoyo et al., 2012).

Results and discussion

Fate of ^{14}C salicylic acid

After an initial short (~4 h) lag phase, salicylic acid underwent rapid mineralization with logistic kinetics that plateaued within 48 hours with a final asymptotic mineralization of ~80% of the ^{14}C -carbon applied (Fig. 1a). Within the first 24 h, the increase in the mineralized fraction was accompanied by a decrease in the extractable ^{14}C from ~70% (at time 0 h) to ~5% of the applied radioactivity (at time 24 h) and a concomitant increase in non-extractable ^{14}C associated with the solid phase (post extraction solids (PES); from 33 to ~50%) and extractable biomass (from <0.1 to ~1.6%) ^{14}C (Fig. 1b). The rapid dissipation of aqueous-extractable salicylic acid after addition to soil is in agreement with Celis et al. (2005) who, for soil slurry incubations receiving 2.76 or 276 $\mu\text{g g}^{-1}$ salicylic acid, reported a reduction of solution-phase salicylic acid concentration to below detection limits or 40% of that applied initially, respectively, over a 24 hour period. The ^{14}C not initially extractable was mostly accounted for in the PES which might be due to rapid sorption of salicylic acid to the soil solid phase, potentially via ligand exchange and formation of bidentate complexes with positively charged metal oxide surfaces (Dubus et al., 2001; Celis et al., 2005; Jagadamma et al., 2012). However, at subsequent time points, the microbial use of salicylic acid C for anabolic (in addition to simultaneous catabolic, i.e. mineralization) processes explains the loss of ^{14}C from the aqueous phase and might explain the additional incorporation of ^{14}C in to the PES through production of non-extractable microbial products and biomass (Nowak et al., 2011). The microbial metabolism of salicylic acid via catechol or gentisic acid to TCA cycle intermediates has been well studied (Karegoudar and Kim, 2000) and the short lag phase prior to rapid mineralization presumably reflects the time required for the induction of salicylic acid catabolic enzymes (Schell, 1985). The final asymptotic amount of ^{14}C mineralized of ~80%

is in agreement with studies that have characterized the mineralization of low concentrations (1 and 1000 ng ml⁻¹) of related aromatic acids (benzoic and phenylacetic) in soil slurries (Amador and Alexander, 1988). The overall mass balance of ¹⁴C recovered across all fractions varied between 103 ± 1.6 % and 86 ± 0.9 % of total applied radioactivity at 0 h and 12 h (Fig. 1c), respectively; losses due to mineralisation during the Nycodenz extraction processing time (4-6 h) are believed to be responsible for the decreased overall recovery seen at 12h and 24h (Fig. 1c).

PSE1284-positive population dynamics and salicylic acid C capture

The PSE1284 hybridising population, which was initially present at ~1.5% of the total FC-detected population increased 7-fold in the 24-h following salicylic acid addition to constitute 9.3 ± 0.2 % of the total FC detected population (Fig 2a and 2b). PSE1284-hybridized populations (n=4 per time point), gated as in the example in Fig. 2a, were recovered by FACS (mean number of cells sorted ranged from 3.7 x 10⁵ to 2.3 x 10⁶, Fig 2b) and the salicylic acid ¹⁴C associated with the sorted cells quantified by scintillation counting. Examining the apportionment of salicylic acid ¹⁴C to PSE1284-positive cells over time (Fig. 3) as a function of the ¹⁴C activity recovered in the total Nycodenz-extracted biomass (also shown in Fig. 1b) reveals an incorporation of 47 ± 4.3% of the salicylic acid-associated radioactivity into the PSE1284-positive biomass over the initial 12 hour period that corresponded to a 4-fold increase in the proportional population size (Fig. 2b). In the subsequent 12 to 24-hour period, PSE1284-positive cells continued to increase their relative abundance (Fig 2b), but their capture of ¹⁴C as a proportion of the ¹⁴C initially added as salicylic acid remained relatively static whilst the Nycodenz-extracted biomass as a whole continued to assimilate ¹⁴C resulting in a decreased

apportionment of salicylic acid ^{14}C to PSE1284-positive cells at 24 hours ($33 \pm 4.2\%$ ^{14}C capture by PSE1284 as a function of the total ^{14}C assimilated). This reduced ^{14}C apportionment might reflect: (i) succession in the assimilation of the added substrate (the PSE1284 population competitively depleted initially high substrate concentrations prior to the utilisation of residual concentrations by a more oligotrophic population); (ii) the turnover of PSE1284-assimilated salicylate ^{14}C and consequent cross-feeding by PSE1284-negative cells. The competitive utilisation of native (potentially primed; Blagodatskaya and Kuzyakov, 2008) soil substrates likely explains how the PSE1284 population could continue to increase in relative abundance in the 12 to 24-hour period (Fig 2b) in the absence of concomitant anabolism of the added ^{14}C source.

As already established, probe PSE1284 targets both *Pseudomonas* and *Burkholderia* spp. It is very likely, given the vast microbial phylogenetic biodiversity known to exist in soils (Lundberg et al., 2012; Peiffer et al., 2013; Chen et al., 2016) that the PSE1284-targetted population represented a small proportion of the diversity present in the studied soil. Against this backdrop, that the PSE1284-positive population was initially able to capture nearly 50% of the carbon that was assimilated in to the extracted biomass suggests specificity/selectivity of salicylic acid for the target population. Plasmid- and chromosomally-mediated biodegradation of salicylic acid has long been associated with members of the genus *Pseudomonas* (e.g. Chakraba.Am, 1972; You et al., 1991; Bosch et al., 1999; Nishi et al., 2000; Sazonova et al., 2008), including *Pseudomonas cepacia* (reclassified as *Burkholderia cepacia* complex; Ramsay et al., 1992) and the C apportionment and response of the PSE1284 population to salicylic acid addition here is compatible with earlier reports that salicylic acid might be used as an “exotic” carbon source to

selectively promote the survival and activity of pseudomonad inoculants in soil (Colbert et al., 1993a; Colbert et al., 1993c; Colbert et al., 1993b). To understand if this selectivity arose because the distribution of the salicylic acid biodegradation trait was conserved within the bacterial groups targeted by the PSE1284 probe, or, through the competitiveness of the PSE1284 population for this substrate against a background of other biodegradative groups, we analysed the taxonomic distribution of putative salicylate hydroxylase enzymes which catalyse the first step in the bacterial degradation of salicylic acid (Supporting Information). This analysis revealed that salicylic acid biodegradation is widely dispersed among bacterial phyla (Supporting Fig. S1) and lower taxonomic levels (Supporting Fig. S2) that are commonly represented in rhizosphere soil and therefore that this trait is not taxonomically conserved. This finding is in agreement with Martiny et al. (2013) who also found that the distribution of traits associated with the assimilation of small organic carbon sources (including p-hydroxy-phenylacetic acid, an aromatic acid) across a range of bacterial phyla to be highly dispersed. Therefore, on the basis of this analysis, it appears that success of the PSE1284 population in exploiting the applied salicylic acid carbon was probably not due to it being the only population in possession of the genetic potential to be able to do so. *Pseudomonas* spp. and *Burkholderia* spp. are well known for their adaptive potential (Silby et al., 2011; Vial et al., 2011) and high competitive ability with strategies, in addition to relatively fast growth rates (Cray et al., 2013), that might allow representatives from these two bacterial genera to outcompete others for salicylic acid carbon. Such strategies might include production of various antibiotic compounds (Raaijmakers and Mazzola, 2012), production of siderophores (dos Santos et al., 2004) and siderophore piracy (Galet et al., 2015).

Evaluation of the ^{14}C -FISH-FACS method

Methods which allow the quantification of C apportionment within the microbial biomass are needed for improved understanding of the relationships between microbial identity, microbial C utilisation traits and C dynamics in soil. Out of the many methods capable of linking biogeochemical processes to specific microbial taxa in the environment, only methods detecting C isotopic incorporation at the whole cell level (rather than biomolecule) level are suitable for the quantification of flows of total substrate carbon through microbial populations. However, as already discussed, current methods which rely on single cell isotopic analysis (by MAR or NanoSIMS) are limited in the number of cells per sample that can be analysed (Clode et al., 2009; Wang et al., 2016). Here, through the application of FACS to sort target cells from the total Nycodenz-extracted cell population we have increased the target cell population that can be analysed from tens of cells to 10^5 to 10^6 cells (Fig. 2b) with modest sorting times of ~ 2 hours. Whilst Nycodenz gradient extraction does not likely achieve exhaustive extraction of bacterial cells from soil (Supporting Table S1; Gougoulias and Shaw, 2012), we know from previous work (Gougoulias and Shaw, 2012) that a diverse community is recovered although with some extraction bias. In relation to this, a key assumption of the method as applied to apportion C flow through microbial communities is that any bias in extraction is constant with sampling time and therefore that changes in the proportional abundance of PSE1284-positive cells and in the substrate ^{14}C they captured are representative of dynamics within the total soil bacterial community.

There have been other published studies that have exploited FACS for the isolation and concentration of marine or lacustrine microbial populations for subsequent isotopic analysis.

However, these studies have used ^{13}C (natural abundance or enriched) and therefore required a custom-built spooling wire microcombustion (SWiM) device coupled to a conventional IRMS (Eek et al., 2007; Hansman and Sessions, 2016) or NanoSIMS (Zimmermann et al., 2015) to detect the $^{12}\text{C}/^{13}\text{C}$ ratios in the resulting cells as regular combustion elemental analysis-coupled IRMS requires sample masses (10–100 $\mu\text{g C}$) which are too large to allow analysis of sorted cells (Hansman and Sessions, 2016). In addition, introduction of C-containing fixatives and probes in to cells during FISH protocols might prove problematic in terms of C contamination and therefore for interpretation of sample $^{12}\text{C}/^{13}\text{C}$ ratios, especially when working at or close to natural abundance levels for ^{13}C (Hansman and Sessions, 2016). In comparison to the use of ^{13}C , the analysis of ^{14}C in sorted cell fractions is relatively straightforward using instrumentation (a scintillation counter) that is common place in research laboratories. Because scintillation quantifies ^{14}C activity, it is not subject to the same sample mass requirements (or C contamination issues) and therefore allows analysis of a (fixed and hybridised) cell number (10^5 - 10^6) that can be realistically achieved by FACS. Here we applied 50 KBq of ^{14}C -carbon to each experimental unit which resulted in detection of ~ 30 -300 Bq ^{14}C -carbon in the FACS-sorted population over the incubation (Fig. 3) which is ~ 60 -600 times the background activity detected in controls not receiving ^{14}C . This sensitivity allowed detection of ^{14}C incorporation and potential cross-feeding over short incubation timecourses (Fig. 3). Whilst we added the test C substrate at a relatively high concentration (50 $\mu\text{g g}^{-1}$) it would be possible to focus on 1000-fold lower concentrations whilst still adding the ^{14}C activity required for detection by increasing the specific activity of the substrate applied (based on specific activities of GBq mmol^{-1} that are typical for ^{14}C -labelled compounds as purchased). Thus, the ^{14}C -FISH-FACS method can be applied for study of the apportionment of substrates applied at concentrations (ng g^{-1}) more

representative of those found in soil, for example, as a result of root exudation (Jones, 1998), or, indeed potentially to the study of the microbial fate of root ^{14}C -exudation *in situ*. Furthermore, the ^{14}C detection in our sorted population of up to ~600 times that of background controls (with 5 min scintillation counting time) points to the capability of the method to be applied to microbial groups that are not such major players in substrate utilisation as the PSE1284 population targeted here. In addition, whilst here we focussed on one PSE1284-positive population, there is the option to target and sort, through the use of FISH probes labelled with different chromophores, multiple bacterial groups.

In terms of the potential limitations of our method, the first relates to the health and safety considerations of working with ^{14}C and in particular the exposure of workers and environment to ^{14}C -labelled bacterial aerosol particles during cell sorting. However, our assessment (see Supporting Information) revealed negligible likely exposure from aerosol creation, even under a worst case scenario that simulated a cell sorter nozzle blockage resulting in an unstable sample stream. Secondly, like all probe-based methods, ^{14}C -FISH-FACS requires prior knowledge of the identity of the population(s) of focus and therefore, on its own, cannot be used as a discovery tool. However, ^{14}C -FISH-FACS might usefully be combined, in sequence, to learn more about the quantitative C cycling role played by hitherto unknown microbial groups that have been discovered by qualitative identity-function methods such as DNA or RNA-SIP. Thirdly, in comparison to other *in situ* probe-based methods (FISH-MAR, FISH-Raman, NanoSIMS-ISH), the quantification of the isotopic incorporation of the sorted cell fraction in bulk precludes the determination of cell-specific information and therefore analysis of functional differences between single cells of the same population. To overcome this, an aliquot of the sorted cells

might be taken for subsequent single cell analysis (for example, by MAR) to get an appreciation of the heterogeneity of metabolism within the microbial population of focus.

Balancing the above-discussed advantages and limitations, we conclude that ^{14}C -FISH-FACS makes a valuable companion to the toolbox of methods capable of quantifying the function of microbial cells *in situ* as demonstrated here in the quantification of the competitive use of salicylic acid by *Pseudomonas* and *Burkholderia* spp. As the Nycodenz protocol used to obtain FACS-ready cell extracts is compatible with monitoring whole system fate (mineralized, extractable, non-extractable) of the added isotope (Fig. 1), including the ^{14}C incorporated by the extractable non-target as well as the target cell population (Fig. 3) over time, we believe our method allows a quantitative dissection of C flow within the microbial biomass black box that has hitherto not been achieved to the same detail.

Experimental Procedures

Rhizosphere Soil

A sandy loam soil (2% organic matter, pH 6.5; sampled from the Rowland Series of the University of Reading Farm, Sonning, UK, NGR SU765765) in pots (10 cm diameter containing 270-300 g soil) was sown with *L. perenne* seed (Herbiseed, UK) at a density of 0.013 g seed cm^{-2} . After growth (15-20 °C) of the *L. perenne* seedlings for 4 weeks, the whole soil content from triplicate pots (defined as rhizosphere soil) was sieved (<2 mm), mixed to form a composite soil and used for timecourse experiments.

¹⁴C-salicylic acid incubation timecourse

Replicate gas-tight glass vials (40 ml) containing rhizosphere soil (15.7 g dry weight basis; moisture content 7.8 % v/w) were spiked with a solution of ¹⁴C-salicylic acid (2-hydroxybenzoic acid-phenyl-UL-¹⁴C) that had been derived from mixing stock solutions of ‘hot’ ¹⁴C-salicylic acid (≥98% purity, Sigma, UK; dissolved in methanol) and ‘cold’ sodium-salicylate (dissolved in ultra-pure water) in order to simultaneously deliver 50 µg salicylic acid per gram of soil (dry weight) and 50 KBq of radioactivity per vial. Controls that received only ‘cold’ salicylic acid (50 µg g⁻¹) were included. Small test-tubes containing a solution of NaOH (2M, 2 ml) to trap ¹⁴C-CO₂ released as a result of mineralization were placed inside the vials on the soil surface and the vials were sealed and incubated at a constant temperature of 26 ± 0.2 °C, under dark conditions. At 0, 12 and 24 h, four replicate vials were destructively sampled for mineralization: the test tubes were removed and the ¹⁴C in the NaOH quantified by liquid scintillation counting (LSC) using a Wallac 1409 DSA liquid scintillation counter (Perkin-Elmer, Boston, MA) with Ultima-Gold (PerkinElmer) as the scintillant in a 1-to-4 (v/v) sample-to-scintillant ratio. The soil in these vials was then used as the basis for soil extractions as described below. In addition, for extended quantification of mineralization only, replicate vials receiving a lower activity of ¹⁴C-salicylic acid (0.5 KBq) were established for sampling of NaOH traps at time points beyond 24 h.

Soil extraction, analysis and ¹⁴C mass balance

The entire soil contents of the vials sampled at 0, 12 and 24 h from the incubation timecourse (above) was slurried in 14.0 ml of phosphate-buffered saline (PBS; 130 mM sodium chloride, 10mM sodium phosphate [pH 7.3]) by vortexing for 2 min with twelve glass beads (6 mm). The

slurry was then subject to low-speed centrifugation ($\sim 600 \times g$, 5 min, ambient temperature) to sediment large soil particles and debris. The resulting supernatant (containing fine soil particles and microbial cells) was removed and retained, and the slurring and low-speed centrifugation step was repeated two additional times by adding 8 ml and 7 ml PBS, sequentially, to the remaining soil pellet. The ^{14}C -activity that remained in the soil pellet (termed Post Extraction Solids ^{14}C (PES- ^{14}C)), was quantified by oxidation (four analytical replicates per sample) using a Perkin Elmer Model 307 combustor after freeze drying to avoid any losses of ^{14}C -salicylic acid due to further mineralisation during drying. The slurry supernatants from the successive extractions were pooled and dispensed to 2 ml microcentrifuge tubes (~ 10 tubes per sample) and concentrated by high speed centrifugation ($16,000 \times g$, 10 min, 4°C). ^{14}C -activity in the resulting supernatants (“extractable ^{14}C ”) was quantified by LSC as described above for the NaOH samples whilst the pellet (containing fine soil particles and microbial cells) was used as the basis for the Nycodenz purification of bacterial cells (from similar-sized soil particles) to prepare samples ready for FISH, flow cytometry and cell sorting as described below. The total ^{14}C -activity recovered in each fraction (extractable ^{14}C , PES- ^{14}C , bacterial biomass- ^{14}C) was calculated as a percentage of total applied radioactivity and summed to give the mass balance in each vial. The mass balance included non-biomass associated ^{14}C that was recovered during the Nycodenz purification, fixation, FISH and washing steps; these fractions were collected, LSC counted and the ^{14}C activity counted as “extractable ^{14}C ”.

Nycodenz purification and fixation of bacterial cells

The main steps of the Nycodenz protocol, as modified from Gougoulis and Shaw (2012) were as follows: the pellets (fine soil particles with bacterial cells) of the initial debris-cleared slurry supernatants were resuspended in PBS (~170 μ l per microcentrifuge tube) by rigorous aspiration through a 200 μ l pipette tip. These resuspensions were pooled to give $2 \times \sim 850$ μ l sub-samples which were then layered onto individual density cushions of 1000 μ l of a 1.3 g/ml Nycodenz (Gentaur, Belgium) solution. Following centrifugation ($16,000 \times g$, 30 min, 4 °C), the Nycodenz/PBS interface containing the final cell suspension (~400 μ l) was removed and diluted twofold with 2 x PBS to provide a purified cell suspension which was harvested by further centrifugation ($16,000 \times g$, 5 min, 4 °C). The two sub-samples were combined to form a 300 μ l composite sample in 1 x PBS to which an equal volume of 96% (wt/vol) ethanol was added. The fixed samples were stored at -20°C for at least 2 h prior to hybridization and for a maximum period of 2 weeks. An aliquot (20 μ l) of the fixed cells was stained with propidium iodide and counted manually under a Brunel epifluorescent microscope to determine the Nycodenz bacterial cell recovery (Supporting Table S1). A further aliquot (30 μ l out of 600 μ l) of the ethanol fixed cells was directly taken for LSC analysis to quantify the radioactivity associated with the total Nycodenz-extracted bacterial fraction. The steps involved in the soil extraction and Nycodenz purification are depicted diagrammatically in the Supporting Information (Fig. S3).

Fluorescence in situ hybridization (FISH)

The following 6-carboxyfluorescein (6-FAM), labeled oligonucleotide probes (Eurofins, MWG-Operon, Ebersberg, Germany) were used: NON-EUB338 [nonspecific for bacteria (Manz et al.,

1992)] and PSE1284 [designed to target *Pseudomonas* spp. (Gunasekera et al., 2003)]. Two aliquots (200 μ l each) of the remaining ethanol fixed cells were washed twice with 1 ml of PBS and resuspended in 180 μ l of prewarmed (48°C) hybridization buffer (0.9 M NaCl, 20 mM Tris-HCl, pH 7.5, 30% formamide, 0.1% sodium dodecyl sulfate) and mixed with 20 μ l either of the specific probe PSE1284 or non-specific probe NONEUB-338 (final concentration: 5 ng μ l⁻¹ for each probe). After incubation for 4 h at 48 °C, the hybridization mixtures were centrifuged (16,000 \times g, 5 min), the cell pellets resuspended in 500 μ l of hybridization buffer and incubated at 48 °C for a further 20 min. The cells were then harvested by centrifugation and incubated in 500 μ l of wash buffer (20 mM Tris-HCl, pH 8.0, 0.9 mM NaCl, 0.1% sodium dodecyl sulfate) for 20 min. Finally, the cells were again harvested by centrifugation and resuspended in 0.2 ml of cold filter-sterilized (0.2 μ m) PBS to await flow cytometric analysis and cell sorting.

Flow cytometry and cell sorting

Flow cytometric analysis and cell sorting were performed with an Influx™ (Becton Dickinson, formerly Cytopeia Inc, Seattle, USA) flow cytometer/cell sorter equipped with a 488-nm solid state laser (Coherent™-Sapphire, output power 200 mW) as described previously (Gougoulas and Shaw, 2012). The Influx™ was also equipped with an extra forward-small size particle detector which allows the detection of particles >0.1 μ m and was calibrated with 0.53 μ m and 3.00 μ m fluorescent microbeads (Saxon Europe Ltd, Kelso, UK). The Influx™ data were acquired as pulse height signals for 100,000 events at a rate of 10,000 to 25,000 events per second with the use of the Spigot™ v.6.1.4 (BD, San Jose-California, USA) software and ~370,000 to 2.3 million events related to the PSE-targeted population were sorted per sample and

LSC counted. Different replicates for each time point were analysed on different days to account for operational variation. During sorting the following parameters were chosen and/or calculated: sheath pressure - 31.7 psi using a 70 µm tip, trigger channel - parallel (small size particle detector), trigger level - 30; drop frequency - 67.9 kHz; piezo amplitude - 2.37-2.51 V; drop delay - 43.8; and, break off point (camera position on drop formation) - 223. Data analysis and graphics were acquired using the FlowJo software package (FlowJo 7.2.4, Tree Star Inc, Ashland, USA).

Acknowledgements

This work was supported by the Biotechnology and Biological Sciences Research Council (BBSRC) in United Kingdom under contract no. BB/F000251/1.

References

Amador, J.A., and Alexander, M. (1988) Effect of humic acids on the mineralization of low concentrations of organic compounds. *Soil Biology & Biochemistry* **20**: 185-191.

Berendsen, R.L., Pieterse, C.M.J., and Bakker, P. (2012) The rhizosphere microbiome and plant health. *Trends in Plant Science* **17**: 478-486.

Blagodatskaya, E., and Kuzyakov, Y. (2008) Mechanisms of real and apparent priming effects and their dependence on soil microbial biomass and community structure: critical review. *Biology and Fertility of Soils* **45**: 115-131.

Bosch, R., Moore, E.R.B., Garcia-Valdes, E., and Pieper, D.H. (1999) NahW, a novel, inducible salicylate hydroxylase involved in mineralization of naphthalene by *Pseudomonas stutzeri* AN10. *Journal of Bacteriology* **181**: 2315-2322.

Celis, R., Real, M., Hermosin, M.C., and Cornejo, J. (2005) Sorption and leaching behaviour of polar aromatic acids in agricultural soils by batch and column leaching tests. *European Journal of Soil Science* **56**: 287-297.

Chakraba. Am (1972) Genetic basis of biodegradation of salicylate in *Pseudomonas* *Journal of Bacteriology* **112**: 815-&.

Chen, L., Brookes, P.C., Xu, J.M., Zhang, J.B., Zhang, C.Z., Zhou, X.Y., and Luo, Y. (2016) Structural and functional differentiation of the root-associated bacterial microbiomes of perennial ryegrass. *Soil Biology & Biochemistry* **98**: 1-10.

Clode, P.L., Kilburn, M.R., Jones, D.L., Stockdale, E.A., Cliff, J.B., Herrmann, A.M., and Murphy, D.V. (2009) In situ mapping of nutrient uptake in the rhizosphere using nanoscale secondary ion mass spectrometry. *Plant Physiology* **151**: 1751-1757.

Colbert, S.F., Hendson, M., Ferri, M., and Schroth, M.N. (1993a) Enhanced growth and activity of a biocontrol bacterium genetically-engineered to utilize salicylate. *Applied and Environmental Microbiology* **59**: 2071-2076.

Colbert, S.F., Schroth, M.N., Weinhold, A.R., and Hendson, M. (1993b) Enhancement of population densities of *Pseudomonas putida* PPG7 in agricultural ecosystems by selective feeding with the carbon source salicylate. *Applied and Environmental Microbiology* **59**: 2064-2070.

Colbert, S.F., Isakeit, T., Ferri, M., Weinhold, A.R., Hendson, M., and Schroth, M.N. (1993c) Use of an exotic carbon source to selectively increase metabolic activity and growth of *Pseudomonas putida* in soil. *Applied and Environmental Microbiology* **59**: 2056-2063.

Cray, J.A., Bell, A.N.W., Bhaganna, P., Mswaka, A.Y., Timson, D.J., and Hallsworth, J.E. (2013) The biology of habitat dominance; can microbes behave as weeds? *Microbial Biotechnology* **6**: 453-492.

dos Santos, V., Heim, S., Moore, E.R.B., Stratz, M., and Timmis, K.N. (2004) Insights into the genomic basis of niche specificity of *Pseudomonas putida* KT2440. *Environmental Microbiology* **6**: 1264-1286.

Dubus, I.G., Barriuso, E., and Calvet, R. (2001) Sorption of weak organic acids in soils: clofencet, 2,4-D and salicylic acid. *Chemosphere* **45**: 767-774.

Eek, K.M., Sessions, A.L., and Lies, D.P. (2007) Carbon-isotopic analysis of microbial cells sorted by flow cytometry. *Geobiology* **5**: 85-95.

Eichorst, S.A., Strasser, F., Woyke, T., Schintlmeister, A., Wagner, M., and Wobken, D. (2015) Advancements in the application of NanoSIMS and Raman microspectroscopy to investigate the activity of microbial cells in soils. *Fems Microbiology Ecology* **91**.

Galet, J., Deveau, A., Hotel, L., Frey-Klett, P., Leblond, P., and Aigle, B. (2015) *Pseudomonas fluorescens* pirates both ferrioxamine and ferricoelichelin siderophores from *Streptomyces ambofaciens*. *Applied and Environmental Microbiology* **81**: 3132-3141.

Geyer, K.M., Kyker-Snowman, E., Grandy, A.S., and Frey, S.D. (2016) Microbial carbon use efficiency: accounting for population, community, and ecosystem-scale controls over the fate of metabolized organic matter. *Biogeochemistry* **127**: 173-188.

Gougoulias, C., and Shaw, L.J. (2012) Evaluation of the environmental specificity of Fluorescence In Situ Hybridization (FISH) using Fluorescence-Activated Cell Sorting (FACS) of probe (PSE1284)-positive cells extracted from rhizosphere soil. *Systematic and Applied Microbiology* **35**: 533-540.

Gougoulias, C., Clark, J.M., and Shaw, L.J. (2014) The role of soil microbes in the global carbon cycle: tracking the below-ground microbial processing of plant-derived carbon for manipulating carbon dynamics in agricultural systems. *Journal of the Science of Food and Agriculture* **94**: 2362-2371.

Gunasekera, T.S., Dorsch, M.R., Slade, M.B., and Veal, D.A. (2003) Specific detection of *Pseudomonas* spp. in milk by fluorescence in situ hybridization using ribosomal RNA directed probes. *Journal of Applied Microbiology* **94**: 936-945.

Gutierrez-Zamora, M.L., and Manefield, M. (2010) An appraisal of methods for linking environmental processes to specific microbial taxa. *Reviews in Environmental Science and Biotechnology* **9**: 153-185.

Hansman, R.L., and Sessions, A.L. (2016) Measuring the in situ carbon isotopic composition of distinct marine plankton populations sorted by flow cytometry. *Limnology and Oceanography-Methods* **14**: 87-99.

Hao, W.Y., Ren, L.X., Ran, W., and Shen, Q.R. (2010) Allelopathic effects of root exudates from watermelon and rice plants on *Fusarium oxysporum* f.sp. *niveum*. *Plant and Soil* **336**: 485-497.

Jagadamma, S., Mayes, M.A., and Phillips, J.R. (2012) Selective sorption of dissolved organic carbon compounds by temperate soils. *Plos One* **7**. DOI: 10.1371/journal.pone.0050434

Jehmlich, N., Vogt, C., Lunsman, V., Richnow, H.H., and von Bergen, M. (2016) Protein-SIP in environmental studies. *Current Opinion in Biotechnology* **41**: 26-33.

Jones, D.L. (1998) Organic acids in the rhizosphere - a critical review. *Plant and Soil* **205**: 25-44.

Karegoudar, T.B., and Kim, C.K. (2000) Microbial degradation of monohydroxybenzoic acids. *Journal of Microbiology* **38**: 53-61.

- Khorassani, R., Hettwer, U., Ratzinger, A., Steingrobe, B., Karlovsky, P., and Claassen, N. (2011) Citramalic acid and salicylic acid in sugar beet root exudates solubilize soil phosphorus. *Bmc Plant Biology* **11**. DOI: 10.1186/1471-2229-11-121
- Lundberg, D.S., Lebeis, S.L., Paredes, S.H., Yourstone, S., Gehring, J., Malfatti, S. et al. (2012) Defining the core *Arabidopsis thaliana* root microbiome. *Nature* **488**: 86-90.
- Lunsmann, V., Kappelmeyer, U., Benndorf, R., Martinez-Lavanchy, P.M., Taubert, A., Adrian, L. et al. (2016) In situ protein-SIP highlights Burkholderiaceae as key players degrading toluene by para ring hydroxylation in a constructed wetland model. *Environmental Microbiology* **18**: 1176-1186.
- Manz, W., Amann, R., Ludwig, W., Wagner, M., and Schleifer, K.H. (1992) Phylogenetic oligodeoxynucleotide probes for the major subclasses of proteobacteria - problems and solutions. *Systematic and Applied Microbiology* **15**: 593-600.
- Martiny, A.C., Treseder, K., and Pusch, G. (2013) Phylogenetic conservatism of functional traits in microorganisms. *Isme Journal* **7**: 830-838.
- Mayali, X., Weber, P.K., Brodie, E.L., Mabery, S., Hoepflich, P.D., and Pett-Ridge, J. (2012) High-throughput isotopic analysis of RNA microarrays to quantify microbial resource use. *Isme Journal* **6**: 1210-1221.
- McGuire, K.L., and Treseder, K.K. (2010) Microbial communities and their relevance for ecosystem models: Decomposition as a case study. *Soil Biology & Biochemistry* **42**: 529-535.
- Miltner, A., Bombach, P., Schmidt-Brucken, B., and Kastner, M. (2012) SOM genesis: microbial biomass as a significant source. *Biogeochemistry* **111**: 41-55.
- Musat, N., Foster, R., Vagner, T., Adam, B., and Kuypers, M.M.M. (2012) Detecting metabolic activities in single cells, with emphasis on nanoSIMS. *Fems Microbiology Reviews* **36**: 486-511.
- Nielsen, U.N., Ayres, E., Wall, D.H., and Bardgett, R.D. (2011) Soil biodiversity and carbon cycling: a review and synthesis of studies examining diversity-function relationships. *European Journal of Soil Science* **62**: 105-116.
- Nishi, A., Tominaga, K., and Furukawa, K. (2000) A 90-kilobase conjugative chromosomal element coding for biphenyl and salicylate catabolism in *Pseudomonas putida* KF715. *Journal of Bacteriology* **182**: 1949-1955.
- Nowak, K.M., Miltner, A., Gehre, M., Schaeffer, A., and Kaestner, M. (2011) Formation and fate of bound residues from microbial biomass during 2,4-D degradation in soil. *Environmental Science & Technology* **45**: 999-1006.
- Peiffer, J.A., Spor, A., Koren, O., Jin, Z., Tringe, S.G., Dangl, J.L. et al. (2013) Diversity and heritability of the maize rhizosphere microbiome under field conditions. *Proceedings of the National Academy of Sciences of the United States of America* **110**: 6548-6553.

Raaijmakers, J.M., and Mazzola, M. (2012) Diversity and Natural Functions of Antibiotics Produced by Beneficial and Plant Pathogenic Bacteria. In *Annual Review of Phytopathology*, Vol 50. VanAlfen, N.K., Leach, J.E., and Lindow, S. (eds). Palo Alto: Annual Reviews, pp. 403-424.

Ramsay, J.R., McEntee, I.D., and Hammond, P.M. (1992) Production and purification of salicylate monoxygenase from *Pseudomonas cepacia* ATCC 29351. *Bioseparation* **2**: 375-383.

Santoyo, G., Orozco-Mosqueda, M.D., and Govindappa, M. (2012) Mechanisms of biocontrol and plant growth-promoting activity in soil bacterial species of *Bacillus* and *Pseudomonas*: a review. *Biocontrol Science and Technology* **22**: 855-872.

Sazonova, O.I., Izmalkova, T.Y., Kosheleva, I.A., and Boronin, A.M. (2008) Salicylate degradation by *Pseudomonas putida* strains not involving the "Classical" nah2 operon. *Microbiology* **77**: 710-716.

Schell, M.A. (1985) Transcriptional control of the nah and sal hydrocarbon-degradation operons by the nahR gene product. *Gene* **36**: 301-309.

Schmidt, S.K., Lipson, D.A., and Raab, T.K. (2000) Effects of willows (*Salix brachycarpa*) on populations of salicylate-mineralizing microorganisms in alpine soils. *Journal of Chemical Ecology* **26**: 2049-2057.

Silby, M.W., Winstanley, C., Godfrey, S.A.C., Levy, S.B., and Jackson, R.W. (2011) *Pseudomonas* genomes: diverse and adaptable. *Fems Microbiology Reviews* **35**: 652-680.

Trivedi, P., Anderson, I.C., and Singh, B.K. (2013) Microbial modulators of soil carbon storage: integrating genomic and metabolic knowledge for global prediction. *Trends in Microbiology* **21**: 641-651.

Vial, L., Chapalain, A., Groleau, M.C., and Deziel, E. (2011) The various lifestyles of the *Burkholderia cepacia* complex species: a tribute to adaptation. *Environmental Microbiology* **13**: 1-12.

Wang, Y., Huang, W.E., Cui, L., and Wagner, M. (2016) Single cell stable isotope probing in microbiology using Raman microspectroscopy. *Current Opinion in Biotechnology* **41**: 34-42.

Wieder, W.R., Allison, S.D., Davidson, E.A., Georgiou, K., Hararuk, O., He, Y.J. et al. (2015) Explicitly representing soil microbial processes in Earth system models. *Global Biogeochemical Cycles* **29**: 1782-1800.

You, I.S., Ghosal, D., and Gunsalus, I.C. (1991) Nucleotide sequence analysis of the *Pseudomonas putida* PPG7 salicylate hydroxylase gene (NahG) and its 3' flanking region. *Biochemistry* **30**: 1635-1641.

Zimmermann, M., Escrig, S., Huebschmann, T., Kirf, M.K., Brand, A., Inglis, R.F. et al. (2015) Phenotypic heterogeneity in metabolic traits among single cells of a rare bacterial species in its natural environment quantified with a combination of flow cell sorting and NanoSIMS. *Frontiers in Microbiology* **6**. DOI: 10.3389/fmicb.2015.00243

Figure Legends

Figure 1. (a) Cumulative mineralization of ^{14}C -salicylic acid ($50 \mu\text{g g}^{-1}$) added to soil sampled from the *L. perenne* rhizosphere over a 132 h timecourse. (b) Partitioning the fate of total applied salicylic acid ^{14}C between mineralized, extractable, post extraction solids (PES) and extracted biomass fractions for the 0-24 hour time points. (c) Mass balance of ^{14}C recovered across all fractions in (b) for the 0-24 hour time points. Data are mean \pm SE (N=4). The ‘extractable’ fraction in the mass balance includes ^{14}C recovered by PBS extraction and in washings during subsequent Nycodenz cell extraction and fixation steps. The following functional form was fitted ($p=0.0003$; $R^2 = 0.98$) to the cumulative mineralization data in (a): $Y = a[1 + (t/t_0)^b]^{-1}$ where Y = cumulative percentage of salicylic acid ^{14}C evolved as $^{14}\text{CO}_2$ and t = time. This function has an asymptote at $a = 80.8 \pm 5\%$.

Figure 2. Flow cytometric (FC) analysis of the response of the PSE1284-6FAM-hybridising population to the addition of salicylic acid ($50 \mu\text{g g}^{-1}$) to *L. perenne* rhizosphere soil. (a) Indicative flow cytometric dot plots where x axis = side light scatter and y axis = green fluorescence per cell in the FITC channel for fixed cells extracted from soil (i,ii) 0-h, (iii) 12-h and (iv) 24-h after salicylic acid addition and targeted with (i) non-specific probe (NON-EUB338-6FAM,) and (ii-iv) probe PSE1284-6FAM. Plots contain 100,000 events. The gated part of the plots indicates the area where the events were sorted from with the numerical value inside the gate depicting the percentage of the total detected events that are in the gate. Graphs shown are indicative of 4 replicates per time point. (b) Mean ($n = 4$; \pm SE) response of the PSE1284-6FAM-hybridising population to the addition of salicylic acid over time (0 -24 h). Data points represent the number of PSE1284-6FAM positive cells as a proportion of the total number of FC detected events (i.e, 100,000 events) whilst the actual number of events sorted as

PSE1284-6FAM positive (the whole aliquot was sorted until the sample was exhausted) are shown above the data points. The correlation between the percentage PSE1284-detected population by FC and the PSE1284-sorted events by FACS had an $R^2 > 0.99$, $p < 0.001$).

Figure 3. Apportionment of salicylic acid ^{14}C to total Nycodenz extracted and PSE1284-sorted bacterial cells over time following addition of ^{14}C -labelled salicylic acid (50 ug/g) to soil. Actual amounts of recovered radioactivity (Bq) are shown on the right Y axis (50,000 Bq was spiked per sample of 15.67 g soil dw). Data are mean \pm SE. Mean values within fraction with a letter in common are not significantly different ($p > 0.05$; Fisher LSD).





