

Condensed tannin changes along the digestive tract in lambs fed with sainfoin pellets or hazelnut skins

Article

Accepted Version

Quijada, J., Drake, C., Gaudin, E., El-Korso, R., Hoste, H. and Mueller-Harvey, I. (2018) Condensed tannin changes along the digestive tract in lambs fed with sainfoin pellets or hazelnut skins. *Journal of Agricultural and Food Chemistry*, 66 (9). pp. 2136-2142. ISSN 1520-5118 doi: <https://doi.org/10.1021/acs.jafc.7b05538> Available at <https://centaur.reading.ac.uk/75832/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

To link to this article DOI: <http://dx.doi.org/10.1021/acs.jafc.7b05538>

Publisher: American Chemical Society

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online

Condensed tannin changes along the digestive tract in lambs fed with sainfoin pellets or hazelnut skins

Jessica Quijada^{†,§*}, Christopher Drake[#], Elodie Gaudin^{†,§,Δ}, Ramzi El-Korso^{†,§}, Hervé Hoste^{†,§},
Irene Mueller-Harvey[#]

[†] INRA, UMR 1225, 23 Chemin des Capelles, Toulouse F-31076, France

[§] Université de Toulouse, ENVT, 23 Chemin des Capelles, Toulouse F-31076, France

[#] School of Agriculture, Policy and Development, University of Reading, , P.O. Box 236, Reading
RG6 6AT, United Kingdom

^Δ MG 2 MIX. La Basse Haye, Châteaubourg 35220, France.

*corresponding author:

E-mail: jessicaquijada7@gmail.com (JQ)

1 **Abstract**

2 The variable anthelmintic efficacy of condensed tannins (CT) against gastrointestinal nematodes
3 may depend on CT concentration, composition or fate along the digestive tract. We analyzed CT
4 concentration and composition by acetone-HCl-butanol and thiolysis coupled to HPLC-MS in
5 digesta and feces of lambs. Lambs had been infected with *Haemonchus contortus* and
6 *Trichostrongylus colubriformis* and received sainfoin pellets and hazelnut skins of contrasting
7 prodelphinidin/procyanidin ratios. The digesta and feces had lower CT concentrations than the
8 original feeds, but similar concentration patterns across the digestive compartments. The changes
9 in assayable CT concentrations between rumen, abomasum and small intestine may be due to
10 complex formation between CT and other dietary components. However, the large CT
11 disappearance (61-85%) from feed to feces could also indicate that CT may have been structurally
12 modified, degraded or absorbed during digestion. Interestingly, there were no changes in the
13 structural features of assayable CT in the digesta.

14 **Keywords:** condensed tannins, nematode, *Onobrychis viciifolia*, *Corylus avellana*, flavan-3-ols,
15 acetone-HCl-butanol, thiolysis, HPLC-MS

16 **Introduction**

17 Tannins are polyphenolic plant compounds and can confer beneficial effects on animal nutrition
18 and health, with anthelmintic (AH) effects being of particular interest.^{1,2} Therefore, tannin-
19 containing resources represent a model to explore the concept of nutraceuticals for controlling
20 gastrointestinal nematodes in ruminants.¹ Proanthocyanidins or condensed tannins (CT) are
21 oligomeric or polymeric flavan-3-ols, where (epi)catechin and (epi)gallocatechin are the most
22 widespread subunits and these give rise to procyanidin (PC) and prodelphinidin (PD) tannins,
23 respectively. In addition, a few plants also contain CT with galloylated flavan-3-ol subunits.³⁻⁵

24 It is often assumed that many of the positive effects of CT in terms of animal health and nutrition
25 are based on their protein binding capacity and possibly also on their antioxidant activities.^{2,6}
26 Formation of CT-protein complexes is thought to cause a shift from urinary to fecal N-excretion,
27 but with a few CT-containing diets this shift can also lead to better dietary protein utilization and,
28 therefore, animal production.^{5,7,8} In addition, dietary CT can also decrease ruminal
29 methanogenesis⁹⁻¹¹ and exert anthelmintic activities.^{1,2,12,13}

30 Our interests focus on the anthelmintic (i.e. antiparasitic) activity of CT against gastrointestinal
31 nematodes both *in vitro* and *in vivo*.^{1,13} Although some *in vitro* and *in vivo* results suggest that CT
32 act via a dose-dependent anthelmintic response,¹⁴⁻¹⁸ CT quantity is not always related to
33 anthelmintic activity.^{19,20} Indeed, recent evidence indicates that CT structural compositions are
34 important for understanding their anthelmintic activities against parasites from cattle,²¹ small
35 ruminants²² and pigs.²³ Of particular interest are polymer size in terms of mean degree of
36 polymerization (mDP) and the composition of monomeric flavan-3-ol subunits (*i.e.* PD/PC ratio),
37 which can modulate their anthelmintic effects.

38 Recent evidence from both *in vitro* and *in vivo* studies suggests that anthelmintic effects vary
39 against gastrointestinal nematode species²⁴ and depend on whether they inhabit the abomasum or
40 the small intestine. Variations with regard to gastrointestinal nematode species have been described
41 *in vitro*. For example, Moreno-Gonzalo et al.^{18,25} evaluated the anthelmintic effect of heather
42 (Ericaceae) extracts on the exsheathment process of *T. circumcincta*, *H. contortus* and *T.*
43 *colubriformis* infective L3 larvae using the larval exsheathment inhibition assay (LEIA). The EC₅₀
44 results showed a higher susceptibility for the intestinal *T. colubriformis* than for the two abomasal
45 species.

46 On the other hand, the effects on gastrointestinal nematodes seem to depend also on the local
47 conditions related either to the host species and/or the local digestive conditions, *e.g.* whether the
48 worms inhabit the stomach or the small intestine. For example, experimentally infected sheep
49 showed a strong anthelmintic effect with quebracho CT against two intestinal species (*Nematodirus*
50 *battus* and *Trichostrongylus colubriformis*) in terms of lower adult worm burden and female
51 fecundity; however, there was no anthelmintic effect against two abomasal species (*Teladorsagia*
52 *circumcincta* and *Haemonchus contortus*).¹⁴ In contrast, the same CT (*i.e.* quebracho) fed to goats
53 reduced the *T. colubriformis* worm burden and *H. contortus* fecundity but there were no changes
54 for *T. circumcincta*.^{26,27}

55 To explain these variations against gastrointestinal nematodes, two hypotheses can be proposed: i)
56 anthelmintic activity stems from a species-specific response or ii) there are differences in CT
57 activity along the digestive tract and the local environmental conditions (*e.g.* pH).^{28,29}

58 For example, with regard to the first hypothesis, when purified CT fractions from 15 different
59 plants were evaluated *in vitro* with the LEIA, Quijada et al.²² observed that nematode species

60 showed different *in vitro* susceptibilities to CT since lower EC₅₀ were recorded for *H. contortus*
61 (more susceptible) than *T. colubriformis*. This also depended on the CT composition. Namely,
62 anthelmintic activity against *H. contortus* (an abomasal species) could be linked to two structural
63 features, mDP-values and PD/PC ratios, whereas for the small-intestinal worm, *T. colubriformis*,
64 only the PD/PC ratio was important. Similar findings on differences in susceptibility between
65 abomasal and intestinal species have also been obtained *in vitro* with gastrointestinal nematodes of
66 cattle.²⁰

67 Up to now, very few studies have addressed the second hypothesis by measuring CT concentrations
68 or activities along the ruminant gut,²⁸⁻³⁰ and no study has compared the effects of CT quality along
69 the gut. Therefore, the present study sought to evaluate the changes of two different CT types from
70 sainfoin plant pellets and hazelnut skins during their passage along the digestive tract of sheep.
71 This study focused i) on CT quantity (concentration) and ii) on CT quality (composition in terms
72 of PD/PC ratios and mDP) in order to assess whether these could explain their *in vivo* anthelmintic
73 activities in lambs, which were experimentally infected with *H. contortus* and *T. colubriformis*.

74 **Materials and Methods**

75 **Trial site**

76 The experiment was carried out at ENVT (National Veterinary School of Toulouse) in the
77 southwest of France (43°35'59'' N, 1°22'41'' E). The facilities hosting the animals and trial
78 performance met and was approved by the French ethical and welfare rules (*Comité d'éthique en*
79 *expérimentation animale* agreement, *Science et Santé Animales SSA N° 115* of December 15,
80 2014). Each group was housed in experimental facilities with concrete floors that had separated
81 boxes of ca. 12 m² each. All animals had ready access to water.

82 **Animals**

83 Twenty-seven 4-month-old lambs of Tarascon breed were used. They had been raised under
84 helminth-free conditions and tests were negative for strongyle nematode infections (by McMaster
85 technique according to Raynaud, 1970) before the start of the study. Diclazuril (Vecoxan®, 2.5
86 mg/mL, Lilly-France, Neuilly-sur-Seine, France) was used, twice at three weekly intervals, at the
87 recommended dose of 1 mg/kg of live weight to prevent coccidian infection. The study was
88 conducted indoors.

89 **Infective larvae**

90 The isolates of either *H. contortus* or *T. colubriformis* were susceptible to anthelmintics. The
91 infective larvae (L3) were cultured from feces of monospecifically infected donor sheep. Larvae
92 were recovered with the Baermann technique and then stored at 4 °C for 1 month (*H. contortus*) or
93 4 months (*T. colubriformis*).

94 **Experimental design**

95 On day 0 (D0), all lambs were orally infected with a single dose of 2000 L3 *H. contortus* and 2000
96 L3 *T. colubriformis*. They had access to *ad libitum* grass hay, mineral block and water and a ration
97 of commercial (tannin-free) pellets. On day 21 (D21) after parasite infection was confirmed by
98 fecal examination, the animals were allocated into three groups of nine lambs, based on
99 experimental diets [hazelnut skin; sainfoin pellets; control (tannin-free) pellets]. The groups were
100 balanced according to sex, live body weight (mean 29.19 ± 2.71 kg), packed cell volume (PCV%
101 = 39.11 ± 2.38) and fecal egg counts (EPG = 1124.1 ± 370.8). From D24 to D28, lambs were
102 allowed to adapt to their diets. During the experimental period (D28 – D57), the rations were
103 adjusted once based on body weight (D34), to meet animal growth requirements. Therefore, from

104 D37 to D44 a second adjustment period was used for the three diets in order to reach an optimal
105 intake level of the two CT-containing diets and to maintain isoproteic and isoenergetic levels in all
106 groups. The condition of the animals was monitored on a daily basis after the infection by checking
107 their feeding and movement behavior and by looking for diarrhea symptoms. Once a week the
108 anemia level was measured (*i.e.* packed cell volume or hematocrit). None of the lambs got severely
109 ill or died during the trial. All lambs were humanely sacrificed under anesthesia, by intravenous
110 injection (3.6 g/lamb) of pentobarbital sodium (Doléthal®, 182.2 mg/mL, Vétquinol S.A.,
111 Magny-Vernois, France) on day D57.

112 **Experimental feeds**

113 Lambs in the experimental group were allocated three different diets. The first group (hazelnut
114 skin) received commercial feed pellets (tannin free) + hazelnut endocarps; the second group
115 (sainfoin) was fed with sainfoin pellets; the third group was the control group and received only
116 commercial, CT-free feed pellets (Passio Ovi Primeur®, Sud Ouest Aliment SOAL, France).
117 During the whole study period (*i.e.* 57 days), all groups received a fattening (total mixed) ration
118 diet, which was isoproteic, isoenergetic and balanced for Ca, P and the Ca:P ratio. Additionally,
119 the two CT-diets (*i.e.* sainfoin pellets and hazelnut skin groups) were fed at equal CT
120 concentrations.

121 **Preparation of digesta and fecal samples**

122 At necropsy, individual digesta samples were retrieved from five lambs (out of nine) per
123 experimental group (*i.e.* sainfoin pellet; hazelnut skin; control). Whole digesta (200 mL) were taken
124 directly from each organ, *i.e.* rumen, abomasum or small intestine (ileum) and fecal samples were

125 collected from the rectum. Each sample was transferred to a 500 mL container and stored at -20
126 °C.

127 The frozen digesta or feces were cooled to -40 °C (-0.5 °C/min) for 2 h (Cryotec, MUT PCCPLS1.5
128 001, France) and freeze-drying was carried out in two phases. Samples were first subjected to a
129 progressive freeze-drying process using the following temperature and pressure program: -30 °C
130 (0.1 °C/min, 0.1 mbar), then at -10 °C (0.2 °C/min, 0.3 mbar) for 19 h 45 min, and finally at -5 °C
131 (0.2 °C/min, 0.15 mbar) until reaching -2 °C. The second phase started when samples had reached
132 -2 °C. They were then kept at 20 °C with a pressure of 0.05 mbar for 15 to 20 h until dry. The
133 freeze-dried digesta or feces were ground in a Retsch impeller SM1 cutting mill (Haan, Germany)
134 to pass a 1 mm sieve and stored at -20 °C until CT analysis.

135 **Condensed tannin analyses**

136 **Chemicals**

137 Hydrochloric acid (37%, analytical reagent grade), acetone (analytical reagent grade), butan-1-ol
138 (standard laboratory reagent grade), acetonitrile (HPLC grade), formic acid (HPLC grade),
139 methanol (HPLC grade) were obtained from Fisher Scientific (Loughborough, UK); benzyl
140 mercaptan (BM) from Sigma-Aldrich (Poole, UK), and ultrapure water (MQ H₂O) from a Milli-Q
141 Plus system (Millipore, Watford, UK).

142 **Tannin analysis by acetone-HCl-butanol assay**

143 The acetone-HCl-butanol assay was described by Grabber et al.³¹ and used with a slight
144 modification as described.²⁸ All samples (sainfoin pellets, control pellets or hazelnut skin, digesta
145 and feces) and a freeze-dried sainfoin sample, which served as an internal laboratory control, were
146 run in triplicate with each batch of samples. . After adding the reagent (10 mL) to the samples (10

147 mg), the tubes were left at room temperature for 1 hour to check for the possibility of flavan-4-ol
148 or flavan-3,4-diol interference. The tubes were then heated at 70 °C for 2.5 hours in the dark. After
149 cooling to room temperature and centrifugation spectra were recorded between 450 and 650 nm on
150 a Jasco V-530 spectrophotometer (Jasco UK, Dunmow, UK). The acetone-HCl-butanol reagent
151 was used as a blank. The absorbance at the peak maximum was determined and converted to CT
152 concentration based on calibration curves derived from a purified prodelphinidin standard, isolated
153 from *Lespedeza cuneata* plants, for sainfoin samples and a purified procyanidin standard, isolated
154 from Tilia flowers, for hazelnut samples.²² The CT concentration was reported as g CT/100 g on a
155 dry weight (DW) basis.

156 **Tannin analysis by thiolysis**

157 The thiolysis reaction was carried out as described previously.³² The reaction products were
158 identified by HPLC-MS analysis^{23,28} and quantified based on peak areas at 280 nm using published
159 flavan-3-ol response factors against taxifolin.^{3,32} This provided information on CT concentration
160 (% CT) and size (mean degree of polymerization, mDP), molar percentages of prodelphinidins
161 (PD) and procyanidins (PC) within CT, and molar percentages of *trans*- vs *cis*-flavan-3-ols (*trans*
162 and *cis*).³ Samples were also analyzed for free flavan-3-ols, but none were detected.

163 **Statistical Analyses**

164 Non-parametric analysis (Kruskal-Wallis and Kolmogorov-Smirnov test) was applied to CT values
165 (CT concentration, mDP, PC, PD, *cis*, *trans*) per sample type (*i.e.* digesta or feces) as determined
166 by each CT assay (acetone-HCl-butanol or thiolysis) and flavan-3-ol terminal and extension units.
167 Comparisons were made between 1) the different diet treatments, and 2) the different segments of
168 the digestive tract within each diet treatment group. All statistical analyses were performed using
169 Systat® 9 software (SPSS Ltd).

170 **Results**

171 **Condensed tannin concentrations in digesta and feces**

172 According to the acetone-HCl-butanol assay, there were no differences ($P > 0.05$) in the CT-
173 concentrations of sainfoin feed pellets and hazelnut skins, *i.e.* 6.5 and 5.1 g CT/100 g DW,
174 respectively (Table 1). As expected the control pellets had no CT. Digesta and fecal samples had
175 significantly lower CT concentrations than the feeds in both the sainfoin- and hazelnut-fed lamb
176 groups (Table 1), *i.e.* from 1.0 to 2.1 g CT/100 g DW. For the lambs of the sainfoin group, these
177 values represented reductions of 84.6 %, 67.7%, 72.4% and 69.2% and for the lambs of the hazelnut
178 group, these CT losses were 78.5%, 66.7%, 76.5% and 60.8% for ruminal, abomasal, small
179 intestinal and fecal samples, respectively. Overall, the CT concentrations showed similar patterns
180 in both groups: slightly higher values were measured in the abomasal and fecal samples, and lower
181 values in the ruminal or small intestinal samples. There were no differences in CT concentrations
182 between the sainfoin and hazelnut groups ($P > 0.05$) but differences were found between the digesta
183 or feces samples within each feed group ($P < 0.05$).

184 In contrast to the acetone-HCl-butanol assay, the thiolysis reaction gave quite different CT
185 concentrations ($P < 0.01$) for the sainfoin pellets (1.7 ± 1.01 g CT/100 g DW) and hazelnut skins
186 (6.3 ± 1.01 g CT/100 g DW) (Table 1). The sainfoin group had the highest CT value in the abomasal
187 digesta (0.7 ± 0.1 g CT/100 g DW), and the hazelnut group in the abomasal and fecal samples
188 (approx. 0.7 ± 0.1 g CT/100 g DW). Thus, apparent CT losses were 85.3%, 58.8%, 76.5% and 76.5%
189 in the sainfoin group, and 92.1%, 88.9%, 93.7% and 88.9% in the hazelnut group in the rumen,
190 abomasum, small intestine and feces compared to the diets, respectively. Differences were found
191 for the CT concentrations measured by thiolysis between the two types of feeds and between the
192 digesta and fecal sample within each feed-group ($P < 0.05$). No differences were recorded between

193 the feed groups when comparing the samples from the same organs ($P > 0.05$). Once, again thiolysis
194 also did not detect any CT in the samples from the control animals.

195 **CT structural features in digesta and feces**

196 Thiolysis also afforded information on the CT composition in terms of molar percentages of
197 prodelphinidins, procyanidins (or PD/PC ratios), *cis*- and *trans*-flavan-3-ols and mean degrees of
198 polymerization (Table 2). The CT in the sainfoin digesta and fecal samples had high percentages
199 of prodelphinidins (*i.e.* rumen 79.5, abomasum 84.1, small intestine 78.7, and feces 72.4%) and
200 *cis*-flavan-3-ols (*i.e.* rumen 87.9, abomasum 91.3, small intestine 87.5, and feces 88.9%), which
201 were similar to the original sainfoin pellets (*i.e.* PD 74.8 and *cis*-flavan-3-ols 85.3%). Due to the
202 low CT concentrations (Table 1), it was not possible to calculate the mDP values in these digesta
203 samples as the peaks of the terminal flavan-3-ol units were too small to be detected. In the hazelnut
204 group, the CT composition was also preserved: hazelnut skins, digesta and fecal samples had high
205 percentages of procyanidins, similar percentages of *cis*- and *trans*-flavan-3-ols and similar mean
206 degrees of polymerization (Table 2).

207 **Discussion**

208 This study was carried out to determine the changes in CT concentrations and compositions during
209 the transit of the sainfoin pellet and hazelnut skin diets in the digestive tract of lambs in order to
210 provide a basis for understating the anthelmintic effects of these diets. Our previous research
211 discovered that gastrointestinal parasites that reside in the abomasum tended to be more sensitive
212 to tannins (*i.e.* lower EC_{50} -values) than parasites that are found in the intestines.²² Lambs were fed
213 with two diets that differed in CT compositions: sainfoin pellets had a high PD/PC ratio (75/25)
214 and hazelnut skins had a low PD/PC ratio (28/72). Samples were taken from along the digestive

215 tract to study CT concentration and compositional changes in the rumen, abomasum, small intestine
216 (ileum) and feces and were compared with the feeds.

217 Given the absence of data on CT changes along the digestive tract, we decided to use two assays
218 that employ different reagents and reaction conditions for the degradation of tannins: the acetone-
219 HCl-butanol reaction uses harsher conditions and is carried out at 70 °C for 2.5 h with 5% HCl and
220 33% water, whereas the thiolysis reaction is milder and takes place at 40 °C for 1 h with <1% HCl
221 in methanol. Previous studies demonstrated that the acetone-HCl-butanol assay can occasionally
222 give higher CT concentrations than the thiolysis assay when plant materials are analyzed.^{2,33,34}

223 **Condensed tannin contents in digesta and feces**

224 There are only a few studies so far that have evaluated changes in CT concentrations in small
225 ruminants and these used a previous, less sensitive, version of the HCl-butanol assay.^{29,30} One
226 recent study also reported thiolysis results for CT concentrations and compositions in digesta from
227 sainfoin-fed cattle, which had been infected with gastro-intestinal nematodes.²⁸ To the best of our
228 knowledge, the current study, therefore, presents for the first time CT concentrations and
229 composition in digesta and feces of lambs. The 60% to 80% decrease of CT concentrations (by
230 acetone-HCl-butanol) from feeds to digesta or feces was comparable to the ¹⁴C-labelled CT losses
231 in sheep of 71.1 - 98.5%.²⁹ Similarly, large decreases in digesta or fecal samples were also
232 described in post-rumen losses in sheep (85 – 86%) and goats (83%).^{29,30}

233 The relatively mild conditions during thiolysis reaction compared to the acetone-HCl-butanol assay
234 may not release all CT from the sample matrix.³² In addition it has also been shown that some CT
235 polymers are resistant to degradation with thiols,^{34,35} which may explain the lower CT
236 concentrations detected by thiolysis than by acetone-HCl-butanol in digesta and feces (Table

237 1).^{28,33,36} Thiolysis also measured much lower CT concentrations than the acetone-HCl-butanol
238 method for the sainfoin pellets (1.7 vs 6.5 g CT/100 g DW) but surprisingly not for the hazelnut
239 skins (6.3 vs 5.1 g CT/100 g DW). The reason for this discrepancy is not clear and will need further
240 investigation; this finding also illustrates the need for using more than one analytical technique
241 when dealing with unusual matrices in order to probe the biological effects of CT.²

242 Despite these differences, both assays revealed a similar pattern (Table 1): the highest CT
243 concentrations were measured in the abomasal samples in both the sainfoin and the hazelnut groups
244 and also in the feces from the hazelnut lamb group. Interestingly, another study that fed sainfoin
245 pellets to cattle also found that CT concentrations were higher in the abomasum (acetone-HCl-
246 butanol: 5.8%; thiolysis 2.3%) than the rumen (acetone-HCl-butanol: 3.0%; thiolysis: 0.5%).²⁸ It is
247 well known that CT bind dietary Rubisco protein optimally at a pH that is close to neutral.² Thus,
248 we hypothesize that dietary proteins are complexed by CT in the rumen (pH 6-7) and released
249 under the acid conditions in the abomasum (pH < 3.5).²⁹ Indeed, the results support this
250 explanation: measured concentrations were highest in the abomasum (Table 1) and a possible
251 explanation could be that these CT were not complexed by proteins and thus remained more
252 accessible and reactive in both assays. In fact, Ramsay et al.³⁴ also noted that benzyl mercaptan in
253 the thiolysis reagent appeared to react preferentially with extractable rather than tightly bound CT.
254 The increased CT concentrations in feces could be due to the combined action of matrix digestion
255 plus bile acids and pH (> 7) that can disrupt CT-protein complexes.³⁰ However, there are also
256 numerous other matrix components with which CT can interact, such as carbohydrates, lipids and
257 intestinal mucosa^{5,37-39} and further work will be needed to establish the interactions between CT
258 and dietary matrix components. Whilst thiolysis appears to preferentially detect extractable CT,^{28,34}
259 the acetone-HCl-butanol assay appears better able to detect bound CT.³⁴

260 However, these results also point to considerable CT modification or degradation in the digestive
261 tract of sheep as pointed out previously with sheep, goat, cattle and pig feeding trials.^{28,29,30,40} If
262 CT were inert, CT concentrations would be expected to increase progressively throughout the tract
263 as dietary matrix components are digested and only the undigestible and non-absorbed components
264 would remain.⁴¹ Mean dry matter digestibilities in sheep are 58% according to a meta-analysis⁴²
265 and, therefore, the CT concentration in feces of sainfoin-fed sheep should have been close to 15%.
266 However, as we could only detect 2% by the acetone-HCl-butanol assay, it would appear that 87%
267 of the CT could no longer be detected. A cattle study that used the same sainfoin diet and acetone-
268 HCl-butanol assay estimated that ca 50% of the CT had disappeared.²⁸ Considerable losses of CT,
269 29% by thiolysis and 17% by acetone-HCl-butanol, were also reported after fermentation of
270 silages³⁴ and from the human digestive tract, where the gut microflora caused extensive losses due
271 to CT metabolism.⁴³

272 **CT structural features in digesta and feces**

273 The CT compositions in Tables 2 and 3 of sainfoin, (mostly prodelphinidins), and hazelnut skins
274 (mostly procyanidins), agree with literature reports.^{32,44,45} Table 3 lists the monomeric subunits that
275 give rise to prodelphinidins (galocatechin and epigallocatechin) and to procyanidins (catechin and
276 epicatechin). Once again, there were no significant changes in these flavan-3-ol compositions
277 between the digesta and the sainfoin feed pellets. The molar composition of these flavan-3-ols
278 decreased as follows: EGC > EC > GC > C, which was in line with the literature.⁴⁵ The flavan-3-
279 ol compositions in the hazelnut skins and the corresponding digesta and fecal samples were also
280 not significantly different (Table 3). However, ca. 5% of the subunits in the hazelnut skins were
281 galloylated, *i.e.* epicatechin gallate (ECg) and epigallocatechin gallate (EGCg), but none of these

282 galloylated subunits could be detected in the digesta or feces, which indicated that the esterified
283 gallic acid may have been cleaved from the CT either by esterases or acids in the gut.

284 It can be concluded that the CT compositional features of PD/PC and *cis/trans* ratios, mean degrees
285 of polymerization, and molar percentages of individual flavan-3-ol subunits were preserved during
286 the digestion in lambs. A similar conclusion was reached after examining the CT composition of
287 ensiled sainfoin.³⁴ These results suggested that CT structures *per se* were not modified during
288 fermentation and digestion - with the exception of esterified gallic acids, which appeared to be
289 cleaved. However, the acetone-HCl-butanol assay measured CT reductions of up to 85% and
290 thiolysis up to 94% in digesta and feces (dry weight basis) compared to the original feeds. These
291 CT decreases suggested that there may be similar processes taking place in the ruminant digestive
292 tract as in the colon of monogastric animals.^{43,47,48} In addition, abomasal digesta samples tended to
293 have the highest levels of assayable CT, which could be due to a matrix effect, as CT tend to bind
294 less strongly at acid pH-values to most proteins.

295 These findings lend support to the hypothesis that CT activity is higher in the abomasum than the
296 intestine, which could explain why CT are more effective against abomasal than intestinal parasite
297 species.⁴⁹ However, our data do not provide support for a species-specific response to CT, despite
298 such evidence from *in vitro* studies with *Haemonchus contortus* (an abomasal species) and
299 *Trichostrongylus colubriformis* (an intestinal species).²² Our results have now revealed that the CT
300 flavan-3-ol subunit composition was preserved along the digestive tract, hence the higher *in vitro*
301 biological activity of prodelphinidins can be expected to be maintained under *in vivo* conditions as
302 long as the overall CT concentration remains sufficiently high.

303 **Abbreviations Used**

304 CT, condensed tannins; PD, prodelphinidins; PC, procyanidins; mDP, mean degree of
305 polymerization; BM, benzyl mercaptan; C, catechin; EC, epicatechin; ECg, epicatechin gallate;
306 EGC, epigallocatechin; EGCg, epigallocatechin gallate; GC, galocatechin.

307 **Acknowledgments**

308 Assistance from Mrs. Fabienne Picard and Dr Vincent Niderkorn (INRA, UMR 1213 Herbivores,
309 Saint-Genès-Campanelle, France) with freeze-drying of digesta samples is deeply appreciated.

310 **Supporting information.** Feed nutritional analyses results for each experimental group are shown
311 in regard to composition, fiber content and nutrition values.

312 **Author's contribution**

313 JQ and HH designed and performed the animal experiments. IMH designed the chemical analyses.
314 EG and REK helped in the animal experiment. JQ, HH and IMH analyzed the data and prepared
315 the manuscript. CD, EG, REK contributed reagents, materials and analysis tools. All authors
316 critically read and approved the final manuscript.

317 **Competing interest**

318 The authors declare that they have no competing interests.

319 **References**

- 320 1. Hoste, H; Torres-Acosta, J.F.J.; Sandoval-Castro, C.A.; Mueller-Harvey, I.; Sotiraki, S.;
321 Louvandini, H.; Thamsborg, S.M.; Terrill, T.H. Tannin containing legumes as a model for
322 nutraceuticals against digestive parasites in livestock. *Vet. Parasitol.* **2015**, 212, 5–17.
- 323 2. Mueller-Harvey, I.; Bee, G.; Dohme-Meier, F.; Hoste, H.; Karonen, M.; Kölliker, R.; Lüscher,
324 A.; Niderkorn, V.; Pellikaan, W.F.; Salminen, J-P.; Skøt, L.; Smith, L.M.J.; Thamsborg, S.M.;
325 Totterdell, P.; Wilkinson, I.; Williams, A.R.; Azuhnwi, B.N.; Baert, N.; Grosse Brinkhaus, A.;
326 Copani, G.; Desrues, O.; Drake, C.; Engström, M.; Fryganas, C.; Girard, M.; Huyen, N.T.;
327 Kempf, K.; Malisch, C.; Mora-Ortiz, M.; Quijada, J.; Ramsay, A.; Ropiak, H.M.; Waghorn,
328 G.C. Benefits of condensed tannins in forage legumes fed to ruminants: importance of structure,
329 concentration and diet composition. *Crop Sci.* **2017**. DOI: 10.2135/cropsci2017.06.0369
- 330 3. Ropiak, H.M.; Ramsay, A.; Mueller-Harvey, I. Condensed tannins in extracts from European
331 medicinal plants and herbal products. *J. Pharm. Biomed. Anal.* **2016**, 121, 225–231.
- 332 4. Hagerman, A. Fifty years of polyphenol-protein complexes. In: Cheynier, V.; Sami-Machado,
333 P.; Quideau, S. (eds) *Recent Advances in Polyphenol Research*. Vol. No. 3. 1st Edition Wiley-
334 Blackwell & Sons, Oxford, UK. **2012**, 364 pp.
- 335 5. Patra, A.K.; Saxena, J. Exploitation of dietary tannins to improve rumen metabolism and
336 ruminant nutrition. *J. Sci. Food Agric.* **2011**, 91, 24–37.
- 337 6. Luciano, G.; Vasta, V.; Monahan, F.J.; López-Andrés, P.; Biondi, L.; Lanza, M.; Priolo, A.
338 Antioxidant status, colour stability and myoglobin resistance to oxidation of *longissimus dorsi*
339 muscle from lambs fed a tannin-containing diet. *Food Chem.* **2011**, 124, 1036–1042.
- 340 7. Buccioni, A.; Pauselli, M.; Viti, C.; Minieri, S.; Pallara, G.; Roscini, V.; Rapaccini, S.;
341 Marinucci, M.T.; Lupi, P.; Conte, G.; Mele, M. Milk fatty acid composition, rumen microbial

- 342 population, and animal performances in response to diets rich in linoleic acid supplemented with
343 chestnut or quebracho tannins in dairy ewes. *J. Dairy Sci.* **2015**, 98, 1145–56.
- 344 8. Waghorn, G.C.; John, A.; Jones, W.T.; Shelton, I.D. Nutritive value of *Lotus corniculatus* L.
345 containing low and medium concentrations of condensed tannins for sheep. *Proc. New Zeal.*
346 *Soc. Anim. Prod.* **1987**, 47, 25–30.
- 347 9. Bhatta, R.; Saravanan, M.; Baruah, L.; Prasad, C.S. Effects of graded levels of tannin-containing
348 tropical tree leaves on *in vitro* rumen fermentation, total protozoa and methane production. *J.*
349 *Appl. Microbiol.* **2015**, 118, 557–564.
- 350 10. Huyen, N. T.; Fryganas, C.; Uittenbogaard, G.; Mueller-Harvey, I.; Verstegen, M.W.A.;
351 Hendriks, W.H.; Pellikaan, W.F. Structural features of condensed tannins affect *in vitro*
352 ruminal methane production and fermentation characteristics. *J. Agr. Sci.* **2016**, 154, 1474–
353 1487.
- 354 11. Liu, H.; Vaddella, V.; Zhou, D. Effects of chestnut tannins and coconut oil on growth
355 performance, methane emission, ruminal fermentation, and microbial populations in sheep. *J.*
356 *Dairy Sci.* **2011**, 94, 6069–6077.
- 357 12. Zhong, R.Z.; Li, H.Y.; Sun, H.X.; Zhou, D.W. Effects of supplementation with dietary green
358 tea polyphenols on parasite resistance and acute phase protein response to *Haemonchus*
359 *contortus* infection in lambs. *Vet. Parasitol.* **2014**, 205, 199–207.
- 360 13. Hoste, H.; Martinez-Ortiz-De-Montellano, C.; Manolaraki, F.; Brunet, S.; Ojeda-Robertos, N.;
361 Fourquaux, I.; Torres-Acosta, J.F.J.; Sandoval-Castro, C.A. Direct and indirect effects of
362 bioactive tannin-rich tropical and temperate legumes against nematode infections. *Vet.*
363 *Parasitol.* **2012**, 186, 18–27.

- 364 14. Athanasiadou, S.; Kyriazakis, I.; Jackson, F.; Coop, R. Direct anthelmintic effects of condensed
365 tannins towards different gastrointestinal nematodes of sheep: *in vitro* and *in vivo* studies. *Vet.*
366 *Parasitol.* **2001**, 99, 205–219.
- 367 15. Brunet, S.; Hoste, H. Monomers of condensed tannins affect the larval exsheathment of
368 parasitic nematodes of ruminants. *J. Agric. Food Chem.* **2006**, 54, 7481–7487.
- 369 16. Brunet, S.; Aufrère, J.; El Babili, F.; Fouraste, I.; Hoste, H. The kinetics of exsheathment of
370 infective nematode larvae is disturbed in the presence of a tannin-rich plant extract (sainfoin)
371 both *in vitro* and *in vivo*. *Parasitology* **2007**, 134, 1253–1262.
- 372 17. Burke, J.M.; Whitley, N.C.; Pollard, D.A.; Miller, J.E.; Terrill, T.H.; Moulton, K.E.; Mosjidis
373 J.A. Dose titration of sericea lespedeza leaf meal on *Haemonchus contortus* infection in lambs
374 and kids. *Vet. Parasitol.* **2011**, 181, 345–349.
- 375 18. Moreno-Gonzalo, J.; Manolaraki, F.; Frutos, P.; Hervás, G.; Celaya, R.; Osoro, K.; Ortega-
376 Mora, L.M.; Hoste, H.; Ferre, I. *In vitro* effect of heather extracts on *Trichostrongylus*
377 *colubriformis* eggs, larvae and adults. *Vet. Parasitol.* **2013**, 197, 586–594.
- 378 19. Naumann, H.D.; Armstrong, S.A.; Lambert, B.D.; Muir, J.P.; Tedeschi, L.O.; Kothmann, M.M.
379 Effect of molecular weight and concentration of legume condensed tannins on *in vitro* larval
380 migration inhibition of *Haemonchus contortus*. *Vet. Parasitol.* **2014**, 199, 93–98.
- 381 20. Novobilský, A.; Stringano, E.; Hayot Carbonero, C.; Smith, L.M.J.; Enemark, H.L.; Mueller-
382 Harvey, I.; Thamsborg, S.M. *In vitro* effects of extracts and purified tannins of sainfoin
383 (*Onobrychis viciifolia*) against two cattle nematodes. *Vet. Parasitol.* **2013**, 196, 532–537.
- 384 21. Novobilský, A.; Mueller-Harvey, I.; Thamsborg, S.M. Condensed tannins act against cattle
385 nematodes. *Vet. Parasitol.* **2011**, 182, 213–220.

- 386 22. Quijada, J.; Fryganas, C.; Ropiak, H.M.; Ramsay, A.; Mueller-Harvey, I.; Hoste, H.
387 Anthelmintic activities against *Haemonchus contortus* or *Trichostrongylus colubriformis* from
388 small ruminants are influenced by structural features of condensed tannins. *J. Agric. Food*
389 *Chem.* **2015**, 63, 6346–6354.
- 390 23. Williams, A.R.; Fryganas, C.; Ramsay, A.; Mueller-Harvey, I.; Thamsborg, S.M. Direct
391 anthelmintic effects of condensed tannins from diverse plant sources against *Ascaris suum*.
392 *PLoS One* **2014**, 9(5): e97053. [DOI: 10.1371/journal.pone.0097053](https://doi.org/10.1371/journal.pone.0097053)
- 393 24. Hoste, H.; Torres-Acosta, J.F.J.; Quijada, J.; Chan-Pérez, J.I.; Dakheel, M.M.; Kommuru, D.S.;
394 Mueller-Harvey, I.; Terrill, T.H. Interactions between nutrition and infections with
395 *Haemonchus contortus* and related gastrointestinal nematodes in small ruminants. In: Gasser,
396 R.B.; von Samson-Himmelstjerna, G. (Ed.), *Haemonchus contortus and Haemonchosis – Past,*
397 *Present and Future Trend.* Elsevier Ltd, **2016**, pp. 239–351.
- 398 25. Moreno-Gonzalo, J.; Manolaraki, F.; Frutos, P.; Hervás, G.; Celaya, R.; Osoro, K.; Ortega-
399 Mora, L.M.; Hoste, H.; Ferre, I. *In vitro* effect of heather (Ericaceae) extracts on different
400 development stages of *Teladorsagia circumcincta* and *Haemonchus contortus*. *Vet. Parasitol.*
401 **2013**, 197:235–243.
- 402 26. Paolini, V.; Bergeaud, J.P.; Grisez, C.; Prevot, F.; Dorchies, P.; Hoste, H. Effects of condensed
403 tannins on goats experimentally infected with *Haemonchus contortus*. *Vet. Parasitol.* **2003**, 113,
404 253–261.
- 405 27. Paolini, V.; Frayssines, A.; Farge, F.; Dorchies, P. Effects of condensed tannins on established
406 populations and on incoming larvae of *Trichostrongylus colubriformis* and *Teladorsagia*
407 *circumcincta* in goats. *Vet. Res.* **2003**, 34, 331–339.

- 408 28. Desrues, O.; Mueller-Harvey, I.; Pellikaan, W.F.; Enemark, H.L.; Thamsborg, S.M. Condensed
409 tannins in the gastrointestinal tract of cattle after sainfoin (*Onobrychis viciifolia*) intake and their
410 possible relationship with anthelmintic effects. *J. Agric. Food Chem.* **2017**, *65*, 1420-1427.
- 411 29. Terrill, T.H.; Waghorn, G.C.; Woolley, D.J.; McNabb, W.C.; Barry, T.N. Assay and digestion
412 of ¹⁴C-labelled condensed tannins in the gastrointestinal tract of sheep. *Br. J. Nutr.* **1994**, *72*,
413 467–477.
- 414 30. Perez-Maldonado, R.A.; Norton, B.W. The effects of condensed tannins from *Desmodium*
415 *intortum* and *Calliandra calothyrsus* on protein and carbohydrate digestion in sheep and goats.
416 *Br. J. Nutr.* **1996**, *76*, 515–533.
- 417 31. Grabber, J.H.; Zeller, W.E.; Mueller-Harvey, I. Acetone enhances the direct analysis of
418 procyanidin- and prodelphinidin-based condensed tannins in lotus species by the butanol-HCl-
419 iron assay. *J. Agric. Food Chem.* **2013**, *61*, 2669–2678.
- 420 32. Gea, A.; Stringano, E.; Brown, R.H.; Mueller-Harvey, I. *In situ* analysis and structural
421 elucidation of sainfoin (*Onobrychis viciifolia*) tannins for high-throughput germplasm
422 screening. *J. Agric. Food Chem.* **2011**, *59*, 495–503.
- 423 33 Azuhnwi, B.N.; Boller, B.; Dohme-Meier, F.; Hess, H.D.; Kreuzer, M.; Stringano, E.; Mueller-
424 Harvey, I. Exploring variation in proanthocyanidin composition and content of sainfoin
425 (*Onobrychis viciifolia*). *J. Sci. Food Agric.* **2013**, *93*, 2102–2109.
- 426 34. Ramsay, A.; Drake, C.; Grosse Brinkhaus, A.; Girard, M.; Copani, G.; Dohme-Meier, F.; Bee,
427 G.; Niderkorn, V.; Mueller-Harvey, I. Sodium hydroxide enhances extractability and analysis
428 of proanthocyanidins in ensiled sainfoin (*Onobrychis viciifolia*). *J. Agric. Food. Chem.* **2015**,
429 *63*, 9471–9479.

- 430 35. Matthews, S.; Mila, I.; Scalbert, A.; Pollet, B.; Lapierre, C.; duPenhoat, C.; Rolando, C.;
431 Donnelly, D.M.X. Method for estimation of proanthocyanidins based on their acid
432 depolymerization in the presence of nucleophiles. *J. Agric. Food Chem.* **1997**, *45*, 1195–1201.
- 433 36. Pérez-Jiménez, J.; Torres, J.L. Analysis of non-extractable polyphenols in foods: the current
434 state of the art. *J. Agric. Food Chem.* **2011**, *59*, 12713–12724.
- 435 37. Bindon, K.A.; Smith, P.A.; Holt, H.; Kennedy, J.A. Interaction between grape-derived
436 proanthocyanidins and cell wall material. 2. Implications for vinification. *J. Agric. Food Chem.*
437 **2010**, *58*, 10736–10746.
- 438 38. Le Bourvellec, C.; Bouchet, B.; Renard, C.M.G.C. Non-covalent interaction between
439 procyanidins and apple cell wall material. Part III: Study on model polysaccharides. *Biochim.*
440 *Biophys. Acta – Gen. Sub.* **2005**, *1725*:10–18.
- 441 39. van Leeuwen, P.; Jansman, A.J.; Wiebenga, J.; Koninkx, J.F.; Mowen, J.M. Dietary effects of
442 faba-bean (*Vicia faba* L.) tannins on the morphology and function of the small-intestinal mucosa
443 of weaned pigs. *Br. J. Nutr.* **1995**, *73*, 31–39.
- 444 40. Choy, Y.Y.; Quifer-Rada, P.; Holstege, D.M.; Frese, S.A.; Calvert, C.C.; Mills, D.A.; Lamuela-
445 Raventos, R.M.; Waterhouse, A.L. Phenolic metabolites and substantial microbiome changes in
446 pig feces by ingesting grape seed proanthocyanidins. *Food Funct.* **2014**, *5*, 2298–2308.
- 447 41. Gedir, J.V.; Sporns, P.; Hudson, R.J. Extraction of condensed tannins from cervid feed and
448 feces and quantification using a radial diffusion assay. *J. Chem. Ecol.* **2005**, *31*, 2761–2773.
- 449 42. Riaz, M.Q.; Südekum, K.H.; Clauss, M.; Jayanegara, A. Voluntary feed intake and digestibility
450 of four domestic ruminant species as influenced by dietary constituents: A meta-analysis. *Livest.*
451 *Sci.* **2014**, *162*, 76–85.

- 452 43. Mena, P.; Calani, L.; Bruni, R.; Del Rio, D. Chapter 6 - Bioactivation of High-Molecular-
453 Weight Polyphenols by the Gut Microbiome. In: *Diet-Microbe Interactions in the Gut* (ed. Rio,
454 K. T. D.), Academic Press, San Diego, **2015**, pp. 73-101.
- 455 44. Del Rio, D.; Calani, L.; Dall'Asta, M.; Brighenti, F. Polyphenolic composition of hazelnut skin.
456 *J. Agric. Food. Chem.* **2011**, 59, 9935–9941.
- 457 45. Wang, Y.; Mc Allister, T.A.; Acharya, S. Condensed tannins in sainfoin: composition,
458 concentration, and effects on nutritive and feeding value of sainfoin forage. *Crop Sci* **2015**, 55,
459 13-22.
- 460 46. Stringano, E.; Hayot Carbonero, C.; Smith, L.M.J.; Brown, R.H.; Mueller-Harvey, I.
461 Proanthocyanidin diversity in the EU “HealthyHay” sainfoin (*Onobrychis viciifolia*) germplasm
462 collection. *Phytochemistry* **2012**, 77, 197–208.
- 463 47. Spiegler, V.; Liebau, E.; Hensel, A. Medicinal plant extracts and plant-derived polyphenols
464 with anthelmintic activity against intestinal nematodes. *Nat. Prod. Rep.* **2017**, 34, 627–643.
- 465 48. Li, M.; Hagerman, A.E. Interactions between plasma proteins and naturally occurring
466 polyphenols. *Curr. Drug Metab.* **2013**, 14, 432-445.49. Arroyo-Lopez, C.; Manolaraki, F.;
467 Saratsis, A.; Saratsis, K.; Stefanakis, A., Skampardonis, V.; Voutzourakis, N.; Hoste, H.;
468 Sotiraki, S. Anthelmintic effect of carob pods and sainfoin hay when fed to lambs after
469 experimental trickle infections with *Haemonchus contortus* and *Trichostrongylus colubriformis*.
470 *Parasite* **2014**, 21, 71-80.

471 **Funding**

472 The research was funded by the European Commission through the PITN-GA-2011-289377
473 “LegumePlus” project.

Table 1. Mean (\pm SD) Concentrations of Condensed Tannin (g CT/100 g DW) Measured Either With the Acetone-HCl-butanol or the Thiolytic Assays in Feeds, Digesta and Fecal Samples from Each Experimental Group (n= 5 lambs).

	Feed	Rumen	Abomasum	Small Intestine	Feces
Acetone-HCl-butanol assay					
Sainfoin pellets group	6.5 \pm 0.3 ^a	1.0 \pm 0.1 ^b	2.1 \pm 0.3 ^{c*}	1.8 \pm 0.3 ^b	2.0 \pm 0.4 ^b
Hazelnut skin group	5.1 \pm 0.1 ^a	1.1 \pm 0.1 ^b	1.7 \pm 0.2 ^{bc}	1.2 \pm 0.1 ^{bc}	2.0 \pm 0.3 ^{c*}
Thiolytic assay					
Sainfoin pellets group	1.7 \pm 0.1 ^{a**}	0.3 \pm 0.1 ^b	0.7 \pm 0.1 ^{c*}	0.4 \pm 0.1 ^{bc}	0.4 \pm 0.1 ^b
Hazelnut skin group	6.3 \pm 0.1 ^{a**}	0.5 \pm 0.1 ^b	0.7 \pm 0.1 ^b	0.4 \pm 0.1 ^b	0.7 \pm 0.1 ^b

** (P < 0.01) indicates significant differences between sainfoin pellets and hazelnut skin feeds; * (P < 0.05)

^{a,b,c} different superscripts within rows indicate significant differences depending on the digestive organs or feces; \pm indicates standard deviations

Table 2. Condensed Tannin Compositions in Digesta or Fecal Samples from Lambs (n= 5)

Fed with either Sainfoin Pellets or Hazelnut Skins.

	mDP	PD/PC %	<i>cis/trans</i>-flavan-3-ols %
Sainfoin pellets	11.5±0.3	74.8/25.2 (±0.5)	85.3/14.7 (±0.1)
Rumen	-	79.5/20.5 (±0.9)	87.9/12.1 (±0.7)
Abomasum	-	84.1/15.9 (±0.50)	91.3/8.7 (±0.5)
Small intestine	-	78.7/21.3 (±1.1)	87.5/12.5 (±0.3)
Feces	-	72.4/27.6 (±1.6)	88.9/11.1 (±1.3)
Hazelnut skin	13.3±0.1	27.9/72.1 (±0.2)	58.4/41.6 (±0.2)
Rumen	14.8±0.7	34.3/65.7 (±1.5)	46.3/53.7 (±1.2)
Abomasum	13.9±0.3	33.4/66.6 (±0.7)	51.3/48.7 (±0.6)
Small intestine	13.8±1.2	33.4/66.6 (±1.7)	46.9/53.1 (±2.3)
Feces	13.2±0.3	18.9/81.1 (±2.4)	48.4/51.6 (±0.6)

Note: there were no significant differences between the different organs.

Abbreviations: mean degree of polymerization (mDP); % refers to molar percentages of procyanidins (PC), prodelphinidins (PD), *cis*- or *trans*- flavan-3-ols (*cis* or *trans*); ± refers to standard deviations

Table 3. Molar Percentages (%) of Terminal and Extension Flavan-3-ol Subunits within CT from Digesta and Fecal Samples Collected from Lambs that Had Been Fed with Sainfoin Pellets or Hazelnut Skins.

	Terminal units (%)				Extension units (%)					
	GC	EGC	C	EC	GC-BM	EGC-BM	C-BM	EC-BM	ECg-BM	EGCg-BM
Sainfoin pellets	2.4±0.1	1.8±0.1	1.9±0.1	2.7±0.1	9.5±0.3	61.2±0.5	0.9±0.1	19.7±0.3	0.0	0.0
Rumen	0.0	0.0	0.0	0.0	10.3±0.5	69.2±1.5	1.7±0.1	18.8±0.9	0.0	0.0
Abomasum	0.0	0.0	1.3±0.0	1.2±0.0	8.3±0.3	75.7±0.7	0.6±0.0	15.4±0.5	0.0	0.0
Small intestine	0.0	0.0	0.0±0.0	1.9±0.0	11.9±0.3	66.7±1.5	0.5±0.0	19.8±1.9	0.0	0.0
Feces	0.0	0.0	2.5±0.2	2.3±0.2	10.1±0.7	62.3±2.2	0.0	24.8±1.1	0.0	0.0
Hazelnut skins	0.0	0.0	7.5±0.1	0.0	12.1±0.1	15.1±0.1	21.2±0.1	39.4±0.3	0.8±0.1	3.9±0.1
Rumen	0.0	0.0	6.8±0.2	0.0	20.1±1.5	14.3±0.3	26.8±0.6	31.9±1.1	0.0	0.0
Abomasum	0.0	0.0	6.5±0.1	0.7±0.1	16.3±0.6	17.2±0.2	25.9±0.6	33.5±0.4	0.0	0.0
Small intestine	0.0	0.0	7.5±0.8	0.0	17.8±0.5	15.6±1.4	27.8±1.7	31.4±1.1	0.0	0.0
Feces	0.0	0.0	7.6±0.2	0.0	10.1±1.7	8.8±0.9	34.0±1.0	39.5±1.3	0.0	0.0

Abbreviations: Gallocatechin (GC), epigallocatechin (EGC), catechin (C), epicatechin (EC), epicatechin gallate (ECg), epigallocatechin gallate (EGCg), benzyl mercaptan adduct (-BM); ± refers to standard deviations

Table of content (TOC)

