

Eating pathology in midlife women: similar or different to younger counterparts?

Article

Accepted Version

Jenkins, P. E. ORCID: <https://orcid.org/0000-0003-1673-2903> and Price, T. (2018) Eating pathology in midlife women: similar or different to younger counterparts? *International Journal of Eating Disorders*, 51 (1). pp. 3-9. ISSN 0276-3478 doi: <https://doi.org/10.1002/eat.22810> Available at <https://centaur.reading.ac.uk/76081/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

Published version at: <http://dx.doi.org/10.1002/eat.22810>

To link to this article DOI: <http://dx.doi.org/10.1002/eat.22810>

Publisher: Wiley

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online



1 Eating pathology in midlife women: similar or different to younger counterparts?

2 Running title: Eating Disorders in Midlife Women

3

4 Paul E. Jenkins*^{1, 2}

5 Tom Price^{1, 3}

6

7 ¹Cotswold House Eating Disorders Service, Oxford Health NHS Foundation Trust,
8 Oxford, United Kingdom.

9 ²School of Psychology, University of Birmingham, Birmingham, United Kingdom.

10 ³Institute of Psychiatry, Psychology and Neuroscience, King's College London,
11 United Kingdom.

12

13 *Present address: School of Psychology and Clinical Language Sciences, University
14 of Reading, RG6 6AL, UK. E-mail: pej106@gmail.com

15

16 Word Count (Abstract): 158

17 Word Count (Main Text): 3438

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22

Abstract

Objective: This study examined potential similarities and differences between women with eating disorders (EDs) in midlife and their younger counterparts.

Method: Seven hundred and three women assessed by a specialist eating disorder service were divided into 3 groups based on age (18 – 25, 25 – 40, and 40+) and compared on a number of clinical and demographic measures. Distribution of ED diagnoses was also examined between groups.

Results: Midlife women were less likely to receive a diagnosis of anorexia nervosa and more likely to receive a diagnosis of binge-eating disorder than their younger counterparts. Duration of illness was longer and age of ED onset later in the midlife group but no differences were seen on measures of global ED pathology, psychosocial impairment, or psychological distress.

Discussion: This study adds to the developing literature concerning EDs in midlife women, although further work is needed to support the findings presented here and to examine profiles of males presenting to treatment centres.

Keywords: middle-age, midlife, diagnosis, age of onset, anorexia nervosa, bulimia nervosa, binge-eating disorder

Acknowledgements

The authors would like to thank Ceri Morgan for assistance in data collection and analysis.

Introduction

1
2 Incidence of eating disorders (EDs) is commonly associated with younger women,
3 perhaps due to the relatively early typical age of onset and the gender disparity
4 frequently seen (e.g., Fairburn & Harrison, 2003). However, studies of women in
5 midlife (a commonly used range is 35 – 55y, although definitions often vary) suggest
6 that around 5% meet diagnostic criteria for an ED, with greater numbers falling within
7 the spectrum of eating disturbance (Mangweth-Matzek et al., 2014; see also
8 Larrañaga, Docet, & García-Mayor, 2012; McGuinness & Taylor, 2016; Marcus et al.,
9 2007; Slof-Op't Landt et al., 2017). Distinct symptoms, such as binge eating and
10 fear of weight gain, have been found to occur in more than 10% of midlife women
11 (e.g., Fairweather-Schmidt, Lee, & Wade, 2015; de Freitas et al., 2008; Marcus et
12 al., 2007) and a number of studies suggest that admission rates of midlife women to
13 ED units have increased markedly in recent decades (e.g., Ackard et al., 2013;
14 Cumella & Kally, 2008).

15 Given these concerning statistics, alongside findings that older age is
16 associated with poorer outcome (e.g., Ackard et al., 2014; Marcus et al., 2007), it is
17 perhaps surprising that there exists little information on the characteristics of midlife
18 women presenting for outpatient ED treatment. Elran-Barak et al. (2015) looked at
19 individuals seeking treatment at one of four sites in the USA, finding that prevalence
20 of bulimia nervosa (BN) was lower in midlife individuals (age ≥ 40 years) than a
21 younger sample (18 – 25 years of age), but that binge-eating disorder (BED) and
22 atypical EDs were more common. Prevalence of anorexia nervosa (AN) did not
23 differ between groups. This study provided important information on diagnostic
24 prevalence, but only reported comparisons between younger and older women
25 within diagnostic groups. Therefore, more information is needed concerning whether

1 midlife women differ from younger groups in degree of ED pathology, for example.
2 Elran-Barak et al. also highlighted the need for more research investigating age of
3 onset and psychological comorbidity, which may have implications for treatment
4 design and provision.

5 Looking specifically at inpatient admissions, contrasting findings were
6 presented by Ackard et al. (2013, Study 1) who found that midlife patients (40 years
7 or older) were more likely to present with BN than their younger counterparts (18 –
8 39y). The same authors also reported on a sample of 164 women presenting for any
9 level of treatment (Study 2), finding few differences between the two groups in terms
10 of self-esteem, depression, anxiety, and eating pathology. However, Ackard and
11 colleagues did find that age of onset was higher in midlife compared to a younger
12 group (21.4y vs. 17.0y) and that duration of illness was also different (26.5y vs. 8.1y)
13 (see also Forman & Davis, 2005). It would be expected, though, that duration of
14 illness would typically be longer in midlife samples seeking treatment as, irrespective
15 of age at onset, midlife women have lived longer than their younger counterparts.

16 It seems, then, that data on whether diagnostic prevalence differs between
17 midlife and younger samples is limited, and at times conflicting. Studies on midlife
18 women, generally considered as those aged over 40 years old (Elran-Barak et al.,
19 2015), have tended to focus on more broadly-defined eating pathology (Slevec &
20 Tiggemann, 2011), with many concerned only with community samples or those
21 presenting for residential treatment. Some data suggest that midlife women report
22 fewer concerns about body shape compared to younger patients but may experience
23 poorer quality of life, and higher levels of psychological distress and interpersonal
24 problems (e.g., Ackard et al., 2014). Other equivocal data concern age of onset,
25 with some studies finding older age of onset (and longer duration of illness) in midlife

1 samples. This contrasts with data suggesting that age of onset in some EDs is
2 decreasing (e.g., Favaro et al., 2009) and might suggest that there is a subgroup of
3 older individuals who develop an ED later in life (i.e., late onset; see Gupta, 1990;
4 Bueno et al., 2014). Such 'late onset' EDs have been variously defined as over 40
5 years old (Kally & Cumella, 2008) or over 25 years old (Bueno et al., 2014),
6 reflecting the view that age of onset is notoriously "difficult to determine with any
7 precision in most cases" (Arcelus, Mitchell, & Wales, 2011, p. 729).

8 Given the small number of studies, it is perhaps unsurprising that some
9 contradictions have arisen. Differences between the studies of Ackard et al. (2013)
10 and Elran-Barak et al. (2015) may have been due to how age groups were split –
11 Ackard et al. compared adults over 40 with those under 40 whereas Elran-Barak et
12 al. looked at three groups (18 – 25, 26 – 39, and 40+), which may lead to
13 contradictions in interpretation, particularly if those aged 18 – 25 present differently
14 to older groups. Furthermore, the upper age was different between the studies and
15 nearly half of those presenting for treatment in the study of Ackard et al. received
16 inpatient treatment, which is unlikely to be representative of many treatment
17 programs in the UK at least (e.g., Robinson, 1993).

18 The lack of information on midlife women led one group of authors to
19 conclude recently that "our understanding of the needs of [middle-aged] women
20 seeking treatment is in its nascence" (Ackard et al., 2013, p. 177). The current study
21 aims to look in detail at individuals in midlife who attend for specialist treatment of an
22 ED. It will present data on the distribution of different ED diagnoses and will
23 compare midlife women with younger women on a number of clinical and
24 demographic measures. Given conflicting methods in existing empirical work, the
25 study will compare three age groups (i.e., 18 – 25, 25 – 40, 40+), using 40 years of

1 age as a common definition of midlife (e.g., Elran-Barak et al., 2015; Kally &
2 Cumella, 2008). The method of dividing the sample into three separate groups was
3 chosen in order to include the full age spectrum, whilst being able to look at
4 differences between clearly-defined age groups that have been previously studied
5 (cf., Elran-Barak et al., 2015). We were also interested in whether age differences
6 existed between those who did and did not attend their initial assessment as this has
7 important implications for accessing treatment. Given limited findings to-date, we
8 hypothesised that BED would be more common in the midlife group. In line with
9 others (Mangweth-Matzek et al., 2014) our clinical experience suggests that midlife
10 women present with similar levels of symptoms and impairment to younger samples,
11 and were interested to see whether this was substantiated in a large, treatment-
12 seeking sample.

13

14

Methods

Participants

16 Participants were referred to an outpatient eating disorders service based in the UK.
17 This service offers specialist, evidence-based treatment to adults with an ED,
18 including guided self-help, cognitive behaviour therapy, and interpersonal
19 psychotherapy. Some individuals referred to the service will subsequently be
20 referred for more intensive treatment (e.g., inpatient care). Between April 2014 and
21 April 2017, 896 outpatient referrals were received, the majority of which came from
22 primary care (n = 641; 74.8%). In line with the study's primary aims, for remaining
23 analyses only those who attended for assessment were included. Men (n = 47;
24 6.3%) were also excluded due to relatively small numbers. Participants were
25 subsequently stratified by age into 3 groups: 18 to 25 years (54.3%; n = 382), 25 to

1 40 years (28.4%, n = 200) and 40 years and above (17.2%, n = 121). The study was
2 approved by the local NHS Quality and Audit Team.

3

4 **Procedures**

5 Individuals met with a qualified clinician (e.g., nurse specialist, psychologist,
6 psychiatrist) who established diagnosis following a semi-structured interview
7 according to criteria for feeding and eating disorders (DSM-5; APA, 2013). Twenty-
8 nine of those who attended assessment (4.1%) did not meet criteria for an ED.
9 Additional information collected included age and self-reported age at onset of ED
10 (although the procedure for identifying the latter was not standardised); duration of
11 illness was calculated as the difference. Of 611 individuals who provided this
12 information, 317 (51.9%) reported having received previous treatment. Weight and
13 height were measured with calibrated scales, used to calculate body mass index
14 (BMI; kg/m²). In advance of this appointment, each individual was sent a self-report
15 questionnaire pack which they were invited to complete although not all individuals
16 did so.

17 The Eating Disorder Examination – Questionnaire (EDE-Q; Fairburn & Beglin,
18 1994) is a 36-item measure assessing behavioural and cognitive symptoms of eating
19 pathology experienced in the last 28 days. The EDE-Q produces a number of
20 behavioural indices (e.g., objective binge eating, self-induced vomiting) as well as
21 four subscales (Restraint, Eating Concern, Shape Concern, Weight Concern).
22 Subscales are rated on a six-point Likert scale, with higher scores indicative of
23 greater symptomatology. A Global score can be calculated from the cognitive items,
24 which provides a general index of ED pathology. Due to strong correlation ($r = 0.88$)
25 between the Shape Concern and Weight Concern subscales, a combined

1 'Weight/Shape Concern' subscale was derived from the mean of all 12 items (e.g.,
2 see Berg, Peterson, Frazier, & Crow, 2012). Cronbach's α in the current sample
3 were as follows: Restraint, 0.82; Eating Concern, 0.77; and Weight/Shape Concern,
4 0.92.

5 The Clinical Impairment Assessment questionnaire (CIA; Bohn & Fairburn,
6 2008) assesses severity of psychosocial impairment attributable to ED symptoms
7 experienced over the last 28 days. The measure has shown good psychometric
8 properties in comparable samples (Jenkins, 2013), and the 16 items are rated on a
9 0–3 scale, with higher scores indicating greater levels of impairment. A cut-off of 16
10 has been suggested to predict ED case status (Bohn et al., 2008). Cronbach's α
11 was 0.92.

12 The Clinical Outcomes in Routine Evaluation-Outcome Measure (CORE-OM;
13 Barkham et al., 2001) is a 34-item scale, which provides a measure of general
14 psychological distress experienced over the previous 7 days. Items are scored from
15 0 to 4 and provide a total score, calculated as the mean of all items multiplied by 10
16 (see Connell et al., 2007). A cut-off of 10 is recommended to indicate clinical
17 significance and its utility in the study of individuals with EDs has been established
18 (Jenkins & Turner, 2014). Cronbach's α was 0.95.

19

20 **Statistical Analyses**

21 χ^2 tests were used to examine differences among groups on categorical variables
22 (e.g., diagnoses, assessment attendance) and continuous data (e.g., age of onset,
23 EDE-Q scores) were analysed with Kruskal-Wallis or Mann-Whitney procedures due
24 to non-normal distribution of data. Significant differences were explored using post

1 hoc Mann-Whitney U tests. Significance level was set at $p < .05$ and all tests were
2 conducted using SPSS Version 22.

3

4

Results

5 Attendance at First Assessment

6 There were no significant differences between those who attended their initial
7 assessment ($n = 750$) and those who did not ($n = 146$) in terms of age (mean (SD) =
8 28.2y (11.1y) vs. 28.6y (11.5y), $U = 52540.500$, $z = -.774$, $p = .439$) or gender
9 (93.7% female vs. 89.7% female, $\chi^2 (1) = 3.047$, $p = .081$).

10

11 Diagnostic Differences

12 As summarised in Table 1, those aged 18 – 25 were more likely to be diagnosed
13 with AN than other groups ($\chi^2 (1) = 8.92$, $p = .003$). Those aged 25 – 40 were
14 equally likely as other groups to be diagnosed with AN ($\chi^2 (1) = 0.24$, $p = .63$), and
15 midlife women were less likely to receive an AN diagnosis ($\chi^2 (1) = 11.28$, $p = .001$).

16 There were no group differences regarding proportions of BN ($\chi^2 (2) = 1.284$, $p =$
17 .53). Those aged 18 – 25 were less likely to be diagnosed with BED than other
18 groups ($\chi^2 (1) = 13.67$, $p = <.001$). Those aged 25 – 40 were equally likely to be
19 diagnosed with BED ($\chi^2 (1) = 0.02$, $p = .90$) as other age groups. Midlife women
20 were more likely to be diagnosed with BED ($\chi^2 (1) = 22.34$, $p = <.001$). There were
21 no group differences regarding proportion of 'other' ED diagnoses ($\chi^2 (2) = 1.451$, $p =$
22 .48).

23

24

INSERT TABLE 1

25

1 **Previous Treatment**

2 Results found no differences between age groups regarding receipt of previous
3 treatment ($\chi^2(2) = 1.69, p = .431$); see Table 2.

4

5 **Markers of Disordered Eating**

6 As shown in Table 2, BMI differed across groups ($H(2) = 17.48, p < .001$), with post
7 hoc analyses finding that the youngest group (18 – 25y) was lower than both other
8 groups, which were statistically equivalent. Duration of illness differed across groups
9 ($H(2) = 207.76, p < .001$), with post hoc analyses finding that all groups differed from
10 each other. Scores on the Global scale of the EDE-Q were equivalent ($H(2) = 3.89,$
11 $p = 0.14$). Weight and Shape Concerns differed across groups ($H(2) = 7.68, p =$
12 0.021), and Mann-Whitney tests found that the youngest group (18 – 25y) had lower
13 scores than the other two groups, which were statistically equivalent.

14

15

INSERT TABLE 2

16

17 **Psychological Distress and Psychosocial Impairment**

18 Scores on the CORE-OM ($H(2) = 3.20, p = 0.202$) and CIA ($H(2) = 2.47, p = 0.292$)
19 were equivalent across groups. Degree of distress and impairment was high within
20 the sample, with 498 of 559 (89.1%) individuals scoring above 10 on the CORE-OM,
21 and 501 of 560 (89.5%) scoring above 16 on the CIA.

22

23 **Age of Onset**

24 Age of onset was different across groups ($H(2) = 10.32, p = .006$), with the youngest
25 group (18 – 25y) reporting a younger age of onset than the other two groups, which

1 were statistically equivalent. Of 307 individuals for whom data were available, only 2
2 (0.7%) were documented as having an ED onset over 40 years of age and 20 (6.5%)
3 over the age of 25. Statistical comparisons between groups were therefore not
4 performed. However, for those with an onset over 25, seven (35.0%) had AN, four
5 (20.0%) BN, and the remainder 'other EDs' (n = 9; 45.0%). There were no
6 individuals with BED with a reported onset after 25 years of age.

7

8

Discussion

9 The current study aimed to describe the characteristics of women in midlife who
10 present for treatment of EDs. Overall, slightly fewer than 20% of individuals
11 assessed for treatment were aged over 40 and, of these, just over one-third
12 presented with atypical or 'other' eating disorders.

13 Comparison with existing studies suggests that the proportion of atypical EDs is
14 similar, at around 30 – 40% (see Ackard et al., 2013; Elran-Barak et al., 2015), but
15 proportions of other diagnoses were more variable. For example, similar numbers of
16 BN were observed between the current study (20.7%) and that of Ackard et al.
17 (2013) (16.7%), with slightly higher rates (29.9%) reported by Elran-Barak et al.
18 (2015). Regarding BED, the results presented here concur with those of Elran-Barak
19 et al. in suggesting that women in midlife are more likely to be diagnosed with BED
20 than younger individuals. Findings regarding AN are more equivocal, partly
21 confounded by the fact that many studies have included inpatient samples, which are
22 likely to report a higher proportion of individuals with AN (e.g., Cumella & Kally,
23 2008). However, the finding reported here that women in midlife are less likely to be
24 diagnosed with AN has been reported previously (Ackard et al., 2014; cf. Elran-
25 Barak et al., 2015).

1 Looking at ED symptoms, results suggest that midlife women present perhaps
2 as more similar than they are different compared to younger groups (see also Perez,
3 Hernandez, Clarke, & Joiner, 2007). Although levels of ED pathology, as measured
4 by the Global scale of the EDE-Q, were similar, the youngest group (18 – 25y)
5 reported lower weight and shape concerns than the older groups, which were
6 equivalent, a finding similar to that of Tiggemann and Stevens (1999) in a sample of
7 180 women aged between 18 – 60. Similar patterns were also evident in BMI and
8 age of onset. The finding that duration of illness is longer (e.g., Cumella & Kally,
9 2008; Forman & Davis, 2005) was supported, suggesting that women in midlife
10 presenting for treatment have lived with the illness for longer than their younger
11 counterparts, which might negatively affect treatment outcome (e.g., Reas,
12 Williamson, Martin, & Zucker, 2000). Although this should not be taken as evidence
13 that older women cannot benefit from treatment, it does suggest that greater efforts
14 should be made to intervene with disordered eating at earlier stages of the illness.
15 The finding of a longer duration of illness, however, is confounded by obvious age
16 differences between groups. Indeed, mean duration of illness (27.4y) of the oldest
17 group exceeded the upper age limit for the youngest group.

18 Many of these findings, however, should be seen in the context of the
19 distribution of ED diagnoses. For example, it is likely that differences in BMI, for
20 example, were strongly related to the higher prevalence of AN in the younger sample
21 and BED in the older samples. However, as BMI increases with age (Deurenberg,
22 Weststrate, & Seidell, 1991), this conclusion cannot be confirmed in the current
23 study. Interestingly, the current study showed that all age groups were equivalent on
24 measures of psychological distress and impairment (cf. Ackard et al., 2014), and

1 scores for the midlife group were in line with other studies of ED samples (e.g.,
2 Jenkins, 2013; Jenkins & Turner, 2014), again suggestive of similarity.

3 Investigating age of onset within EDs was a secondary aim of the study,
4 although this is difficult to determine accurately (Arcelus et al., 2011) and the
5 procedure was not standardised in the current study. Given these caveats, relatively
6 small numbers of 'late onset' cases were seen, with smaller proportions than
7 previous work (e.g., Bueno et al., 2014; Kimura et al., 2007). Of particular note,
8 there were no cases of 'late onset' BED in this large sample, although this has been
9 documented (e.g., Beck, Casper, & Andersen, 1996) and more work is required in
10 this area given what little is known (Bueno et al., 2014). In addition to possibly being
11 related to diagnostic distribution within the different age groups, findings may also
12 reflect different referral, identification, and assessment practices across time periods.
13 Thus, the possibility that cohort effects influenced findings represents a limitation of
14 the study, which nonetheless provides a cross-sectional view of eating pathology
15 and impairment across the age range.

16 The current study adds to the literature in a number of ways. Firstly, it
17 presents information from a large sample of women regarding prevalence of ED
18 diagnoses and symptoms, showing similar figures to existing work (e.g., Ackard et
19 al., 2014). However, it also goes beyond this by including a large sample of women
20 presenting for outpatient treatment and looking at both diagnostic breakdown (e.g.,
21 Elran-Barak et al., 2015) and wider symptoms. The study presents data on three
22 distinct age groups, perhaps revealing interesting similarities amongst those aged
23 over 25. Finally, it reviewed age of onset, although found few individuals reporting
24 illness onset beyond 25 years of age, which warrants further research. Some
25 shortcomings and limitations should be noted. As men were omitted from this study,

1 this perhaps lends support to the idea that “men aged midlife and beyond might be
2 the most stigmatized group to suffer from an eating disorder” (Reas & Stedal, 2015,
3 p. 254). The study was limited by use of retrospective review of routinely-referred
4 patients within an outpatient setting in the UK, particularly as results may differ
5 across settings (e.g., inpatients, or primary care). As sociocultural factors have been
6 associated with presentation of different EDs (e.g., Hoek et al., 1995), further studies
7 might seek to replicate the findings reported here. In addition, a large proportion of
8 individuals indicated that they had previously received treatment for an ED, although
9 this proportion is similar to other tertiary centres in the UK (e.g., Treasure, Schmidt,
10 Troop, Tiller, Todd, & Turnbull, 1996) and clinics in the US (e.g., Crow, Mussell,
11 Peterson, Knopke, & Mitchell, 1999). Differences with previous work might reflect
12 this methodological aspect but could also be related to the diagnostic criteria used
13 (i.e., DSM-IV vs. DSM-5).

14 Results of the study argue for continued attention to reducing barriers to
15 accessing treatment and identifying cases as early as possible (e.g., Reas et al.,
16 2000). A recent study highlighted the importance of adequate healthcare provision
17 with regards to women in midlife (Micali et al., 2017), and thus the present findings
18 serve as a continued reminder that EDs occur across the age range. Further studies
19 might look at whether EDs in older individuals are being correctly identified, and
20 whether current psychological therapies are sufficiently personalised to address the
21 needs of all those presenting for ED treatment. Given the similarities across groups
22 noted here, it likely that many core strategies will remain relevant and effective for
23 women in midlife, but age-related differences may warrant consideration.

24 In summary, this study suggests that midlife women are more similar than
25 different to their younger counterparts, although older women appear more likely to

- 1 present with BED. Degree of global eating pathology, ED-related psychosocial
- 2 impairment, and psychological distress remained similar across the age range.
- 3 Further work in this area is needed, particularly concerning age of onset and
- 4 inclusion of male samples.

References

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23

American Psychiatric Association. (2013). *Diagnostic and statistical manual of mental disorders* (5th ed.). Washington, DC.

Ackard, D. M., Richter, S., Frisch, M. J., Mangham, D., & Cronemeyer, C. L. (2013). Eating disorder treatment among women forty and older: Increases in prevalence over time and comparisons to young adult patients. *Journal of Psychosomatic Research*, *74*(2), 175-178. DOI: 10.1016/j.jpsychores.2012.10.014.

Ackard, D. M., Richter, S., Egan, A., & Cronemeyer, C. (2014). Poor outcome and death among youth, young adults, and midlife adults with eating disorders: an investigation of risk factors by age at assessment. *International Journal of Eating Disorders*, *47*(7), 825-835. DOI: 10.1002/eat.22346.

Arcelus, J., Mitchell, A. J., Wales, J., & Nielsen, S. (2011). Mortality rates in patients with anorexia nervosa and other eating disorders: A meta-analysis of 36 studies. *Archives of General Psychiatry*, *68*(7), 724-731. DOI: 0.1001/archgenpsychiatry.2011.74.

Barkham, M., Margison, F., Leach, C., Lucock, M., Mellor-Clark, J., Evans, C., . . . McGrath, G. (2001). Service profiling and outcomes benchmarking using the CORE-OM: toward practice-based evidence in the psychological therapies. *Journal of Consulting and Clinical Psychology*, *69*(2), 184-196.

Beck, D., Casper, R., & Andersen, A. (1996). Truly late onset of eating disorders: a study of 11 cases averaging 60 years of age at presentation. *International Journal of Eating Disorders*, *20*(4), 389-395.

- 1 Berg, K. C., Peterson, C. B., Frazier, P., & Crow, S. J. (2012). Psychometric
2 evaluation of the Eating Disorder Examination and Eating Disorder
3 Examination-Questionnaire: a systematic review of the literature.
4 *International Journal of Eating Disorders, 45*(3), 428-438. DOI:
5 10.1002/eat.20931.
- 6 Bohn, K., & Fairburn, C.G. (2008). The clinical impairment assessment questionnaire
7 (CIA 3.0). In: Fairburn, C.G. (Ed.), *Cognitive Behaviour Therapy and Eating*
8 *Disorders*. Guilford Press, New York, 315-317.
- 9 Bohn, K., Doll, H. A., Cooper, Z., O'Connor, M., Palmer, R. L., & Fairburn, C. G.
10 (2008). The measurement of impairment due to eating disorder
11 psychopathology. *Behaviour Research and Therapy, 46*(10), 1105-1110.
12 DOI: 10.1016/j.brat.2008.06.012.
- 13 Bueno, B., Krug, I., Bulik, C. M., Jiménez-Murcia, S., Granero, R., Thornton, L., . . .
14 Fernández-Aranda, F. (2014). Late onset eating disorders in Spain: Clinical
15 characteristics and therapeutic implications. *Journal of Clinical Psychology,*
16 *70*(1), 1-17. DOI: 10.1002/jclp.22006.
- 17 Connell, J., Barkham, M., Stiles, W. B., Twigg, E., Singleton, N., Evans, O., Miles, J.
18 N. V. (2007). Distribution of CORE–OM scores in a general population,
19 clinical cut-off points and comparison with the CIS–R. *British Journal of*
20 *Psychiatry, 190*(1), 69-74. DOI: 10.1192/bjp.bp.105.017657.
- 21 Crow, S., Mussell, M. P., Peterson, C., Knopke, A., & Mitchell, J. (1999). Prior
22 treatment received by patients with bulimia nervosa. *International Journal of*

- 1 *Eating Disorders*, 25, 39-44. DOI: 10.1002/(SICI)1098-
2 108X(199901)25:1<39::AID-EAT5>3.0.CO;2-W
- 3 Cumella, E. J., & Kally, Z. (2008). Profile of 50 women with midlife-onset eating
4 disorders. *Eating Disorders*, 16(3), 193-203. DOI:
5 10.1080/10640260802016670
- 6 de Freitas, S. R., Appolinario, J. C., Souza A. d. M., & Sichieri, R. (2008).
7 Prevalence of binge eating and associated factors in a Brazilian probability
8 sample of midlife women. *International Journal of Eating Disorders*, 41(5),
9 471-478. DOI: 10.1002/eat.20530.
- 10 Deurenberg, P., Weststrate, J. A., & Seidell, J. C. (1991). Body mass index as a
11 measure of body fatness: age- and sex-specific prediction formulas. *British*
12 *Journal of Nutrition*, 65(2), 105 – 114.
- 13 Elran-Barak, R., Fitzsimmons-Craft, E. E., Banyamini, Y., Crow, S. J., Peterson, C.
14 B., Hill, L. J., Crosby, R. D., Mitchell, J. E., & Le Grange, D. (2015). Anorexia
15 nervosa, bulimia nervosa, and binge eating disorder in midlife and beyond.
16 *The Journal of Nervous and Mental Disease*, 203(8), 1-8. DOI:
17 10.1097/NMD.0000000000000333.
- 18 Fairburn, C.G., & Beglin, S.J. (1994). Assessment of eating disorders: interview or
19 self-report questionnaire? *International Journal of Eating Disorders*, 16(4),
20 363–370.
- 21 Fairburn, C. G., & Harrison, P. J. (2003). Eating disorders. *The Lancet*, 361(9355),
22 407-416. DOI: 10.1016/s0140-6736(03)12378-1.

- 1 Favaro, A., Caregaro, L., Tenconi, E., Bosello, R., & Santonastaso, P. (2009). Time
2 trends in age at onset of anorexia nervosa and bulimia nervosa. *Journal of*
3 *Clinical Psychiatry, 70*(12), 1715-21. DOI: 10.4088/JCP.09m05176blu.
- 4 Fairweather-Schmidt, A. K., Lee, C., & Wade, T. D. (2015). A longitudinal study of
5 midage women with indicators of disordered eating. *Developmental*
6 *Psychology, 51*(5), 722-729. DOI: 10.1037/dev0000011.
- 7 Forman, M. E., & Davis, W. N. (2005) Characteristics of Middle-Aged Women in
8 Inpatient Treatment for Eating Disorders. *Eating Disorders, 13*(3), 231-243,
9 DOI: 10.1080/10640260590932841.
- 10 Gupta, M. A. (1990). Fear of aging: a precipitating factor in late onset anorexia
11 nervosa. *International Journal of Eating Disorders, 9*(2), 221-224. DOI:
12 10.1002/1098-108X(199003)9:2<221::AID-EAT2260090213>3.0.CO;2-K.
- 13 Hoek, H. W., Bartelds, A. I. M., Bosveld, J. J. F., van der Graaf, Y., Limpens, V. E.
14 L., Maiwald, M., & Spaaij, C. J. K. (1995). Impact of urbanization on detection
15 rates of eating disorders. *The American Journal of Psychiatry, 152*(9), 1272-
16 1278.
- 17 Jenkins, P. E. (2013). Psychometric validation of the Clinical Impairment
18 Assessment in a UK eating disorder service. *Eating Behaviours, 14*(2), 241-
19 243. DOI: 10.1016/j.eatbeh.2012.12.001.
- 20 Jenkins, P. E., & Turner, H. M. (2014). An investigation into the psychometric
21 properties of the CORE-OM in patients with eating disorders. *Counselling*
22 *and Psychotherapy Research, 14*(2), 102-110. DOI:
23 10.1080/14733145.2013.782057.

- 1 Kally, Z., & Cumella, E. J. (2008). 100 midlife women with eating disorders: A
2 phenomenological analysis of etiology. *The Journal of General Psychology*,
3 135(4), 359–377. DOI: 10.3200/GENP.135.4.359-378.
- 4 Kimura, H., Tonoike, T., Muroya, T., Yoshida, K., Ozaki, N. (2007). Age of onset has
5 limited association with body mass index at time of presentation for anorexia
6 nervosa: Comparison of peak-onset and late-onset anorexia nervosa groups.
7 *Psychiatry and Clinical Neuroscience*, 61(6), 646-50.
- 8 Larrañaga, A., Docet, M. F., & García-Mayor, R. V. (2012). High prevalence of
9 eating disorders not otherwise specified in north western Spain: population-
10 based study. *Social Psychiatry and Psychiatric Epidemiology*, 47(10), 1669-
11 73. DOI: 10.1007/s00127-012-0473-1.
- 12 McGuinness, S., & Taylor, J. E. (2016). Understanding body image dissatisfaction
13 and disordered eating in midlife adults. *New Zealand Journal of Psychology*,
14 45(1), 4-12.
- 15 Mangweth-Matzek, B., Hoek, H. W., Rupp, C. I., Lackner-Seifert, K., Frey, N.,
16 Whitworth, A. B., Pope, H. G., Jr., & Kinzl J. (2014). Prevalence of eating
17 disorders in middle-aged women. *International Journal of Eating Disorders*,
18 47(3), 320-324. DOI: 10.1002/eat.22232.
- 19 Marcus, M. D., Bromberger, J. T., Wei, H.-L., Brown, C., & Kravitz, H. M. (2007).
20 Prevalence and selected correlates of eating disorder symptoms among a
21 multiethnic community sample of midlife women. *Annals of Behavioral*
22 *Medicine*, 33(3), 269-277.

- 1 Micali, N., Martini, M. G., Thomas, J. J., Eddy, K. T., Kothari, R., Russell, E., . . .
2 Treasure, J. (2017). Lifetime and 12-month prevalence of eating disorders
3 amongst women in mid-life: a population-based study of diagnoses and risk
4 factors. *BMC Medicine*, *15*, 12. DOI: 10.1186/s12916-016-0766-4
- 5 Perez, M., Hernandez, A., Clarke, A., & Joiner, T. E., Jr. (2007). Analysis of bulimic
6 symptomatology across age and geographic locations. *Eating Behaviors*,
7 *8*(1), 136-142.
- 8 Reas, D. L., & Stedal, K. (2015). Eating disorders in men aged midlife and beyond.
9 *Maturitas*, *81*(2), 248-255. DOI: 0.1016/j.maturitas.2015.03.004.
- 10 Reas, D. L., Williamson, D. A., Martin, C. K., & Zucker, N. L. (2000). Duration of
11 illness predicts outcome for bulimia nervosa: a long-term follow-up study.
12 *International Journal of Eating Disorders*, *27*, 428-434. DOI:
13 10.1002/(SICI)1098-108X(200005)27:4<428::AID-EAT7>3.0.CO;2-Y.
- 14 Robinson, P. (1993). Treatment for eating disorders in the United Kingdom: Part I. A
15 survey of specialist services. *Eating Disorders Review*, *1*(1), 4–9. DOI:
16 10.1002/erv.2400010103.
- 17 Slevec, J., & Tiggemann, M. (2011). Media Exposure, Body Dissatisfaction, and
18 Disordered Eating in Middle-aged Women: A Test of the Sociocultural Model
19 of Disordered Eating. *Psychology of Women Quarterly* *35*(4) 617-627. DOI:
20 10.1177/0361684311420249.
- 21 Slof-Op't Landt, M. C. T., van Furth, E. F., van Beijsterveldt, C. E. M., Bartels, M.,
22 Willemsen, G., de Geus, E. J., Ligthart, L., & Boomsma, D. I. (2017).
23 Prevalence of dieting and fear of weight gain across ages: a community

1 sample from adolescents to the elderly. *International Journal of Public Health*.

2 DOI: 10.1007/s00038-017-0948-7.

3 Tiggemann, M., & Stevens, C. (1999). Weight concern across the life-span:

4 Relationship to self-esteem and feminist identity. *International Journal of*

5 *Eating Disorders*, 26(1), 103 – 106.

6 Treasure, J., Schmidt, U., Troop, N., Tiller, J., Todd, G., & Turnbull, S. (1996).

7 Sequential treatment for bulimia nervosa incorporating a self-care manual.

8 *British Journal of Psychiatry*, 168, 94-98. DOI: 10.1192/bjp.168.1.94

1 Table 1. Distribution of diagnoses across age groups

Diagnoses, n	Age Group			Total Sample (n = 703)	χ^2
	18 – 25y (n = 382)	25 – 40y (n = 200)	40+ (n = 121)		
AN	162 (42.4%)	72 (36.0%)	29 (24.0%)	263 (37.6%)	13.584**
BN	88 (23.0%)	52 (26.0%)	25 (20.7%)	165 (23.5%)	1.284
BED	8 (2.1%)	10 (5.0%)	16 (13.2%)	34 (4.8%)	24.744**
Other	111 (29.1%)	59 (29.5%)	42 (34.7%)	212 (30.2%)	1.451
No ED	13 (3.4%)	7 (3.5%)	9 (7.4%)	29 (4.1%)	4.059

2 Note: **p < .01; AN = anorexia nervosa; BN = bulimia nervosa; BED = binge-eating
3 disorder

1 Table 2. Demographic and clinical characteristics of women with EDs

Demographics, mean (SD)	Age Group			Total Sample	Kruskal-Wallis	
	18 – 25y (n = 382) ^a	25 – 40y (n = 200) ^b	40+ (n = 121) ^c		χ^2	Mann-Whitney
BMI	19.16 (4.11)	21.30 (6.38)	24.13 (10.69)	20.59 (6.56)	17.476**	a < b, a < c, b = c
Age of onset	16.05 (2.91)	18.11 (5.70)	20.16 (10.09)	17.15 (5.35)	10.315**	a < b, a < c, b = c
DOI	4.31 (3.27)	12.97 (6.82)	27.36 (11.16)	9.63 (9.65)	207.756**	a < b < c
EDE-Q Global	3.90 (1.40)	4.15 (1.26)	4.15 (1.31)	4.01 (1.35)	3.822	ns
WSC	4.29 (1.51)	4.59 (1.37)	4.62 (1.40)	4.42 (1.46)	7.483*	a < b, a < c, b = c
CIA	31.40 (11.77)	32.52 (10.39)	33.87 (10.40)	32.11 (11.20)	2.431	ns
CORE-OM	18.73 (7.28)	19.54 (7.14)	20.31 (8.09)	19.21 (7.39)	3.253	ns
Previous treatment, <i>n</i>	178 (51.9%)	82 (48.8%)	57 (57.0%)	317 (51.9%)	1.685	-

2 Note: Numbers differ across groups as not all individuals completed all measures

3 **p* < .05, ***p* < .01; BMI = body mass index; DOI = duration of illness; EDE-Q =

4 Eating Disorder Examination – Questionnaire; WSC = Weight and Shape Concern;

- 1 CIA = Clinical Impairment Assessment questionnaire; CORE-OM = Clinical
- 2 Outcomes in Routine Evaluation-Outcome Measure