

# *Impact of polystyrene microplastics on Daphnia magna mortality and reproduction in relation to food availability*

Article

Published Version

Creative Commons: Attribution 4.0 (CC-BY)

Open access

Aljaibachi, R. and Callaghan, A. ORCID:  
<https://orcid.org/0000-0002-2731-3352> (2018) Impact of polystyrene microplastics on Daphnia magna mortality and reproduction in relation to food availability. PeerJ, 6. e4601. ISSN 2167-8359 doi: 10.7717/peerj.4601 Available at <https://centaur.reading.ac.uk/76166/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

Published version at: <https://peerj.com/>

To link to this article DOI: <http://dx.doi.org/10.7717/peerj.4601>

Publisher: PeerJ Inc

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

[www.reading.ac.uk/centaur](http://www.reading.ac.uk/centaur)

**CentAUR**

Central Archive at the University of Reading

Reading's research outputs online



# Impact of polystyrene microplastics on *Daphnia magna* mortality and reproduction in relation to food availability

Rana Aljaibachi and Amanda Callaghan

Ecology and Evolutionary Biology Section, School of Biological Sciences, University of Reading, Reading, United Kingdom

## ABSTRACT

Microplastics (MPs) in the environment continue to be a growing area of concern in terms of acute and chronic impacts on aquatic life. Whilst increasing numbers of studies are providing important insights into microparticle behaviour and impacts in the marine environment, a paucity of information exists regarding the freshwater environment. This study focusses on the uptake, retention and the impact of 2  $\mu$ m polystyrene MPs in the freshwater cladoceran *Daphnia magna* in relation to food intake (algae *Chlorella vulgaris*), with MP size chosen to approximately match the cell size of the algae. *Daphnia* were exposed to varied concentrations of MPs and algae. When exposed to a single concentration of MPs *Daphnia* almost immediately ate them in large quantities. However, the presence of algae, even at low concentrations, had a significant negative impact on MP uptake that was not in proportion to relative availability. As MP concentrations increased, intake did not if algae were present, even at higher concentrations of MPs. This suggests that *Daphnia* are selectively avoiding eating plastics. Adult *Daphnia* exposed to MPs for 21 days showed mortality after seven days of exposure in all treatments compared to the control. However significant differences were all related to algal concentration rather than to MP concentration. This suggests that where ample food is present, MPs have little effect on adults. There was also no impact on their reproduction. The neonate toxicity test confirmed previous results that mortality and reproduction was linked to availability of food rather than MP concentrations. This would make sense in light of our suggestion that *Daphnia* are selectively avoiding eating microplastics.

Submitted 18 November 2017

Accepted 21 March 2018

Published 18 April 2018

Corresponding author  
Amanda Callaghan,  
a.callaghan@reading.ac.uk

Academic editor  
Chris Vulpe

Additional Information and  
Declarations can be found on  
page 15

DOI 10.7717/peerj.4601

© Copyright  
2018 Aljaibachi and Callaghan

Distributed under  
Creative Commons CC-BY 4.0

OPEN ACCESS

**Subjects** Ecology, Ecotoxicology, Environmental Impacts

**Keywords** Microplastics, Eco-toxicology, *Daphnia magna*, Chronic toxicity, *Chlorella vulgaris*, Polystyrene, Life history

## INTRODUCTION

Plastics are used extensively worldwide since they are cheap, easy to manufacture and have properties that allow them to replace natural products, including wood, stone and glass (Cole et al., 2011). Around 90% of the world's plastics are low- or high-density polyethylene (PE), polypropylene (PP), polyvinyl chloride (PVC), polystyrene (PS) or polyethylene terephthalate (PET) (Andrady & Neal, 2009). Few of these are released into

the environment without some form of modification (additives) that can potentially leach into the water and many have properties that allow the adsorption of hydrophobic pollutants, so continuing their persistence and spread throughout the water (Andrady & Neal, 2009). Enormous amounts of plastic enter aquatic environments from the land, with an estimated 4.8–12.7 million tons of plastic entering oceans per year (Jambeck et al., 2015). Visible, larger plastic fragments pose well known risks to marine life and environments, but there is an increasing awareness of the impact of microplastics (MPs) which are defined as plastics less than 5 mm in size (Moore, 2008; Cole et al., 2011; Wright, Thompson & Galloway, 2013). MPs can be generated from the degradation of larger pieces but many are manufactured specifically, for example, for use in cosmetics products such as facial scrubs (Napper et al., 2015). Ingestion of MPs has been demonstrated for many marine organisms and there is considerable evidence to suggest that they are transferred up between different trophic levels (Eriksson & Burton, 2003). MPs in freshwater ecosystems (Besseling et al., 2017) are less studied and yet an increasingly important environmental issue (Wagner, Engwall & Hollert, 2014; Dris et al., 2015). MPs enter freshwater bodies through land-based sources or wastewater treatment plants in addition to the potential degradation of large plastic particles (Mason et al., 2016). Cosmetic products such as facial scrubs, toothpaste or body wash are a primary source of MPs with up to 100,000 MPs released into wastewater in a single use (Napper et al., 2015). Very recently, environmental lobbying in the UK has resulted in a ban of MPs in personal beauty products which will come into force in 2018 as well as a greater awareness of the issues. However, this does not extend to other sources of microplastic (MP) pollution. These include synthetic fibres in clothing (e.g., fleece) following machine washing (Fossi et al., 2014) as well as resin pellets used in plastics manufacture. A survey of the River Thames (UK) shoreline revealed 1–4 mm MPs at concentrations of 22–297 particles/L (Horton et al., 2016). The most dominant MPs were fibres, but in one site downstream of a storm drain, receiving urban runoff, many of the plastics were derived from thermoplastic road-surface marking paints (Horton et al., 2016). In the US effluent from wastewater treatment plants was shown to contain an average of  $0.05 \pm 0.024$  MP particles/L (Mason et al., 2016) and effluent feeding into lakes generated an average of  $0.79 \pm 0.88$  mg/L to  $1.56 \pm 1.64$  mg/L MPs (Lasee et al., 2017). Likewise, a survey of beach sediments from the subalpine Lake Garda in Italy revealed high concentrations of plastics including polystyrene (45.6%), polyethylene (43.1%) and polypropylene (9.8%) as well particles (9–500  $\mu\text{m}$ ) of polyamide and polyvinylchloride (Imhof et al., 2013a).

Recent investigations have argued that aquatic organisms have a limited ability to distinguish between food and MPs (Von Moos, Burkhardt-Holm & Köhler, 2012; Fossi et al., 2014). For example, the copepod *Corvus typicus* could not differentiate between algae and 20.6  $\mu\text{m}$  MPs (Cole et al., 2013). Similar results were found when *Acartia clausi* and *Calanus pacificus nauplii* were exposed to MPs in the presence of food (Cole et al., 2013). These factors have led to increasing concerns regarding MPs and have contributed to a recent increase in studies on the impact of MPs on marine and freshwater environments.

Studies on freshwater invertebrate species including *Lumbriculus variegatus*, *Daphnia magna*, *Notodromas monacha*, *Potamopyrgus antipodarum*, and *Gammarus pulex*, have

confirmed that ingestion of MPs occurs (Imhof et al., 2013b). Ingestion and elimination of polyamide fibres and polystyrene MPs were demonstrated in the freshwater amphipod *Gammarus fossarum* and impacted on food assimilation (Blarer & Burkhardt-Holm, 2016).

Various studies have begun to look at the impact of MPs on growth and reproduction in the model ecotoxicology organism *Daphnia*. *Daphnia magna* (Cladocera, Crustacea) is a freshwater filter feeder with the ability to uptake and ingest small suspended particles from the water (Ebert, 2005). *Daphnia* usually feed on algae, although they are capable of feeding on bacteria and can consume particles between 1–70 µm in size (Ebert, 2005). *Daphnia* are said to be unable to distinguish between particles size and quality (DeMott, 1986), which implies a lack of selection and likely ingestion of MPs. Tiny carboxylate polystyrene MPs (20 and 1,000 nm) can cross the *Daphnia* gut epithelium and accumulate in lipid storage droplets (Rosenkranz et al., 2009). Bioaccumulation of these nano-polystyrenes is associated with a negative impact on the growth, mortality and reproduction of *D. magna* (Besseling et al., 2014). MP fibres (ground polyethylene terephthalate) from textiles were toxic to unfed *D. magna* but no mortality occurred when animals had been fed (Jemec et al., 2016). Degraded macro plastics were also toxic to *D. magna*, increasing inter-brood period and decreasing reproduction at high concentrations whereas responses to cosmetic MPs found no such effect and effects were restricted to the level of nutrition (Ogonowski et al., 2016). This contrasts with the effect of polyethylene MPs where ingestion of high concentrations of 1 µm MPs led to the immobilisation of *D. magna* (Rehse, Kloas & Zarfl, 2016).

The aim of the present study was to add to the growing body of evidence of the impact of MPs through a thorough investigation of the impact of 2 µm micro-polystyrene on *D. magna* on reproduction, growth and mortality. This size was chosen since it is a similar size to the alga *Chlorella vulgaris* which is used to feed *Daphnia*. This work was undertaken at a point where few had studied the impact of MPs on *Daphnia* and freshwater ecosystems. We hypothesized that *Daphnia* would be unable to differentiate between MPs and algae and that the uptake of both algae and MPs would be equivalent. Given an inability to distinguish between food and non-food, we predicted that the reduction in food ingested would have a significant impact on the fitness of the *D. magna*. Longer term impacts would also be dependent on excretion of MPs as well as general toxicity. We measured the impact on growth and reproduction using standard ecotoxicology 21 days' life history tests following OECD guideline 211 (OECD, 2012).

## MATERIALS AND METHODS

### *Daphnia magna* and *Chlorella vulgaris* culture

*Daphnia magna* were obtained from the Water Research Centre (WRC, Medmenham, UK) and cultured at the University of Reading for more than ten years prior to this experiment. Full details of culturing methods are given in (Hooper et al., 2006). *Daphnia* were maintained in Organization for Economic Co-operation and Development (OECD) reconstituted water (media) and fed yeast and *C. vulgaris* var Viridis following the methods of (Hayashi et al., 2008). New cultures of *Daphnia* were prepared with 15 neonates in 1,200 ml beakers filled with OECD media (the progeny of these neonates are the first brood).

Juveniles were removed regularly from the culture and the media was changed once a week. The third brood produced by the original 15 neonates were used for experiments.

### Preparation of MPs

MPs used for uptake and depuration experiments were supplied by Sigma-Aldrich, Dorset, UK, (Lot no. MKBQ9691: batch no. 1001856699) as 2  $\mu\text{m}$  carboxylate-modified polystyrene, fluorescent yellow-green (excitation 470 nm; emission 505 nm), density 1.050g/cm<sup>3</sup>. MPs were stored as a stock suspension (2.5 mg mL<sup>-1</sup>) in distilled water and mixed using a vortex prior to dilutions. MPs used for toxicity tests were also supplied by Sigma-Aldrich, Dorset, UK, (Lot no. BCBN6954V: batch no.78452) as 2  $\mu\text{m}$  non-fluorescent polystyrene microplastics. These were stored as an aqueous suspension of 10% solids.

### Microplastic uptake

Adult (18 days) *D. magna* were placed individually in glass beakers filled with 50 ml media and starved for 24 h. They were then exposed to  $1.46 \times 10^2$  mg/L 2  $\mu\text{m}$  MPs with and without algae (calculated based on carbon  $1.00 \times 10^{-1}$  mg/L) of *C. vulgaris*. Four replicates of each treatment were exposed for 15, 30, 60, 120 and 240 min in daylight conditions. at  $20 \pm 2$  °C. Following exposure *Daphnia* were washed with deionized water to remove MPs that had adhered to the carapace, dried on tissue and placed in an Eppendorf tube and stored at  $-20$  °C.

### Microplastic depuration

In order to evaluate depuration of MPs, 20 *Daphnia* from both treatments (see previous section) were exposed for one hour then transferred into clean media for 15, 30, 60, 120 and 240 min (replicated four times). *Daphnia* were washed twice with deionized water for one minute before freezing at  $-20$  °C.

Frozen *Daphnia* were placed individually into a 1.5 ml Eppendorf tube filled with 500  $\mu\text{l}$  distilled water and homogenised by crushing the *Daphnia* using a glass Kontes Pellet Pestle (Fisher Sciences, Loughborough, UK) for one minute. A further 500  $\mu\text{l}$  distilled water was added to wash the pestle. A 0.2 ml aliquot of the homogenate was filtered onto a black background nucleopore track-etched membrane (Whatman, Kent, UK)  $<0.2$   $\mu\text{m}$ , by using a glass vacuum filter holder connected to a manual air pump. The membrane was examined under the epi-fluorescent microscope (Zeiss Axioskop) under (20 $\times$ ) to count the fluorescent MPs in each treatment.

### Microplastic visual assessment

Four adult *Daphnia* from each treatment for each experiment were observed under an epi-fluorescent microscope (Carl Zeiss Axioskop, Wetzlar, Germany), at (10 $\times$ ) magnification with the main focus on the gut system. Images were taken through a blue filter (excitation 450–490 nm) to differentiate the MPs from algae which fluoresced under a green filter (excitation 510–560 nm) (Figs. S1–S20).

**Table 1** Concentrations (mg/L) of MPs and algae added to each treatment to study the uptake of microplastics by *Daphnia magna*.

Treatments	Volume μl	Algae concentration (mg/L)	MPs concentrations (mg/L)
Control MP only	50	0	$6.93 \times 10^{-4}$
	100	0	$1.39 \times 10^{-3}$
	200	0	$2.77 \times 10^{-3}$
	400	0	$5.54 \times 10^{-3}$
	600	0	$8.31 \times 10^{-3}$
	800	0	$1.11 \times 10^{-2}$
Algae = MP	50	$5.00 \times 10^{-2}$	$6.93 \times 10^{-4}$
	100	$1.00 \times 10^{-1}$	$1.39 \times 10^{-3}$
	200	$2.00 \times 10^{-1}$	$2.77 \times 10^{-3}$
	400	$4.00 \times 10^{-1}$	$5.54 \times 10^{-3}$
	600	$6.00 \times 10^{-1}$	$8.31 \times 10^{-3}$
	800	$8.00 \times 10^{-1}$	$1.11 \times 10^{-2}$
Algae > MP	50	$5.00 \times 10^{-2}$	$1.39 \times 10^{-3}$
	100	$1.00 \times 10^{-1}$	$1.39 \times 10^{-3}$
	200	$2.00 \times 10^{-1}$	$1.39 \times 10^{-3}$
	400	$4.00 \times 10^{-1}$	$1.39 \times 10^{-3}$
	600	$6.00 \times 10^{-1}$	$1.39 \times 10^{-3}$
	800	$8.00 \times 10^{-1}$	$1.39 \times 10^{-3}$
MP > Algae	50	$1.00 \times 10^{-1}$	$6.93 \times 10^{-4}$
	100	$1.00 \times 10^{-1}$	$1.39 \times 10^{-3}$
	200	$1.00 \times 10^{-1}$	$2.77 \times 10^{-3}$
	400	$1.00 \times 10^{-1}$	$5.54 \times 10^{-3}$
	600	$1.00 \times 10^{-1}$	$8.31 \times 10^{-3}$
	800	$1.00 \times 10^{-1}$	$1.11 \times 10^{-2}$

### Differential microplastic uptake under varying food regime

Adult *D. magna* were exposed individually for 60 min to a different combination of MPs and algae concentrations (Table 1). Each treatment was replicated six times and assigned to a randomized block design (where replicates were randomly situated on the laboratory bench to eliminate the contribution of factors other than treatment as an experimental error). Following exposure, *Daphnia* were washed twice with deionized water for one minute before freezing at  $-20^{\circ}\text{C}$ . MPs were quantified as described earlier.

### Chronic toxicity tests—adults

*Daphnia* (18 days old from the third brood) were placed individually into glass beakers filled with 50 ml media and exposed to one of eight treatments (replicated five times) (Table 2). Media and concentrations were renewed three times per week. In all treatments, life history traits (survival and reproduction) were monitored for 21 days. Neonates were counted daily and removed. Animals unable to swim after gentle stirring for 15 s were considered dead. The experiment was run under laboratory conditions at  $20 \pm 2^{\circ}\text{C}$ , and 16 h light :8 h dark.

**Table 2** Concentrations (mg/L) of MPs and algae added to each treatment to study chronic toxicity in *Daphnia magna*.

Treatments	Algae concentration (mg/L)	MPs (mg/L)
Control low algae	$1.00 \times 10^{-1}$	0
Control high algae	$8.00 \times 10^{-1}$	0
Algae = MP (low)	$1.00 \times 10^{-1}$	$1.39 \times 10^{-3}$
Algae = MP (high)	$8.00 \times 10^{-1}$	$1.11 \times 10^{-2}$
Algae > MP	$8.00 \times 10^{-1}$	$1.39 \times 10^{-3}$
MP > Algae	$1.00 \times 10^{-1}$	$1.11 \times 10^{-2}$

### Chronic toxicity test—neonates

A standard chronic toxicity test was conducted in compliance with OECD guideline 211 (OECD, 2012). Five third brood neonates (<24 h) old were placed in glass beakers, and exposed to different treatments in 50 ml media (Table 2). Media and concentrations were renewed three times a week and life history traits (survival, growth and reproduction) were monitored for 21 days. Body length (the area from the top of the head to the base of the tail spine) was measured under a stereomicroscope every other day. The experiment was run at  $20 \pm 2$  °C, with 16 h light: 8 h dark.

### Statistical methods

Data were checked for normality using the Shapiro–Wilk test, SPSS 21.0 (SPSS Incorp., Chicago, IL, USA). The uptake and depuration experiments were normally distributed data and analysed by ANOVA whereas T-tests were conducted to compare between 15 and 240 min.

For the reproduction experiment the Kolmogorov–Smirnov test showed non-normally distributed data. Data were analysed using the Wald chi-square test, which is based on the linearly independent pairwise comparisons among the estimated marginal means of the offspring. T-tests were used to compare the mean reproduction between treatments with and without MPs.

Growth rate data was normally distributed and analysed by UNIANOVA. A post-hoc pairwise comparison was undertaken with t-tests to measure the effect of MPs on the growth rate.


The mortality tests were normally distributed. Analysis was carried out using Minitab V. 17, general linear model and a probit analysis conducted, and the response curve for concentrations was made using a scatterplot.

## RESULTS

### Uptake of MPs

*Daphnia* that were fed with algae plus MPs ate significantly fewer MPs compared to those with just MPs ( $F_{1,30} = 50.702$ ,  $P < 0.001$ ) (Fig. 1) (Tables S1–S2). *Daphnia* treated with MPs and algae not only ate fewer MPs than *Daphnia* without algae, but there was a significant reduction in bead uptake over time ( $F_{4,19} = 5.771$ ,  $P = 0.005$ ). There were





**Figure 1** Uptake of 2  $\mu\text{m}$  polystyrene MPs by *Daphnia magna* exposed to MPs only ( $1.46 \times 10^2$  mg/L) or MPs with algae ( $1.46 \times 10^2$  mg/L and  $1.00 \times 10^{-1}$  mg/L) over 240 min. Each point represents the mean  $\pm$  the standard error.

Full-size [DOI: 10.7717/peerj.4601/fig-1](https://doi.org/10.7717/peerj.4601/fig-1)


significantly fewer MPs in *Daphnia* fed algae after 240 min compared to the number after 15 min'  $t$  (d.f. 6) = 2.5,  $P = 0.042$ . Although *Daphnia* without algae ingested significantly more MPs over time than those fed algae, ingestion over time did not increase significantly ( $F_{4,19} = 1.244$ ,  $P = 0.335$ ). The number of MPs ingested after 15 min did not significantly vary from the number of MPs ingested after 240 min  $t$  (d.f. 6) = -1.2,  $P = 0.27$  (See Figs. S1–S10).

## Depuration of MPs

As before, *Daphnia* fed algae contained significantly fewer MPs compared to those without algae ( $F_{1,30} = 976.162$ ,  $P < 0.001$ ) (Fig. 2) (Tables S3–S4). Bead excretion in *Daphnia* fed algae did not vary over time ( $F_{4,19} = 1.006$ ,  $P = 0.435$ ). There were no significant differences in bead counts after 240 min compared to the number after 15 min  $t$  (d.f.6) = 0.978,  $P = 0.366$ . However, the unfed *Daphnia* excreted a significant number of MPs over time compared to fed *Daphnia* ( $F_{4,19} = 5.452$ ,  $P = 0.006$ ). There were significantly fewer MPs in *Daphnia* after 240 min compared to the number after 15 min  $t$  (d.f.6) = 3.5,  $P = 0.013$  (Figs. S11–S20).


## Differential microplastic uptake under varying food regime

Four treatment regimes, which varied in the amount of either MPs, algae or both MPs and algae resulted in significant differences in the amount of MPs ingested ( $F_{3,120} = 114.899$ ,  $P < 0.001$ ) (Fig. 3). In unfed *Daphnia* treated with MPs (MPX), the internal bead count increased significantly as the concentration of MPs increased ( $F_{5,120} = 12.849$ ,  $P < 0.001$ ). Where a low amount of food was introduced (MP>Algae) ingested MPs dropped significantly ( $F_{3,120} = 20.788$ ,  $p < 0.001$ ) but did not change with increasing



**Figure 2** Excretion of 2  $\mu\text{m}$  polystyrene MPs from the gut of *Daphnia magna* exposed to MPs only ( $1.46 \times 10^2$  mg/L) or MPs with algae ( $1.46 \times 10^2$  mg/L and  $1.00 \times 10^{-1}$  mg/L) over 240 min. Each point represents the mean  $\pm$  the standard error.


Full-size [DOI: 10.7717/peerj.4601/fig-2](https://doi.org/10.7717/peerj.4601/fig-2)



**Figure 3** Uptake of 2  $\mu\text{m}$  polystyrene MPs by *Daphnia magna* with and without algae in various volumes ( $\mu\text{l}$ ) (see Table 1 for actual concentrations). Each point represents the mean  $\pm$  the standard error.

Full-size [DOI: 10.7717/peerj.4601/fig-3](https://doi.org/10.7717/peerj.4601/fig-3)

concentrations of MPs ( $F_{5,120} = 1.207$ ,  $P < 0.310$ ). An equal dose of MPs and algae (MP = Algae) was likewise significantly lower than MPX and did not increase as both concentrations increased ( $F_{5,120} = 1.131$ ,  $P < 0.348$ ). Where *Daphnia* were treated with a fixed low concentration of MPs and algal concentrations increased (Algae > MP), the number of MPs decreased significantly ( $F_{5,120} = 4.242$ ,  $P < 0.001$ ).



**Figure 4** Mortality of *Daphnia magna* expressed as a function of time after chronic exposure to MPs under high and low food conditions for 21 days. Asterisks denote overlap between two treatments.

Full-size [DOI: 10.7717/peerj.4601/fig-4](https://doi.org/10.7717/peerj.4601/fig-4)

**Table 3** Mean  $\pm$  standard error (S.E.) of lethal time (LT10, LT50 and LT90) of adult *Daphnia magna* exposed to different concentrations (mg/L) of MPs and algae.


Treatments	Concentrations		Lethal time %		
	Algae mg/L	MPs mg/L	LT <sub>10</sub> $\pm$ SE	LT <sub>50</sub> $\pm$ SE	LT <sub>90</sub> $\pm$ SE
Algae (low)	$1.00 \times 10^{-1}$	–	$7.33 \pm 0.825$	$12.47 \pm 0.69$	$17.6 \pm 0.85$
Algae (high)	$8.00 \times 10^{-1}$	–	$39.4 \pm 0$	$44.5 \pm 0$	$49.9 \pm 50$
Algae = MP (low)	$1.00 \times 10^{-1}$	$1.39 \times 10^{-3}$	$11.22 \pm 0.79$	$16.36 \pm 0.69$	$21.5 \pm 0.87$
Algae = MP (high)	$8.00 \times 10^{-1}$	$1.11 \times 10^{-2}$	$12.35 \pm 0.76$	$17.48 \pm 0.70$	$22.6 \pm 0.92$
Algae > MP	$8.00 \times 10^{-1}$	$1.39 \times 10^{-3}$	$39.40 \pm 0$	$44.54 \pm 0$	$49.68 \pm 0$
MP > Algae	$1.00 \times 10^{-1}$	$1.11 \times 10^{-2}$	$4.95 \pm 0.85$	$10.09 \pm 0.70$	$15.23 \pm 0.83$

### Mortality test—adults

Mortality rates varied significantly between treatments ( $F_{5,618} = 43.38$ ,  $P < 0.001$ ) (Fig. 4). All treatments had significant mortality compared to the control (algae  $8.00 \times 10^{-1}$  mg/L) ( $F_{5,618} = 48.97$ ,  $P < 0.001$ ). A pairwise comparison of mortality in *Daphnia* with restricted food ( $1.00 \times 10^{-1}$  mg/L algae) plus or minus  $1.39 \times 10^{-3}$  mg/L MPs was significant  $t$  (d.f. 1) =  $-3.99$ ,  $P < 0.001$  and with ample food ( $8.00 \times 10^{-1}$  mg/L algae) plus or minus  $1.11 \times 10^{-2}$  mg/L MPs was highly significant  $t$  (d.f. 1) =  $4.59$ ,  $P < 0.001$  since there was no mortality in the treatment without MPs (Table 3).

### Reproduction test—adults

A 21 day reproduction test on adult *Daphnia* resulted in no significant differences in the mean number of offspring between treatments (Fig. 5) ( $X^2(5, N = 30) = 4.62$ ,  $P = 0.463$ ). There was no significant effect on the mean reproduction between treatments with low algal concentration ( $1.00 \times 10^{-1}$  mg/L) and those with low algal concentrations with the addition of  $1.39 \times 10^{-3}$  mg/L of MPs  $t$  (d.f. 8) =  $0.971$ ,  $P = 0.36$ . Similarly, there were no significant differences in mean reproduction between *Daphnia* treated with high algal



**Figure 5** Effects of combinations of high and low MPs and algae concentrations on the mean number of offspring on *Daphnia magna*. Error bars indicate  $\pm$  95% confidence intervals and asterisks denote significant differences compared to the control  $p < 0.001$ .

Full-size [DOI: 10.7717/peerj.4601/fig-5](https://doi.org/10.7717/peerj.4601/fig-5)


concentrations ( $8.00 \times 10^{-1}$  mg/L) and those with high algal concentrations with the addition of  $1.11 \times 10^{-2}$  mg/L of MPs  $t$  (d.f. 8) = 0.067,  $P = 0.948$ .

### Mortality rate—neonate

In neonates, there was a significant difference in the percentage mortality between treatments ( $F_{5,618} = 26.86$ ,  $P < 0.001$ ) (Fig. 6). The percentage mortality of *Daphnia* fed low algal concentrations and MPs was significantly higher than that of those treated with the same algal concentrations but no MPs  $t$  (d.f. 5) = 4.24,  $P < 0.001$ . In contrast, MPs had no impact on mortality when fed to *Daphnia* on a high algal concentration  $t$  (d.f. 5) = 4.51,  $P = 0.776$ . Calculated lethal times (LT<sub>10</sub>, LT<sub>50</sub> and LT<sub>90</sub>) are presented in Table 4.

### Reproduction test—neonate

The reproduction rate was significantly different between adults that had been treated since they were neonates with high and low algae concentrations (Fig. 7) ( $X^2(5, N = 28) = 618$ ,  $P > 0.001$ ), this was because all treatments that included ( $8.00 \times 10^{-1}$  mg/L) algae had significantly higher reproduction compared to those on ( $1.00 \times 10^{-1}$  mg/L). There were no significant differences in reproduction between identical food regimes with and without MPs (low algae concentration ( $1.00 \times 10^{-1}$  mg/L) in the presence of MPs ( $1.39 \times 10^{-3}$



**Figure 6** Mortality of neonate *Daphnia magna* after exposure to different treatments of MPs and algae over 21 days. Asterisks denote overlap between two treatments.

Full-size [DOI: 10.7717/peerj.4601/fig-6](https://doi.org/10.7717/peerj.4601/fig-6)

**Table 4** Mean  $\pm$  standard error of lethal time (LT10, LT50 and LT90) of neonate *Daphnia magna* exposed to different concentrations (mg/L) of MPs and algae.

Treatments	Concentrations		Lethal time %		
	Algae mg/L	MPs mg/L	LT10 $\pm$ SE	LT50 $\pm$ SE	LT90 $\pm$ SE
Algae (low)	$1.00 \times 10^{-1}$	–	$7.71 \pm 0.39$	$14.23 \pm 0.35$	$20.75 \pm 0.42$
Algae (high)	$8.00 \times 10^{-1}$	–	$9.02 \pm 0.4$	$15.54 \pm 0.36$	$22.06 \pm 0.44$
Algae = MP (low)	$1.00 \times 10^{-1}$	$1.39 \times 10^{-3}$	$4.98 \pm 0.41$	$11.51 \pm 0.35$	$18.03 \pm 0.41$
Algae = MP (high)	$8.00 \times 10^{-1}$	$1.11 \times 10^{-2}$	$9.03 \pm 0.4$	$15.55 \pm 0.36$	$22.07 \pm 0.44$
Algae > MP	$8.00 \times 10^{-1}$	$1.39 \times 10^{-3}$	$13.07 \pm 0.42$	$19.59 \pm 0.42$	$26.11 \pm 0.52$
MP > Algae	$1.00 \times 10^{-1}$	$1.11 \times 10^{-2}$	$5.57 \pm 0.4$	$12.09 \pm 0.34$	$18.61 \pm 0.4$


mg/L)  $t$  (d.f. 7) = 1.63,  $p$  = 0.146, high algal concentration ( $8.00 \times 10^{-1}$  mg/L) in addition to MPs ( $1.11 \times 10^{-2}$  mg/L)  $t$  (d.f. 8) = 0.46,  $P$  = 0.652).

## Growth rate

*Daphnia* body length increased significantly in all treatments over 21 days compared to the initial size ( $F_{5,274} = 166.8$ ,  $P < 0.001$ ) (Fig. 8). *Daphnia* growth was lower in animals exposed to low algae concentrations ( $1.00 \times 10^{-1}$  mg/L) compared to those given high algae concentrations ( $8.00 \times 10^{-1}$  mg/L)  $t$  (d.f. 278) =  $-14.5$ ,  $P < 0.001$ . MPs significantly reduced growth in both algal food regimes ( $1.00 \times 10^{-1}$  mg/L,  $F_{2,126} = 3.009$ ,  $P = 0.05$ ); ( $8.00 \times 10^{-1}$  mg/L,  $F_{2,149} = 0.63$ ,  $P = 0.05$ ).

## DISCUSSION


Previous studies have suggested that *D. magna* find it hard to distinguish between MP and food particles in the media (Wiedner & Vareschi, 1995). This study was therefore designed to look at the interaction between MP ingestion and food intake, with MP size chosen to approximately match the cell size of the algae. Algae concentrations were chosen based on



**Figure 7** *Daphnia magna* reproduction (neonate production) after 21 days' exposures to a range of MP and algae treatments (algae (low), algae (high), Algae = MP(low), Algae = MP(high), Algae > MP, MP > Algae). Error bars indicate  $\pm$  95% confidence intervals and asterisks denote significant differences compared to the control  $p < 0.001$ .

Full-size [DOI: 10.7717/peerj.4601/fig-7](https://doi.org/10.7717/peerj.4601/fig-7)

the minimum and maximum normal daily feeding of *Daphnia* (Hooper *et al.*, 2006), while MPs concentrations were chosen to be close to the algae concentrations volume/volume. When exposed to a single concentration of MPs *Daphnia* almost immediately ate them in large quantities. Previous studies have demonstrated that *Daphnia* will feed on polystyrene, beating their appendages at a constant rate regardless of the food concentration (Pavlaki *et al.*, 2014). The same study also found that plastics of different sizes (1.1  $\mu\text{m}$  and 5.7  $\mu\text{m}$ ) were ingested in proportion to their concentration (Pavlaki *et al.*, 2014). This could explain the fact that the amount eaten was reduced when food (algae) was present, which suggests a simple competition for uptake. This hypothesis was further tested by mixing concentrations of algae and MPs. Here there was a direct correlation between MP uptake and concentration so that *Daphnia* ate more MPs as more were available. However, the presence of algae, even at low concentrations, had a significant negative impact on MP uptake. Although MP concentrations increased, intake did not if algae were present, even at higher concentrations of MPs. A high concentration of algae significantly reduced the uptake of a low concentration of MPs so that they were not ingested in direct proportion. Therefore, unlike the study looking at the uptake of different sized polystyrene, there is no evidence that MPs are eaten in proportion to their concentration when algae are present. This implies selectivity (Pavlaki *et al.*, 2014). Previous studies have demonstrated that



**Figure 8** Effect of 21 days' exposure to different combinations of MPs and algae (Algae (low), algae (high), Algae = MP(low), Algae = MP(high), Algae > MP, MP > Algae) on body length of *Daphnia magna*. Each point represents the mean of five replicates  $\pm$  standard error.

Full-size [DOI: 10.7717/peerj.4601/fig-8](https://doi.org/10.7717/peerj.4601/fig-8)

*Daphnia* are not blindly filtering particles from the water, since particles smaller than the mesh size of their appendages are still eaten (Pavlaki et al., 2014). The uptake of particles is strongly influenced by surface chemistry (negative charges increase uptake) and wettability (which can be reduced by the presence of surfactants to decrease uptake) (Pavlaki et al., 2014). There is also some evidence that *Daphnia* can discriminate artificial from natural particles by taste for instance selecting to feed on phytoplankton rather than clay particles (DeMott, 1986). A previous study noted that MPs tended to form aggregates with algae, thereby effectively increasing the food particle size, with a reduction in MP ingestion when algae were present (Long et al., 2015). There was no evidence of aggregation in any of the experiments presented here pre-feeding.

Where no food was offered, the number of MPs in the *Daphnia* gut remained relatively stable with no significant difference between the number in the gut at the beginning and the end of the experiment. However, there was a significant reduction in MPs over time when food was present.

The excretion of particles was further investigated by measuring the depuration of MPs following a 1 h exposure to MPs followed by clean OECD water for 15, 30, 60, 120 and 240 min. The results showed that the number of MPs in the *Daphnia* decreased over time, suggesting excretion. This has been shown before, with small MPs clearing more rapidly than larger ones (Rosenkranz et al., 2009). However, in the individuals exposed to MPs and food particles, the number of MPs in the gut hardly decreased over time. It is not clear why this is the case and it could be that MPs and food aggregate in the gut, making it hard

to excrete MPs. Additionally, it could be that *Daphnia* is re-ingesting the plastic particles after excretion into the media.

*Daphnia* exposed to MPs for 21 days showed mortality after seven days of exposure in all treatments compared to the controls. However, treatments with high MP concentrations and low food levels had the highest mortality compared to the control with low food concentrations. Treatments with high MPs and ample food showed lower mortality compared to the control with low food concentrations. This suggests that where ample food is present, MPs have little effect on adults.

There was also no impact on their reproduction. Again, food levels were more important than MPs concentrations. Similar results had been found recently by [Ogonowski et al. \(2016\)](#), when exposed *Daphnia* to primary MPs or kaolin with low and high food concentrations, results showed life history traits more related to food concentration rather than MPs.

The neonate toxicity test confirmed previous results that mortality was linked to availability of food rather than MP concentrations. *Daphnia* exposed to high food concentrations show higher survival despite MP treatment and reproduction was dramatically decreased in treatments with low food concentrations. Only in the treatment with low food concentrations and low doses of MPs could we see a potential impact of MP with no reproduction. This issue had been investigated previously by exposing the *Daphnia* to metal and different food concentrations which shows that chronic toxicity is linked more to food availability rather than metals; however, that metal get more toxic when there is low food level ([Pavlaki et al., 2014](#)).

Growth rate was more effected by food concentrations rather than MPs, *Daphnia* treated with high food concentrations grow are twice as large as those with low food availability. Similar results were obtained elsewhere with both *Daphnia* and marine isopods *Idotea emarginata* ([Hamer et al., 2014](#); [Ogonowski et al., 2016](#)). This is in contrast to work published on the effect of ~70 nm nano-polystyrenes (Nano-PS) on both *Daphnia* and algae (*Scenedesmus obliquus*) ([Besseling et al., 2014](#)). Nano-PS negatively impacted population growth and reduced chlorophyll concentrations in the algae. *Daphnia* exposed to Nano-PS had reduced body sizes and the numbers and body size of neonates were lower. Nano-PS also caused high numbers of neonate malformations. However the difference here, apart from the size, is that the authors didn't use pristine polystyrene but aged the nano-PS with the algae for 5 days ([Besseling et al., 2014](#)). Their comparison between aged and pristine nano-PS demonstrated that pristine MPs may not represent the full impact of the exposure.

## CONCLUSIONS

Our research was designed to determine the effect of 2  $\mu\text{m}$  MPs on *Daphnia magna* in the presence of algae *Chlorella vulgaris*. This was an experimental approach and was not intended to reflect environmental concentrations of MPs. There is no accurate measurement of 2  $\mu\text{m}$  MPs in freshwater environment and this particular size of plastic can be generated from either a primary source such as cosmetics, or from the degradation of large plastic particles ([Connors, Dyer & Belanger, 2017](#)).



The uptake of MPs decreased in the presence of algae and excretion of MPs reduced. The concentration of MPs ingested did not increase with concentration when algae were available which indicates that the *Daphnia* is selectively eating the algae rather than MPs. Chronic toxicity tests (mortality and reproduction rate) found no toxic effect after a 96 h of exposure although seven days of exposure to high concentration of MPs increased mortality. Life history traits of neonates (mortality, reproduction and growth rate) was mainly linked to food concentrations rather than MPs that could confirm *Daphnia* select the food particles rather than MPs.

The study presented here was undertaken to look at the impact of the MPs themselves and as such our results have been obtained with clean MPs that have not been exposed to any contaminants. Environmental MPs are likely to mix with other contaminants, some of which could bind to them and alter their toxicity. Therefore, a future direction of research should include investigating the toxicity of MPs collected from the aquatic environment or in mixtures with known freshwater pollutants such as pesticides.

## ADDITIONAL INFORMATION AND DECLARATIONS

### Funding

The authors received no funding for this work.

### Competing Interests

The authors declare there are no competing interests.

### Author Contributions

- Rana Aljaibachi conceived and designed the experiments, performed the experiments, analyzed the data, contributed reagents/materials/analysis tools, prepared figures and/or tables, authored or reviewed drafts of the paper, approved the final draft.
- Amanda Callaghan conceived and designed the experiments, authored or reviewed drafts of the paper, approved the final draft.

### Data Availability

The following information was supplied regarding data availability:

The raw data are provided in the [Supplemental File](#).

### Supplemental Information

Supplemental information for this article can be found online at <http://dx.doi.org/10.7717/peerj.4601#supplemental-information>.

## REFERENCES

- Andrady AL, Neal MA. 2009. Applications and societal benefits of plastics. *Philosophical Transactions of the Royal Society B: Biological sciences* **364**:1977–1984 DOI [10.1098/rstb.2008.0304](https://doi.org/10.1098/rstb.2008.0304).

- Besseling E, Quik JTK, Sun M, Koelmans AA. 2017. Fate of nano- and microplastic in freshwater systems: a modeling study. *Environmental Pollution* 220:540–548 DOI 10.1016/j.envpol.2016.10.001.
- Besseling E, Wang B, Lürling M, Koelmans AA. 2014. Nanoplastic affects growth of *S. obliquus* and reproduction of *D. magna*. *Environmental Science & Technology* 48:12336–12343 DOI 10.1021/es503001d.
- Blarer P, Burkhardt-Holm P. 2016. Microplastics affect assimilation efficiency in the freshwater amphipod *Gammarus fossarum*. *Environmental Science and Pollution Research* 23:23522–23532 DOI 10.1007/s11356-016-7584-2.
- Cole M, Lindeque P, Fileman E, Halsband C, Goodhead R, Moger J, Galloway TS. 2013. Microplastic ingestion by zooplankton. *Environmental Science and Technology* 47:6646–6655 DOI 10.1021/es400663f.
- Cole M, Lindeque P, Halsband C, Galloway TS. 2011. Microplastics as contaminants in the marine environment: a review. *Marine Pollution Bulletin* 62:2588–2597 DOI 10.1016/j.marpolbul.2011.09.025.
- Connors KA, Dyer SD, Belanger SE. 2017. Advancing the quality of environmental microplastic research. *Environmental Toxicology and Chemistry* 36:1697–1703 DOI 10.1002/etc.3829.
- DeMott WR. 1986. The role of taste in food selection by freshwater zooplankton. *Oecologia* 69:334–340 DOI 10.1007/BF00377053.
- Dris R, Imhof H, Sanchez W, Gasperi J, Galgani F, Tassin B, Laforsch C. 2015. Beyond the ocean: contamination of freshwater ecosystems with (micro-) plastic particles. *Environmental Chemistry* 12:539–550 DOI 10.1071/EN14172.
- Ebert D. 2005. *Ecology, epidemiology, and evolution of parasitism in Daphnia*. US: National Center for Biotechnology Information.
- Eriksson C, Burton H. 2003. Origins and biological accumulation of small plastic particles in fur seals from Macquarie Island. *AMBIO: A Journal of the Human Environment* 32:380–384 DOI 10.1579/0044-7447-32.6.380.
- Fossi MC, Coppola D, Bains M, Giannetti M, Guerranti C, Marsili L, Panti C, De Sabata E, Clò S. 2014. Large filter feeding marine organisms as indicators of microplastic in the pelagic environment: the case studies of the Mediterranean basking shark (*Cetorhinus maximus*) and fin whale (*Balaenoptera physalus*). *Marine Environmental Research* 100:17–24 DOI 10.1016/j.marenvres.2014.02.002.
- Hamer J, Gutow L, Kohler A, Saborowski R. 2014. Fate of microplastics in the marine isopod *Idotea emarginata*. *Environmental Science & Technology* 48:13451–13458.
- Hayashi Y, Heckmann LH, Callaghan A, Sibly RM. 2008. Reproduction recovery of the crustacean *Daphnia magna* after chronic exposure to ibuprofen. *Ecotoxicology* 17:246–251 DOI 10.1007/s10646-008-0191-3.
- Hooper HL, Connon R, Callaghan A, Maund SJ, Liess M, Duquesne S, Hutchinson TH, Moggs J, Sibly RM. 2006. The use of image analysis to estimate population growth rate in *Daphnia magna*. *Journal of Applied Ecology* 43:828–834 DOI 10.1111/j.1365-2664.2006.01180.x.

- Horton AA, Svendsen C, Williams RJ, Spurgeon DJ, Lahive E. 2016. Large microplastic particles in sediments of tributaries of the River Thames, UK—Abundance, sources and methods for effective quantification. *Marine Pollution Bulletin* 114:218–226 DOI 10.1016/j.marpolbul.2016.09.004.
- Imhof HK, Ivleva NP, Schmid J, Niessner R, Laforsch C. 2013a. Contamination of beach sediments of a subalpine lake with microplastic particles. *Current Biology* 23:1–15 DOI 10.1016/j.cub.2013.09.001.
- Imhof HK, Ivleva NP, Schmid J, Niessner R, Laforsch C. 2013b. Contamination of beach sediments of a subalpine lake with microplastic particles. *Current Biology* 23:R867–R868 DOI 10.1016/j.cub.2013.09.001.
- Jambeck JR, Geyer R, Wilcox C, Siegler TR, Perryman M, Andrady A, Narayan R, Lavender K. 2015. Plastic waste inputs from land into the ocean. *Science* 347:768–770 DOI 10.1017/CBO9781107415386.010.
- Jemec A, Horvat P, Kunej U, Bele M, Kržan A. 2016. Uptake and effects of microplastic textile fibers on freshwater crustacean *Daphnia magna*. *Environmental Pollution* 219:201–209 DOI 10.1016/j.envpol.2016.10.037.
- Lasee S, Mauricio J, Thompson WA, Karnjanapiboonwong A, Kasumba J, Subbiah S, Morse AN, Anderson TA. 2017. Microplastics in a freshwater environment receiving treated wastewater effluent. *Integrated Environmental Assessment and Management* 13:528–532 DOI 10.1002/ieam.1915.
- Long M, Moriceau B, Gallinari M, Lambert C, Huet A, Raffray J, Soudant P. 2015. Interactions between microplastics and phytoplankton aggregates: impact on their respective fates. *Marine Chemistry* 175:39–46 DOI 10.1016/j.marchem.2015.04.003.
- Mason SA, Garneau D, Sutton R, Chu Y, Ehmman K, Barnes J, Fink P, Papazissimos D, Rogers DL. 2016. Microplastic pollution is widely detected in US municipal wastewater treatment plant effluent. *Environmental Pollution* 218:1045–1054 DOI 10.1016/j.envpol.2016.08.056.
- Moore CJ. 2008. Synthetic polymers in the marine environment: a rapidly increasing, long-term threat. *Environmental Research* 108:131–139 DOI 10.1016/j.envres.2008.07.025.
- Napper IE, Bakir A, Rowland SJ, Thompson RC. 2015. Characterisation, quantity and sorptive properties of microplastics extracted from cosmetics. *Marine Pollution Bulletin* 99:178–185 DOI 10.1016/j.marpolbul.2015.07.029.
- Ogonowski M, Schür C, Å Jarsén, Gorokhova E. 2016. The effects of natural and anthropogenic microparticles on individual fitness in *Daphnia magna*. *PLOS ONE* 11:e0155063 DOI 10.1371/journal.pone.0155063.
- Organisation for Economic Co-operation and Development (OECD). 2012. *Daphnia magna* reproduction test. In: *OECD guidelines for the testing of chemicals—effects on biotic systems: section 2*, 23 DOI 10.1787/9789264070127-en.
- Pavlaki MD, Ferreira ALG, Soares AMVM, Loureiro S. 2014. Changes of chemical chronic toxicity to *Daphnia magna* under different food regimes. *Ecotoxicology and Environmental Safety* 109:48–55 DOI 10.1016/j.ecoenv.2014.07.039.

- Rehse S, Kloas W, Zarfl C. 2016.** Short-term exposure with high concentrations of pristine microplastic particles leads to immobilisation of *Daphnia magna*. *Chemosphere* **153**:91–99 DOI [10.1016/j.chemosphere.2016.02.133](https://doi.org/10.1016/j.chemosphere.2016.02.133).
- Rosenkranz PHR, Chaudhry QAC, Stone VIS, Ernandes TEFF. 2009.** A comparison of nanoparticle and fine particle uptake by *Daphnia magna*. *Environmental Toxicology and Chemistry* **28**:2142–2149 DOI [10.1897/08-559.1](https://doi.org/10.1897/08-559.1).
- Von Moos N, Burkhardt-Holm P, Köhler A. 2012.** Uptake and effects of microplastics on cells and tissue of the blue mussel *Mytilus edulis* L. after an experimental exposure. *Environmental Science & Technology* **46**:11327–11335 DOI [10.1021/es302332w](https://doi.org/10.1021/es302332w).
- Wagner M, Engwall M, Hollert H. 2014.** Editorial: (Micro)Plastics and the environment. *Environmental Sciences Europe* **26**:16 DOI [10.1186/s12302-014-0016-3](https://doi.org/10.1186/s12302-014-0016-3).
- Wiedner C, Vareschi E. 1995.** Evaluation of a fluorescent microparticle technique for measuring filtering rates of *Daphnia*. *Hydrobiologia* **302**:89–96 DOI [10.1007/BF00027033](https://doi.org/10.1007/BF00027033).
- Wright SL, Thompson RC, Galloway TS. 2013.** The physical impacts of microplastics on marine organisms: a review. *Environmental pollution* **178**:483–92 DOI [10.1016/j.envpol.2013.02.031](https://doi.org/10.1016/j.envpol.2013.02.031).