

Carbon-13 cross-polarization magic-angle spinning nuclear magnetic resonance (CPMAS NMR) for measuring proanthocyanidin content and procyanidin to prodelphinidin ratio in sainfoin (Onobrychis viciifolia) tissues

Article

Accepted Version

Fryganas, C., Drake, C., Ropiak, H. M., Mora Ortiz, M., Smith, L. M. J., Mueller-Harvey, I. and Kowalczyk, R. (2018) Carbon-13 cross-polarization magic-angle spinning nuclear magnetic resonance (CPMAS NMR) for measuring proanthocyanidin content and procyanidin to prodelphinidin ratio in sainfoin (*Onobrychis viciifolia*) tissues. *Journal of Agricultural and Food Chemistry*, 66 (16). pp. 4073-4081. ISSN 0021-8561 doi: <https://doi.org/10.1021/acs.jafc.8b01215> Available at <https://centaur.reading.ac.uk/76550/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

To link to this article DOI: <http://dx.doi.org/10.1021/acs.jafc.8b01215>

Publisher: American Chemical Society

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online

1 **Carbon-13 Cross-Polarization Magic-Angle Spinning**
2 **Nuclear Magnetic Resonance (CPMAS NMR) for**
3 **Measuring Proanthocyanidin Content and Procyanidin to**
4 **Prodelphinidin Ratio in Sainfoin (*Onobrychis viciifolia*)**
5 **Tissues**

6

7 Christos Fryganas,^{†,‡,*} Christopher Drake,[†] Honorata M. Ropiak,^{†,§} Marina Mora-Ortiz,[#]8 Lydia M. J. Smith,[⊥] Irene Mueller- Harvey,[†] and Radoslaw M. Kowalczyk,^{⊥,*}

9

10 [†] School of Agriculture, Policy and Development, University of Reading, P.O. Box 236, 1

11 Earley Gate, Reading RG6 6AT, United Kingdom

12 [#] Department of Food and Nutritional Sciences, University of Reading, Whiteknights

13 Campus, P.O. Box 226, Reading RG6 6AP, United Kingdom

14 [⊥] National Institute of Agricultural Botany, Huntingdon Road, Cambridge CB3 0LE, United

15 Kingdom

16 [⊥] Chemical Analysis Facility, School of Chemistry, Food and Pharmacy, University of

17 Reading, P.O. Box 224, Whiteknights Campus, Reading RG6 6AD, United Kingdom

18

19 ^{*} Corresponding author (Tel.: + 44 (0)118 378 4756; Email: r.m.kowalczyk@reading.ac.uk)

20

[‡] CF Present address: Biobased Chemistry & Technology, Wageningen University, P.O. Box 17, 6700 AA, Borse Weilanden 9, Wageningen, The Netherlands

[§] HMR Present address: Pirbright Institute, Ash Road, Pirbright, Woking GU24 0NF, United Kingdom

21 **ABSTRACT**

22 A procedure based on ^{13}C CPMAS NMR was developed to study procyanidins (PCs) and
23 prodelphinidins (PDs) directly in milled sainfoin plant tissues. Blackcurrant and *Tilia* samples
24 enabled reference spectra of purified proanthocyanidin (PA) fractions, crude extracts and
25 milled plant tissues, with characteristic resonances at 155, 144 and 132 ppm. PC/PD ratios
26 were estimated from the I_{132}/I_{155} intensity ratio and differed by 2.5 to 5.9% compared to
27 thiolysis data. Normalization to the 155 ppm signal intensity from reference spectra enabled
28 analysis of PA contents with an error of ca 8 g PAs/100 g plant tissue. The procedure
29 estimates the lignin contribution and allows for a correction of the PA content. In six sainfoin
30 accessions, estimated PA contents, were 1.6- to 20.8-fold higher than the thiolysis and 1.4- to
31 2.6-fold higher than the HCl-butanol-acetone results. Method differences may reflect the
32 presence of unextractable, possibly high molecular weight PAs in sainfoin.

33

34 **KEYWORDS:** *condensed tannins; CPMAS NMR; thiolysis; HCl-butanol-acetone assay;*
35 *extractable and non-extractable proanthocyanidins; sainfoin;*

36

37

38

39

40

41

42

43

44

45

46 INTRODUCTION

47 Given the beneficial effects of proanthocyanidins (PAs) on human and animal
48 nutrition and health,^{1,2} there is a need for analytical methods capable of providing
49 information on the true PA content and their composition in plants. However, their analysis is
50 challenging as many plants synthesize complex polymeric mixtures of PAs (Figure 1).^{3,4}
51 Extractable PAs are typically analyzed after degradation in the presence of nucleophiles
52 followed by chromatographic separation of the reaction products and mass spectrometric
53 analysis; this gives data on PA content, flavan-3-ol subunit composition, mean degree of
54 polymerization (mDP) and procyanidin/prodelphinidin (PC/PD) and *cis/trans* flavan-3-ol
55 ratios.⁵⁻⁸ Extraction, however, is often incomplete as many plants also contain large quantities
56 of unextractable PAs, which can lead to underestimation of total contents.⁹⁻¹¹ Therefore, *in*
57 *planta* PA depolymerization with HCl-butanol-acetone, thiolysis or phloroglucinolysis has
58 been used to estimate extractable plus unextractable PAs; but, incomplete reaction, side
59 reactions or degradation-resistant PAs continue to pose challenges.^{3,4, 12-14} Of particular
60 concern is the fact that the HCl-butanol-acetone reagent often, but not always, gives higher
61 PA contents than thiolytic degradation;^{15, 16} although opposite results have also been found
62 with galloylated PAs (unpublished results). Questions remain, therefore, about the true PA
63 contents in plants and foods. Currently, little is known about the quantities and composition
64 of unextractable PAs and alternative methods deserve evaluation.^{10, 17}

65 In contrast to the above degradation methods, nuclear magnetic resonance (NMR) has
66 the potential to provide direct evidence and accurate results for total, i.e. extractable plus
67 unextractable, PAs. Solution-state ¹³C NMR spectroscopy has been used to determine the
68 relative purities,^{18, 19} PC/PD¹⁸⁻²¹ and *cis/trans* flavan-3-ol ratios^{18, 20, 21} and mDP values of
69 extracted PAs.^{20, 21} Whilst two dimensional gel-state NMR proved useful for assessing the
70 presence and composition of PAs in *Lotus* tissues,²² our proof-of-concept study showed that

71 gel NMR spectra of sainfoin (*Onobrychis viciifolia*) tissues had signal resolution and matrix
72 interference problems.

73 Solid-state NMR experiments have been used for the direct analysis of plant materials
74 and require neither extraction nor derivatization of PAs.²³⁻²⁵ Cross-polarization magic-angle
75 spinning (CPMAS) NMR techniques have previously been used to evaluate the efficiency of
76 PA extraction in pine barks,²³ softwood and barks from quebracho²⁵ and humus²⁶ as well as
77 to characterize PAs in dietary fiber samples.²⁷ In addition, differences in fingerprint spectra
78 were used to distinguish between PA transformations of leather tannings²⁸ and during fungal
79 degradation.²⁹ However, significant problems with poor resolution and frequent overlaps of
80 multiple signals prevented any in depth quantitative estimates of the PA content and PC/PD
81 ratios.

82 Sainfoin is a PA-containing forage legume, which is of interest for its nutraceutical
83 and environmental properties. It engenders high voluntary intakes,^{30, 31} has excellent nutritive
84 value,^{30, 31} lowers *in vitro* greenhouse gas emissions,³² prevents bloat in ruminants,³³ lowers
85 gastrointestinal nematodes,³⁴ and can enhance the unsaturated fatty acid composition of meat
86 and milk products.^{35, 36} The work presented here used six sainfoin accessions as a case study
87 for complex PA mixtures.^{37, 38} Such complexity may have been the main reason for the
88 observed discrepancies between the thiolysis and HCl-butanol-acetone assay results and still
89 requires detailed experimental confirmation.³⁹ Therefore, the aim of this study was to develop
90 a procedure based on a ¹³C CPMAS NMR technique for estimating PA content and
91 composition directly, without chemical modification in different sainfoin accessions and to
92 compare the results to thiolysis and HCl-butanol-acetone data.

93

94 MATERIALS AND METHODS

95 **Plant Samples.** Blackcurrant (*Ribes nigrum*) leaves were collected from Hildred PYO farm
96 (Goring-on-Thames, UK). *Tilia* (*Tilia* spp.) flowers were purchased from Flos (Mokrsko,
97 Poland) as described by Ropiak et al.⁸ Sainfoin (*Onobrychis viciifolia*) leaves and stems,
98 from six accessions, were provided by the National Institute of Agricultural Botany
99 (Cambridge, UK). Blackcurrant leaves and sainfoin plants were lyophilized or air-dried and
100 subsequently ground to pass a 1 mm sieve and ball-milled. Fine powders of ball-milled
101 blackcurrant leaves and *Tilia* flowers were also mixed on a weight basis (100/0, 70/30, 50/50,
102 30/70 and 0/100) for NMR analysis and the development of the procedure to elucidate PC/PD
103 ratios.

104

105 **Chemicals and Reagents.** Hydrochloric acid (36%), formic acid, acetic acid, butan-1-ol,
106 HPLC-grade acetone, HPLC-grade methanol, HPLC-grade dichloromethane, HPLC-grade
107 hexane, HPLC-grade acetonitrile and ammonium chloride were purchased from Fisher
108 Scientific Ltd (Loughborough, UK). (±) – Dihydroquercetin (98%) was from Apin Chemicals
109 (Abingdon, UK). Benzyl mercaptan and acetone-*d*₆ (99.9%) were supplied by Sigma-Aldrich
110 (Poole, UK). Deuterium oxide (D₂O) was from CK Isotopes (Ibstock, UK). Sephadex LH-20
111 was purchased from GE Healthcare (Little Chalfont, UK). Deionized water was purified in a
112 Milli-Q system (Millipore, Watford, UK).

113

114 **Proanthocyanidin Extraction and Purification.** Finely ground plant tissue (50 g) was
115 weighed into a conical flask. Acetone/water (500 mL, 7:3, v/v) was added and the mixture
116 was vigorously stirred for 1 h. The mixture was filtered under vacuum. The filtrate was
117 further extracted with dichloromethane (250 mL) to remove lipids and chlorophyll.
118 Polyphenols were concentrated in the aqueous phase with a rotary evaporator below 37.5 °C
119 to yield the “crude extract”.

120 This crude extract was diluted with deionized water (2 L) and filtered under vacuum.
121 The filtrate was loaded on a 400 x 65 mm i.d., glass column which had been packed with 70 x
122 65 mm i.d., Sephadex LH-20 resin and was equipped with a sintered-glass frit.⁴⁰ The column
123 was then washed with deionized water (2 L) to elute sugars and low molecular mass
124 phenolics.⁴⁰ Gravity elution with acetone/water (3:7, 1:1 and 8:2, v/v; 1 L each) yielded three
125 PA fractions from each plant source; the first 200 mL of each eluate was discarded and PA
126 fractions 1, 2 and 3 were collected in the next 500, 300, and 300 mL, respectively. The
127 organic solvent was removed on a rotary evaporator below 37.5 °C and the PA fractions were
128 frozen, freeze-dried and stored at -20 °C. Fractions 2 that eluted with 1:1 acetone/water were
129 designated as “purified PA fractions”. Various aliquots from the purified blackcurrant and
130 *Tilia* PA fractions (152:0, 36:83, 45:38, 63:25, 0:157 mg) were also mixed, in order to
131 provide a range of nominal PC/PD values of 0/100, 30/70, 50/50, 70/30 and 100/0, for NMR
132 analysis. The calculated PC and PD contents within purified PA fraction (in g PCs or
133 PDs/100 g purified PA fraction) were based on the PC and PD contents from thiolysis (*Tilia*:
134 97.4 g PCs/100 g of purified PA fraction and blackcurrant: 94.9 g PDs/100 g of purified PA
135 fraction).

136

137 **Thiolysis of Purified Proanthocyanidins.** Thiolysis reactions on purified PA fractions that
138 eluted with acetone/water (1:1 v/v) were performed according to Novobilský et al.⁴¹ Purified
139 PA fractions (4 mg) were weighed into a 10 mL screw-capped vial, dissolved in methanol
140 (1.5 mL) and acidified with HCl (0.5 mL, 3.3% in methanol, v/v). This was followed by
141 addition of benzyl mercaptan (50 µL) and the reaction mixture was stirred in a water-bath for
142 1 h at 40 °C. The reaction was quenched by adding ultrapure water (2.5 mL) to the mixture at
143 room temperature. The analysis and quantification of thiolysis reaction products were

144 performed with reverse-phase HPLC according to Gea et al.⁵ The operating conditions and
145 parameters of HPLC analysis were described by Williams et al.⁴²

146

147 ***In planta* Thiolysis of Sainfoin Proanthocyanidins.** Milled freeze-dried sainfoin plant
148 material (200 mg) was weighed into a 10 mL screw-capped vial, dissolved in methanol (2
149 mL) and acidified with HCl (1 mL, 3.3% in methanol, v/v). This was followed by addition of
150 benzyl mercaptan (100 μ L) and the mixture was stirred in a water-bath for 1 h at 40 °C. The
151 reaction was stopped by transferring the vials into an ice-bath and 1% formic acid in
152 ultrapure water (9 mL) was added at room temperature. The samples were centrifuged (3000
153 rpm, 5 min) and 1 mL of the mixtures added to HPLC vials for LC-MS analysis.

154 Samples (5 μ L) were injected into an 1100 series HPLC system (Agilent
155 Technologies, Stockport, UK) connected to a 150 x 3 mm i.d., 5 μ m, ACE super C₁₈ column
156 (Hichrom, Theale, UK) fitted to a 150 x 3 mm i.d., 5 μ m, ACE EXCEL Ultra-Inert UHPLC
157 compatible guard cartridge (Hichrom, Theale, UK). The column temperature was set at 60
158 °C. The HPLC system consisted of a G1379A degasser, a G1312A binary pump, a G1313A
159 ALS autoinjector, a G1314A VWD UV detector and a G1316A column oven and an 1100
160 series MSD API-ES mass spectrometer (Agilent Technologies, Waldbronn, Germany). Data
161 were acquired and processed with ChemStation software (version A 10.01 Rev. B.01.03).
162 The flow rate was 0.4 mL/min using formic acid (1%) in water containing 100 mg/L
163 ammonium chloride (solvent A) and HPLC-grade acetonitrile (solvent B). The thiolysis
164 reaction products eluted with the following gradient: 0-7 min, 2.5% B; 7-15 min, 2.5-5% B;
165 15-22 min, 5-10% B; 22-40 min, 10-40% B; 40-45 min, 40-100% B; 45-49, 100-2.5% B; 49-
166 60, 2.5% B. Mass spectra were recorded in the negative ionization scan mode from *m/z* 100-
167 1000. The mass spectrometer operating conditions were as follows: 3000 V for capillary
168 voltage, nebulizer gas pressure at 35 psi, drying gas at 12 mL/min and dry heater temperature

169 at 350 °C. Flavan-3-ols and their benzyl mercaptan adducts were identified by their retention
170 times, ultraviolet-visible (UV) spectra recorded at 280 nm and molecular masses and they
171 were quantitated using published response factors against dihydroquercetin.^{5,8}

172

173 ***In planta* Sainfoin Proanthocyanidin Analysis with the HCl-Butanol-Acetone Assay.** The
174 HCl-butanol-acetone assay followed the procedure of Grabber et al.²² with minor
175 modifications. Briefly, lyophilized sainfoin plant tissue was weighed (10 mg) into a 10 mL
176 screw-capped vial. The reagent mixture was prepared by mixing ammonium ferric sulfate
177 (150 mg) in ultrapure water (3.3 mL), hydrochloric acid (12 M, 5 mL), butan-1-ol (42 mL)
178 and acetone (50 mL). An aliquot of the reagent (10 mL) was added to the sainfoin sample.
179 The sample was left at room temperature (1 h) to check for the presence of flavan-4-ols or
180 flavan-3,4-diols as these generate false positives.⁴³ Tubes were then stirred and heated at 70
181 °C for 2.5 h in the dark. Samples were left to cool to room temperature and centrifuged (3000
182 rpm, 1 min). Absorbance of the supernatant was recorded by scanning from 450-650 nm
183 using a V530 spectrophotometer (Jasco, Great Dunmow, UK). The HCl-butanol-acetone
184 reagent was used as a blank and all samples were run in duplicate. A sainfoin sample of
185 known PA content and composition was used for quality control purposes.

186

187 **Analysis of Proanthocyanidins by ¹³C Cross-Polarization Magic-Angle Spinning**

188 **Nuclear Magnetic Resonance.** The solid-state CPMAS NMR spectra were recorded on a
189 Bruker Avance III 500 MHz spectrometer (Bruker Biopsin, Rheinstetten, Germany) operating
190 at carbon-13 Larmor frequency of 125.78 MHz (11.75T). Crude extracts and purified PA
191 fractions from *Tilia* and blackcurrant and ball-milled *Tilia*, blackcurrant and sainfoin plant
192 tissue samples (~80 mg) were compressed into standard 4 mm zirconia rotors prior to
193 analysis. A standard bore Bruker MAS probe was used and rotors were spun at 10 kHz rate.

194 The proton 90° pulse width was $3.7 \mu\text{s}$ at a power level of 32 W. The variable amplitude CP
195 ramp (90-100) was used with a contact time of 1 ms. In total, 4096 signal transients were
196 averaged into each spectrum with a 6 s relaxation delay at ambient temperature. All spectra
197 were referenced to an external adamantane signal (at 38.5 ppm with respect to TMS) as a
198 secondary reference. Dipolar dephasing experiments were run with identical parameters to
199 ^{13}C CPMAS NMR experiments and dephasing filter time (tdd) of $45 \mu\text{s}$ was optimized to
200 attenuate the signals of all protonated carbons.

201 To accurately extract intensity of all signals, each experimental spectrum was fitted
202 with a calculated spectrum that was constructed from the sum of signals with a Gaussian line
203 shape. Numbers and positions of signals in the calculated spectrum corresponded directly to
204 the number and positions of signals in the experimental spectrum. The width, amplitude and
205 position of each signal in the calculated spectrum was optimized by minimizing the
206 discrepancy between the experimental and calculated spectra using an “in house” written
207 Matlab routine based on the least square method. A number of initial parameters were tested
208 and the sets with the smallest discrepancy led to final results with a narrow range of the
209 fitting error. Once the width and the amplitude of the fitted signal at 155 ppm were known,
210 the intensity was calculated.

211 The PA content of sainfoin plant tissue samples was estimated by comparing the
212 signal intensity of the reference signal I_R at 155 ppm (from *Tilia* with 95 g PAs/100 g of
213 purified PA fraction, determined by thiolysis) with the intensity I_S of the 155 ppm signal in
214 the investigated sample.

215 The calculation of experimental error involved three factors: (i) the non-strictly
216 quantitative character of the solid-state NMR experiments because of the cross-polarization
217 signal enhancement (and possibly the miscalibration of the relaxation parameters); (ii) the

218 small variation in setting hardware parameters, such as tuning and matching of the probe; and
219 (iii) the numerical fitting of the spectra.

220 The recording of 2-3 spectra of the same sample with slightly different settings
221 regarding the first two factors (including contact time and relaxation delay) resulted in an
222 error of ca 2%. The discrepancies of the spectra acquired with different parameters
223 introduced an additional error of ca 2% after the fitting process. This accounted for a total
224 error of ca 4% for the intensity measurements. The PA content was calculated by comparing
225 the reference signal intensity to the intensity of the sample of interest. Therefore, the total
226 experimental error (ca 8% or 8 g PAs/100 g of plant tissue) derived from the sum of the
227 intensity errors between reference and sample intensity.

228 The influence of the spinning sidebands on the signal intensity was assessed by
229 comparing the 155 ppm signal intensity recorded at 10, 12 and 15 kHz spinning rates for both
230 crude extract and sainfoin plant tissue samples. Only small differences of less than 1%, well
231 within the experimental error were observed. At 10 kHz spinning rate, the spinning
232 sidebands of the signal at 155 ppm were not fully removed and decreased the intensity by ca
233 12%. However, the identical anisotropy of the chemical shift of C5, C7 and C8a carbons in
234 purified PA fraction and plant tissue PAs is expected (due to the same molecular structure)
235 and by normalizing the intensity of the 155 ppm signal from the sample of interest to the high
236 purity reference with 95 g PAs/100 g of purified PA fraction from *Tilia*, the contribution of
237 these factors to the intensity error could be neglected.

238 However, it should be noted that ^{13}C CPMAS NMR is not strictly quantitative and,
239 therefore, it is unlikely but not impossible that the discrepancies could have exceeded to a
240 small extent the range of the estimated experimental error.

241

242 **RESULTS AND DISCUSSION**

243 **Assignment of Signals in the Solid-State NMR Spectrum.** The ^{13}C CPMAS NMR spectra
244 of the purified PA fractions from *Tilia* and blackcurrant consisted of approximately ten
245 resolved resonances each centered between 170 and 0 ppm and showed several PA
246 fingerprint signals (Figure 2B). Characteristic resonances at 155, 144 and 132 ppm were
247 clearly observed in the spectra of all three sample types, i.e. purified PA fractions, crude
248 extracts and plant tissues (Figure 3). The signal at 155 ppm originated from the non-
249 protonated carbons C5, C7, C8a of the flavan-3-ol structure (Figure 2A).^{25, 27, 44} The signal at
250 144 ppm was from the resonance of the non-protonated carbons C3', C5' of blackcurrant PAs
251 and the non-protonated carbons C3', C4' of *Tilia* PAs.^{24, 25, 44} The most distinctive signal at
252 132 ppm originated from the non-protonated carbons C1', C4' in blackcurrant PAs but only
253 from the non-protonated carbon C1' in *Tilia* PAs.^{24, 25, 44}

254 In order to confirm these assignments, dipolar dephasing solid-state NMR
255 experiments were performed. The experimental parameters were optimized to detect only
256 signals from non-protonated carbons (Figure 2B). The resulting data confirmed our initial
257 identification and were in good agreement with spectra in reports on pecan nutshell PDs⁴⁵ and
258 *Photinia* leaf PCs.⁴⁴ Weak signals of not completely removed spinning sidebands were
259 detected in the 80-40 ppm region. Some very weak contributions from the protonated carbons
260 were also visible in the 40-20 ppm region and could have been caused by a small
261 miscalibration of parameters for the dephasing filter.

262 The resonances from protonated carbons were spread across the 120-0 ppm region
263 and showed line broadening and overlap, especially for the crude extracts and milled plant
264 tissues (Figures 2B and 3). This was likely due to interferences from other plant components
265 such as lignin, pectin, cutin, cellulose and hemicellulose as previously reported in crude
266 extracts and milled plant samples.^{25, 28, 44} However, our signal assignments of the purified PA

267 fractions from *Tilia* and blackcurrant were consistent with the much better resolved ^1H - ^{13}C
268 HSQC solution NMR spectra of the same samples.

269

270 **Estimation of Proanthocyanidin Content in the Presence of Lignin.** Accurate
271 quantification of PAs in plant matrices is of great interest and several techniques have been
272 used including solid-state NMR. The region between 160-120 ppm is understood to be the
273 most suitable despite some possible lignin or lignin-PA complex contributions especially in
274 milled plant samples.^{25, 44} In particular, Wawer et al.²⁷ postulated that resonances at 155 and
275 144 ppm in solid-state NMR dipolar dephased spectra from *Aronia* and blackcurrant fiber
276 powders could include responses from C3, 5-OMe and C4-OH carbons of lignin. However,
277 this interference was on a smaller scale (i.e. there were no significant changes to the signal
278 pattern) compared to resonances below 120 ppm (Figure 3) where the appearance changed
279 completely between purified PA fractions and milled plant tissues.

280 The characteristic signals at 155, 144 and 132 ppm in the spectra of the purified PA
281 fractions could only have derived from PAs as large polymeric lignin molecules were not
282 expected to be soluble in acetone/water.^{27, 46} Our strict, two step purification process also
283 assured that the low molecular weight phenolics or lignin fragments which may have been
284 present in small quantities in crude extracts were predominantly removed in the purified PA
285 fractions. The intensities of the 132 ppm signal spinning side bands at 211 ppm (left spinning
286 side band) and 53 ppm (right spinning side band) were exactly the same (0.02% difference)
287 which proves that there is no, or only negligible, lignin contribution to the NMR spectra of
288 the purified PA fractions. Figure 3 shows the comparison of the normalized amplitude of
289 these characteristic signals between purified PA fractions, crude extracts and milled plant
290 tissues. However, it was not possible to estimate the contribution of impurities or plant matrix
291 components to the intensity because currently there is no technique that can provide

292 completely accurate data on PA concentrations in the plant tissues. However, a contribution
293 of lignin in this region of the spectrum was expected between 155 and 144 ppm as
294 postulated,²⁷ and it should affect the width of the resonances due to overlap. Therefore,
295 measuring the change of the signal width in purified PA fractions, crude extracts and milled
296 plant tissues could provide an estimate of lignin interference. In our case, the estimated
297 increase in the width of the 155 and 144 ppm signals was <15% for crude extracts and <25%
298 for plant tissues, and corresponded well to the increased intensity of the lignin signal at 53
299 ppm, which was especially evident for the *Tilia* plant tissues (Figure 3B).

300 This simple estimate showed that the lignin interference had to be taken into account
301 as a correction to the PA content in plant samples. To test the accuracy of this assumption,
302 the content of extractable PAs in the crude extract from blackcurrant was estimated with ¹³C
303 CPMAS NMR and compared to the thiolysis result. The PA content determined by NMR
304 without correction was 39.7 g PAs/100 g crude extract and with the 15% reduction was 33.7
305 g PAs/100 g crude extract (with an error of ± 8 g PAs/100 g). The latter value was in
306 excellent agreement with the thiolysis result of 29.2 g PAs/100 g crude extract. This
307 demonstrates that both techniques can be used for the analysis of extractable PAs.

308 Although this simple calculation may take into account a lignin contribution, other
309 factors could also influence the width of the resonance line, e.g. relaxation and mobility
310 issues can arise from large polymeric molecules²⁵ and small variation in the shimming
311 quality.⁴⁷ Further, one needs to be aware that the thiolysis and HCl-butanol-acetone assays
312 have their own limitations, especially when applied directly to plant tissues.

313 Despite problems arising from possible plant matrix interferences in the milled plant
314 tissues, it may be concluded that the 155 ppm signal of the purified PA fractions was a
315 suitable marker to estimate PA contents in unknown milled plant tissues. Therefore, the PA
316 content of sainfoin plant tissue samples was estimated by comparing the intensity of a

317 reference signal, I_R , of the purified PA fraction (95 g PAs/100 g as determined by thiolytic)
318 from *Tilia* and the intensity of the I_S signal of the investigated sample.²³ The content error
319 was estimated to be ca 8 g PAs/100 g plant tissue.

320 The PA content of selected sainfoin plant tissues determined by ^{13}C CPMAS NMR
321 (without and with a 25% reduction for the lignin correction) was then compared to results
322 from the thiolytic and the HCl-butanol-acetone assays (Table 1). Selected spectra are shown
323 in Figure 4. Clearly, solid-state NMR consistently gave the largest PA content in all sainfoin
324 plant tissue samples and thiolytic gave the smallest. Most HCl-butanol-acetone values were
325 higher than the thiolytic data¹⁶ and accounted for ca 50% of the NMR data after lignin
326 correction. The results varied greatly between the sainfoin accessions for all methods, but it is
327 worth pointing out that the three Zeus samples (S1, S2 and S6) gave consistent results with
328 the NMR and HCl-butanol-acetone but not the thiolytic technique.

329 The above results illustrate the complexity of determining PA content directly in
330 plants. Our NMR results can be regarded as an upper limit of PA content compared to the
331 other techniques (Table 1) and is more likely to measure previously undetected unextractable
332 PAs because NMR does not require any chemical modification of the sample. Solid-state
333 NMR has already proved useful for estimating differences in PA contents within species, i.e.
334 for samples with similar PA compositions.^{25, 28} Our approach to account for a lignin
335 contribution could improve the way NMR could be used for PA measurements more
336 generally, including for the first time full quantitative elucidation. The validity of our
337 approach is illustrated by the analysis of the amplitude of the lignin signal at 53 ppm for
338 samples S1, S3 and S4 (Figure 4). The strongest lignin signal was detected in the S1 sample
339 and the weakest and approximately equal signals in the S3 and S4 samples. Had lignin
340 interference been the predominant factor in the measurement of PA content then S1 should
341 have had more PAs than the S3 and S4 samples. As evident from the 155 and 145 ppm

342 signals this was not the case as the S3 sample with the smallest lignin contribution had the
343 largest PA content.

344 There are several possible reasons for the differences between the NMR and the
345 thiolysis or HCl-butanol-acetone data. Whilst our results account for a lignin contribution at
346 155 ppm on the basis of the reference *Tilia* and blackcurrant samples, they do not rule out
347 contributions from other matrix components in the investigated sainfoin plant tissues.^{24, 27}
348 Secondly, solid-state NMR experiments are not strictly quantitative because the efficiency of
349 the cross-polarization depends on proton abundance in the neighboring environment, the
350 molecular dynamics and the relaxation behavior for improving the ¹³C signal.^{25, 28} These
351 factors are difficult to fully control in NMR experiments and are unlikely to be the same for
352 all carbons, and will also vary between different plant matrices. Therefore, it is possible that
353 the proton environment of the more complex and possibly larger sainfoin PAs (mDP of 12 to
354 27) could have resulted in a larger signal enhancement (and consequently PA content)
355 compared to the purified *Tilia* PAs (mDP of 8), but it is unlikely that this factor alone
356 increased the NMR signal by 100% (Table 1).

357 Another explanation could arise from the limitations of the thiolysis and HCl-butanol-
358 acetone assays in quantitating PAs directly in plant tissues.^{3, 4} It has been shown that longer
359 reaction times during thiolysis with benzyl mercaptan resulted in considerably higher
360 concentrations.⁵ In addition, the existence of thiolysis-resistant PAs has also been reported.⁹
361 The previous versions of the HCl-butanol assay often underestimated total PA content when
362 applied directly to plant or extracted residue samples, possibly because of incomplete
363 interaction with bound PAs.¹² However, the present HCl-butanol-acetone assay used a recent
364 modification by Grabber et al.,²² who demonstrated that inclusion of acetone achieved
365 complete degradation of unextractable PAs from two *Lotus* species. Nevertheless, it is
366 possible that sainfoin may have more diverse PA mixtures than *Lotus*,⁴⁸ which could explain

367 the observed differences between the NMR, thiolytic and HCl-butanol-acetone results.
368 Overall, the important point here is that the NMR results suggested that NMR allows
369 accounting for all extractable and unextractable PAs, whereas the thiolytic and HCl-butanol-
370 acetone assays may be underestimating PA contents.

371

372 **Estimation of Mean Degree of Polymerization.** The mean degree of polymerization (mDP)
373 can be obtained by integrating the C3 signals from PA extension units at 73 ppm and the
374 corresponding signals of PA terminal units at 67 ppm in solution NMR.^{20, 21, 26} However,
375 significant signal overlap from various plant constituents in milled plant tissues²⁶ and
376 consequently poor resolution of that region in the ¹³C CPMAS NMR spectra (Figure 3)
377 prevented estimation of mDP values.

378

379 **Estimation of procyanidin/prodelphinidin ratios.** The procedure to estimate PC to PD
380 ratios within PAs was based on their different hydroxylation patterns, which were clearly
381 distinguishable in the 140-100 ppm region of the ¹³C CPMAS NMR spectra (Figure 2B). The
382 resonance at 132 ppm was assigned to C1' and C4' carbons of PDs from blackcurrant, but in
383 *Tilia* PCs this resonance was assigned only to C1' carbon and had only half the intensity of
384 the PDs signal.^{44, 45} This difference in the 132 ppm signal intensity directly reflected the PD
385 and PC composition because the PA contents were comparable (i.e. blackcurrant 87 g
386 PAs/100 g of purified PA fraction; and *Tilia* 95 g PAs/100 g of purified PA fraction); the
387 molar percentages obtained by thiolytic were 94.9% PDs and 97.4% PCs for the purified PA
388 fractions, from blackcurrant and *Tilia* respectively.

389 This observation was validated by thiolytic and ¹H-¹³C HSQC solution NMR
390 experiments on artificial samples with the nominal PC/PD values equal to 100/0, 70/30,

391 50/50, 30/70 and 0/100, prepared by mixing purified PA fractions from *Tilia* and
392 blackcurrant.

393 However, the solution NMR methods, such as used by Czochanska et al.²⁰ and Zeller
394 et al.,²³ to estimate PC/PD ratios, which rely on the high resolution of the solution NMR
395 spectra could not be transferred directly to the solid-state NMR experiments. The
396 experimental resolution of the recorded solid-state NMR spectra was in the order of 12.5 Hz
397 (or 0.1 ppm) but the typical linewidth was ca 550 Hz (or 4.5 ppm), as expected for large
398 polymeric molecules. In addition to that, the spectra of milled plant tissues show significant
399 overlap with other signals from the plant matrix below 110 ppm, which prevents any
400 meaningful PC/PD analysis (Figures 3 and 5A). Therefore, it was impossible to calculate
401 PC/PD ratio on the basis of the C2' - C6' carbons as used in solution NMR experiments.

402 The experiment on the artificial mixtures of purified PA fractions with nominal
403 PC/PD values was repeated using the solid-state NMR technique. The intensity ratios
404 between signals at 132 and 155 ppm were selected as a calibration standard (to minimize the
405 experimental errors) because they were expected to vary linearly between a value of 1/3 or ca
406 0.33 for pure PCs, and 2/3 or ca 0.67 for pure PDs reflecting the number of carbons
407 contributing to each signal (Figures 2B and 5A). The resulting I_{132}/I_{155} experimental ratios
408 were within the theoretical limits (Figure 5B) and showed excellent linear dependence when
409 plotted as a function of the nominal PC/PD values.

410 However, to extend the procedure to estimate the unknown PC/PD ratios of milled
411 sainfoin tissues, it was necessary to repeat the experiment on the artificial mixtures but this
412 time these were prepared on a plant weight (mg) basis using the milled *Tilia* and blackcurrant
413 tissues. This allowed us to assess and overcome several problems, such as imprecisions in the
414 nominal PC/PD values and plant matrix interference, which varied between different plants
415 and could obscure any direct calculation of the unknown PC/PD ratio using only milled *Tilia*

416 or blackcurrant plant tissues. It was assumed that the nominal PC/PD values (again equal to
417 100/0, 70/30, 50/50, 30/70 and 0/100) in these plant tissues would be the same as in the
418 purified PA fraction mixtures. A similar linear dependence of the I_{132}/I_{155} on the nominal
419 PC/PD values was indeed observed for the milled plant tissues compared to the purified PA
420 fraction mixtures (Figure 5B). The large offset between both curves shows that there is a
421 possible interference of the plant matrix. However, both curves showed a similar slope, which
422 confirmed that extraction and purification had no effect on the PC/PD ratio in *Tilia* and
423 blackcurrant mixtures, and that the matrix effect on intensity was almost uniform across these
424 PC/PD ratios. Therefore, the gradient of the I_{132}/I_{155} intensity ratio as a function of nominal
425 PC/PD values could be regarded as a universal tool for setting up a calibration curve to
426 estimate unknown PC/PD ratios in other plant samples such as sainfoin.

427 Table 1 shows a comparison between the PC content in sainfoin samples from the ^{13}C
428 CPMAS NMR spectra and the thiolysis data. It can be seen that NMR estimated much higher
429 PC contents in purified PA fractions than thiolysis in four of the eight samples, but had a
430 rather low accuracy with an estimated experimental error of ± 16 mg PCs/100 mg purified PA
431 fraction. The biggest contributor to the estimated error was the uncertainty in the fitting of the
432 132 ppm signal, which overlapped with other signals from the plant matrices (Figures 4 and
433 6A). This was not surprising given the large offset already observed for the I_{132}/I_{155} intensity
434 ratio dependence on nominal PC/PD values between purified PA fractions and milled plant
435 tissues (Figure 5B). However, the degree and pattern of overlap was much larger and variable
436 in the sainfoin compared to the reference *Tilia* and blackcurrant plant tissues (Figure 5A).
437 This problem was particularly evident when comparing the NMR spectra of the S3 and S1
438 tissues (Figure 4); and was also highlighted in Figure 6, which compared the NMR spectra of
439 sainfoin leaves and stems. Obviously, it was only possible in the case of the fairly resolved
440 structure to extract the correct intensity of the signal at 132 ppm from the spectrum.

441 Therefore, a better resolution of the NMR spectra will be required to improve the accuracy of
442 the estimated PC molar percentage or PC/PD ratio beyond a qualitative assessment.

443 However, the comparison of sainfoin leaf and stem spectra also demonstrated the
444 potential of the ^{13}C CPMAS NMR approach to gain a rapid qualitative assessment (Figure 6).
445 The stem spectra showed a very different pattern of resonances in the region between 160 -
446 120 ppm. This was most likely due to poorer resolution of signals at 144 and 132 ppm in
447 stems, which tend to have higher lignin content than leaves, and also corresponded to lower
448 PA content in stems than leaves³⁷ and a greater contribution of plant matrix interferences.

449 This study reported a first attempt to develop a procedure based on ^{13}C CPMAS NMR
450 technique for the direct screening of PAs in sainfoin plant tissues. By using contrasting
451 signature PA spectra from blackcurrant and *Tilia* reference samples, a calibration for PAs was
452 developed based on signal intensities at 155 and 132 ppm. There were surprisingly large
453 differences between the PA content obtained by the thiolysis or HCl-butanol-acetone assays
454 and ^{13}C CPMAS NMR results. These differences could be due to interfering plant matrix
455 components, size and mobility differences in reference and sainfoin PAs, and/or presence of
456 unextractable PAs in sainfoin plant tissues. It is very unlikely that various sizes and mobility
457 differences would influence the cross-polarization enhancement to such an extent that it could
458 account for the observed PA content differences. A simple correction to account for the
459 matrix interference based on the linewidth difference between the purified PA fraction and
460 milled plant tissue has been used. However, even with the applied correction the differences
461 between the ^{13}C CPMAS NMR and thiolysis or HCl-butanol assays indicated that the two
462 degradation methods may not detect the entire PA present in some plants. Therefore, only ^{13}C
463 CPMAS NMR provided full information on both extractable and unextractable PAs in
464 sainfoin plant tissues.

465 In conclusion, this newly developed procedure based on ^{13}C CPMAS NMR technique
466 has proved to be a useful tool for estimating the upper limit of PA content directly in whole
467 plants and could be of interest for probing the bioactivities of unextractable PAs.

468

469 **ABBREVIATIONS USED:** PA, proanthocyanidin; PC, procyanidin; PD, Prodelphinidin;
470 ^{13}C CPMAS NMR, carbon-13 cross-polarization magic-angle spinning nuclear magnetic
471 resonance; mDP, mean degree of polymerization;

472

473 **ACKNOWLEDGMENTS:** We thank Hildred PYO farm for permission to collect the
474 blackcurrant leaf samples. Use of the NMR spectrometer at the Chemical Analysis Facility
475 (CAF), University of Reading is gratefully acknowledged.

476

477 **AUTHOR INFORMATION:** *Alternate Corresponding author (Tel.: +31 (0)317486721;
478 Email: christos.fryganas@wur.nl).

479

480 **FINANCIAL SUPPORT:** This work was supported by a European Union Marie Curie
481 training network (PITN-GA-2011-289377, “LegumePlus”).

482

483 **CONFLICTS OF INTEREST:** The authors declare no conflict of interest.

484

485 **SUPPORTING INFORMATION DESCRIPTION:** ^1H - ^{13}C Heteronuclear Single Quantum
486 Coherence (HSQC) Nuclear Magnetic Resonance Analysis of the Purified PA Fraction
487 Mixtures; Chromatogram of the Thiolytic Reaction Products from the Blackcurrant Purified
488 PA Fraction; Chromatogram of the Thiolytic Reaction Products from the *Tilia* Purified PA
489 Fraction.

490 **References**

- 491 (1) Cheynier, V.; Tomas-Barberan, F. A.; Yoshida, K. Polyphenols: From plants to a variety
492 of food and nonfood uses. *J. Agric. Food Chem.* **2015**, *63*, 7589-7594.
- 493 (2) Mueller-Harvey, I. Unravelling the conundrum of tannins in animal nutrition and health.
494 *J. Sci. Food Agric.* **2006**, *86*, 2010-2037.
- 495 (3) Hummer, W.; Schreier, P. Analysis of proanthocyanidins. *Mol. Nutr. Food Res.* **2008**, *52*,
496 1381-1398.
- 497 (4) Schofield, P.; Mbugua, D. M.; Pell, A. N. Analysis of condensed tannins: A review. *Anim.*
498 *Feed Sci. Technol.* **2001**, *91*, 21-40.
- 499 (5) Gea, A.; Stringano, E.; Brown, R. H.; Mueller-Harvey, I. *In situ* analysis and structural
500 elucidation of sainfoin (*Onobrychis viciifolia*) tannins for high-throughput germplasm
501 screening. *J. Agric. Food Chem.* **2011**, *59*, 495-503.
- 502 (6) Karonen, M.; Leikas, A.; Loponen, J.; Sinkkonen, J.; Ossipov, V.; Pihlaja, K. Reversed-
503 phase HPLC-ESI/MS analysis of birch leaf proanthocyanidins after their acidic degradation
504 in the presence of nucleophiles. *Phytochem. Anal.* **2007**, *18*, 378-386.
- 505 (7) Meagher, L. P.; Lane, G.; Sivakumaran, S.; Tavendale, M. H.; Fraser, K. Characterization
506 of condensed tannins from *Lotus* species by thiolytic degradation and electrospray mass
507 spectrometry. *Anim. Feed Sci. Technol.* **2004**, *117*, 151-163.
- 508 (8) Ropiak, H. M.; Ramsay, A.; Mueller-Harvey, I. Condensed tannins in extracts from
509 European medicinal plants and herbal products. *J. Pharm. Biomed. Anal.* **2016**, *121*, 225-
510 331.
- 511 (9) Matthews, S.; Mila, I.; Scalbert, A.; Donnelly, D. M. X. Extractable and non-extractable
512 proanthocyanidins in barks. *Phytochemistry* **1997**, *45*, 405-410.

- 513 (10) Perez-Jimenez, J.; Arranz, S.; Saura-Calixto, F. Proanthocyanidin content in foods is
514 largely underestimated in the literature data: An approach to quantification of the missing
515 proanthocyanidins. *Food Res. Int.* **2009**, *42*, 1381-1388.
- 516 (11) Ramsay, A.; Drake, C.; Grosse Brinkhaus, A.; Girard, M.; Copani, G.; Dohme-Meier, F.;
517 Bee, G.; Niderkorn, V.; Mueller-Harvey, I. Sodium hydroxide enhances extractability and
518 analysis of proanthocyanidins in ensiled sainfoin (*Onobrychis viciifolia*). *J. Agric. Food*
519 *Chem.* **2015**, *63*, 9471-9479.
- 520 (12) Makkar, H. P. S.; Gamble, G.; Becker, K. Limitation of the butanol-hydrochloric acid-
521 iron assay for bound condensed tannins. *Food Chem.* **1999**, *66*, 129-133.
- 522 (13) Callemien, D.; Guyot, S.; Collin, S. Use of thiolysis hyphenated to RP-HPLC-ESI-
523 MS/MS for the analysis of flavanoids in fresh lager beers. *Food Chem.* **2008**, *110*, 1012-
524 1018.
- 525 (14) Wayne McGraw, G.; Steynberg, J. P.; Hemingway, R. W. Condensed tannins: A novel
526 rearrangement of procyanidins and prodelphinidins in thiolytic cleavage. *Tetrahedron Lett.*
527 **1993**, *34*, 987-990.
- 528 (15) Klongsiriwet, C. Phytochemical analysis of a willow germplasm collection and its
529 pharmacological activities. Ph.D. Thesis, The University of Reading, UK, June 2016.
- 530 (16) Azuhnwi, B. N.; Boller, B.; Dohme-Meier, F.; Hess, H. D.; Kreuzer, M.; Stringano, E.;
531 Mueller-Harvey, I. Exploring variation in proanthocyanidin composition and content of
532 sainfoin (*Onobrychis viciifolia*). *J. Sci. Food Agric.* **2013**, *93*, 2102-2109.
- 533 (17) Hellstrom, J. K.; Mattila, P. H. HPLC determination of extractable and unextractable
534 proanthocyanidins in plant materials. *J. Agric. Food Chem.* **2008**, *56*, 7617-7624.
- 535 (18) Zeller, W. E.; Ramsay, A.; Ropiak, H. M.; Frygas, C.; Mueller-Harvey, I.; Brown, R.
536 H.; Drake, C.; Grabber, J. H. ^1H - ^{13}C HSQC NMR spectroscopy for estimating

- 537 procyanidin/prodelphinidin and *cis/trans*-flavan-3-ol ratios of condensed tannin samples:
538 Correlation with thiolysis. *J. Agric. Food Chem.* **2015**, *63*, 1967-1973.
- 539 (19) Hellstrom, J.; Sinkkonen, J.; Karonen, M.; Mattila, P. Isolation and structure elucidation
540 of procyanidin oligomers from Saskatoon berries (*Amelanchier alnifolia*). *J. Agric. Food*
541 *Chem.* **2007**, *55*, 157-164.
- 542 (20) Czochanska, Z.; Foo, L. Y.; Newman, R. H.; Porter, L. J. Polymeric proanthocyanidins -
543 stereochemistry, structural units, and molecular-weight. *J. Chem. Soc., Perkin Trans. 1* **1980**,
544 2278-2286.
- 545 (21) Qaadan, F.; Nahrstedt, A.; Schmidt, M.; Mansoor, K. Polyphenols from *Ginkgo biloba*.
546 *Sci. Pharm.* **2010**, *78*, 897-907.
- 547 (22) Grabber, J. H.; Zeller, W. E.; Mueller-Harvey, I. Acetone enhances the direct analysis of
548 procyanidin- and prodelphinidin-based condensed tannins in *Lotus* species by the butanol-
549 HCl-iron assay. *J. Agric. Food Chem.* **2013**, *61*, 2669-2678.
- 550 (23) Grigsby, W. J.; Hill, S. J.; McIntosh, C. D. NMR estimation of extractables from bark:
551 Analysis method for quantifying tannin extraction from bark. *J. Wood Chem. Technol.* **2003**,
552 *23*, 179-195.
- 553 (24) Parfitt, R. L.; Newman, R. H. ¹³C NMR study of pine needle decomposition. *Plant Soil*
554 **2000**, *219*, 273-278.
- 555 (25) Reid, D. G.; Bonnet, S. L.; Kemp, G.; van der Westhuizen, J. H. Analysis of commercial
556 proanthocyanidins. Part 4: Solid state ¹³C NMR as a tool for *in situ* analysis of
557 proanthocyanidin tannins, in heartwood and bark of quebracho and acacia, and related
558 species. *Phytochemistry* **2013**, *94*, 243-248.
- 559 (26) Lorenz, K.; Preston, C. M. Characterization of high-tannin fractions from humus by
560 carbon-13 cross-polarization and magic-angle spinning nuclear magnetic resonance. *J.*
561 *Environ. Qual.* **2002**, *31*, 431-436.

- 562 (27) Wawer, I.; Wolniak, M.; Paradowska, K. Solid state NMR study of dietary fiber powders
563 from aronia, bilberry, black currant and apple. *Solid State Nucl. Magn. Reson.* **2006**, *30*, 106-
564 113.
- 565 (28) Romer, F. H.; Underwood, A. P.; Senekal, N. D.; Bonnet, S. L.; Duer, M. J.; Reid, D.
566 G.; van der Westhuizen, J. H. Tannin fingerprinting in vegetable tanned leather by solid state
567 NMR spectroscopy and comparison with leathers tanned by other processes. *Molecules* **2011**,
568 *16*, 1240-1252.
- 569 (29) Gamble, G. R.; Akin, D. E.; Makkar, H. P.; Becker, K. Biological degradation of tannins
570 in sericea lespedeza (*Lespedeza cuneata*) by the white rot fungi *Ceriporiopsis subvermispora*
571 and *Cyathus stercoreus* analyzed by solid-state ¹³C nuclear magnetic resonance spectroscopy.
572 *Appl. Environ. Microbiol.* **1996**, *62*, 3600-3604.
- 573 (30) Hayot Carbonero, C.; Mueller-Harvey, I.; Brown, T. A.; Smith, L. Sainfoin (*Onobrychis*
574 *viciifolia*): A beneficial forage legume. *Plant Genet. Resour.* **2011**, *9*, 70-85.
- 575 (31) Wang, Y. X.; McAllister, T. A.; Acharya, S. Condensed tannins in sainfoin:
576 Composition, concentration, and effects on nutritive and feeding value of sainfoin forage.
577 *Crop Sci.* **2015**, *55*, 13-22.
- 578 (32) Chung, Y. H.; Mc Geough, E. J.; Acharya, S.; McAllister, T. A.; McGinn, S. M.;
579 Harstad, O. M.; Beauchemin, K. A. Enteric methane emission, diet digestibility, and nitrogen
580 excretion from beef heifers fed sainfoin or alfalfa. *J. Anim. Sci.* **2013**, *91*, 4861-4874.
- 581 (33) McMahon, L. R.; McAllister, T. A.; Berg, B. P.; Majak, W.; Acharya, S. N.; Popp, J. D.;
582 Coulman, B. E.; Wang, Y.; Cheng, K. J. A review of the effects of forage condensed tannins
583 on ruminal fermentation and bloat in grazing cattle. *Can. J. Plant Sci.* **2000**, *80*, 469-485.
- 584 (34) Manolaraki, F.; Sotiraki, S.; Stefanakis, A.; Skampardonis, V.; Volanis, M.; Hoste, H.
585 Anthelmintic activity of some Mediterranean browse plants against parasitic nematodes.
586 *Parasitology* **2010**, *137*, 685-696.

- 587 (35) Girard, M.; Dohme-Meier, F.; Silacci, P.; Ampuero Kragten, S.; Kreuzer, M.; Bee, G.
588 Forage legumes rich in condensed tannins may increase n-3 fatty acid levels and sensory
589 quality of lamb meat. *J. Sci. Food Agric.* **2016**, *96*, 1923-1933.
- 590 (36) Girard, M.; Dohme-Meier, F.; Wechsler, D.; Goy, D.; Kreuzer, M.; Bee, G. Ability of 3
591 tanniferous forage legumes to modify quality of milk and Gruyère-type cheese. *J. Dairy Sci.*
592 **2016**, *99*, 205-220.
- 593 (37) Malisch, C. S.; Luscher, A.; Baert, N.; Engstrom, M. T.; Studer, B.; Fryganas, C.; Suter,
594 D.; Mueller-Harvey, I.; Salminen, J. P. Large variability of proanthocyanidin content and
595 composition in sainfoin (*Onobrychis viciifolia*). *J. Agric. Food Chem.* **2015**, *63*, 10234-
596 10242.
- 597 (38) Stringano, E.; Hayot Carbonero, C.; Smith, L. M.; Brown, R. H.; Mueller-Harvey, I.
598 Proanthocyanidin diversity in the EU 'Healthyhay' sainfoin (*Onobrychis viciifolia*)
599 germplasm collection. *Phytochemistry* **2012**, *77*, 197-208.
- 600 (39) Hixson, J. L.; Bindon, K. A.; Smith, P. A. Evaluation of direct phloroglucinolysis and
601 colorimetric depolymerization assays and their applicability for determining condensed
602 tannins in grape marc. *J. Agric. Food Chem.* **2015**, *63*, 9954-9962.
- 603 (40) Brown, R. H.; Mueller-Harvey, I.; Zeller, W. E.; Reinhardt, L.; Stringano, E.; Gea, A.;
604 Drake, C.; Ropiak, H. M.; Fryganas, C.; Ramsay, A.; Hardcastle, E. E. Facile purification of
605 milligram to gram quantities of condensed tannins according to mean degree of
606 polymerization and flavan-3-ol subunit composition. *J. Agric. Food Chem.* **2017**, *65*, 8072-
607 8082.
- 608 (41) Novobilský, A.; Mueller-Harvey, I.; Thamsborg, S. M. Condensed tannins act against
609 cattle nematodes. *Vet. Parasitol.* **2011**, *182*, 121-220.

- 610 (42) Williams, A. R.; Fryganas, C.; Ramsay, A.; Mueller-Harvey, I.; Thamsborg, S. M.
611 Direct anthelmintic effects of condensed tannins from diverse plant sources against *Ascaris*
612 *suum*. *PLoS One* 2014, 9 (5), No. e97053, DOI: 10.1371/journal.pone.0097053.
- 613 (43) Watterson, J. J.; Butler, L. G. Occurrence of an unusual leucoanthocyanidin and absence
614 of proanthocyanidins in sorghum leaves. *J. Agric. Food Chem.* **1983**, 31, 41-45.
- 615 (44) Newman, R.; Porter, L. Solid state ¹³C-NMR studies on condensed tannins. In *Plant*
616 *Polyphenols*, Hemingway, R.; Laks, P., Eds. Springer US: 1992; Vol. 59, pp 339-347.
- 617 (45) Preston, C. M.; Sayer, B. G. What's in a nutshell: An investigation of structure by
618 carbon-13 cross-polarization magic-angle spinning nuclear magnetic resonance spectroscopy.
619 *J. Agric. Food Chem.* **1992**, 40, 206-210.
- 620 (46) Schuerch, C. The solvent properties of liquids and their relation to the solubility,
621 swelling, isolation and fractionation of lignin. *J. Am. Chem. Soc.* **1952**, 74, 5061-5067.
- 622 (47) Kim, H.; Ralph, J. Solution-state 2D NMR of ball-milled plant cell wall gels in DMSO-
623 d₆/pyridine-d₅. *Org. Biomol. Chem.* **2010**, 8, 576-591.
- 624 (48) Marais, J. P.; Mueller-Harvey, I.; Brandt, E. V.; Ferreira, D. Polyphenols, condensed
625 tannins, and other natural products in *Onobrychis viciifolia* (sainfoin). *J. Agric. Food Chem.*
626 **2000**, 48, 3440-3447.
- 627
- 628
- 629
- 630
- 631
- 632
- 633
- 634

FIGURE CAPTIONS:

Figure 1. A generic proanthocyanidin structure depicting flavan-3-ol subunits.

Figure 2. (A) Schematic representation of a proanthocyanidin structure consisting of a prodelphinidin subunit (top) and a procyanidin subunit (bottom), and (B) ^{13}C CPMAS NMR spectra (black solid line) of purified prodelphinidin fraction from blackcurrant leaves (upper) and purified procyanidin fraction from *Tilia* flowers (lower). The assignment of the resonances is consistent with the labelled carbon positions of the proanthocyanidin structure including resolved terminal (4t) and internal (4i) carbons. The grey lines depict the corresponding dipolar dephased solid-state NMR spectra, which were optimized to detect only non-protonated carbons. The asterisks mark the spinning side-bands.

Figure 3. Comparison of ^{13}C CPMAS NMR spectra of milled plant tissues (black line), acetone/water crude extracts (green line) and purified proanthocyanidin fractions (blue line) from (A) blackcurrant leaves and (B) *Tilia* flowers. Black arrows depict the signals at 155 and 132 ppm. All spectra were normalized to the amplitude of the 155 ppm signal to evaluate interferences from other plant components in the area of interest (160-120 ppm, highlighted in grey color).

Figure 4. Comparison of ^{13}C CPMAS NMR spectra of sainfoin plant (A) S3, (B) S1 and (C) S4 tissues. The insets show superimposed regions for comparison of the 155 and 145 ppm proanthocyanidin signals and the 53 ppm lignin signal.

Figure 5. (A) Comparison of the selected regions of ^{13}C CPMAS NMR spectra recorded for mixtures with nominal PC/PD values of 100/0, 70/30, 50/50, 30/70 and 0/100, prepared by mixing purified PA fractions (bottom) and milled plant tissues (top) of *Tilia* (PC-rich) and blackcurrant (PD-rich). The red arrows reflect the change in the 132 ppm signal intensity as function of the nominal PC/PD value and a small change in chemical shift. The increase in the color intensity of lines (light grey to black) denotes the increase in the PC/PD value. The black arrows depict other signals that also reflect spectral differences but are less useful due to signal overlaps. All signals were normalized to the amplitude of the signal at 155 ppm (red dot). (B) The relation between the intensity ratio I_{132}/I_{155} and nominal PC/PD values for the calibration mixtures of purified PA fractions and milled plant tissues. The dots denote the experimental points and the red lines show fitted linear dependence.

Figure 6. Fingerprint ^{13}C CPMAS NMR spectra from sainfoin (A) leaves and (B) stems. Both spectra were fitted using the sum of Gaussian shape lines and the result of the fit is superimposed on the original spectrum (red dotted lines). The grey lines below show Gaussian lines used to fit the corresponding region of the spectrum. The signals from this region were used to estimate PA content and PC/PD ratios and are shown in blue.

Table 1. Proanthocyanidin (PA) Contents and Procyanidin (PC) Contents within Purified PA Fraction Determined by ^{13}C CPMAS NMR, the HCl-Butanol-Acetone Assay and Thiolytic-HPLC Analysis.

Sainfoin plant tissue	Accession	PA (g/100 g of plant tissue)				PC (mg/100 mg of purified PA fraction)	
		NMR	NMR (-25%) ^a	HCl-butanol	Thiolytic	NMR	Thiolytic
S1	Zeus	16 ^b	12	5.0 ^c	1.5 ^d	20 ^b	23 ^d
S2	Zeus	16	12	5.3	3.4	30	18
S3	Cholderton	21	16	8.8	10.0	50	19
S4	Hampshire	14	10	7.1	5.2	30	19
S5	Ambra	18	14	<i>nt</i> ^d	5.2	30	19
S6	Zeus	15	11	4.3	1.7	10	18
S7	Perly	14	11	5.1	0.5	20	18
S8 ^e	Cotswold Common	19	14	<i>nt</i>	4.2	20	19

^a Correction for lignin interference; ^b experimental error was ± 8 g/100 g of plant tissue for the PA content, and ± 16 mg PCs/100 mg of purified PA fraction for the PC content within purified PA fraction; ^c experimental error was ± 0.5 g/100 g of plant tissue for the PA content; ^d experimental error was ± 0.6 g/100 g of plant tissue for the PA content, and ± 5 mg PCs/100 mg of purified PA fraction for the PC content within purified PA fraction; ^d *nt*: not tested; ^e sample S8 was analyzed only once.

Figure 1.




Figure 2.




Figure 3.




Figure 4.




Figure 5.





Figure 6.



GRAPHIC FOR TABLE OF CONTENTS: