

The North Atlantic Waveguide and Downstream Impact Experiment

Article

Accepted Version

Creative Commons: Attribution 4.0 (CC-BY)

Schäfler, A., Craig, G., Wernli, H., Arbogast, P., Doyle, J. D., McTaggart-Cowan, R., Methven, J. ORCID: <https://orcid.org/0000-0002-7636-6872>, Rivière, G., Ament, F., Boettcher, M., Bramberger, M., Cazenave, Q., Cotton, R., Crewell, S., Delanoë, J., Dörnbrack, A., Ehrlich, A., Ewald, F., Fix, A., Grams, C. M., Gray, S. L. ORCID: <https://orcid.org/0000-0001-8658-362X>, Grob, H., Groß, S., Hagen, M., Harvey, B. ORCID: <https://orcid.org/0000-0002-6510-8181>, Hirsch, L., Jacob, M., Kölling, T., Konow, H., Lemmerz, C., Lux, O., Magnusson, L., Mayer, B., Mech, M., Moore, R., Pelon, J., Quinting, J., Rahm, S., Rapp, M., Rautenhaus, M., Reitebuch, O., Reynolds, C. A., Sodemann, H., Spengler, T., Vaughan, G., Wendisch, M., Wirth, M., Witschas, B., Wolf, K. and Zinner, T. (2018) The North Atlantic Waveguide and Downstream Impact Experiment. *Bulletin of the American Meteorological Society*, 99 (8). pp. 1607-1637. ISSN 1520-0477 doi: 10.1175/BAMS-D-17-0003.1 Available at <https://centaur.reading.ac.uk/76719/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

To link to this article DOI: <http://dx.doi.org/10.1175/BAMS-D-17-0003.1>

Publisher: American Meteorological Society


All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online



AMERICAN METEOROLOGICAL SOCIETY

Bulletin of the American Meteorological Society

EARLY ONLINE RELEASE

This is a preliminary PDF of the author-produced manuscript that has been peer-reviewed and accepted for publication. Since it is being posted so soon after acceptance, it has not yet been copyedited, formatted, or processed by AMS Publications. This preliminary version of the manuscript may be downloaded, distributed, and cited, but please be aware that there will be visual differences and possibly some content differences between this version and the final published version.

The DOI for this manuscript is doi: 10.1175/BAMS-D-17-0003.1

The final published version of this manuscript will replace the preliminary version at the above DOI once it is available.

If you would like to cite this EOR in a separate work, please use the following full citation:

Schäfler, A., G. Craig, H. Wernli, P. Arbogast, J. Doyle, R. McTaggart-Cowan, J. Methven, G. Rivière, F. Ament, M. Boettcher, M. Bramberger, Q. Cazenave, R. Cotton, S. Crewell, J. Delanoë, A. Dörnbrack, A. Ehrlich, F. Ewald, A. Fix, C. Grams, S. Gray, H. Grob, S. Groß, M. Hagen, B. Harvey, L. Hirsch, M. Jacob, T. Kölling, H. Konow, C. Lemmerz, O. Lux, L. Magnusson, B. Mayer, M. Mech, R. Moore, J. Pelon, J. Quinting, S. Rahm, M. Rapp, M. Rautenhaus, O. Reitebuch, C. Reynolds, H. Sodemann, T. Spengler, G. Vaughan, M. Wendisch, M. Wirth, B. Witschas, K. Wolf, and T. Zinner, 2018: The North Atlantic Waveguide and Downstream Impact Experiment. Bull. Amer. Meteor. Soc. doi:10.1175/BAMS-D-17-0003.1, in press.



The North Atlantic Waveguide and Downstream Impact Experiment

Andreas Schäfler, George Craig, Heini Wernli, Philippe Arbogast, James D. Doyle,
 Ron McTaggart-Cowan, John Methven, Gwendal Rivière, Felix Ament, Maxi Boettcher,
 Martina Bramberger, Quitterie Cazenave, Richard Cotton, Susanne Crewell, Julien Delanoë,
 Andreas Dörnbrack, André Ehrlich, Florian Ewald, Andreas Fix, Christian M. Grams,
 Suzanne L. Gray, Hans Grob, Silke Groß, Martin Hagen, Ben Harvey, Lutz Hirsch, Marek Jacob,
 Tobias Kölling, Heike Konow, Christian Lemmerz, Oliver Lux, Linus Magnusson, Bernhard Mayer,
 Mario Mech, Richard Moore, Jacques Pelon, Julian Quinting, Stephan Rahm, Markus Rapp,
 Marc Rautenhaus, Oliver Reitebuch, Carolyn A. Reynolds, Harald Sodemann, Thomas Spengler,
 Geraint Vaughan, Manfred Wendisch, Martin Wirth, Benjamin Witschas, Kevin Wolf, Tobias Zinner

Affiliations: SCHÄFLER, BRAMBERGER, DÖRNBRACK, EWALD, FIX, GROß, HAGEN,
 LEMMERZ, LUX, RAHM, RAPP, REITEBUCH, WIRTH AND WITSCHAS — Deutsches Zentrum
 für Luft- und Raumfahrt, Oberpfaffenhofen, Germany; CRAIG, GROB, KÖLLING, MAYER AND
 ZINNER — Ludwig-Maximilians-Universität, München, Germany; BOETTCHER, GRAMS AND
 WERNLI — ETH Zürich, Switzerland; ARBOGAST — CNRM, Météo-France/CNRS, Toulouse,
 France; DOYLE AND REYNOLDS — Naval Research Laboratory, Monterey, California;
 MCTAGGART-COWAN — Environment and Climate Change Canada, Dorval, Canada; GRAY,
 HARVEY AND METHVEN — Department of Meteorology, University of Reading, Reading, United
 Kingdom, UK; RIVIÈRE — LMD/IPSL, ENS/PSL Research University/CNRS, Paris, France;
 CREWELL, JACOB AND MECH — Universität Köln, Köln, Germany; CAZENAVE, DELANOË
 AND PELON — Laboratoire ATmosphère, Milieux, et Observations Spatiales (LATMOS),
 Guyancourt, France; COTTON — Met Office, Exeter, United Kingdom; EHRLICH, WENDISCH
 AND WOLF — Universität Leipzig, Leipzig, Germany; HIRSCH — Max Planck Institute for
 Meteorology, Hamburg, Germany; AMENT AND KONOW — Universität Hamburg, Hamburg,
 Germany and Max Planck Institute for Meteorology, Hamburg, Germany; MAGNUSSON —
 European Centre for Medium-Range Weather Forecasts, Reading, United Kingdom; MOORE —
 Norwegian Meteorological Institute / University of Oslo, Oslo, Norway; QUINTING — ETH Zürich,
 Switzerland and Monash University, Clayton, Australia; RAUTENHAUS — Technische Universität
 München, Garching, Germany; SODEMANN AND SPENGLER — Geophysical Institute, University
 of Bergen and Bjerknes Centre for Climate Research, Bergen, Norway; VAUGHAN — National
 Centre for Atmospheric Science, University of Manchester, Manchester, United Kingdom

Corresponding author:

Dr. Andreas Schäfler, Institut für Physik der Atmosphäre, Deutsches Zentrum für Luft- und Raumfahrt
 (DLR), Oberpfaffenhofen, 82234 Weßling, Germany, email: andreas.schäfler@dlr.de

Capsule. Multi-aircraft and ground-based observations were made over the North Atlantic in fall 2016 to investigate the importance of diabatic processes for midlatitude weather.

Abstract. The North Atlantic Waveguide and Downstream Impact Experiment (NAWDEX) explored the impact of diabatic processes on disturbances of the jet stream and their influence on downstream high-impact weather through the deployment of four research aircraft, each with a sophisticated set of remote-sensing and in situ instruments, and coordinated with a suite of ground-based measurements. A total of 49 research flights were performed, including, for the first time, coordinated flights of the four aircraft; the German High Altitude and Long Range Research Aircraft (HALO), the Deutsches Zentrum für Luft- und Raumfahrt (DLR) Dassault Falcon 20, the French Service des Avions Français Instrumentés pour la Recherche en Environnement (SAFIRE) Falcon 20, and the British Facility for Airborne Atmospheric Measurements (FAAM) BAe 146. The observation period from 17 Sep to 22 Oct 2016 with frequently occurring extratropical and tropical cyclones was ideal to investigate midlatitude weather over the North Atlantic. NAWDEX featured three sequences of upstream triggers of waveguide disturbances, their dynamic interaction with the jet stream, subsequent development, and eventual downstream weather impact on Europe. Examples are presented to highlight the wealth of phenomena that were sampled, the comprehensive coverage and the multi-faceted nature of the measurements. This unique dataset forms the basis for future case studies and detailed evaluations of weather and climate predictions to improve our understanding of diabatic influences on Rossby waves and downstream impact of weather systems affecting Europe.

Progress in understanding the processes controlling midlatitude weather is one of the factors that have contributed to a continuous improvement in the skill of medium-range weather forecasts in recent decades (Thorpe 2004; Richardson et al. 2012; Bauer et al. 2015). Additionally, numerical weather prediction (NWP) has undergone a revolution in recent years, with the development and widespread use of ensemble prediction systems (EPS) to represent forecast uncertainty (Bauer et al. 2015). However, the short-term prediction of high-impact weather (HIW) events (e.g., strong winds and heavy precipitation), and the medium-range prediction of extratropical cyclones, including their tracks and intensity, is still a major challenge (e.g., Frame et al. 2015). Recent research on midlatitude weather has focused on quantifying model errors and predictability, and in particular on investigating the role of diabatic processes such as those related to clouds and radiation, whose interaction with the dynamics of the flow must be understood and represented more accurately in models in order to further improve forecast quality.

Detailed observations are needed to characterize the weather systems and embedded physical processes across a range of spatial and temporal scales that encompass cloud microphysical variability and Rossby waves. In September and October 2016, the North Atlantic Waveguide and Downstream Impact Experiment (NAWDEX) made new multi-scale observations in the North Atlantic basin from eastern Canada to western Europe. Weather features expected to be associated with forecast errors were extensively probed, providing a high-quality set of observations that are not assimilated routinely and thus can be used for validation of the NWP systems.

The fall season was chosen for the experiment because diabatic processes are particularly active due to relatively high sea surface temperatures and the jet stream is intensifying as the high latitudes cool. Many of the weather phenomena central to the growth of disturbances on the jet stream and midlatitude predictability are active in fall, such as extratropical cyclones with intense fronts and warm conveyor belts (WCBs), carrying air from the oceanic boundary

layer into ridges at the tropopause level. There is also the possibility of North Atlantic tropical cyclones (TCs) recurving poleward into midlatitudes and undergoing extratropical transition (ET) – a process known to be associated with low predictability (Harr et al. 2008). Coherent mesoscale depressions of the tropopause, known as tropopause polar vortices (TPVs, Cavallo and Hakim 2010; Kew et al. 2010), can disturb the jet stream if they move equatorward from the Arctic.

NAWDEX contributes to the World Weather Research Program (WWRP) and its High Impact Weather project (Jones and Golding 2015), and aims to provide the observational foundation for further investigating cloud diabatic processes and radiative transfer in North Atlantic weather systems, which will form the basis for future improvements in the prediction of HIW over Europe.

THE ROLE OF DIABATIC PROCESSES. Weather in Europe strongly depends on the life-cycle of Rossby waves that propagate along the slowly varying part of the North Atlantic jet stream (Martius et al. 2010). The strong meridional potential vorticity (PV) gradient associated with the jet stream serves as a waveguide for propagating Rossby waves. Frequently, small disturbances in the jet entrance region over eastern North America grow in baroclinic weather systems and evolve into large-amplitude features in the European sector (Schwierz et al. 2004). Figure 1 portrays an idealized North Atlantic flow situation that could result in HIW in the form of high winds and heavy precipitation over northern Europe. In addition to Rossby waves amplifying through baroclinic instability, diabatic processes are able to modify upper-tropospheric PV at the level of the midlatitude jet stream, which impacts the wavelength and amplitude of the downstream Rossby wave development (e.g., Massacand et al. 2001; Knippertz and Martin 2005; Grams et al. 2011; Teubler and Riemer 2016).

The majority of precipitation and cloud diabatic processes in extratropical cyclones occur within a coherent airstream known as the WCB. It carries warm, moist air from the low-level

warm sector of a cyclone to the ridge at tropopause level within 1-2 days (Browning et al. 1973; Carlson 1980; Wernli and Davies 1997). The boundary layer humidity in the inflow of WCBs (Region 1 in Fig. 1) can impact the outflow height of WCBs (Schäfler and Harnisch 2015). For some WCBs, the inflow region coincides with a filament of strong horizontal water vapor transport, a so-called “atmospheric river”, which can contribute to intense rain in the midlatitudes (Lavers and Villarini 2013). During the ascent of WCBs (Region 2 in Fig. 1), latent heating due to cloud microphysical processes, embedded convection and turbulent fluxes influence the level of the outflow layer, the direction taken by outflow air masses, and the shape of the upper-level ridge (Martínez-Alvarado et al. 2014; Joos and Forbes 2016). The latent heating in WCBs is strong both in the early phase of the ascent when condensation dominates and later when mixed-phase clouds are formed and vapor deposition on ice crystals and snow becomes important (Joos and Wernli 2012).

The effect of the heating on the PV structure is to produce a positive PV anomaly in the lower troposphere (Wernli and Davies 1997), which influences the structure and evolution of midlatitude surface cyclones (e.g., Kuo et al. 1991; Davis et al. 1993; Binder et al. 2016). Above the level of maximum latent heating, PV is reduced by cloud diabatic processes leading to negative PV anomalies in the upper-tropospheric WCB outflow region (Wernli 1997; Pomroy and Thorpe 2000; Madonna et al. 2014; Methven 2015). The divergent outflow winds (Region 3 in Fig. 1) tend to amplify the upper-level downstream ridge and to intensify the jet stream by strengthening the PV gradient (Archambault et al. 2013). If the outflow layer is higher, the negative PV anomaly is stronger and more air mass enters the anticyclonic branch of the WCB flowing into the downstream ridge (Grams and Archambault 2016). In addition, a sharp peak in longwave radiative cooling near the tropopause, associated with a step change in water vapor, creates a reinforcement of the positive PV anomaly in upper-level troughs (Chagnon et al. 2013) and plays a key role in maintaining and strengthening TPVs (Cavallo and Hakim 2012).

Diabatic processes also play a key role in weather systems that act as triggers to disturb the midlatitude waveguide. Recurving TCs undergoing ET (Jones et al. 2003) can enhance the anticyclonic and divergent flow at upper levels, excite and amplify Rossby waves and cause downstream forecast errors, as well as HIW events (e.g., Agusti-Panareda et al. 2004; Harr et al. 2008; Riemer and Jones 2010). Radiatively maintained TPVs, which are positive PV anomalies above the tropopause, can disturb the Rossby waveguide from polar latitudes.

Rossby wave breaking leads to PV filamentation, forming smaller-scale PV anomalies such as PV streamers and cut-off vortices. They form frequently over the eastern North Atlantic and Europe (e.g., Wernli and Sprenger 2007), and several studies have reported their relevance for triggering HIW, in particular heavy precipitation (e.g., Martius et al. 2006; Chaboureaud and Claud 2006; Grams and Blumer 2015). Synoptic wave breaking events are also important for the large-scale flow itself as they reinforce weather regimes such as blocking ridges (Michel and Rivière 2011; Spensberger and Spengler 2014). Blocks are also strongly influenced by diabatic processes in air masses ascending from the lower troposphere (Pfahl et al. 2015).

Disturbances of the waveguide and associated errors can amplify and propagate downstream, and may cause significant forecast errors over Europe (Madonna et al. 2015; Martínez-Alvarado et al. 2016) (Region 4 in Fig. 1). In NWP models, diabatic processes such as those associated with convection, cloud microphysics and radiation are represented by parameterizations of varying degrees of fidelity and may contain both systematic and random errors that influence forecast skill. A distinct Rossby wave pattern associated with poleward transport of warm and moist air over the eastern US and strong diabatic activity has been identified as a common precursor 6 days before the worst forecast busts over Europe (Rodwell et al. 2013). Upscale error growth experiments in numerical models show that the growth of small-scale perturbations is initially confined to regions where condensation is occurring, with the regions of large error amplitude gradually expanding to affect the

synoptic-scale weather pattern (Zhang et al. 2007; Selz and Craig 2015). Doyle et al. (2014) found forecasts of an extratropical cyclone with severe impact in western Europe to be very sensitive to the initial low-level moisture, which influenced the moisture supply in a WCB. At upper levels, global NWP models fail to maintain a sufficiently sharp tropopause, showing a decrease in sharpness with forecast lead time (Gray et al. 2014). This influence on the waveguide can have major implications for the representation of the downstream propagation and amplification of Rossby waves in NWP (Harvey et al. 2015) and the associated prediction of HIW.

Previous studies using measurements to study the influence of diabatic processes on the Rossby waveguide have been primarily based on routinely collected observations by operational meteorological services. These observations rely largely on satellite data, which models predominantly assimilate in cloud-free areas, and on sparse in situ measurements, all of which are combined in the data assimilation system using model forecasts as a background estimate. This approach to studying diabatic processes has significant limitations since these processes tend to be strongest in cloudy and precipitating regions, which are particularly challenging for both observation and modeling systems. The processes associated with diabatic heating are characterized by a high-degree of small-scale variability, particularly in the vertical (e.g., sharp vertical gradients of cloud microphysical processes and their interactions with radiative forcing), which are typically poorly resolved by satellite and conventional in situ observations. Furthermore, rapid error growth and systematic model errors lead to large errors in the background forecast in precisely these regions, which are poorly characterized by error covariance matrices based on climatology and/or sampling using an ensemble of limited size. A field campaign has the potential to address some of these difficulties by deploying specialized observing systems with high resolution and the ability to measure both in and around clouds.

NAWDEX was proposed with the overarching hypothesis that diabatic processes have a major influence on the jet stream structure, the downstream development of Rossby waves, and eventually HIW. Specific science goals were formulated (Table 1), which require observations of moisture advection in the boundary layer and of the vertical distribution of stability, water vapor, liquid droplets and ice crystals. These observations will be used to investigate spatial variability within clouds and the implications for diabatic processes. Detailed wind measurements in the layer of the divergent outflow of the WCB are needed to investigate the interaction of diabatically modified air masses with the upper-level jet. This includes observations of horizontal and vertical gradients of wind, temperature and humidity as well as hydrometeors in clouds. Accordingly, high-resolution cross-sections of wind, temperature and humidity from the lower stratosphere down to the surface, inside and outside of clouds, are the central observational requirement, which are not available from conventional observations.

EXPERIMENTAL DESIGN AND OBSERVATIONS. The need for a new field experiment emerged from a series of campaigns coordinated by the World Meteorological Organization's program THORPEX (The Observing-System Research and Predictability Experiment, Parsons et al. 2017). This series includes the Atlantic THORPEX Regional Campaign (ATReC, Rabier et al. 2008), Winter Storm Reconnaissance (WSR, e.g., Szunyogh et al. 2000), the THORPEX Pacific Asian Regional Campaign (T-PARC, Weissmann et al. 2011), and the Convective and Orographically induced Precipitation Study/European THORPEX Regional Campaign (COPS/ETReC 2007, Wulfmeyer et al. 2011), which all focused on the impact of additional observations on improving forecast accuracy. This idea was pioneered by the Fronts and Atlantic Storm Track Experiment (FASTEX) in 1997 (Joly et al. 1999) where the concept of targeting observations using sensitive area calculations was introduced. The synthesis of these campaigns and data assimilation experiments denying observations in data rich areas showed that the impact of targeted observations on global

forecast systems is weaker than originally anticipated, although they improve forecasts on average (see review of Majumdar (2016)). At the same time, as discussed above, evidence was growing that forecast errors often originate in regions where diabatic processes are strong, and observation and modeling systems are least reliable. This provided the motivation for a new campaign, NAWDEX, that rather than targeting regions of forecast sensitivity, instead focused on observing the processes that are thought to be most uncertain in NWP models.

Diabatic processes are difficult to measure directly, but can be constrained via their observable effects on the structure and evolution of weather systems. In the decade before THORPEX, detailed diagnostic case studies using aircraft measurements (e.g., ERICA (Experiment on Rapidly Intensifying cyclones over the Atlantic, Hadlock and Kreitzberg 1988) and FASTEX) had already shown that diabatic processes, in particular diabatic heating and cooling, can impact the large-scale dynamics via PV modification (Neiman et al. 1993; Pomroy and Thorpe 2000). However, the processes are difficult to accurately quantify since they depend on fine-scale structures (e.g., large gradients) in the water vapor and cloud fields and are influenced by transport and mixing over a wide range of spatial and temporal scales throughout the lifetime of the cyclone (recall Fig. 1). Two single-aircraft field campaigns organized within THORPEX explored how aircraft observations could be used to accurately constrain the impact of diabatic heating in midlatitude cyclones. DIAMET (DIAbatic influence on Mesoscale structures in ExTratropical storms; Vaughan et al. 2015) made airborne in situ measurements of liquid droplets and ice crystals and used them to infer the microphysical processes acting, their positions relative to mesoscale structures (such as fronts and PV anomalies), and their role in the weather system dynamics. Although the observations were limited to the 1-D aircraft flight path, they provided a basis for the modeling studies of Dearden et al. (2014), who obtained estimates of heating rates from various microphysical processes represented by a Lagrangian model initialized with in situ observations of size

distributions for cloud droplets and ice particles. However, using a model to extrapolate the measurement information in space and time represents an additional source of uncertainty in the quantification of the processes. The THORPEX-NAWDEX-Falcon project (Schäfler et al. 2014) attempted to constrain this uncertainty by carrying out in situ observations of clouds, humidity and wind in ascending WCBs, and trying to re-sample the same air masses at a later time to obtain a Lagrangian estimate of integrated diabatic effects. NAWDEX was conceived to expand upon the design of these previous experiments by combining high resolution remote sensing and in situ instrumentation to provide accurate measurements of atmospheric structures including strong gradients, using multiple aircraft to sample air masses at different stages of the WCB ascent and advection along the tropopause.

To allow these observations to be related to the development of weather forecast errors, NAWDEX employed four research aircraft and ground-based stations spanning the northern part of the North Atlantic with the aim of observing the processes influencing the development of disturbances to the North Atlantic waveguide across the Atlantic. This includes upstream triggering of disturbances on the waveguide by phenomena with strong latent heat release, the continuous effects of clouds and radiation near the tropopause, the dynamical interactions between large-scale disturbances, and the potential impact on weather over Europe from the Mediterranean to Scandinavia.

Airborne platforms and payload. NAWDEX employed four research aircraft, the German High Altitude and Long Range Research Aircraft (HALO) and the Deutsches Zentrum für Luft- und Raumfahrt (DLR) Dassault Falcon 20, the French Service des Avions Français Instrumentés pour la Recherche en Environnement (SAFIRE) Falcon 20, and the British Facility for Airborne Atmospheric Measurements (FAAM) BAe 146. FAAM operated from the UK and HALO and the two Falcon aircraft from Keflavik, Iceland, in an area covering the North Atlantic, north of 45° N, and northern and central Europe. The payloads were chosen to

observe the required profiles of wind, temperature, moisture and cloud properties, and in the case of FAAM, in situ cloud microphysics.

The strategy was to deploy HALO with its extended range to observe moisture transport and diabatic processes in weather systems upstream of Iceland that impact the midlatitude waveguide. HALO is a modified Gulfstream G-550 ultra-long-range business jet with a maximum flight range of about 10000 km and a maximum endurance of 10 hours (Krautstrunk and Giez 2012; Wendisch et al. 2016), which allows accessing remote regions over the central North Atlantic that are not accessible by other European research aircraft. The high ceiling of almost 15 km in combination with a sophisticated remote-sensing payload (see sidebar and Table 2) allow HALO to fly above the main commercial aircraft routes and to probe features of interest from above. The two Falcon aircraft, with a maximum range of 3000 km, a maximum endurance of about 4 hours and a ceiling up to 12 km, aimed to observe the approaching cyclones and evolving jet streams close to Iceland. The DLR Falcon was equipped with two wind lidar systems and the SAFIRE Falcon with a remote-sensing payload for clouds and winds (sidebar and Table 2). The FAAM BAe 146, with a maximum endurance of 5 hours and a ceiling of 10 km, was equipped with a range of in situ instrumentation for meteorological, cloud and chemical measurements together with a downward-pointing aerosol lidar and passive spectral radiometers. Its flights from East Midlands, UK, were aimed at observing microphysics and turbulence in WCBs and the structure of the jet stream.

HALO, SAFIRE and the FAAM aircraft were equipped with dropsonde dispensers to measure air temperature, wind and humidity profiles. Global NWP centers could access the dropsonde data from HALO and SAFIRE via the Global Telecommunication System in near real-time. The potential for coordinated application of the various instruments on board multiple aircraft was realized through specific instrument-driven science goals (Table 1 and sidebar). Table 2 indicates which of the research aims listed in Table 1 are addressed by each instrument.

In parallel to NAWDEX, the NOAA (National Oceanic and Atmospheric Administration) SHOUT (Sensing Hazards with Operational Unmanned Technology) campaign took place in the tropical and subtropical western North Atlantic. SHOUT utilized the unmanned NASA Global Hawk aircraft with a suite of remote-sensing platforms and dropsondes to study the impact of the observations on TC forecasts. During the campaign, a tropical storm (TS) moved into the midlatitudes and underwent ET, providing an unprecedented scientific opportunity to observe the interaction of such a system with the jet stream using a combination of upstream flights with the SHOUT Global Hawk and downstream flights with NAWDEX aircraft.

[Place Sidebar 1 here: Active remote-sensing observations for future satellite missions Aeolus and EarthCARE]

Airborne observations. NAWDEX observations took place in the North Atlantic basin between 17 Sep and 22 Oct 2016. Figure 2 shows the tracks of the 47 research flights of the four aircraft, together amounting to 205 flight hours. Performing research flights over the North Atlantic is complicated because of the dense trans-Atlantic air traffic. Commercial airliners are tightly staggered along predefined flight routes, the so-called North Atlantic Tracks (NATs), between altitudes of 9 and 12 km. Operating research aircraft beneath the NATs offers high flexibility for the flight planning; however, the base height of the NATs is often too low to observe the tropopause and jet-related maximum wind speeds. Furthermore the location of the NATs changes from day to day, depending on the forecast wind situation. Height changes and the release of dropsondes from high altitudes are not possible in the NAT area. The requirement of air traffic control (ATC) authorities to have detailed flight plans 2-3 days in advance created challenging circumstances in weather situations with reduced predictability, i.e., in situations with large changes between subsequent forecasts. Therefore, NAWDEX combined modern forecasting tools including ensemble and adjoint-based

diagnostics, and new visualization techniques to incorporate forecast uncertainty in the planning process (see sidebar on forecast products).

HALO covered large parts of the central and eastern North Atlantic and reached flight distances up to 7150 km (~9 h). The flights were performed either at altitudes between 11.5 and 14.2 km above the NATs for remote-sensing observations or at ~8 km to release dropsondes beneath the NATs. The two Falcon aircraft remained in radar-controlled air spaces near Greenland, Iceland and the UK. The FAAM BAe 146 flights were north and west of the UK. A total of 289 dropsondes were released (Fig. 3a).

[Sidebar 2: Forecast products to investigate forecast uncertainty]

The research flights occurred within 13 Intensive Observation Periods (IOPs), which were consecutively numbered and had a duration of 1-6 days. Each IOP was associated with a particular weather system development and addressed one or more NAWDEX science objectives (Table 3). For easier communication, the IOPs were given names, which either correspond to the cyclone naming of the Free University of Berlin or the National Hurricane Center, or were invented by the NAWDEX team (Table 3). Some IOPs overlap in time when different weather systems were observed simultaneously by the different aircraft.

To exploit instrument synergies and enable direct instrument comparisons, coordinated flights were performed, i.e., the same air mass was near-simultaneously probed by different aircraft on common flight legs. In total, 16 coordinated legs, with a total flight time of 14.5 h and a distance of about 10000 km, were achieved. The longest coordinated leg with the SAFIRE Falcon and HALO on 14 Oct 2016 had a distance of 1365 km (1.8 h). On two occasions the coordination involved three aircraft: HALO and the two Falcons flew together for ~30 min (~300 km) on 9 Oct between the UK and Iceland, and on 14 Oct, FAAM, HALO and the SAFIRE Falcon had a common leg between the Faroe Islands and Scotland (55 min, 570 km).

Ground-based facilities and observations. During several IOPs additional ground-based observations were taken to complement the aircraft operations and to enhance the temporal and spatial coverage of routine observations. In total 589 additional radiosondes from 40 stations in 14 countries were launched (Fig. 3b and Table 4). Of these launches, 253 were achieved through the cooperation of national meteorological agencies in the European Meteorological Services Network (EUMETNET), complemented by additional radiosondes from Iceland, UK, France and Norway. Launches from land stations or commercial ships were requested daily depending on the predicted evolution of weather systems. Furthermore, two additional radiosondes were launched daily during the campaign from six stations in eastern Canada, upstream of the main NAWDEX area (336 in total).

Special ground-based observations were conducted in Iceland, the UK and France (Fig. 3b). At Keflavik airport, a radiosonde facility was set up by DLR to increase the frequency of the operational soundings. In cases of orographically induced gravity waves (GWs), large balloons were launched to reach altitudes up to 42 km. Also in Keflavik, a Doppler cloud radar BASTA (Delanoë et al. 2016) allowed several comparisons with its airborne counterpart on board the SAFIRE Falcon during overflights. In the UK, a mesosphere–stratosphere–troposphere (MST) radar, Raman lidar, and radiosondes were operated at Capel Dewi (Wales), together with another MST radar wind profiler at South Uist (Scotland). Additionally, the MST radar at Andøya, Norway, measured tropospheric winds upon request. Two observational sites were active in France during the campaign. The site in Lannion (Brittany) operated a wind profiler, the BASTA Doppler cloud radar and a GPS station. The Site Instrumental de Recherche par Télédétection Atmosphérique (SIRTA) near Paris (Haefelin et al. 2005) operated radar and lidars, and launched radiosondes.

METEOROLOGICAL CONDITIONS. The fall of 2016 was a favorable period for observing midlatitude weather over the North Atlantic. The average synoptic situation for the

campaign period was characterized by an increased frequency of relevant weather systems compared to climatology (Fig. 4). One of the most prominent features was a long-lasting blocking high and surface anticyclone covering large parts of Scandinavia (Fig. 4a). Extratropical cyclones occurred more frequently than normal south of Iceland and Greenland (Fig. 4b), in the core area of airborne NAWDEX observations. Consistent with the increased frequency of cyclones relative to climatology, the WCB frequency (Fig. 4c) shows increased activity over large parts of the North Atlantic. During the campaign, a succession of events with poleward transport of warm air and ascent of low-PV air into the upper troposphere was observed that appeared to strengthen the downstream anticyclonic anomaly. Most midlatitude cyclones (Fig. 4d) approached Iceland from the southwest, which was favorable for reaching them with Falcon flights from Keflavik. Only a small fraction of the extratropical cyclones moved into central and northern Europe. Six TS occurred during NAWDEX. Ian (12-16 Sep), Julia (13-16 Sep), Karl (14-25 Sep) and Lisa (19-25 Sep) did not exceed TS strength, while Matthew (29 Sep-9 Oct) and Nicole (4-18 Oct) were classified as major hurricanes. Ian, Karl and Nicole underwent ET and moved far into the midlatitudes. TPVs originating over the Canadian polar region were observed twice when they moved southward over the Davis Strait and interacted with the midlatitude waveguide.

North Atlantic weather regimes during NAWDEX show Scandinavian blocking to be the dominant regime (blue line in Fig. 5a), corresponding to the anomalous anticyclone activity over northern Scandinavia (Fig. 4a). In late September the block decayed and a short period with a positive North Atlantic Oscillation (NAO) prevailed before the Scandinavian blocking pattern was again established.

A broad measure of forecast quality during NAWDEX is provided by the anomaly correlation coefficient (ACC) of the mid-tropospheric geopotential height pattern over the eastern North Atlantic as predicted by the ECMWF Integrated Forecasting System (IFS) in fall 2016 (Fig. 5b). Periods of increased 120-h forecast errors and high spread in the ensemble

forecasts are evident, and four of these periods were directly relevant to NAWDEX. Three periods of reduced forecast skill (23 to 27 Sep, 29 Sep to 3 Oct and 5 to 10 Oct) occurred during NAWDEX and two periods of the four were accompanied by a weather regime transition (Fig. 5a). Forecast uncertainty was high on 26 Sep during the onset of a positive NAO phase, and on 1 Oct during the return to the Scandinavian blocking regime. High uncertainty also occurred prior to the campaign, for forecasts initialized between 10 and 14 Sep, again covering a regime transition to Scandinavian blocking. This period affected NAWDEX as it complicated the planning of the transfer flight to Keflavik (IOP 1) five days later.

The progression of weather systems across the North Atlantic during NAWDEX can be conveniently described as a storyline characterized by *upstream triggers*, their dynamic *interaction* with the jet stream, *subsequent development* of disturbances, and downstream *weather impacts* over Europe. Three such sequences occurred completely within the NAWDEX period, and their timespan is indicated by dark grey shading in Fig. 5. In each case, low predictability was found in 5-day forecasts for the eastern North Atlantic initialized within the *trigger* stage (marked by a drop in forecast skill in Fig. 5b), while the final *impact* stage was associated with significant changes in the weather over Europe at the verification time 5 days later. The snapshots for each sequence in Fig. 6 show that the interaction of the trigger disturbance with the waveguide featured intensification of a surface cyclone, with a diabatic contribution consistent with the first three regions identified in the conceptual model presented in Fig. 1. However, the subsequent development and impact stages differed markedly, with the pattern of low PV in the downstream ridge affecting weather even further downstream than suggested by region 4 of Fig. 1. The temporal continuity between the snapshots in Fig. 6 is shown by labeling several coherent long-lived features (identified in the caption). Prominent ridges (R1-R9) along the North Atlantic waveguide are identified as

northward excursions of the jet stream (and the PV gradient). Since each ridge is characterized by low PV air, the associated flow tends to be anticyclonic.

Sequence A is *triggered* by TS Karl leaving the subtropics and moving northwards into the midlatitudes (Fig. 6, panel A.1). Large ensemble spread and changes between consecutive forecast runs showed that the subsequent evolution was very sensitive to uncertainties in the location and timing of the interaction of Karl with ridge R2, the trough upstream, and the associated weak surface cyclone (not shown). The *interaction* that took place was a merging of Karl with a low-level cyclone, leading to rapid re-intensification and the formation of a cyclonic hook at tropopause level separating ridge R2 from the new ridge R3 (Fig. 6 A.2). The ridge-building is intensified by diabatically produced low PV in the WCB outflow. Hence in the subsequent *development*, the jet stream is unusually strong on its southern flank, forming a jet streak that propagates ahead from Karl reaching Scotland the following day (Fig. 6 A.3). The *impact* on European weather occurs through the formation of a new cyclone Walpurga (W in Fig. 6 A.4), which develops to the west of ridge R3 helping to amplify it. Moisture laden air on the western flank of ridge R3 is drawn around the subtropical high. During IOP 5, HALO observed the moist boundary layer in this atmospheric river type flow that extends to Norway where it causes heavy, persistent rainfall, similar to the case studied in Sodemann and Stohl (2013).

Sequence B begins as sequence A ends, in a southwesterly flow situation with a long PV streamer that formed through the merger of the trough west of R3 and the large cut-off feature C (Fig. 6 A.3-A.4). The *trigger* for this sequence appears to follow from the vortex roll-up of the streamer through shear instability, resulting in a new cut-off over Newfoundland (V in Fig. B.1), which then *interacts* and merges with a large-scale trough west of R5 advancing rapidly from the northwest. Note that ridge R5 and its upstream trough wrap up cyclonically during the development so that the trough catches up with the cut-off to the south of R5. The tropopause was very low just in the very center of this system, which therefore has been

named the “stalactite cyclone” (St in Fig. 6 B.2). In the *development* stage, a second cyclone (F in Fig. 6 B.3) intensified rapidly between ridge R6 and the trough to its west. The poleward moving air in R6 crossed Iceland and reinforced the anticyclonic anomaly formed by ridge R5 of the stalactite cyclone. The *impact* of the sequence comes not as a classical severe weather event, but through the establishment of a strong blocking anticyclone over Northern Europe, which persisted for the next two weeks.

Sequence C begins with two upstream *triggers*. A TPV originating in the Canadian Arctic is carried rapidly southeastward on the poleward flank of the jet stream (T in Fig. 6 B.4 and C.1). It is hypothesized that the TPV locally enhanced the cyclonic circulation about the tip of the large-scale trough (T in Fig. 6 C.1), which eventually wrapped cyclonically over Iceland (Fig. 6 C.2). At the same time the remnants of cut-off C appear to be associated with the emergence of a small surface cyclone, which has been named “Sanchez” (S in Fig. 6 C.1). The European dipole block (cf. Rex et al. 1950) is well established at this time so that the ridge R8 and the cyclonic PV anomaly over Iceland are held stationary and a PV filament forms in the deformation region on their western side. The filament is unstable and experiences vortex roll-up, forming three tropopause-level cyclonic vortices. The key *interaction* in this sequence occurs as the low-level cyclone Sanchez passes the southernmost cut-off, but then phase-locks with the central cut-off resulting in baroclinic intensification (S in Fig. 6 C.2). As the sequence *develops*, the resulting cut-off cyclone progresses slowly eastwards (Fig. 6 C.3) and is responsible for some of the most dramatic high *impact* weather during NAWDEX, with heavy precipitation and flooding across southern France and northwestern Italy in the southerly flow ahead of it (Fig. 6 C.4). But this HIW is not the only significant outcome of Sequence C. Returning to stage C.2, ridges R8 and R9 are similar in horizontal extent, but the tropopause is much higher above R9 than R8 (not shown) with the result that the anticyclonic circulation induced by R9 is stronger and R8 is stretched out meridionally between R6 and R9 (Fig. 6 C.3). As NAWDEX draws to a close, the ridge R9

extends rapidly into the Arctic, reinforcing the block and forming a PV anomaly in the shape of the Icelandic character Þ (the first letter of Þor - pronounced Thor - the ancient Norse god of HIW).

It is important to note that the large-amplitude ridge building leading to the Thor block is not R8, which developed as part of the interaction phase of Sequence C, but rather R9, associated with a second cyclone that develops to the west (and which may play a role in the cut-off of Sanchez). Indeed for all three sequences, the development stage leading to weather impacts over Europe appears to be associated with a second cyclone that forms in an environment modified by the interaction of the trigger disturbance with the midlatitude flow, and whose development is difficult to predict because of the low predictability of the environment.

HIGHLIGHTS OF NAWDEX. Observations in NAWDEX were organized in IOPs that focused on key weather systems involved in the longer sequences (Table 5). Several of these IOPs are unprecedented in terms of the phenomena that were sampled or the comprehensive coverage and multi-faceted nature of the measurements. While the analysis of the data is just beginning, a first impression of the results can be obtained from four highlights that illustrate the unique sets of multi-platform and multi-instrument observations that were obtained.

Extratropical transition of TS Karl. The evolution of TS Karl is the central feature of Sequence A discussed above and IOP 4. It was the first extratropical transition sequence that has been observed with research aircraft through at all stages of development, including TS status, ET, re-intensification with impacts on jet stream strength, moisture transport and downstream HIW (Table 5). By flying over the TS and its northwestern flank twice, the SHOUT Global Hawk observed the development stage that occurred far south of the midlatitude jet stream on 22/23 Sep (Fig. 7a), and the ET phase on 24/25 Sep (Fig. 7b). On 26 Sep, HALO observed the interaction with the waveguide and re-intensification phase of the

storm by flying over the cyclone center (Fig. 7c), WCB ascent, the low-valued PV air in the WCB outflow and the dry intrusion (not shown). When Karl moved rapidly towards Scotland, decaying in strong horizontal shear on 27 Sep, IOP 5 focused on the intense jet streak at tropopause level and the strong moisture transport along the equatorward side of the jet with a combination of HALO, FAAM and DLR Falcon flights (Fig. 7d).

IOP 4 will contribute to answer several of the posed research questions (see Table 1 and 3). The large number of dropsonde and special radiosonde measurements that were assimilated into operational forecasts in real-time will provide a basis for observational impact and predictability studies. Detailed airborne remote-sensing observations will allow examination of the role of diabatic processes and their representation in numerical models. Both the synergies of the instruments and the storm-following observational strategy give unprecedented information about this intense and long-lived cyclone and a unique opportunity to analyze forecast error growth due to in situ processes vs. downstream propagation.

Cloud physics in a WCB. IOP 3 focused on observing the vertical cloud structure and cloud microphysical processes in a WCB that was related to the midlatitude cyclone Vladiana south of Iceland and west of Scotland on 23 Sep 2016 (Fig. 7a and Table 5). The WCB transported moist air northeastward just west of the UK as indicated by the low-valued PV air in the upper troposphere (Fig. 8a).

HALO first stayed beneath the NATs at altitudes of ~8 km on the way to the southwestern-most point of the flight (white circle) to begin the first of three sections across the WCB. On this leg to Ireland, 12 dropsondes were released before HALO climbed to ~13 km in Irish airspace. Over northern Ireland, HALO and FAAM joined to perform coordinated remote-sensing and in situ observations of the WCB. HALO measured the WCB by remote-sensing from above while FAAM performed 4 in situ legs at different altitudes to measure

cloud-microphysical parameters inside the WCB. After the coordinated leg, HALO crossed the WCB a third time and observed the outflow of the WCB between Scotland and Iceland.

Figure 8 focuses on the first and second crossing of the WCB. The WALES lidar measured water vapor profiles throughout the troposphere and lower stratosphere in the absence of clouds (Fig. 8b). At the western side of the cross-sections, where HALO was located on the stratospheric side of the waveguide, the post-frontal troposphere was cloud-free except for boundary layer clouds reaching up to 2 km. The water vapor shows high variability, which portrays the dynamically modulated transport of moisture related to cyclone Vladiana. On both crossings of the waveguide, one west-to-east and one east-to-west, a tilted dry layer is visible at altitudes between 5 and 9 km (1110-1125 and 1305-1325 UTC), related to a dry intrusion west of the low-level cold front. The wedge-shaped moist layer on top is associated with high moisture values in the WCB outflow. The second crossing at high altitudes depicts a strong vertical moisture gradient, on top of the elevated moist layer, that marks the tropopause and extends further east into the area where WCB clouds reach high altitudes. A decrease of the tropopause height is detected towards the west on the second leg. The radar shows two vertically (~11.5 km) and horizontally (~400 km) extended and coherent clouds (Fig. 8c) representing the double crossing of the WCB. In between, i.e., on the eastern side of the WCB, cloud tops are lower and the clouds are intermittent. The sharp vertical gradient in radar reflectivity at about 3 km altitude marks the melting layer.

On the second transect the FAAM aircraft performed in situ measurements on flight legs beneath HALO (Figs. 8b,c). HALO met FAAM at the beginning of its second WCB leg (purple diamond marker in Fig. 8c) where FAAM started its lowest leg at about 3 km altitude, just above the melting layer, with subsequent legs at 4, 6 and 7.5 km. The in situ observations show that both mixed-phase and ice-only clouds were encountered during the low-level run, but during the high-level runs only ice was observed. The ice water content (IWC) in the WCB shows maximum values of 0.4 g m^{-3} on the lowest two legs (Fig. 8d). Ice images show

large differences in the form of the particles at different altitudes. On the lowest leg, large aggregates (~6 mm) dominate close to the freezing level, while on the highest level higher concentrations of small irregularly shaped crystals (< 1 mm) prevail.

HALO also observed the interaction of Vladiana's WCB outflow with the jet stream in coordination with the DLR Falcon (not shown). IOP 3 contributes to all research aims (Table 3) and future work on the cloud microphysics observations will investigate, e.g., the correlation of increased IWC with particularly high radar reflectivity. Data from liquid and ice particle size distributions will be used to improve the retrieval of cloud properties from the HALO remote-sensing instruments. Overall, this is a unique set of comprehensive and complementary airborne observations of a WCB, its embedded microphysical processes and its outflow interaction with the jet stream.

Wind observations in the jet stream and outflow of a WCB. Figure 6 (panel B.2) shows the stalactite cyclone that formed previously via merging of two near-surface vorticity maxima with a very intense, small-scale, upper-level PV anomaly south of Newfoundland (not shown). The rapid development of the cyclone occurred in the mid-Atlantic between 30 Sep and 2 Oct. On 2 Oct (IOP 6), a coordinated flight of the DLR and SAFIRE Falcons observed WCB ascent and outflow when the stalactite cyclone was most intense (Fig. 9a). The aircraft flew together to intersect the jet stream on the northwestern edge of ridge R5 wrapping cyclonically around the stalactite cyclone. On a common leg between Iceland and Greenland both aircraft crossed the jet stream (Fig. 9b) and made complementary wind observations (Figs. 9c,d). The DWL lidar on the DLR Falcon observed two wind maxima up to 50 m s^{-1} in cloud-free regions and in optically thin cirrus in the WCB outflow. Complementarily, the SAFIRE radar observed in-cloud winds in the region of WCB ascent in the mid and lower troposphere. Only in dry and aerosol-poor air masses over Greenland, i.e., on the stratospheric side of the waveguide, the combination of both instruments provides poor data coverage. The

SAFIRE Falcon released 9 dropsondes when crossing the jet stream, yielding further profiles of winds, temperature and moisture.

Future research on IOP 6 will be mainly dedicated to predictability issues associated with the blocking formation downstream of the cyclone. The block formed at the time when a loss of predictability in the ECMWF forecasts occurred (Fig. 5b). Winds measured by the two aircraft will help to characterize the role of the WCB outflow in the ridge building. The observed high winds and strong vertical gradients were repeatedly observed on flights across the jet stream with observed maxima up to 80 m s^{-1} and often related to strong vertical wind speed gradients up to $30 \text{ m s}^{-1} \text{ km}^{-1}$. A unique aspect of this example is the benefit of coordinated flights with complementary instruments to address one the key objectives of NAWDEX (Table 1 and 3) – observing the strong wind shear and PV gradients near a WCB outflow.

HIW related to cut-off cyclone Sanchez. Cut-off Sanchez was initiated in the central North Atlantic and reached southern Europe between 12 and 14 Oct 2016 (Fig. 6, Seq. C). On its leading edge moisture was advected northward (Fig. 10a) on 13 Oct when it triggered heavy precipitation and strong winds over France and Italy. The 24-hour accumulated precipitation in the Herault region reached $\sim 250 \text{ mm}$ (Figs. 10b), and wind gusts exceeding 100 km h^{-1} were observed along the French Mediterranean coast (Figs. 10c). As in typical Cevenol episodes, strong southerlies brought warm and moist air from the Mediterranean Sea toward the Massif Central and caused heavy orographic precipitation over the mountain ranges of the Cevennes. Upper-level cut-offs like Sanchez are known to be favorable synoptic conditions for triggering convective mesoscale events (Nuissier et al. 2008), which were intensively studied during the recent international field campaign HyMex (Ducrocq et al. 2016). Part of the air masses responsible for the HIW subsequently reached as far north as the SIRTa site near Paris, causing precipitation in the afternoon of 13 Oct (Fig. 10d). This episode illustrates

one of the key NAWDEX aims (Table 1) to investigate how HIW events over Europe are associated with complex waveguide dynamics (in this case the formation of a PV cut-off) over the upstream North Atlantic. The combination of the ground-based data with NAWDEX observations both from aircraft and from the many additional radiosondes taken during IOP 9 will enable detailed studies of the forecast sensitivity of HIW to upstream initial condition errors.

SUMMARY. NAWDEX was the first field experiment with synergistic airborne and ground-based observations from the entrance to the exit region of the climatological storm track, taken to investigate the role of diabatic processes in altering jet stream disturbances, their development, and effects on HIW downstream.

Because of the operational limitations on research aircraft flights over the North Atlantic and Europe and the need for high-resolution profile measurements of thermal and dynamic properties, NAWDEX focused on airborne remote-sensing observations and the deployment of multiple aircraft. Advanced instruments for remote sensing of wind, water vapor and cloud properties provide an integrated picture of the atmospheric structure in regions where diabatic processes were active, from the synoptic to sub-kilometer scale. The unique combination of the four aircraft and the first deployment of HALO in a campaign focusing on midlatitude dynamics allowed observations in large parts of the North Atlantic. Often, the same weather system could be sampled at different stages of its development, and the interaction of successive weather systems have been observed following the NAWDEX storyline. Additional ground-based observations and an enhanced density of operational radiosonde releases yielded very high coverage with high-resolution vertical profiles from the ground to the lower stratosphere. The region with enhanced atmospheric profiling extended from eastern Canada to most parts of Europe. The coverage and fidelity of the resulting observations will enable future studies to estimate diabatic heating through the use of models and diagnostics

constrained by the NAWDEX observations, particularly in situations when the atmospheric flow is especially sensitive to small changes in diabatic heating. Over the 13 IOPs it was possible to address all original campaign objectives (Table 1 and 3). Table 5 lists a number of particular highlights and “firsts” that have drawn the attention of the NAWDEX scientists.

The success of the observational campaign was possible because of the favorable meteorological conditions, with many cyclones and WCBs in the vicinity of Iceland. Importantly, the NAWDEX period contained episodes of reduced predictability, indicating that uncertainties originating in the estimated atmospheric state and model formulation grew rapidly. The suggestion that these uncertainties spread via their impact on the lifecycle of a “second cyclone” forming to the west, rather than through a process of downstream development, shows that NAWDEX has the potential to make an important contribution to the study of predictability of midlatitude weather and the representation of uncertainty in EPSs. Since there were also episodes of HIW in Europe connected to disturbances of the North Atlantic waveguide, NAWDEX also is a unique opportunity to explore HIW predictability.

To the best of our knowledge, the NAWDEX period provides the most complete set of combined wind, humidity, temperature and cloud profile observations of the North Atlantic jet stream yet accomplished. This dataset will form the basis of detailed case studies and the evaluation of weather and climate prediction models for many years. The widespread coverage of high-resolution multi-variate cross-sections across the jet stream and weather systems developing from one side of the North Atlantic to the other enables examination of the whole chain of processes from the triggering of disturbances on the waveguide to the ultimate impact on weather systems affecting Europe.

ACKNOWLEDGMENTS. Many international institutions contributed to the implementation of NAWDEX and the results presented in this overview article. NAWDEX was only possibly due to the close cooperation of many colleagues including the planning team, the instrument groups and the flight operations (Appendix A). The HALO and DLR Falcon campaign received funding from DLR (project “Klimarelevanz von atmosphärischen Spurengasen, Aerosolen und Wolken” KliSAW), ETH Zurich, NRL Monterrey, the German Science Foundation (DFG; within SPP1294 HALO and SFB/TRR165 Waves to Weather), the European Facility for Airborne Research (EUFAR, project NAWDEX Influence) and the European Space Agency (ESA, providing funds related to the preparation of Aeolus (WindVal II, contract No. 4000114053/15/NL/FF/gp) and EarthCARE (EPATAN, contract No. 4000119015/16/NL/CT/gp)). Special thanks are due to the Max Planck Institute Hamburg for sharing the HALO payload during the NARVAL-II and NAWDEX campaigns in summer and fall 2016 and for their general support of NAWDEX. We are grateful to the DLR flight experiments team; in particular the colleagues of flight operations for the careful preparation and the outstanding support in Keflavik. The authors are grateful to the HALO and Falcon pilots and the technical and sensor team from DLR flight operations for excellent support prior and during NAWDEX. The SAFIRE Falcon contribution to NAWDEX received direct funding from IPSL, Météo-France, INSU-LEFE, EUFAR-NEAREX and ESA (EPATAN, contract No. 4000119015/16/NL/CT/gp). The UK funding for the FAAM aircraft flights, dropsondes and additional radiosondes was provided by the Met Office. Special thanks are due to the SHOUT mission for their open collaboration which enabled coordinated observations of tropical storm Karl. Observations at Capel Dewi were funded by the National Centre for Atmospheric Science. We thank the European Meteorological Service Network (EUMETNET) for funding additional radiosondes and for providing access to the data. Environment and Climate Change Canada provided funding for enhanced radiosonde frequency during the campaign. The Icelandic Meteorological Office and DLR are thanked

665 for providing additional radiosondes from Iceland. We thank ECMWF for providing access to
666 data in the framework of the Support Tool for HALO Missions (SPDEHALO) special project.
667 We gratefully acknowledge many discussions with US colleagues, in particular Chris Davis,
668 Pat Harr and Heather Archambault, during the scientific preparation of NAWDEX. We thank
669 Michael Sprenger (ETH Zurich) for preparing the data for Fig. 4. MB and CMG acknowledge
670 funding from the Swiss National Science Foundation (Projects 200020_165941 and
671 PZ00P2_148177, respectively). JDD and CAR acknowledge the support of the U.S. Navy
672 Chief of Naval Research through the NRL Base Program PE 0601153N, and the ONR PE
673 0602435N.

674 **APPENDIX A: The NAWDEX Team**

675 Table A1

REFERENCES

- Agusti-Panareda, A., C. D. Thorncroft, G. C. Craig, and S. L. Gray, 2004: The extratropical transition of hurricane Irene (1999): A potential-vorticity perspective. *Quart. J. Roy. Meteor. Soc.*, **130**, 1047–1074.
- Archambault, H. M., L. F. Bosart, D. Keyser, and J. M. Cordeira, 2013: A climatological analysis of the extratropical flow response to recurving western North Pacific tropical cyclones. *Mon. Wea. Rev.*, **141**, 2325–2346.
- Bauer, P., A. Thorpe, and G. Brunet, 2015: The quiet revolution of numerical weather prediction. *Nature*, **525**, 47–55.
- Binder, H., M. Boettcher, H. Joos, and H. Wernli, 2016: The role of warm conveyor belts for the intensification of extratropical cyclones in Northern Hemisphere winter. *J. Atmos. Sci.*, **73**, 3997–4020.
- Brogniez, G., C. Pietras, M. Legrand, P. Dubuisson, and M. Haeffelin, 2003: A high-accuracy multiwavelength radiometer for in situ measurements in the thermal infrared. Part II: Behavior in field experiments. *J. Atmos. Oceanic Technol.*, **20**, 1023–1033.
- Browning, K. A., M. E. Hardman, T. W. Harrold, C. W. Pardoe, 1973: Structure of rainbands within a mid-latitude depression. *Quart. J. Roy. Meteor. Soc.*, **99**, 215–231.
- Bruneau, D., J. Pelon, F. Blouzon, J. Spatazza, P. Genau, G. Buchholtz, N. Amarouche, A. Abchiche, and O. Aouji, 2015: 355nm high spectral resolution airborne lidar LNG: system description and first results. *Appl. Opt.*, **54**, 8776–8785.
- Carlson, T. N., 1980: Airflow through midlatitude cyclones and the comma cloud pattern. *Mon. Wea. Rev.*, **108**, 1498–1509.
- Cavallo, S. M., and G. J. Hakim, 2010: Composite structure of tropopause polar cyclones. *Mon. Wea. Rev.*, **138**, 3840–3857.

700 Chaboureaud, J.-P., and C. Claud, 2006: Satellite-based climatology of Mediterranean cloud
 701 systems and their association with large-scale circulation. *J. Geophys. Res.*, **111**, D01102,
 702 doi: 10.1029/2005JD006460.

703 Chagnon, J., S. L. Gray, and J. Methven, 2013: Diabatic processes modifying potential
 704 vorticity in a North Atlantic cyclone. *Quart. J. Roy. Meteor. Soc.*, **139**, 1270–1282.

705 Davis, C. A., M. T. Stoelinga, and Y.-H. Kuo, 1993: The integrated effect of condensation in
 706 numerical simulations of extratropical cyclogenesis. *Mon. Wea. Rev.*, **121**, 2309–2330.

707 Delanoë, J., A. Protat, O. Jourdan, J. Pelon, M. Papazzoni, R. Dupuy, J.-F. Gayet, and C.
 708 Jouan, 2013: Comparison of airborne in-situ, airborne radar-lidar, and spaceborne radar-
 709 lidar retrievals of polar ice cloud properties sampled during the POLARCAT campaign. *J.*
 710 *Atmos. Oceanic Technol.*, **30**, 57–73.

711 Delanoë, J., A. Protat, J.-P. Vinson, W. Brett, C. Caudoux, F. Bertrand, J. Parent du Chatelet,
 712 R. Hallali, L. Barthes, M. Haeffelin, and J.-C. Dupont, 2016: BASTA: A 95-GHz FMCW
 713 Doppler Radar for Cloud and Fog Studies. *J. Atmos. Oceanic Technol.*, **33**, 1023–1038.

714 Doyle, J. D., C. Amerault, C. A. Reynolds, and P. A. Reinecke, 2014: Initial condition
 715 sensitivity and predictability of a severe extratropical cyclone using a moist adjoint. *Mon.*
 716 *Wea. Rev.*, **142**, 320–342.

717 Ducrocq, V., S. Davolio, R. Ferretti, C. Flamant, V. H. Santaner, N. Kalthoff, E. Richard, and
 718 H. Wernli, 2016: Introduction to the HyMeX Special Issue on ‘Advances in understanding
 719 and forecasting of heavy precipitation in the Mediterranean through the HyMeX SOP1
 720 field campaign’. *Quart. J. Roy. Meteor. Soc.*, **142**, 1–6.

721 Ehrlich, A., E. Bierwirth, M. Wendisch, J.-F. Gayet, G. Mioche, A. Lampert, and J.
 722 Heintzenberg, 2008: Cloud phase identification of Arctic boundary-layer clouds from
 723 airborne spectral reflection measurements: test of three approaches. *Atmos. Chem. Phys.*, **8**,
 724 7493–7505

725 ESA European Space Agency, 2008: ADM-Aeolus Science Report, ESA SP-1311, ISBN 978-
 726 92-9221-404-3, 121 pp.

727 Ewald, F., T. Kölling, A. Baumgartner, T. Zinner, and B. Mayer, 2015: Design and
 728 characterization of specMACS, a multipurpose hyperspectral cloud and sky imager. *Atmos.*
 729 *Meas. Tech.*, **9**, 2015–2042.

730 Frame, T. H. A., J. Methven, N. M. Roberts, and H. A. Titley, 2015: Predictability of frontal
 731 waves and cyclones. *Wea. Forecasting*, **30**, 1291-1302.

732 Grams, C. M., H. Wernli, M. Boettcher, J. Campa, U. Corsmeier, S. C. Jones, J. H. Keller, C.
 733 J. Lenz, and L. Wiegand, 2011: The key role of diabatic processes in modifying the upper
 734 tropospheric wave guide: a North Atlantic case-study. *Quart. J. Roy. Meteor. Soc.*, **137**,
 735 2174–2193.

736 Grams, C. M., and S. R. Blumer, 2015: European high-impact weather caused by the
 737 downstream response to the extratropical transition of North Atlantic Hurricane Katia
 738 (2011). *Geophys. Res. Lett.*, **42**, 8738–8748.

739 Grams, C. M., and H. M. Archambault, 2016: The key role of diabatic outflow in amplifying
 740 the midlatitude flow: a representative case study of weather systems surrounding western
 741 North Pacific extratropical transition. *Mon. Wea. Rev.*, **144**, 3847–3869.

742 Gray, S. L., C. Dunning, J. Methven, G. Masato, and J. Chagnon, 2014: Systematic model
 743 forecast error in Rossby wave structure. *Geophys. Res. Lett.*, **41**, 2979–2987.

744 Haeffelin, M., L. Barthès, O. Bock, C. Boitel, S. Bony, D. Bouniol, H. Chepfer, M. Chiriaco,
 745 J. Cuesta, J. Delanoë, P. Drobinski, J.-L. Dufresne, C. Flamant, M. Grall, A. Hodzic, F.
 746 Hourdin, F. Lapouge, Y. Lemaître, A. Mathieu, Y. Morille, C. Naud, V. Noël, W. O'Hirok,
 747 J. Pelon, C. Pietras, A. Protat, B. Romand, G. Scialom, and R. Vautard, 2005: SIRTa, a
 748 ground-based atmospheric observatory for cloud and aerosol research. *Annales*
 749 *Geophysicae*, **23**, 1–23.

750 Hadlock, R., and C. W. Kreitzberg, 1988: The Experiment on Rapidly Intensifying Cyclones
751 over the Atlantic (ERICA) field study: Objectives and plans. *Bull. Amer. Meteor. Soc.*, **69**,
752 1309-1320.

753 Harr, P. A., D. Anwender, and S. C. Jones, 2008: Predictability associated with the
754 downstream impacts of the extratropical transition of tropical cyclones: Methodology and a
755 case study of Typhoon Nabi (2005). *Mon. Wea. Rev.*, **128**, 2613–2633.

756 Harvey, B., J. Methven, and M. Ambaum, 2016: Rossby wave propagation on potential
757 vorticity fronts with finite width. *J. Fluid Mech.*, **794**, 775–797.

758 Illingworth, A.J., H.W. Barker, A. Beljaars, M. Ceccaldi, H. Chepfer, N. Clerbaux, J. Cole, J.
759 Delanoë, C. Domenech, D.P. Donovan, S. Fukuda, M. Hidakata, R.J. Hogan, A.
760 Huenerbein, P. Kollias, T. Kubota, T. Nakajima, T.Y. Nakajima, T. Nishizawa, Y. Ohno,
761 H. Okamoto, R. Oki, K. Sato, M. Satoh, M.W. Shephard, A. Velázquez-Blázquez, U.
762 Wandinger, T. Wehr, and G. van Zadelhoff, 2015: The EarthCARE Satellite: The Next
763 Step Forward in Global Measurements of Clouds, Aerosols, Precipitation, and Radiation.
764 *Bull. Amer. Meteor. Soc.*, **96**, 1311–1332.

765 Joos, H., and H. Wernli, 2012: Influence of microphysical processes on the potential vorticity
766 development in a warm conveyor belt: a case-study with the limited-area model COSMO.
767 *Quart. J. Roy. Meteor. Soc.*, **138**, 407–418.

768 Joos, H., and R. Forbes, 2016: Impact of different IFS microphysics on a warm conveyor belt
769 and the downstream flow evolution. *Quart. J. Roy. Meteor. Soc.*, **142**, 2727–2739.

770 Joly, A., K. A. Browning, P. Bessemoulin, J.-P. Cammas, G. Caniaux, J.-P. Chalon, S. A.
771 Clough, R. Dirks, K. A. Emanuel, L. Eymard, R. Gall, T. D. Hewson, P. H. Hildebrand, D.
772 Jorgensen, F. Lalaurette, R. H. Langland, Y. Lemaître, P. Mascart, J. A. Moore, P. Olag.
773 Persson, F. Roux, M. A. Shapiro, C. Snyder, Z. Toth and R. M. Wakimoto, 1999:

774 Overview of the field phase of the fronts and Atlantic Storm-Track EXperiment (FASTEx)
775 project. *Quart. J. Roy. Meteor. Soc.*, **125**, 3131–3163.

776 Jones, S. C., P. A. Harr, J. Abraham, L. F. Bosart, P. J. Bowyer, J. L. Evans, D. E. Hanley, B.
777 N. Hanstrum, R. E. Hart, F. Lalaurette, M. R. Sinclair, R. K. Smith, and C. Thorncroft,
778 2003: The Extratropical Transition of Tropical Cyclones: Forecast Challenges, Current
779 Understanding, and Future Directions. *Wea. Forecasting*, **18**, 1052–1092.

780 Jones, S. C., and B. Golding, 2015: High Impact Weather Implementation Plan. Available:
781 https://www.wmo.int/pages/prog/arep/wwrp/new/high_impact_weather_project.html

782 Kew, S. F., M. Sprenger, and H. C. Davies, 2010: Potential vorticity anomalies of the
783 lowermost stratosphere: A 10-yr winter climatology. *Mon. Wea. Rev.*, **138**, 1234–1249.

784 Knippertz, P., and J. E. Martin, 2005: Tropical plumes and extreme precipitation in
785 subtropical and tropical West Africa. *Quart. J. Roy. Meteor. Soc.*, **131**, 2337–2365.

786 Krautstrunk, M., and A. Giez, 2012: The transition from FALCON to HALO era airborne
787 atmospheric research. In: *Atmospheric Physics: Background – Methods – Trends Research*
788 *Topics in Aerospace*, U. Schumann, Ed., Springer-Verlag Berlin, 604–629. Heidelberg,
789 609–624, ISBN978-3-642-30182-7.

790 Kuo, Y. H., M. A. Shapiro, and E. G. Donall, 1991: The interaction between baroclinic and
791 diabatic processes in a numerical simulation of a rapidly intensifying extratropical marine
792 cyclone. *Mon. Wea. Rev.*, **119**, 368–384.

793 Lavers, D. A., and G. Villarini, 2013: The nexus between atmospheric rivers and extreme
794 precipitation across Europe. *Geophys. Res. Lett.*, **40**, 3259–3264.

795 Madonna, E., H. Wernli, H. Joos, and O. Martius, 2014: Warm conveyor belts in the ERA-
796 Interim dataset (1979–2010). Part I: Climatology and potential vorticity evolution. *J.*
797 *Climate*, **27**, 3–26.

798 Madonna, E., M. Boettcher, C. M. Grams, H. Joos, O. Martius, and H. Wernli, 2015:
799 Verification of North Atlantic warm conveyor belt outflows in ECMWF forecasts. *Quart.*
800 *J. Roy. Meteor. Soc.*, **141**, 1333–1344.

801 Majumdar, S. J., 2016: A review of targeted observations. *Bull. Amer. Meteor. Soc.*, **97**, 2287-
802 2303.

803 Marksteiner, U., O. Reitebuch, S. Rahm, I. Nikolaus, C. Lemmerz, and B. Witschas, 2011:
804 Airborne direct-detection and coherent wind lidar measurements along the east coast of
805 Greenland in 2009 supporting ESA's Aeolus mission. *Proc. of SPIE Remote Sensing*,
806 **8182**, Prague, 19-23 September, 81810J-1 – 81820J-8.

807 Martínez-Alvarado, O., H. Joos, J. Chagnon, M. Boettcher, S. L. Gray, R. S. Plant, J.
808 Methven, and H. Wernli, 2014: The dichotomous structure of the warm conveyor belt.
809 *Quart. J. Roy. Meteor. Soc.*, **140**, 1809–1824.

810 Martínez-Alvarado, O., E. Madonna, S. L. Gray, and H. Joos, 2016: A route to systematic
811 error in forecasts of Rossby waves. *Quart. J. Roy. Meteor. Soc.*, **142**, 196–210.

812 Martius, O., E. Zenklusen, C. Schwierz, and H. C. Davies, 2006: Episodes of Alpine heavy
813 precipitation with an overlying elongated stratospheric intrusion: A climatology. *Int. J.*
814 *Climatol.*, **26**, 1149–1164.

815 Martius, O., C. Schwierz, and H. C. Davies, 2010: Tropopause-level waveguides. *J. Atmos.*
816 *Sci.*, **67**, 866–879.

817 Massacand, A. C., H. Wernli, and H. C. Davies, 2001: Influence of upstream diabatic heating
818 upon an alpine event of heavy precipitation. *Mon. Wea. Rev.*, **129**, 2822–2828.

819 Mech, M., E. Orlandi, S. Crewell, F. Ament, L. Hirsch, M. Hagen, G. Peters, and B. Stevens,
820 2014: HAMP – the microwave package on the High Altitude and LOng range research
821 aircraft (HALO). *Atmos. Meas. Tech.*, **7**, 4539–4553.

822 Methven, J., 2015: Potential vorticity in warm conveyor belt outflow. *Quart. J. Roy. Meteor.*
823 *Soc.*, **141**, 1065–1071.

824 Michel, C., and G. Rivière, 2011: A link between Rossby wave breakings and weather regime
825 transitions. *J. Atmos. Sci.*, **68**, 1730–1748.

826 Michelangeli, P., R. Vautard, and B. Legras, 1995: Weather regimes: Recurrence and quasi
827 stationarity. *J. Atmos. Sci.*, **52**, 1237–1256.

828 Nuissier, O., V. Ducrocq, D. Ricard, C. Lebeaupin, and S. Anquetin, 2008: A numerical study
829 of three catastrophic precipitating events over southern France. I: Numerical framework
830 and synoptic ingredients. *Quart. J. Roy. Meteor. Soc.*, **134**, 111–130.

831 Parsons, D.B., M. Beland, D. Burridge, P. Bougeault, G. Brunet, J. Caughey, S.M. Cavallo,
832 M. Charron, H.C. Davies, A.D. Niang, V. Ducrocq, P. Gauthier, T.M. Hamill, P.A. Harr,
833 S.C. Jones, R.H. Langland, S.J. Majumdar, B.N. Mills, M. Moncrieff, T. Nakazawa, T.
834 Paccagnella, F. Rabier, J. Redelsperger, C. Riedel, R.W. Saunders, M.A. Shapiro, R.
835 Swinbank, I. Szunyogh, C. Thorncroft, A.J. Thorpe, X. Wang, D. Waliser, H. Wernli, and
836 Z. Toth, 2017: THORPEX research and the science of prediction. *Bull. Amer. Meteor.*
837 *Soc.*, **98**, 807–830.

838 Pfahl, S., C. Schwierz, M. Croc-Maspoli, C.M. Grams, and H. Wernli, 2015: Importance of
839 latent heat release in ascending air streams for atmospheric blocking, *Nature Geosci.*, **8**,
840 610–614.

841 Pomroy, H. R., and A. J. Thorpe, 2000: The evolution and dynamical role of reduced upper-
842 tropospheric potential vorticity in intensive observing period one of FASTEX. *Mon. Wea.*
843 *Rev.*, **128**, 1817–1834.

844 Protat A., J. Pelon, J. Testud, N. Grand, P. Delville, P. Laborie, J.-P. Vinson, D. Bouniol, D.
845 Bruneau, H. Chepfer, J. Delanoe, M. Haeffelin, V. Noel, and C. Tinel, 2004: Le projet

846 RALI : Combinaison d'un radar nuage et d'un lidar pour l'étude des nuages faiblement
847 précipitants, *La Météorologie*, n° 47, 23–33.

848 Rabier F., P. Gauthier, C. Cardinali, R. Langland, M. Tsyrlnikov, A. Lorenc, P. Steinle, R.
849 Gelaro, K. Koizumi, 2008: An update on THORPEX-related research in data assimilation
850 and observing strategies. *Nonlin. Proc. Geophys.*, **15**, 81–94.

851 Rautenhaus, M., G. Bauer, and A. Dörnbrack, 2012. A web service based tool to plan
852 atmospheric research flights. *Geosci. Model Dev.*, **5**, 55–71.

853 Rautenhaus, M., M. Kern, A. Schäfler, and R. Westermann, 2015a: Three-dimensional
854 visualization of ensemble weather forecasts - Part 1: The visualization tool Met.3D
855 (version 1.0). *Geosci. Model Dev.*, **8**, 2329–2353.

856 Rautenhaus, M., C. M. Grams, A. Schäfler, and R. Westermann, 2015b: Three-dimensional
857 visualization of ensemble weather forecasts - Part 2: Forecasting warm conveyor belt
858 situations for aircraft-based field campaigns. *Geosci. Model Dev.*, **8**, 2355–2377.

859 Reitebuch, O., C. Lemmerz, E. Nagel, U. Paffrath, Y. Durand, M. Endemann, F. Fabre, and
860 M. Chaloupy, 2009: The airborne demonstrator for the direct-detection Doppler wind lidar
861 ALADIN on ADM-Aeolus: I. Instrument design and comparison to satellite instrument. *J.*
862 *Atmos. Oceanic Technol.*, **26**, 2501–2515.

863 Rex, D. F., 1950: Blocking action in the middle troposphere and its effect upon regional
864 climate. I. An aerological study of blocking action. *Tellus*, **2**, 196–211.

865 Richardson, D., J. Bidlot, L. Ferranti, A. Ghelli, T. Haiden, T. Hewson, M. Janousek, F.
866 Prates, and F. Vitart, 2012: Verification statistics and evaluations of ECMWF forecasts in
867 2011-2012. ECMWF Technical Memorandum, 688.

868 Riemer, M., and S. C. Jones, 2010: The downstream impact of tropical cyclones on a
869 developing baroclinic wave in idealized scenarios of extratropical transition. *Quart. J. Roy.*
870 *Meteor. Soc.*, **136**, 617–637.

871 Rodwell, M. J., L. Magnusson, P. Bauer, P. Bechtold, M. Bonavita, C. Cardinali, M.
 872 Diamantakis, P. Earnshaw, A. Garcia-Mendez, L. Isaksen, E. Källén, D. Klocke, P. Lopez,
 873 T. McNally, A. Persson, F. Prates, and N. Wedi,, 2013: Characteristics of occasional poor
 874 medium-range weather forecasts for Europe. *Bull. Amer. Meteor. Soc.*, **94**, 1393–1405.
 875 Schäfler, A., and F. Harnisch, 2015: Impact of the inflow moisture on the evolution of a warm
 876 conveyor belt. *Quart. J. Roy. Meteor. Soc.*, **141**, 299–310.
 877 Schäfler, A., M. Boettcher, C. M. Grams, M. Rautenhaus, H. Sodemann, and H. Wernli, 2014:
 878 Planning of aircraft measurements within a warm conveyor belt. *Weather*, **69**, 161–166.
 879 Schwierz, C., S. Dirren, and H. C. Davies, 2004: Forced waves on a zonally aligned jet
 880 stream. *J. Atmos. Sci.*, **61**, 73–87.
 881 Selz, T., and G. C. Craig, 2015: Upscale error growth in a high-resolution simulation of a
 882 summertime weather event over Europe. *Mon. Wea. Rev.*, **143**, 813–827.
 883 Sodemann, H., and A. Stohl, 2013: Moisture origin and meridional transport in Atmospheric
 884 Rivers and their association with multiple cyclones, *Mon. Wea. Rev.*, **141**, 2850–2868.
 885 Spensberger, C., and T. Spengler, 2014: A new look at deformation as a diagnostic for large-
 886 scale flow. *J. Atmos. Sci.*, **71**, 4221–4234.
 887 Sprenger, M., G. Fragkoulidis, H. Binder, M. Croci-Maspoli, P. Graf, C. M. Grams, P.
 888 Knippertz, E. Madonna, S. Schemm, B. Skerlak, and H. Wernli, 2017: Global
 889 climatologies of Eulerian and Lagrangian flow features based on ERA-Interim reanalyses.
 890 *Bull. Amer. Meteor. Soc.*, **98**, 1739–1748.
 891 Szunyogh, I., Z. Toth, R. E. Morss, S. J. Majumdar, B. J. Etherton, and C. H. Bishop, 2000:
 892 The effect of targeted dropsonde observations during the 1999 Winter Storm
 893 Reconnaissance Program. *Mon. Wea. Rev.*, **128**, 3520–3537.
 894 Teubler, F., and M. Riemer, 2016: Dynamics of Rossby Wave Packets in a Quantitative
 895 Potential Vorticity–Potential Temperature Framework. *J. Atmos. Sci.*, **73**, 1063–1081.

896 Thorpe, A. J., 2004: Weather forecasting: A centenary perspective. *Weather*, **59**, 332–335.

897 Vaughan, G., J. Methven, D. Anderson, B. Antonescu, L. Baker, T.P. Baker, S.P. Ballard,
898 K.N. Bower, P.R. Brown, J. Chagnon, T.W. Choularton, J. Chylik, P.J. Connolly, P.A.
899 Cook, R.J. Cotton, J. Crosier, C. Dearden, J.R. Dorsey, T.H. Frame, M.W. Gallagher, M.
900 Goodliff, S.L. Gray, B.J. Harvey, P. Knippertz, H.W. Lean, D. Li, G. Lloyd, O. Martínez–
901 Alvarado, J. Nicol, J. Norris, E. Öström, J. Owen, D.J. Parker, R.S. Plant, I.A. Renfrew,
902 N.M. Roberts, P. Rosenberg, A.C. Rudd, D.M. Schultz, J.P. Taylor, T. Trzeciak, R. Tubbs,
903 A.K. Vance, P.J. van Leeuwen, A. Wellpott, and A. Woolley, 2015: Cloud banding and
904 winds in intense European cyclones: results from the DIAMET project. *Bull. Amer.*
905 *Meteor. Soc.*, **96**, 249–265.

906 Weissmann, M., R. Busen, A. Dörnbrack, S. Rahm, and O. Reitebuch, 2005: Targeted
907 observations with an airborne wind lidar. *J. Atmos. Oceanic Technol.*, **22**, 1706–1719.

908 Weissmann, M., F. Harnisch, C. Wu, P. Lin, Y. Ohta, K. Yamashita, Y. Kim, E. Jeon, T.
909 Nakazawa, and S. Aberson, 2011: The Influence of Assimilating Dropsonde Data on
910 Typhoon Track and Midlatitude Forecasts. *Mon. Wea. Rev.*, **139**, 908–920.

911 Wendisch, M., D. Müller, D. Schell, and J. Heintzenberg, 2001: An air-borne spectral
912 albedometer with active horizontal stabilization. *J. Atmos. Oceanic Technol.*, **18**, 1856–
913 1866.

914 Wendisch, M., U. Pöschl, M. O. Andreae, L. A. T. Machado, R. Albrecht, H. Schlager, D.
915 Rosenfeld, S. T. Martin, A. Abdelmonem, A. Afchine, A. Araujo, P. Artaxo, H. Aufmhoff,
916 H. M. J. Barbosa, S. Borrmann, R. Braga, B. Buchholz, M. A. Cecchini, A. Costa, J.
917 Curtius, M. Dollner, M. Dorf, V. Dreiling, V. Ebert, A. Ehrlich, F. Ewald, G. Fisch, A. Fix,
918 F. Frank, D. Fütterer, C. Heckl, F. Heidelberg, T. Hüneke, E. Jäkel, E. Järvinen, T. Jurkat,
919 S. Kanter, U. Kästner, M. Kenntner, J. Kesselmeier, T. Klimach, M. Knecht, R. Kohl, T.
920 Kölling, M. Krämer, M. Krüger, T. C. Krisna, J. V. Lavric, K. Longo, C. Mahnke, A. O.
921 Manzi, B. Mayer, S. Mertes, A. Minikin, S. Molleker, S. Münch, B. Nillius, K.

Pfeilsticker, C. Pöhlker, A.-E. Roiger, D. Rose, D. Rosenow, D. Sauer, M. Schnaiter, J. Schneider, C. Schulz, R. A. F. de Souza, A. Spanu, P. Stock, D. Vila, C. Voigt, A. Walser, D. Walter, R. Weigel, B. Weinzierl, F. Werner, M. A. Yamasoe, H. Ziereis, T. Zinner, M. Zöger, 2016: The ACRIDICON-CHUVA campaign: Studying tropical deep convective clouds and precipitation over Amazonia using the new German research aircraft HALO. *Bull. Am. Meteor. Soc.*, **97**, 10, 1885–1908.

Wernli, H., and H. C. Davies, 1997: A Lagrangian-based analysis of extratropical cyclones. I. The method and some applications. *Quart. J. Roy. Meteor. Soc.*, **123**, 467–489.

Wernli, H., 1997: A Lagrangian-based analysis of extratropical cyclones. II. A detailed case-study. *Quart. J. Roy. Meteor. Soc.*, **123**, 1677–1706.

Wernli, H., and M. Sprenger, 2007: Identification and ERA15 climatology of potential vorticity streamers and cut-offs near the extratropical tropopause. *J. Atmos. Sci.*, **64**, 1569–1586.

Wirth, M., A. Fix, P. Mahnke, H. Schwarzer, F. Schrandt, and G. Ehret, 2009: The airborne multi-wavelength water vapor differential absorption lidar WALES: system design and performance. *Appl. Phys. B*, **96**, 201–213.

Witschas, B., S. Rahm, A. Dörnbrack, J. Wagner, and M. Rapp, 2017: Airborne wind lidar measurements of vertical and horizontal winds for the investigation of orographically induced gravity waves. *J. Atmos. Oceanic Technol.*, **34**, 1371–1386.

Wulfmeyer, V., and Co-authors, 2011: The Convective and Orographically induced Precipitation Study (COPS): The scientific strategy, the field phase, and research highlights. *Quart. J. Roy. Meteor. Soc.*, **137**, 3–30.

Zhang, F., N. Bei, R. Rotunno, C. Snyder, and C. C. Epifanio, 2007: Mesoscale predictability of moist baroclinic waves: Convection-permitting experiments and multistage error growth dynamics. *J. Atmos. Sci.*, **64**, 3579–3594.

Sidebar 1: Active remote-sensing observations for future satellite missions Aeolus and EarthCARE. HALO and the SAFIRE and DLR Falcon aircraft were equipped with remote-sensing instruments that are specifically relevant for the future Earth Explorer satellite missions EarthCARE (Illingworth et al. 2015) and Aeolus (ESA 2008) of the European Space Agency (ESA). NAWDEX observations, through coordinated flights of multiple aircraft and of aircraft with satellite overpasses, provide data from comparable airborne instruments for the preparation and future validation of these satellite instruments.

HALO was equipped with the High Spectral Resolution (HSRL, 532 nm) and water vapor Differential Absorption Lidar (DIAL) WALES, the HALO Microwave Package (HAMP) with a 35.2 GHz cloud radar and microwave radiometers, the cloud spectrometer (specMACS) and the visible to near-infrared SMART instrument (Table 2). The French Falcon was equipped with the RADar-Lidar (RALI, Protat et al. 2004) payload consisting of the 94 GHz cloud radar RASTA and the UV High Spectral Resolution lidar LNG (Table 2). These aircraft provide the most complete instrumentation package available at the European level to mimic upcoming EarthCARE measurements and thus provide valuable data for preparing the EarthCARE mission and for future validation. Coordinated flights with both aircraft as well as CALIPSO-Cloudsat underpasses during NAWDEX delivered independent measurements for testing EarthCARE L2 algorithms at different wavelengths and for performing a first rehearsal of the validation/calibration for EarthCARE.

Figure SB1a illustrates the complementary character of lidar and radar measurements taken during the HALO research flight on 1 Oct 2016. Optically thin ice clouds at cloud top are only visible in the lidar measurements (green marked curtain), while optically thicker cloud regions are only visible in cloud radar measurements (red marked curtain).

The DLR Falcon was equipped with a Doppler wind lidar (DWL) payload consisting of the A2D direct-detection DWL and a 2- μ m scanning coherent/heterodyne detection DWL. The A2D is the prototype of the satellite-borne wind lidar instrument on Aeolus and provides

973 range-resolved line-of-sight wind speeds with high data coverage by exploiting both
974 molecular and particulate backscatter return. With a view to the pre-launch activities for the
975 upcoming Aeolus mission, NAWDEX offered the opportunity to extend the A2D dataset and
976 to perform wind measurements in dynamically complex scenes, including strong wind shear
977 and varying cloud conditions, as well as multiple instrument calibrations, which are a
978 prerequisite for accurate wind retrieval. RALI on board the SAFIRE Falcon complemented
979 the A2D instrument with wind measurements in clouds and aerosol-rich layers.

980 Figure SB1b shows collocated wind observations from the A2D and the 2- μ m DWL from
981 a flight of the DLR Falcon east of Iceland on 4 Oct 2016. The good vertical coverage, limited
982 only by a dense cloud layer, is achieved by combining complementary information from both
983 aerosol backscatter (A2D Mie channel and 2- μ m DWL) and molecular backscatter (A2D
984 Rayleigh channel).

Sidebar 2: Forecast products to investigate forecast uncertainty. NAWDEX focused on weather phenomena that are poorly represented in NWP, so a strong effort to estimate forecast uncertainty was essential for the planning of the IOPs. Deterministic forecasts from the European Centre for Medium-Range Weather Forecasts (ECMWF), the UK Met Office, the Naval Research Laboratory (NRL), Météo France, the Icelandic Meteorological Office (IMO) and the Danish Meteorological Institute (DMI) were available. Additionally, ensemble forecasts from the ECMWF, Met Office (MOGREPS-G) and Météo France (PEARP short-range ensemble) played an essential role.

Each day a standard set of synoptic charts and tailored weather products (e.g. PV on isentropic surfaces and WCB trajectories) were produced using a common map projection and pre-defined cross sections. Ensemble diagnostics of mean and spread of several variables, as well as tailored ensemble forecast products for NAWDEX-relevant features (e.g., WCB and cyclone frequencies, and tropopause height) were created. These forecast products were provided via web sites. In addition, an interactive web interface allowed the flight planning team to compute backward and forward trajectories from planned flight tracks, facilitating the planning of flights to attempt Lagrangian re-sampling of air masses.

Flight planning typically requires cross-section information, e.g., to obtain an accurate picture of tropopause height, winds speeds and cloud layers, and to assess forecast uncertainties along hypothetical flight routes. The NAWDEX community had access to special flight planning tools that allowed an interactive visualization of forecast products. Central to forecasting and flight planning operations was the "Mission Support System" (MSS; Rautenhaus et al. 2012). In addition, the interactive 3D forecast tool "Met.3D" (Rautenhaus et al. 2015a) provided specialized forecast products. Two workstations were set up at the operation center in Keflavik to run Met.3D and enable the novel ensemble forecasting workflow described in Rautenhaus et al. (2015b). Ensemble forecasts by ECMWF could be interactively analyzed in combined 2D/3D depictions. WCB trajectories and derived

probabilities of WCB occurrence could be combined with additional forecast information. The ability of Met.3D to interactively navigate the ensemble data proved particularly useful, facilitating analysis of the uncertainty for features such as the predicted tropopause position.

Figure SB2 shows an example of forecast products used for planning the IOP 4 flight. The +60 h deterministic IFS forecast shows ex-TS Karl as a deep surface cyclone south of Greenland (Fig. SB2a) with cyclonically wrapped PV contours resulting from an advection of low-PV air to upper levels in the outflow of a WCB (not shown). High WCB probabilities with two distinct maxima north and east of Karl indicate that the location of the tropopause and WCB outflow is predicted with high certainty (Fig. SB2b). Images from Met.3D (Fig. SB2c,d) show the relation between the jet stream, WCB and tropopause in the ECMWF ensemble mean along cross sections intersecting the waveguide and the WCB east of the surface cyclone. A cross section with ensemble mean PV (Fig. SB2c) shows a low tropopause north of the jet (depicted by an isosurface of wind speed), whereas a high tropopause appears to the south. This coincides with high probabilities of WCB (Fig. SB2d). WCB trajectories of a selected ensemble member show two distinct branches (Fig. SB2d). One branch wraps cyclonically around the cyclone and features a lower outflow compared to the second branch, which follows anticyclonic pathways at higher elevations, contributing to the elevated WCB probability maximum there. Real-time adjoint products from COAMPS (Doyle et al. 2014) were used to identify regions of initial condition sensitivity. At 12 UTC 24 Sep, the maximum moisture sensitivity is located in the low- to mid-levels and positioned along the eastern portion of TS Karl (Fig. SB2e). The adjoint sensitivity is computed using a kinetic energy response function located in a box ($450 \times 600 \text{ km}^2$ in the horizontal and extending from the surface to 700 hPa) centered on the ascending WCB at the 48 h forecast time at 12 UTC 26 Sep when the IOP 4 flights were planned. Optimal perturbations derived from the adjoint sensitivity show an increase of wind speeds from 30 m s^{-1} to over 45 m s^{-1} in the WCB

1036 highlighting the importance of the mid-level moisture associated with Karl (48 h prior) for the
1037 intensification of the WCB.

Table 1: NAWDEX research aims and science goals. Region numbers refer to Fig. 1			
Aim Nr	Topic	Science Goals	Region
1	Moisture structure in the boundary layer	<ul style="list-style-type: none"> • Characterization of low-level moisture in atmospheric rivers and WCB inflow regions • Investigation of impact of low-level moisture on downstream weather evolution 	1,(2,3),4
2	Mixed phase and cirrus clouds	<ul style="list-style-type: none"> • In situ and remote sensing measurements of cloud properties and meteorological parameters during WCB ascent and outflow • Comparison of observations and models to quantify latent and radiative heating/cooling in and below WCB • Role of slantwise ascent vs. embedded convection in WCB • Characterization of vertical moisture gradient and cirrus structure in WCB outflow, and effects on radiation 	2, 3
3	Potential vorticity	<ul style="list-style-type: none"> • Quantitative estimate of PV from observations • Verification of PV structures, PV gradients and jet stream winds in numerical models • Structure of negative PV anomalies in WCB outflows and upper tropospheric ridges • Role of divergent outflow of WCBs for ridge amplification • Spatial distribution of turbulence in the free atmosphere and relationship to jet stream and PV structures 	3
4	Tropopause waveguide, predictability and consequences for HIW	<ul style="list-style-type: none"> • Relevance of amplifying small errors at tropopause level for uncertainty in surface weather downstream • Influence of observations within and outside of diabatically active regions on the predictability of downstream HIW 	3, 4
5	Instrument-driven aims	<ul style="list-style-type: none"> • Comparison of measured radiances and retrieved cloud optical properties between SMART-HALO and specMACS • Cloud regime characterization in midlatitude cyclones and analysis of model representation at different resolutions • Radiometer retrieval development for profiles and hydrometeor paths using instrument synergies • Validation of Aeolus calibration and wind retrieval algorithms • Intercomparison of wind and aerosol products from different instruments on DLR and SAFIRE Falcon • First test of EarthCARE calibration and validation strategy 	2,3

Table 2: Aircraft and instrumentation for NAWDEX and contribution to research aims (Table1)

Aircraft	Instruments	Measured and derived properties	Aim Nr
HALO	HAMP (HALO Microwave Package) Microwave radiometer with 26 channels spanning the frequency range from 22 to 183 GHz, and Ka-band (35.2 GHz) cloud radar (Mech et al. 2014)	Radiometers: Integrated water vapor, temperature and humidity profiles, liquid and ice water path Radar: profiles of radar reflectivity, depolarization ratio, vertical velocity	2,3,5
	WALES (Water Vapor Lidar Experiment in Space): Four-wavelength Differential Absorption (DIAL) and High Spectral Resolution Lidar (HSRL) (Wirth et al. 2009)	Profiles of water vapor, backscatter coefficient lidar/color ratio, particle linear depolarization ratio, particle extinction coefficient	1,2,4,5
	SMART (Spectral Modular Airborne Radiation Measurement System): Passive cloud spectrometer (Wendisch et al., 2001; Ehrlich et al., 2008)	Spectral nadir radiance, spectral upward and downward irradiance (300-2200 nm), cloud top albedo, cloud thermodynamic phase, cloud optical thickness, effective radius, cloud cover / statistics	2,4,5
	specMACS (Cloud spectrometer of the Munich Aerosol Cloud Scanner): Imaging cloud spectrometer plus 2D RGB camera (+/-35° fov) (Ewald et al. 2015)	Spectral radiance (400-2500 nm), push-broom imaging at nadir and +/- 17° across track, cloud thermodynamic phase, liquid and ice optical thickness, particle size, cloud cover	2,4,5
	Bahamas (Basic HALO Measurements and Sensor System)	In situ observations of pressure, temperature, wind, humidity, TAS aircraft position, attitude, heading, altitude	3,4,5
	Drosondes Vaisala RD94	Temperature, humidity and wind profiles	1,2,3,4,5
DLR Falcon	A2D (ALADIN Airborne demonstrator): direct-detection DWL (Reitebuch et al. 2009, Marksteiner et al. 2011)	Profiles of line-of-sight wind speed and aerosol/cloud layers(20° off-nadir)	3,5
	2-µm scanning coherent/heterodyne detection DWL (Weissmann et al. 2005; Witschas et al. 2017)	Vertical profiles of line-of-sight wind speed, horizontal wind vectors, and aerosol/cloud layers	3,5
	Basic in situ measurements	In situ observations of pressure, temperature, wind, humidity, TAS aircraft position, attitude, heading, altitude	3,4,5
SAFIRE Falcon	RASTA (RAdar SysTem Airborne):	Doppler velocity and reflectivity from three antennas (including spectral width), cloud and	2,3,5

	95-GHz Doppler cloud radar (Delanoë et al 2013)	precipitation microphysics (ice and liquid water content), dynamics (horizontal and vertical wind)	
	LNG (Leandre New Generation): high-spectral-resolution lidar (Bruneau et al. 2015)	Three-wavelength (1064, 532, and 355 nm) backscatter lidar with polarization analysis at 355 nm, High Spectral Resolution capability including Doppler measurement, based on a Mach–Zehnder Interferometer, at 355 nm. Radiative properties and dynamics of cloud and aerosol	2,3,5
	CLIMAT infrared radiometer (Brogniez et al. 2003)	Radiances measured simultaneously in three narrowband channels centered at 8.7, 10.8, and 12.0 μm	2,4,5
	Dropsondes Vaisala RD94	Temperature, humidity and wind profiles	1,2,3,4,5
	Aircraft in situ measurements	In situ observations of pressure, temperature, wind, humidity, TAS aircraft position, attitude, heading, altitude	3,4,5
FAAM BAe 146	In situ temperature, Buck CR-2 and WVSS-2 hygrometers, two turbulence probes	Temperature, humidity, and wind and turbulent fluxes	2,3
	PCASP (aerosol size probe), CDP (scattering cloud droplet probe), CIP-15, CIP-100 (cloud imaging probes)	Cloud particle size spectrum: 2 μm -6 mm diameter, cloud droplet spectrum: 3-50 μm	2,5
	Nevzorov hot wire probe	Ice/Liquid water content	2,5
	TECO 49C UV analyser, Aerolaser AL5002, Los Gatos Fast Greenhouse Gas analyser	O ₃ , CO, CH ₄ , CO ₂	2
	Lidar: downward-pointing Leosphere ALS450 (355 nm, scattering and depolarization)	Position of different atmospheric layers below the aircraft (clear air, aerosols, cloud tops)	2
	ISMAR (International Sub-Millimetre Airborne Radiometer)	Passive radiometer with polarization and multiple channels (118, 243 (V/H), 325, 424, 448, 664 (V/H) and 874 GHz (V/H)) (IOP 11 only)	5
	MARSS (Microwave Airborne Radiometer Scanning System)	Scanning microwave radiometer operating at AMSU-B channels 16-20 (89-183GHz) and pointing both upward and downward (IOP 11 only)	5
	Dropsondes Vaisala RD94	Temperature, humidity and wind profiles	1,2,3,4,5

Table 3: IOPs, key weather systems and associated flights together with the number of dropsondes from all aircraft. As some of the long-range flights of HALO were related to different weather systems, dropsondes were assigned to the respective IOP. Aims numbered according to Table 1 show contribution to NAWDEX science goals.

IOP	Period	Key Weather systems	Date	HALO	DLR Falcon	SAFIRE Falcon	FAAM BAe 146	Drops	Aim Nr
1	16-17 Sep	Outflow of ex-TC Ian , low predictability case	17 Sep	RF01	RF01/02			10	2,3,4,5
2	21-22 Sep	WCB ascent and outflow of extratropical cyclone Ursula	21 Sep	RF02	RF03			14	2,3,4,5
3	23-25 Sep	WCB ascent of extratropical cyclone Vladiana	23 Sep	RF03	RF04		RF01 (B980)	32	1,2,3,4,5
4	22-28 Sep	Re-intensification phase of ex-TS Karl and jet streak forming downstream	26 Sep	RF04				25	2,3,4,5
			27 Sep	RF05	RF05		RF02 (B981)	22	
5	26-29 Sep	Strong WV transport of extratropical cyclone Walpurga leading to HIW in Scandinavia	27 Sep	RF05				20	1,3,4
6	1-5 Oct	Stalactite cyclone and low predictability over Europe	1 Oct	RF06		RF05		3	2,3,4,5
			2 Oct		RF07	RF06/07		9	
7	4-5 Oct	Strong extratropical cyclone originating as frontal wave near Newfoundland	4 Oct		RF08/09	RF08		5	2,3,4
			5 Oct			RF09		4	
8	6-9 Oct	TPV near Newfoundland and downstream forming cyclone	6 Oct	RF07				20	4,5
			7 Oct			RF10		7	
			9 Oct	RF08	RF10	RF11/12		9	
			10 Oct			RF13		6	
9	9-14 Oct	PV cut-off cyclone Sanchez & downstream impact over the Mediterranean	9 Oct	RF08					2,3,4,5
			10 Oct	RF09				20	
10	12-15 Oct	Formation and extension of tropopause ridge Thor and the Scandinavian anticyclone	11 Oct			RF14		4	3,4,5
			12 Oct			RF15		8	
			13 Oct	RF10		RF16		26	
			15 Oct	RF12				12	
11	14 Oct	Radar and lidar mission for instrument comparisons and satellite underflights	14 Oct	RF11		RF17/18	RF03 (B984)	15	5
12	15 Oct	TPV over Baffin Island	15 Oct	RF12					4, 5
13	18 Oct	PV streamer over UK	18 Oct	RF13	RF13/14			16	2,3
		Instrument and calibration flights	28 Sep		RF06				5
			15 Oct		RF11/12				
			16 Oct			RF19		2	
			22 Oct		RF15/16				

Table 4: NAWDEX IOPs and periods of increased ground-based observation activities.		
IOP	Additional Observations	Period
1	Radiosondes from UK, Torshavn and Iceland for a temporal sequence of the arrival of outflow of ex-TC Ian as it extends northeastwards.	16-17 Sep
2	Radiosondes from UK, Iceland and eastern Greenland for a time series during arrival and passage of cyclone Ursula.	21-22 Sep
3	Radiosondes from northern UK to observe rapidly intensifying frontal cyclone Vladiana with strong WCB and ridge building.	23-25 Sep
4	Radiosondes around the northern North Atlantic and Scandinavia to observe the structure and evolution of ex-TS Karl and to observe GWs above Iceland at the jet stream. Jet streak maximum passes directly above MST radar wind profiler on South Uist, Scotland.	26-28 Sep
5	Radiosondes in UK and southern Scandinavia to observe the strong water vapor transport and related HIW. Passage of jet stream over Capel Dewi.	27-29 Sep
6	Radiosondes northwest of Iceland to observe ridge building in relation to the stalactite cyclone. Radiosondes over southern Europe to observe a cut-off downstream. Radiosondes at Iceland to observe GWs in the stratosphere.	1-5 Oct
8	Radiosondes over Iceland and eastern Greenland to observe WCB ascent and cyclone structure. Observation of orographic GWs above Iceland.	6-9 Oct
9	Radiosondes from the western Mediterranean, at Capel Dewi and at SIRTa to observe cut-off Sanchez and related HIW. Passage of outflow from Sanchez over MST radar at Capel Dewi. Radiosondes above Iceland to observe strong GW activity in the stratosphere.	10-14 Oct
8,10	Radiosondes over North Atlantic to obtain a time series of the vertical structure of ridge Thor. MST radar wind observations at Andøya, Norway.	10-15 Oct
11	Radiosondes at SIRTa to observe the downstream impact.	15-16 Oct

Table 5: NAWDEX observational highlights		
IOP	Period	Specific aspects of the observations
3	23-25 Sep	Coordinated flights to observe the cloud structure and cloud physics in the WCB ascent related to cyclone Vladiana and the interaction of the WCB outflow with the jet stream.
4	22-28 Sep	First ever observations of a TS from tropical phase and ET (SHOUT observations) through midlatitude re-intensification, jet streak formation, ridge enhancement and HIW over Scandinavia (NAWDEX observations)
5	26-29 Sep	Large-scale strong moisture transport in an atmospheric river type flow upstream of cyclone Walpurga causing HIW over Scandinavia
6	1-5 Oct	Lowest predictability case with observation of the WCB ascent and outflow of the stalactite cyclone and the subsequent influence in the onset of the European block.
8/12	26-29 Sep, 15 Oct	First ever airborne observation of temperature, wind and moisture structure of two TPV events in a phase when they interacted with the midlatitude waveguide
9	9-14 Oct	Roll-up of the positive PV filament giving rise to mesocyclone Sanchez connected to HIW in France and Italy
10	12-15 Oct	Low PV ridge builds and extends into the Arctic reinforcing the anticyclonic part of the block. Profile observations characterizing the low PV anomaly structure.
11	14 Oct	Coordination of three aircraft and joint underflight of the Calipso/Cloudsat satellite path to exploit instrument synergies of radar, lidar and radiometer instruments

Table A1. Overview of the NAWDEX team members and their roles in the campaign.			
Organization	Country	Participants	Role
Monash University	Australia	Julian Quinting	Flight planning team
Environment and Climate Change Canada (ECCC)	Canada	Ron McTaggart-Cowan	PI Canadian radiosondes, science team
Division Technique, INDU	France	Frédéric Blouzon	RALI team
Institut Pierre Simon Laplace	France	Jean-Charles Dupont	Coordinator of radiosonde launches at SIRTa
Laboratoire d'Aérodynamique	France	Jean-Pierre Chaboureau	Flight planning team
Laboratoire de Météorologie Dynamique	France	Gwendal Rivière	Science team, flight planning team
Laboratoire Atmosphère, Milieux et Observations Spatiales France	France	Julien Delanoë Jacques Pelon	Science team, flight planning team
		Christophe Caudoux, Quitterie Cazenave, Abdenour Irbah, Mathilde Van Haecke	RALI team (airborne radar-lidar)
Météo France	France	Philippe Arbogast	Science team, flight planning team
		Jean-Marie Donier	UHF radar at Lannion
SAFIRE	France	Jean-Christophe Canonici	SAFIRE coordinator
		Dominique Duchanoy, Guillaume Seurat	Falcon pilots
		Hubert Bellec, Nelly Geil, David Mourlas, Thierry Perrin, Frédéric Pouvesle	Falcon technical support
Deutsches Zentrum für Luft- und Raumfahrt (DLR), Flight Experiments	Germany	Andreas Minikin, Robert Uebelacker, Katrin Witte	HALO and Falcon project management
		Stefan Hempe, Frank Probst	HALO and Falcon operations
		Steffen Gamsa, Michael Grossrubatscher, Stefan Grillenbeck, Philipp Weber, Roland Welser, Matthias Wiese	HALO and Falcon pilots
		Volker Dreiling, Andreas Giez, Christian Mallaun, Martin Zöger	HALO and Falcon sensor and data team
		Michael Kettenberger,	HALO technical support

		Thomas Leder, Florian Gebhardt, Christoph Grad, Stephan Storhas, David Woudsma	
Deutsches Zentrum für Luft- und Raumfahrt (DLR), Institute for Atmospheric Physics	Germany	Andreas Schäfler	Mission coordinator, science team, flight planning team
		Axel Amediek, Andreas Fix, Silke Groß, Manuel Gutleben, Martin Wirth	WALES team
		Christian Lemmerz, Oliver Lux, Uwe Marksteiner, Engelbert Nagel, Stephan Rahm, Oliver Reitebuch, Benjamin Witschas	Wind lidar team
		Florian Ewald, Martin Hagen	HAMP team
		Martina Bramberger, Alenka Senika	Radiosonde team at Keflavik, Flight planning team
Karlsruhe Institute of Technology	Germany	Pila Bossmann, Enrico Di Muzio, Florian Pantillion	Flight planning team
Max Planck Institute for Meteorology (MPI-M) Hamburg	Germany	Björn Brüggemann, David Hellmann, Lutz Hirsch, Friedhelm Jansen, Marcus Klingebiel	HAMP team
Technical University Munich	Germany	Marc Rautenhaus	Flight planning team
University of Cologne	Germany	Susanne Crewell, Lisa Dirks, Marek Jacob, Mario Mech	HAMP team
University of Hamburg, Max Planck Institute for Meteorology (MPI-M)	Germany	Felix Ament, Heike Konow	HAMP team
University of Leipzig	Germany	Tim Carlsen, André Ehrlich, Manfred Wendisch, Kevin Wolf,	SMART team
University of Mainz	Germany	Marlene Baumgart, Christian Euler, Paolo Ghinassi, Michael Riemer, Volkmar Wirth	Flight planning team
University of Munich	Germany	George Craig	HALO mission PI, science team, flight planning team
		Florian Baur, Lotte Bierdel, Christian Keil, Julia Mack, Tobias Selz	Flight planning team

		Hans Grob, Lucas Höppler, Tobias Kölling, Bernhard Mayer, Tobias Zinner	specMACS team
Norwegian Meteorological Institute	Norway	Rich Moore	NEAREX team
University of Bergen	Norway	Harald Sodemann, Thomas Spengler	NEAREX team
National Institute of Research and Development for Optoelectronics	Romania	Dragos Ene	Flight planning team
ETH Zurich	Switzerland	Heini Wernli	Science team, flight planning team
		Roman Attinger, Hanin Binder, Maxi Boettcher, Bas Crezee, Christian Grams, Jacopo Riboldi	Flight planning team
University of Bern	Switzerland	Matthias Röthlisberger	Flight planning team
Met Office	UK	Richard Cotton, Stuart Fox	FAAM aircraft
NCAS and University of Manchester	UK	Bogdan Antonescu, Hugo Ricketts, Geraint Vaughan	Capel Dewi observations, FAAM aircraft
University of Reading and NCAS	UK	Suzanne Gray, John Methven	FAAM aircraft, NAWDEX- Influence PIs, science team, flight planning team
		Ben Harvey, Jacob Maddison, Oscar Martínez-Avarado, Leo Saffin	Flight planning team
Naval Research Laboratory Monterey, CA and Washington, DC	USA	James Doyle	Science team, flight planning team
		Stephen Eckermann, Carolyn Reynolds	Flight planning team
EUMETNET		Susanne Hafner, Stefan Klink	EUMETNET coordinators
ESA		Dirk Schüttemeyer	ESA campaign section

List of Figures

Fig. 1. Schematic of an idealized weather situation during NAWDEX. The blue line marks the location of the waveguide with a strong isentropic PV gradient separating stratospheric (blue background, $PV > 2$ PVU) from tropospheric air (white background). The jet stream (dark blue arrows) follows the waveguide. Surface lows develop below the leading edge of upper-level positive PV anomalies (black lines indicate sea level pressure and dark blue and red lines surface cold and warm fronts, respectively). Grey shaded areas indicate clouds related to ascending WCBs (yellow arrows). Purple arrows mark divergent outflow at the tropopause. The four green boxes outline the main regions of interest, i.e., the inflow (1), ascent (2) and outflow (3) of WCBs, as well as a region of expected downstream impact (4).

Fig. 2. Tracks of consecutively numbered research flights (RFs) of (a) HALO (97 flight hours during 13 RFs), (b) DLR Falcon (51 flight hours during 16 RFs), (c) SAFIRE Falcon (42 flight hours during 15 RFs; Flights 01 to 04 were made over France just before the campaign to calibrate the instruments), and (d) FAAM BAe 146 (15 flight hours during 3 RFs). The inset in (b) shows the DLR Falcon transfer flights and two flights to the Mediterranean at the end of the campaign.

Fig. 3. (a) Dropsondes launched from HALO (red dots, 191 dropsondes), SAFIRE Falcon (green dots, 59 dropsondes) and FAAM BAe 146 (blue dots, 39 dropsondes). (b) Ground-based observation sites during NAWDEX: Canadian radiosonde stations (red dots), European radiosonde stations that performed only operational ascents (blue dots) and those with requested additional radiosonde launches (green dots), and six sites with additional profile observations (black diamonds).

Fig. 4. (a-c) Synoptic-scale conditions during NAWDEX. All panels are based on ECMWF ERA-Interim data (1979-2016) and show in colors deviations of the campaign period in 2016 from the mid-September to mid-October climatology. (a-c) frequencies (all in %) of (a) surface anticyclones, (b) surface cyclones, and (c) WCB. The tracking method is explained in Sprenger et al. (2017). Black contours show the ERA-Interim 37-year climatological mean; (d) best track data (HURDAT2 of the National Hurricane Center) of six tropical cyclones (red sections classified as hurricane, orange as TS and blue as extratropical storm), and cyclone tracks during the NAWDEX period (light green lines) and before/after the campaign period (dark green lines).

Fig. 5. (a) Weather regime indices following the definition of Michel and Rivière (2011): Scandinavian blocking (blue line), positive NAO (red line), negative NAO (green line), and Atlantic ridge (yellow line), identified with a k-means clustering approach (Michelangeli et al. 1995). (b) Time series of ECMWF IFS anomaly correlation coefficient (ACC) for geopotential height at 500 hPa over an area 35 to 75°N and 60 to 0°W for a forecast lead time of +120 h (shown at the initial time of the forecast): IFS deterministic forecast (black line), ensemble mean (red line), 50% of the ensemble members (orange area) and all members (yellow area). (c) NAWDEX IOPs as indicated in Table 3 (red bars). Light grey box depicts the NAWDEX campaign period and the dark grey boxes mark the duration of weather sequences as shown in Figure 6.

Fig. 6. Three sequences that illustrate the NAWDEX storyline from trigger to interaction, development and HIW in Europe (based on ECMWF IFS operational analyses). All panels display PV on 325 K (PV < 2 PVU in white, 2 PVU \geq PV < 5 PVU in red, 5 PVU \geq PV < 8 PVU in orange, PV \geq 8 PVU in yellow), contours of wind speeds (grey contours; 60, 70 and 80 m s⁻¹) and MSLP (blue contours; interval 10 hPa). Some long-lived, coherent features are labelled to enable links from one frame to the next. K refers to TS Karl; W, F, S and St mark midlatitude cyclones observed by NAWDEX; C labels a tropopause level cut-off that persists for 10 days; R1-R9 refer to the prominent ridges along the North Atlantic waveguide, identified as northward excursions of the jet stream and the isentropic PV gradient; T marks a TPV.

Fig. 7. ECMWF IFS operational analyses of wind speed (color shading), 2 PVU (green line) at 325 K and MSLP (blue contours, in hPa) at (a) 12 UTC 23 Sep 2016 with Global Hawk flight track (black line, from 2120 UTC 22 Sep to 2100 UTC 23 Sep), (b) 12 UTC 25 Sep 2016 with Global Hawk flight track (black line, from 1820 UTC 24 Sep to 1715 UTC 25 Sep), (c) 12 UTC 26 Sep 2016 with HALO flight track (black line, from 10 to 19 UTC), and (d) 12 UTC 27 Sep 2016 with HALO (black line, 1130 to 2030 UTC), FAAM (pink line, 0800 to 1230 UTC) and DLR Falcon (purple line, 0930 to 1330 UTC) flight tracks.

Fig. 8. WCB observations on 23 Sep 2016 (IOP 3): (a) ECMWF IFS operational analysis of PV at 325 K (shading) and MSLP (black contours, in hPa) at 12 UTC 23 Sep 2016 with HALO flight track from Iceland (gray / green line; green part corresponds to the section shown in (a) and (b)) and FAAM flight track (gray / black line; black part

corresponds to track in (a) and (b)). The circle and diamond markers indicate the start and end positions of the latitudinal WCB cross-sections. (b) WALES DIAL water vapor mixing ratio (colors) and (c) HAMP radar reflectivity with HALO flight track (green line), FAAM flight track (thick black line, lowest leg was flown first) and dropsonde release positions (thin black lines). Only the part of the FAAM flight track with a spatial collocation to HALO is shown and both aircraft started at the same time, but had a time lag of ~2.5 h at the end of the last upper-most FAAM leg. (d) Ice water content as observed along the FAAM flight track. Differences between the flight tracks in (a,b) and (d) result from interpolation of FAAM position to the closest HALO observation in (a). The longitude axis in (d) was reversed to align with the time axis of the HALO flight track.

Fig. 9. Jet stream observations on 2 Oct 2016 (IOP 6): (a) ECMWF IFS operational analysis of PV at 320 K (shading) and MSLP (black contours, in hPa), and (b) 300 hPa wind speed (colors) and geopotential height (black contours, in dam) at 06 UTC 2 Oct 2016. (a) and (b) include flight tracks of the DLR Falcon (light green line) and SAFIRE Falcon (dark green line). The coordinated part of the flight from east to west shown in panels (c) and (d) is marked with the purple line. (c) DLR Falcon 2- μ m DWL wind speeds (colors) and (d) SAFIRE radar-derived wind speeds (colors). Grey area in (c) and (d) marks the topography of Greenland.

Fig.10. (a) ERA Interim moisture fluxes at 850 hPa (arrows, shadings shows magnitude) and surface pressure (black contours, hPa) at 12 UTC 13 Oct 2016. The red star indicates the location of the SIRTA surface observation site. (b) Daily accumulated precipitation (mm) and (c) daily maximum of instantaneous surface wind in southern France on 13 Oct 2016 from the high-resolution climatological network of Météo France surface weather stations. The black areas in (b) and (c) mark topography of the French Pyrenees, the Massif Central and the French Alps. (d) Reflectivity and Doppler velocity (approximately equal to terminal fall speed) at the 25 m resolution of the radar BASTA at SIRTA on 13 Oct 2016.

Fig. SB1. (a) Collocated observations of the vertical cloud structure below HALO, based on lidar (backscatter, along green part of the flight) and radar (radar reflectivity along red line). The underlying true color image was acquired by MODIS Aqua near the time of the flight (Global Imagery Browse Services (GIBS), operated by the NASA/GSFC/Earth Science Data and Information System). (b) Collocated wind

observations on board the DLR Falcon using the A2D direct-detection wind lidar (along the blue line) and the 2- μ m coherent DWL (along the red line). The latter is horizontally displaced from the actual flight track for clarity (Background picture: © 2017 Google).

Fig. SB2. NAWDEX forecast products for 12 UTC 26 Sep 2016 (lead time +60 h): (a) ECMWF IFS deterministic forecast of PV on 330 K (color shading) and mean sea level pressure (MSLP, in hPa). (b) WCB column probabilities of occurrence (color shading in %), derived from ECMWF ensemble (Schäfler et al. 2014; Rautenhaus et al. 2015b). Black line indicates location of cross section in Met.3D visualizations in (c) and (d). (c) Isosurface of ensemble mean wind of 60 m s⁻¹ (color indicates pressure on isosurface in hPa) and MSLP (black surface contours). The cross section shows ensemble mean PV (color shading) and potential temperature (black contours). Colored lines represent WCB trajectories of ensemble member 22, starting at 06 UTC 25 Sep 2016 (colored by pressure). The black vertical poles have been added to aid spatial perception; they are labeled with pressure in hPa. (d) WCB trajectories as in (c) but from a different viewpoint and combined with a cross section showing 3D WCB probabilities (color shading, in %), ensemble mean potential temperature (black contours) and the 2 PVU isoline (red contour). (e) COAMPS adjoint 48 h forecast moisture sensitivity at 850 hPa [color shading, increments every 0.2 m² s⁻² (g kg⁻¹)⁻¹] and 850 hPa geopotential heights (contours every 30 m) valid at 12 UTC 24 Sep (initial forecast time).




Fig. 1. Schematic of an idealized weather situation during NAWDEX. The blue line marks the location of the waveguide with a strong isentropic PV gradient separating stratospheric (blue background, $PV > 2$ PVU) from tropospheric air (white background). The jet stream (dark blue arrows) follows the waveguide. Surface lows develop below the leading edge of upper-level positive PV anomalies (black lines indicate sea level pressure and dark blue and red lines surface cold and warm fronts, respectively). Grey shaded areas indicate clouds related to ascending WCBs (yellow arrows). Purple arrows mark divergent outflow at the tropopause. The four green boxes outline the main regions of interest, i.e., the inflow (1), ascent (2) and outflow (3) of WCBs, as well as a region of expected downstream impact (4).




Fig. 2. Tracks of consecutively numbered research flights (RFs) of (a) HALO (97 flight hours during 13 RFs), (b) DLR Falcon (51 flight hours during 16 RFs), (c) SAFIRE Falcon (42 flight hours during 15 RFs; Flights 01 to 04 were made over France just before the campaign to calibrate the instruments), and (d) FAAM BAe 146 (15 flight hours during 3 RFs). The inset in (b) shows the DLR Falcon transfer flights and two flights to the Mediterranean at the end of the campaign.




Fig. 3. (a) Dropsondes launched from HALO (red dots, 191 dropsondes), SAFIRE Falcon (green dots, 59 dropsondes) and FAAM BAe 146 (blue dots, 39 dropsondes). (b) Ground-based observation sites during NAWDEX: Canadian radiosonde stations (red dots), European radiosonde stations that performed only operational ascents (blue dots) and those with requested additional radiosonde launches (green dots), and six sites with additional profile observations (black diamonds).




Fig. 4. (a-c) Synoptic-scale conditions during NAWDEX. All panels are based on ECMWF ERA-Interim data (1979-2016) and show in colors deviations of the campaign period in 2016 from the mid-September to mid-October climatology. (a-c) frequencies (all in %) of (a) surface anticyclones, (b) surface cyclones, and (c) WCB. The tracking method is explained in Sprenger et al. (2017). Black contours show the ERA-Interim 37-year climatological mean; (d) best track data (HURDAT2 of the National Hurricane Center) of six tropical cyclones (red sections classified as hurricane, orange as TS and blue as extratropical storm), and cyclone tracks during the NAWDEX period (light green lines) and before/after the campaign period (dark green lines).




Fig. 5. (a) Weather regime indices following the definition of Michel and Rivière (2011): Scandinavian blocking (blue line), positive NAO (red line), negative NAO (green line), and Atlantic ridge (yellow line), identified with a k-means clustering approach (Michelangeli et al. 1995). (b) Time series of ECMWF IFS anomaly correlation coefficient (ACC) for geopotential height at 500 hPa over an area 35 to 75°N and 60 to 0°W for a forecast lead time of +120 h (shown at the initial time of the forecast): IFS deterministic forecast (black line), ensemble mean (red line), 50% of the ensemble members (orange area) and all members (yellow area). (c) NAWDEX IOPs as indicated in Table 3 (red bars). Light grey box depicts the NAWDEX campaign period and the dark grey boxes mark the duration of weather sequences as shown in Figure 6.




Fig. 6. Three sequences that illustrate the NAWDEX storyline from trigger to interaction, development and HIW in Europe (based on ECMWF IFS operational analyses). All panels display PV on 325 K ($PV < 2$ PVU in white, $2 \text{ PVU} \leq PV < 5 \text{ PVU}$ in red, $5 \text{ PVU} \leq PV < 8$

1214 PVU in orange, $PV \geq 8$ PVU in yellow), contours of wind speeds (grey contours; 60, 70 and
1215 80 m s^{-1}) and MSLP (blue contours; interval 10 hPa). Some long-lived, coherent features are
1216 labelled to enable links from one frame to the next. K refers to TS Karl; W, F, S and St mark
1217 midlatitude cyclones observed by NAWDEX; C labels a tropopause level cut-off that persists
1218 for 10 days; R1-R9 refer to the prominent ridges along the North Atlantic waveguide,
1219 identified as northward excursions of the jet stream and the isentropic PV gradient; T marks a
1220 TPV.




Fig. 7. ECMWF IFS operational analyses of wind speed (color shading), 2 PVU (green line) at 325 K and MSLP (blue contours, in hPa) at (a) 12 UTC 23 Sep 2016 with Global Hawk flight track (black line, from 2120 UTC 22 Sep to 2100 UTC 23 Sep), (b) 12 UTC 25 Sep 2016 with Global Hawk flight track (black line, from 1820 UTC 24 Sep to 1715 UTC 25 Sep), (c) 12 UTC 26 Sep 2016 with HALO flight track (black line, from 10 to 19 UTC), and (d) 12 UTC 27 Sep 2016 with HALO (black line, 1130 to 2030 UTC), FAAM (pink line, 0800 to 1230 UTC) and DLR Falcon (purple line, 0930 to 1330 UTC) flight tracks.




Fig. 8. WCB observations on 23 Sep 2016 (IOP 3): (a) ECMWF IFS operational analysis of PV at 325 K (shading) and MSLP (black contours, in hPa) at 12 UTC 23 Sep 2016 with HALO flight track from Iceland (gray / green line; green part corresponds to the section shown in (a) and (b)) and FAAM flight track (gray / black line; black part corresponds to track in (a) and (b)). The circle and diamond markers indicate the start and end positions of the latitudinal WCB cross-sections. (b) WALES DIAL water vapor mixing ratio (colors) and (c) HAMP radar reflectivity with HALO flight track (green line), FAAM flight track (thick black line, lowest leg was flown first) and dropsonde release positions (thin black lines). Only the part of the FAAM flight track with a spatial collocation to HALO is shown and both aircraft started at the same time, but had a time lag of ~2.5 h at the end of the last upper-most FAAM leg. (d) Ice water content as observed along the FAAM flight track. Differences between the flight tracks in (a,b) and (d) result from interpolation of FAAM position to the closest HALO observation in (a). The longitude axis in (d) was reversed to align with the time axis of the HALO flight track.




Fig. 9. Jet stream observations on 2 Oct 2016 (IOP 6): (a) ECMWF IFS operational analysis of PV at 320 K (shading) and MSLP (black contours, in hPa), and (b) 300 hPa wind speed (colors) and geopotential height (black contours, in dam) at 06 UTC 2 Oct 2016. (a) and (b) include flight tracks of the DLR Falcon (light green line) and SAFIRE Falcon (dark green line). The coordinated part of the flight from east to west shown in panels (c) and (d) is marked with the purple line. (c) DLR Falcon 2- μ m DWL wind speeds (colors) and (d) SAFIRE radar-derived wind speeds (colors). Grey area in (c) and (d) marks the topography of Greenland.




Fig.10. (a) ERA Interim moisture fluxes at 850 hPa (arrows, shadings shows magnitude) and surface pressure (black contours, hPa) at 12 UTC 13 Oct 2016. The red star indicates the location of the SIRTa surface observation site. (b) Daily accumulated precipitation (mm) and (c) daily maximum of instantaneous surface wind in southern France on 13 Oct 2016 from the high-resolution climatological network of Météo France surface weather stations. The black areas in (b) and (c) mark topography of the French Pyrenees, the Massif Central and the French Alps. (d) Reflectivity and Doppler velocity (approximately equal to terminal fall speed) at the 25 m resolution of the radar BASTA at SIRTa on 13 Oct 2016.




Fig. SB1. (a) Collocated observations of the vertical cloud structure below HALO, based on lidar (backscatter, along green part of the flight) and radar (radar reflectivity along red line). The underlying true color image was acquired by MODIS Aqua near the time of the flight (Global Imagery Browse Services (GIBS), operated by the NASA/GSFC/Earth Science Data and Information System). (b) Collocated wind observations on board the DLR Falcon using the A2D direct-detection wind lidar (along the blue line) and the 2-μm coherent DWL (along the red line). The latter is horizontally displaced from the actual flight track for clarity (Background picture: © 2017 Google).




Fig. SB2. NAWDEX forecast products for 12 UTC 26 Sep 2016 (lead time +60 h): (a) ECMWF IFS deterministic forecast of PV on 330 K (color shading) and mean sea level pressure (MSLP, in hPa). (b) WCB column probabilities of occurrence (color shading in %), derived from ECMWF ensemble (Schäfler et al. 2014; Rautenhaus et al. 2015b). Black line indicates location of cross section in Met.3D visualizations in (c) and (d). (c) Isosurface of ensemble mean wind of 60 m s^{-1} (color indicates pressure on isosurface in hPa) and MSLP (black surface contours). The cross section shows ensemble mean PV (color shading) and potential temperature (black contours). Colored lines represent WCB trajectories of ensemble member 22, starting at 06 UTC 25 Sep 2016 (colored by pressure). The black vertical poles have been added to aid spatial perception; they are labeled with pressure in hPa. (d) WCB trajectories as in (c) but from a different viewpoint and combined with a cross section showing 3D WCB probabilities (color shading, in %), ensemble mean potential temperature (black contours) and the 2 PVU isoline (red contour). (e) COAMPS adjoint 48 h forecast moisture sensitivity at 850 hPa [color shading, increments every $0.2 \text{ m}^2 \text{ s}^{-2} (\text{g kg}^{-1})^{-1}$] and 850 hPa geopotential heights (contours every 30 m) valid at 12 UTC 24 Sep (initial forecast time).