

*Molecular characterization and  
phylogenetic analysis of betasatellite  
molecules associated with okra yellow  
vein mosaic disease in Sri Lanka*

Article

Accepted Version

Jeyeaseelan, T. C., Jeayaseelan, E. C., De Costa, D. M. and Shaw, M. W. (2018) Molecular characterization and phylogenetic analysis of betasatellite molecules associated with okra yellow vein mosaic disease in Sri Lanka. *Tropical Plant Pathology*, 43 (5). pp. 468-472. ISSN 1983-2052 doi: <https://doi.org/10.1007/s40858-018-0225-1> Available at <http://centaur.reading.ac.uk/77431/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

To link to this article DOI: <http://dx.doi.org/10.1007/s40858-018-0225-1>

Publisher: Springer International

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

[www.reading.ac.uk/centaur](http://www.reading.ac.uk/centaur)

## **CentAUR**

Central Archive at the University of Reading

Reading's research outputs online

## Title Page

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20

**Title:**

Molecular characterization and phylogenetic analysis of beta-satellite molecules associated with okra yellow vein mosaic disease in Sri Lanka

**Authors:** Tharmila C. Jeyaseelan<sup>1,4</sup>, Emmanuel C. Jeyaseelan<sup>1</sup>, Devika M. De Costa<sup>2,4</sup>, Michael W. Shaw<sup>3</sup>

1- Department of Botany, Faculty of Science, University of Jaffna, Jaffna, JA 40000, Sri Lanka.

2- Department of Agricultural Biology, Faculty of Agriculture, University of Peradeniya, Peradeniya, KY 20400, Sri Lanka.

3- School of Agriculture, Policy and Development, University of Reading, Reading, RG6 6BZ, United Kingdom.

4- Plant Protection Board of Study, Postgraduate Institute of Agriculture, University of Peradeniya, Peradeniya, KY 20400, Sri Lanka.

Corresponding author: Tharmila C. Jeyaseelan

E-mail address: [siva.tharmi3@gmail.com](mailto:siva.tharmi3@gmail.com)

[stharmila@jfn.ac.lk](mailto:stharmila@jfn.ac.lk)

**Abstract**

21 In the past few years, okra growth and yield has been greatly affected in Sri Lanka by a  
22 disease called okra yellow vein mosaic (OYVM). The disease is caused by begomovirus and  
23 associated beta-satellites. The associated beta-satellite molecule determines the development  
24 and severity of the disease. Therefore, knowledge of the sequence and diversity of the beta-  
25 satellite molecules associated with OYVM disease would assist choice of virus isolates or  
26 trial locations to use when selecting resistant varieties of okra. The present study aimed to  
27 characterize the beta-satellite DNA sequences associated with OYVM disease in Sri Lanka  
28 and determine their phylogenetic relationships. Beta-satellite DNA of six virus isolates from  
29 widely separated geographical locations were sequenced and compared with already reported  
30 begomovirus beta-satellites. The beta-satellite molecules have features common to  
31 begomovirus beta-satellite DNAs: a conserved nonanucleotide TAATATTAC, a coding  
32 sequence for the protein  $\beta$ C1, an adenine rich region and a satellite conserved region. er  
33 Nucleotide diversity among the isolates was relatively low ( $\pi = 0.027$ ). Recombination events  
34 were detected at a specific region in the genome of all the isolates. The isolates shared  
35 sequence identity >96 % with bhendi yellow vein beta-satellites reported from India. This is  
36 further supported by phylogenetic analysis.

37

38 **Key words:**

39 *Abelmoschus esculentus*, Begomovirus, Plant virus, Virus recombination

40

## 41 **Introduction**

42 Yellow vein mosaic (YVM) disease of Okra (*Abelmoschus esculentus*; local names Bhendi /  
43 Vendi / Bandakka) causes heavy economic losses, among pests and diseases that affect the  
44 yield and fruit quality of okra throughout the okra cultivating regions of the world (Pun et al.  
45 2005). Leaves of infected plants are characterized by alternate green and yellow patches, vein  
46 clearing and a conspicuous yellow network of vein. In severe cases the chlorosis may extend  
47 to the inter-veinal area and may result in complete yellowing of leaves with stunting of plants  
48 (Taware et al. 2010). The disease is caused by a geminivirus called okra yellow vein mosaic  
49 virus (OYVMV) (Jose and Usha 2003) which belongs to the genus *Begomovirus* (Fauquet and  
50 Stanley 2005). In *Begomovirus*, single stranded, circular genomic DNA is encapsulated in  
51 quasi-isometric (“geminata”) virions and transmitted by whitefly (*Bemisia tabaci*) (Briddon  
52 et al. 2008).

53 Most OYVMV isolates are monopartite viruses with a single strand DNA (DNA A) of helper  
54 begomovirus which associated with a single strand beta-satellite molecule. The beta-satellites  
55 have a gene which codes a protein known as  $\beta$ C1 (Mansoor et al. 2003). It is essential for the  
56 development of typical symptoms in infected okra plants (Jose and Usha 2003), suppression  
57 of post-transcriptional gene silencing and up-regulation of viral DNA levels *in planta*  
58 (Briddon et al. 2008). Many isolates of the virus and associated satellite molecules have been  
59 characterized in various okra growing countries (Ghosh et al. 2008; Taware et al. 2010;  
60 Venkataravanappa et al. 2011). Recent studies have revealed that the OYVM disease can be  
61 caused by a bipartite begomovirus species known as Bhendi yellow vein mosaic Delhi virus  
62 [BYVDV-IN (India: Delhi: okra)]. This has both DNA-A and DNA-B molecules, but not  
63 beta-satellites, in okra plants which nonetheless have typical symptoms (Venkataravanappa et  
64 al. 2012).

65

66 In Sri Lanka cultivated okra varieties such as TV8, MI5 and MI7 are greatly affected by the  
67 disease. Cultivation of the variety TV8 (locally known as ‘Paal Vendi’), a popular variety  
68 with a great market value in North and East parts of Sri Lanka, is vanishing because of  
69 widespread disease. The variety ‘Haritha’ was recommended as resistant by the Department  
70 of Agriculture, but in recent seasons the variety has suffered a high disease incidence,  
71 especially in Northern Sri Lanka. So far, studies on OYVM disease in Sri Lanka have mainly  
72 focusing on screening of resistant okra varieties, determination of disease incidence. More  
73 recent studies have tried to identify the causative agent, based on partial amplification of virus  
74 DNA by PCR (Senevirathna et al. 2016).

75 These studies have found that an okra plant showing yellow vein mosaic disease may carry  
76 mixed infections with Bhendi yellow vein mosaic virus (BYVMV) DNA-A, Okra enation leaf  
77 curl virus (OELCuV) DNA-A, beta- and alpha-satellites (Priyavathi et al. 2016). This  
78 complex mix of virus genetic materials associated with OYVM disease suggests that  
79 identification and characterization of the causative virus and associated satellite molecules  
80 may help explain why OYVM disease has become more prevalent and severe recently. In  
81 2016, we identified the causative agent of OYVM disease of the same samples as the  
82 begomovirus OYVMV based on whole genome sequencing of the virus isolates (Tharmila et  
83 al. 2017). The present study focused on genome sequence, organisation and phylogenetic  
84 relationship of beta-satellite molecules associated with OYVM disease in Sri Lanka.

85 Leaf samples of asymptomatic okra plants and okra plants showing yellow vein mosaic  
86 symptoms were collected from six different widely separated locations across Sri Lanka  
87 between May and July in 2015 (Table 1). In each location, three symptomatic and three non-  
88 symptomatic leaf samples were collected from three different farmer’s field or home gardens,  
89 while the okra plants were at their post flowering stage. Total DNA was extracted from both

90 non-symptomatic and symptomatic okra leaf samples using a DNA extraction protocol  
91 (Ghosh et al. 2009) with some modifications.

92 The extracted DNAs were PCR amplified with specific primers for the begomovirus genome.  
93 Presence of begomovirus in symptomatic samples was confirmed by PCR amplification of  
94 begomovirus DNA-A using a pair of degenerate primers (Deng et al. 1994). Full length beta-  
95 satellite DNA was amplified with a pair of universal primers  $\beta 01$  and  $\beta 02$  as described by  
96 Briddon et al. (2002). Six different PCR products of beta-satellite DNA, one from each  
97 location sampled, were cloned and sequenced. The PCR products were purified using a spin  
98 column PCR purification kit (NBS Biological, Huntingdon, Cambridgeshire) and cloned into  
99 pCR®2.1 vector using a TA cloning kit (Invitrogen, USA) as described in the manufacturers'  
100 protocols. The clones were sequenced by a commercial automated Sanger sequencing service  
101 (Source Bioscience, UK). The complete nucleotide sequences of the beta-satellites were  
102 deposited in GenBank database; accession numbers are given in Table 1.

103 Identity searches for the sequences were carried out by using the BLAST $n$  program available  
104 in the NCBI. Sequence alignments were performed using MUSCLE (Edgar 2004) and  
105 pairwise identity scores were calculated with the Species Demarcation Tool (SDT) (Muhire et  
106 al. 2014). The details of sequences retrieved from GenBank database are given in  
107 Supplementary Table 1. Phylogenetic analysis was performed in MEGA 7 (Kumar et al.  
108 2016) using the maximum likelihood algorithm with 1000 bootstrap replications. The mean  
109 pairwise number of nucleotide differences per site ( $\pi$ ) was estimated using DnaSP v. 5.10  
110 (Rozas et al. 2003). Putative parental viruses and recombination breakpoints were determined  
111 using Recombination Detection Program (RDP) v. 4.0 (Martin et al. 2015). Alignments were  
112 analysed using default settings for the different methods and statistical significance was  
113 defined as a  $P$  value less than the Bonferroni-corrected cut-off of 0.05.

114 PCR-mediated amplification of DNA extracts of all symptomatic samples with specific  
115 primers for begomovirus DNA-A and beta-satellite yielded products with an approximate  
116 length of 0.52 kb and 1.3 kb respectively (Data not shown). Amplifications from non-  
117 symptomatic plants were uniformly negative with all primers. This shows the frequent  
118 association of helper virus and beta-satellite with symptomatic plants.

119 The complete nucleotide sequences of the six beta-satellites ranged between 1318 and 1369  
120 bp in size (Table 1). The sequences showed all the features typical of a beta-satellite (Bridson  
121 et al. 2008), including a single ORF in a complementary sense strand which encodes a 118  
122 amino acid protein and an adenine rich region. They also contain a nonanucleotide stem-loop  
123 structure (TAATATTAC) and a satellite conserved region (SCR).

124  
125 SDT based pairwise alignment shows that the Sri Lankan okra yellow vein mosaic beta-  
126 satellites (OYVMB) had between 93.4 % to 99.2 % sequence identity. The most divergent  
127 isolate OYVMB [LK:Kan:15] had 93.4 % to 94.0 % sequence identity with the rest of the Sri  
128 Lankan OYVMB. The above isolates shared > 91 % of identity with some bhendi yellow vein  
129 beta-satellites (BYVB), bhendi yellow vein mosaic beta-satellites (BYVMB) and  
130 OYVMB[IN:Aur:08] from India. This classifies the sequences as coming from the same  
131 species, according to the recently proposed beta-satellite species demarcation threshold of  
132 91% ([https://talk.ictvonline.org/files/ictv\\_official\\_taxonomy](https://talk.ictvonline.org/files/ictv_official_taxonomy_updates_since_the_8th_report/m/plant-official/6689)  
133 [updates\\_since\\_the\\_8th\\_report/m/plant-official/6689](https://talk.ictvonline.org/files/ictv_official_taxonomy_updates_since_the_8th_report/m/plant-official/6689)). Sequence identity of Sri Lankan  
134 OYVMB isolates was < 91% with sequences of okra leaf curl beta-satellites (OLCuB), bhendi  
135 yellow vein India beta-satellites (BYVIB) and croton yellow vein mosaic beta-satellite  
136 (CrYVMB [IN\_Bhu\_OYBHU\_06]), all of which were also reported to be associated with  
137 okra yellow vein mosaic disease in India (Venkataravanappa et al. 2011).—Beta satellite



138 isolates associated with other begomovirus diseases in Sri Lanka had even lower identity (<70  
139 %) with the Sri Lankan OYVMB isolates.

140

141 Nucleotide diversity ( $\pi$ ) was measured in three subpopulations of beta-satellite isolates; Sri  
142 Lankan OYVMB (n=6), all the beta-satellite isolates reported from Sri Lanka (n=11) and  
143 some selected beta-satellites associated with okra (n=20). The lowest nucleotide diversity was  
144 noticed in Sri Lankan OYVMB ( $\pi=0.027$ ) and the highest nucleotide diversity was in the total  
145 beta-satellite isolates reported from Sri Lanka ( $\pi=0.245$ ).

146

147 Briddon et al. (2003) reported two major clusters of beta-satellites corresponding to hosts in the  
148 Malvaceae and non-Malvaceae. The Sri Lankan OYVMB clustered phylogenetically with  
149 isolates from the Malvaceae and were completely separated from previously published beta-  
150 satellite sequences from Sri Lanka, as expected (data not shown). Analysis of OYVMB and  
151 selected beta-satellites associated with okra/bhendi clearly showed the close clustering of Sri  
152 Lankan OYVMB with already reported Indian BYVB and BYVMB rather than okra enation  
153 leaf curl beta-satellites (OELCuB) and OLCuB (Figure 1). The isolates OYVMB [LK:Put:15]  
154 and OYVMB [LK:Kan:15] clustered separately from the rest of the Sri Lankan OYVMB  
155 isolates.

156 Genetic recombination is an important process in the evolution of viruses and it is frequent in  
157 begomoviruses and inter-species recombination is the major factor behind the emergence of  
158 new begomovirus species (Lefeuvre et al. 2007). We looked for recombination events among  
159 27 full-length genomes of different beta satellites associated with okra and other plants. A  
160 recombination event was detected in the genome of all the test OYVMB isolates. The  
161 recombination break point was detected immediately upstream of A-rich region and SCR  
162 region. The major and minor parents were inferred to be ageratum yellow vein beta-satellite

163 (AYVB[LK:Age:03] ) and *Leucas zeylanica* yellow vein beta-satellite (LZYVB[LK:Leu:06])  
164 respectively (Table 2). These are associated with common weeds, *Ageratum* spp. and *Leucas*  
165 *zeylanica* respectively, in Sri Lanka. The isolate OYVMB[LK:Kan:15] showed an additional  
166 recombination event a few nucleotides downstream of the previous recombination.

167 OYVMB[IN:Aur:08] and AYVB[LK:Age:03] were inferred to be the minor and major  
168 parents, respectively. One more recombination event was detected in the analysis, where the  
169 isolates OYVMB[LK:Jaf:15] and OYVMB[LK:Kan:15] were parents of the recombinant  
170 isolate OELCuB[IN:Jal-4/okr:15].

171 ~~In India, the beta-satellite molecules associated with okra segregated into four groups~~  
172 ~~(Venkataravanappa et al. 2011): okra leaf curl beta-satellites (OLCuB); bhendi yellow vein~~  
173 ~~beta-satellites (BYVB); bhendi yellow vein India beta-satellites (BYVIB); and a surprising~~  
174 ~~distinct group called croton yellow vein mosaic beta-satellite (CrYVMB). In present study,~~  
175 ~~the Sri Lankan beta-satellite isolates, except OYVMB[LK:Kan:15], were more closely related~~  
176 ~~to BYVB (>96 %) than rest of the three groups.~~

177 Begomovirus may range over long distances with little differentiation in infectivity and  
178 genome sequence. As expected, Sri Lankan isolates showed great similarities with Indian  
179 isolates. The countries are close together and agricultural products are frequently exchanged,  
180 so the isolates that associated with Sri Lankan OYVM disease might have arrived from India.  
181 Sequence diversity among the Sri Lankan OYVM beta-satellites was lower than in previously  
182 published isolates from India, possibly because the country is smaller and has less  
183 geographical and environmental variation than India, or because the disease invaded recently.

#### 184 **Acknowledgement**

185 The authors are grateful to Dr. George Gibbings and Dr. Ihsan Ullah for technical assistance in  
186 Molecular genetics laboratory, School of Agriculture, Policy and Development, University of

187 Reading, U.K. The works in Sri Lanka was supported by University Grants Commission,  
188 Government of Sri Lanka under the project: UGC/DRIC/PG/2015(ii)/JFN/01 is duly  
189 acknowledged.

190

## 191 **References**

192 Briddon R, Bull S, Mansoor S, Amin I, Markham G (2002) Universal primers for the PCR-  
193 Mediated amplification of DNA  $\beta$ . *Molecular Biotechnology* 20: 315–318.

194 Briddon R, Brown J, Moriones E, Stanley J, Zerbini M, Zhou X, Fauquet CM (2008)  
195 Recommendations for the classification and nomenclature of the DNA-  $\beta$  satellites of  
196 begomoviruses. *Archives of Virology* 153: 763–781.

197 Deng D, Mcgrath PF, Robinson DJ, Harrison BD (1994) Detection and differentiation of  
198 whitefly-transmitted geminiviruses in plants and vector insects by the polymerase chain  
199 reaction with degenerate primers. *Annals of Applied Biology* 125: 327–336.

200 Edgar RC (2004) MUSCLE: multiple sequence alignment with high accuracy and high  
201 throughput. *Nucleic Acids Research* 32:1792–1797.

202 Fauquet CM, Stanley J (2005) Revising the way we conceive and name viruses below the  
203 species level : A review of geminivirus taxonomy calls for new standardized isolate  
204 descriptors. *Archives of Virology* 150: 2151–2179.

205 Ghosh R, Paul S, Ghosh SK, Roy A (2009) An improved method of DNA isolation suitable  
206 for PCR-based detection of begomoviruses from jute and other mucilaginous plants.  
207 *Journal of Virological Methods* 159: 34–39.

- 208 Ghosh R, Paul S, Palit P, Das S, Das A, Acharyya S, Mir JI, Ghosh SK, Roy A (2008)  
209 Molecular characterization and computational analysis of coat protein gene of an east  
210 Indian isolate of Bhendi yellow vein mosaic virus and associated DNA-beta satellite.  
211 Indian Journal of Virology 19: 173–182.
- 212 Jose J, Usha R (2003) Bhendi yellow vein mosaic disease in India is caused by association of  
213 a DNA beta satellite with a begomovirus. Virology 305: 310–317.
- 214 Kumar S, Stecher G, Tamura K (2016) MEGA7 : Molecular evolutionary genetics analysis  
215 version 7 . 0 for bigger datasets. Molecular Biology and Evolution 33: 1870–1874.
- 216 Lefeuvre P, Martin DP, Hoareau M, Naze F, Delatte H, Thierry M, Varsani A, Becker N,  
217 Reynaud B, Lett JM (2007) Begomovirus ‘melting pot’ in the south-west Indian Ocean  
218 islands: molecular diversity and evolution through recombination. Journal of General  
219 Virology 88:3458–3468.
- 220 Mansoor S, Briddon RW, Zafar Y, Stanley J (2003) Geminivirus disease complexes : an  
221 emerging threat. Trends in Plant Science 8: 128–134.
- 222 Martin DP, Murrell B, Golden M, Khoosal A, Muhire B (2015) RDP4: Detection and analysis  
223 of recombination patterns in virus genomes. Virus Evolution 1:vev003
- 224 Muhire B, Varsani A, Martin DP (2014) SDT: A virus classification tool based on pairwise  
225 sequence alignment and identity calculation. PLoS One 9:e108277.
- 226 Priyavathi P, Kavitha V, Gopal P (2016) Complex nature of infection associated with yellow  
227 vein mosaic disease in Bhendi (*Abelmoschus esculentus*). Current Science 111: 1511–  
228 1515.

- 229 Pun KB, Doraiswamy S, Balasubramanian G (2005) Prediction of whitefly population and  
230 Okra yellow vein mosaic virus disease incidence in okra. Indian Journal of Virology  
231 16: 19-23.
- 232 Rozas J, Sanchez-Delbarrio JC, Messeguer X, Rozas R (2003) DnaSP, DNA polymorphism  
233 analyses by the coalescent and other methods. Bioinformatics 19:2496–2497.
- 234 Senevirathna HMSI, Wasala SK, Senanayake DMJB, Weerasekara D (2016) Characterization  
235 and detection of yellow vein disease of okra (*Abelmoschus esculentus* (L.) Moench) in  
236 Sri Lanka. Tropical Agricultural Research 27: 360–369.
- 237 Taware S, Thengane R, Harke S, Taware A (2010) Genome sequence analysis of DNA-A  
238 from okra yellow vein mosaic virus isolates. International Journal of Current Research 8:  
239 96–107.
- 240 Tharmila CJ, Jeyaseelan EC, Ihsan U, Wetten AC, De Costa DM, Shaw MW (2017) First  
241 report on association of okra yellow vein mosaic virus with yellow vein mosaic  
242 disease of okra (*Abelmoschus esculentus*) in Sri Lanka. Plant Disease 101:1335.
- 243 Venkataravanappa V, Reddy CNL, Jalali S, Reddy MK (2012) Molecular characterization of  
244 distinct bipartite begomovirus infecting bhendi (*Abelmoschus esculentus* L .) in India.  
245 Virus Genes 44: 522–535.
- 246 Venkataravanappa V, Reddy CNL, Swaranalatha P, Jalali S, Briddon RW, Reddy MK (2011)  
247 Diversity and phylogeography of begomovirus- associated beta satellites of okra in India.  
248 Virology Journal 8: 555.
- 249

250 **Table 1** Characteristic features of beta-satellites isolated from OYVMD from different locations in Sri Lanka

Samples collected districts in Sri Lanka	Beta-Satellites	Abbreviation	Accession number in NCBI GenBank	Length of the DNA (bp)	GenBank sequence showing highest sequence identity in nucleotide BLAST search	
Vavuniya	Okra yellow vein mosaic beta-satellite [Sri Lanka: Vavuniya: 2015]	OYVMB [LK:Vav:15]	KX174318	1318	Bhendi yellow vein India beta-satellite [India:Jalna:Jal:2009] (KJ462078)	96%
Jaffna	Okra yellow vein mosaic beta-satellite [Sri Lanka: Jaffna: 2015]	OYVMB [LK:Jaf:15]	KX174319	1334	Bhendi yellow vein India beta-satellite [India :Coimbatore:Co:2009] (KJ462077)	98%
Kandy	Okra yellow vein mosaic beta-satellite [Sri Lanka: Kandy: 2015]	OYVMB [LK:Kan:15]	KX174320	1369	Bhendi yellow vein mosaic beta-satellite [India:Guntur:OY112:2006] (GU111969)	94%
Matara	Okra yellow vein mosaic beta-satellite [Sri Lanka: Matara: 2015]	OYVMB [LK:Mat:15]	KX174321	1351	Bhendi yellow vein mosaic beta-satellite [India:Madurai:MKU-1:2014] (KR068483)	98%
Trincomalee	Okra yellow vein mosaic beta-satellite	OYVMB [LK:Tri:15]	KX174322	1320	Bhendi yellow vein India beta-satellite [India:Jalna:Jal:2009] (KJ462078)	96%

	[Sri Lanka: Trincomalee: 2015]					
Puttalam	Okra yellow vein mosaic beta-satellite [Sri Lanka: Puttalam: 2015]	OYVMVB [LK:Put:15]	KX174323	1351	Bhendi yellow vein mosaic beta-satellite [India:Aurangabad:OY165:2006] (GU111977)	97%

252 **Table 2** Putative recombination events detected within beta-satellites associated with okra and beta-satellites reported from Sri Lanka, based on  
 253 full-length beta-satellite sequences

Event	Breakpoints		Recombinant	Parents		Methods <sup>1</sup>	P-Value <sup>2</sup>
	Begin	End		Major	Minor		
1	1076	1316	OYVMB[LK:Vav:15]	AYVB[LK:Age:03]	LZYVB[LK:Leu:06]	RGBMCS <sub>3</sub>	3.16E-06
2	1092	1332	OYVMB[LK:Jaf:15]	AYVB[LK:Age:03]	LZYVB[LK:Leu:06]	RGBMCS <sub>3</sub>	3.16E-06
3	1154	1367	OYVMB[LK:Kan:15]	AYVB[LK:Age:03]	LZYVB[LK:Leu:06]	RGBMCS <sub>3</sub>	3.16E-06
4	1109	1349	OYVMB[LK:Mat:15]	AYVB[LK:Age:03]	LZYVB[LK:Leu:06]	RGBMCS <sub>3</sub>	3.16E-06
5	1078	1318	OYVMB[LK:Tri:15]	AYVB[LK:Age:03]	LZYVB[LK:Leu:06]	RGBMCS <sub>3</sub>	3.16E-06
6	1109	1349	OYVMB[LK:Put:15]	AYVB[LK:Age:03]	LZYVB[LK:Leu:06]	RGBMCS <sub>3</sub>	3.16E-06
7	998	1127	OYVMB[LK:Kan:15]	OYVMB[IN:Aur:08]	AYVB[LK:Age:03]	RGBMCS <sub>3</sub>	5.93E-11
8	52	812	OELCuB[IN:Jal-4/okr:15]	OYVMB[LK:Jaf:15]	OYVMB[LK:Kan:15]	RGBMCS <sub>3</sub>	2.95E-15

254 <sup>1</sup> *R* RDP, *G* GeneConv, *B* Bootscan, *M* MaxChi, *C* Chimera, *S* SisScan, *3* 3SEQ

255 <sup>2</sup> The reported *P* values are for the methods by underline, and they are the lowest *P* values calculated for the region in question

256

257



258 **Figure legends**

259 **Fig. 1** Colour coded pairwise identity matrix generated from 27 different beta-satellite DNAs,  
260 including 6 beta-satellites described in this work, 5 beta-satellites associated with different  
261 begomoviral diseases in Sri Lanka, 14 different beta-satellites associated with okra/bhendi  
262 and 2 other beta-satellites associated with chilli and cotton. See Supplementary Table 1 for  
263 details on the compared sequences. Each coloured cell represents a percentage identity score  
264 between two sequences. The coloured key indicates the correspondence between pairwise  
265 identities and the colours displayed in the matrix.

266 **Fig. 2** Molecular Phylogenetic analysis by Maximum Likelihood method

267 The evolutionary history was inferred by using the Maximum Likelihood method based on  
268 the General Time Reversible model. The tree with the highest log likelihood (-10077.7604) is  
269 shown. The percentage of trees in which the associated taxa clustered together is shown next  
270 to the branches. Initial tree(s) for the heuristic search were obtained automatically by applying  
271 Neighbor-Join and BioNJ algorithms to a matrix of pairwise distances estimated using the  
272 Maximum Composite Likelihood (MCL) approach, and then selecting the topology with  
273 superior log likelihood value. A discrete Gamma distribution was used to model evolutionary  
274 rate differences among sites (5 categories (+G, parameter = 0.9163)). The tree is drawn to  
275 scale, with branch lengths measured in the number of substitutions per site. The analysis  
276 involved 30 nucleotide sequences; Indian and Sri Lankan isolates are denoted IN and LK  
277 respectively within the parentheses following the isolate code, . All positions containing gaps  
278 and missing data were eliminated. There were a total of 806 positions in the final dataset.  
279 Evolutionary analyses were conducted in MEGA7.

280

281