

*Grass silage particle size when fed with or without maize silage alters performance, reticular pH and metabolism of Holstein-Friesian dairy cows*

Article

Accepted Version

Figure 1

Tayyab, U., Wilkinson, R. G., Charlton, G. L., Reynolds, C. K. ORCID: <https://orcid.org/0000-0002-4152-1190> and Sinclair, L. A. (2019) Grass silage particle size when fed with or without maize silage alters performance, reticular pH and metabolism of Holstein-Friesian dairy cows. *Animal*, 13 (3). pp. 524-532. ISSN 1751-7311 doi: <https://doi.org/10.1017/S1751731118001568> Available at <https://centaur.reading.ac.uk/77459/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

To link to this article DOI: <http://dx.doi.org/10.1017/S1751731118001568>

Publisher: Cambridge University Press

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

[www.reading.ac.uk/centaur](http://www.reading.ac.uk/centaur)

**CentAUR**

Central Archive at the University of Reading

Reading's research outputs online

