

On the numerical approximation of p -biharmonic and ∞ -biharmonic functions

Article

Accepted Version

Pryer, T. and Katzourakis, N. (2019) On the numerical approximation of p -biharmonic and ∞ -biharmonic functions. Numerical Methods for Partial Differential Equations, 35 (1). pp. 155-180. ISSN 1098-2426 doi: 10.1002/num.22295 Available at <https://centaur.reading.ac.uk/77597/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

To link to this article DOI: <http://dx.doi.org/10.1002/num.22295>

Publisher: Wiley

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online

ON THE NUMERICAL APPROXIMATION OF p -BIHARMONIC AND ∞ -BIHARMONIC FUNCTIONS

NIKOS KATZOURAKIS AND TRISTAN PRYER

ABSTRACT. The ∞ -Bilaplacian is a *third order* fully nonlinear PDE given by

$$\Delta_\infty^2 u := (\Delta u)^3 |\mathbf{D}(\Delta u)|^2 = 0.$$

In this work we build a numerical method aimed at quantifying the nature of solutions to this problem which we call ∞ -Biharmonic functions. For fixed p we design a mixed finite element scheme for the pre-limiting equation, the p -Bilaplacian

$$\Delta_p^2 u := \Delta(|\Delta u|^{p-2} \Delta u) = 0.$$

We prove convergence of the numerical solution to the weak solution of $\Delta_p^2 u = 0$ and show that we are able to pass to the limit $p \rightarrow \infty$. We perform various tests aimed at understanding the nature of solutions of $\Delta_\infty^2 u$ and we prove convergence of our discretisation to an appropriate weak solution concept of this problem, that of \mathcal{D} -solutions.

1. INTRODUCTION AND THE ∞ -BILAPLACIAN

Let $\Omega \subset \mathbb{R}^d$ be an open and bounded set. For a given function $u : \Omega \rightarrow \mathbb{R}$ we denote the gradient of u as $Du : \Omega \rightarrow \mathbb{R}^d$ and its Hessian $\mathbf{D}^2 u : \Omega \rightarrow \mathbb{R}^{d \times d}$ and Laplacian $\Delta u : \Omega \rightarrow \mathbb{R}$. The p -Bilaplacian

$$(1.1) \quad \Delta_p^2 u := \Delta(|\Delta u|^{p-2} \Delta u) = 0$$

is a fourth order elliptic partial differential equation (PDE) which is a nonlinear generalisation of the Bilaplacian. Such problems typically arise from areas of elasticity, in particular, the nonlinear case can be used as a model for travelling waves in suspension bridges [27, 18]. It is a fourth order analogue to its second order sibling, the p -Laplacian, and as such it is useful as a prototypical nonlinear fourth order problem.

The efficient numerical simulation of general fourth order problems has attracted growing interest. A conforming approach to this class of problems would require the use of C^1 finite elements, the Argyris element for example [7, Section 6]. From a practical point of view the approach presents difficulties, in that the C^1 finite elements are difficult to design and complicated to implement, especially when working in three spatial dimensions. Other possibilities include discontinuous Galerkin methods, which form a class of nonconforming finite element method. If $p = 2$ we have the special case that the (2-)Bilaplacian, $\Delta^2 u = 0$, is linear. It has been well studied in the context of both C^1 finite elements [7] and discontinuous Galerkin methods; for example, the papers [26, 13] study the use of h - k dG finite elements (where k here means the local polynomial degree as opposed to the usual convention which is p) applied to the (2-)Bilaplacian. Alternative methods do exist, including those of virtual element type [33, 6] and recovered element type [15]. In addition to this, the classical work of [3] proposed mixed methods for the linear problem whose analysis was based on the mesh-dependent norms in [2]. The numerical approximation of p -Bilaplacian (quasi-linear, fourth order) type PDEs is relatively untouched. To the authors' knowledge, the only known work is [31] where a discontinuous Galerkin method based on a variational principle was derived and was shown to converge under minimal regularity. However, no rates of convergence were proven.

In this work we propose a method based on C^0 -mixed finite elements very much in the spirit of [3]. We rewrite the minimisation problem in mixed formulation and prove that the method converges under minimal regularity of the solution. In addition, using an inf-sup condition inspired by [29, 14, 12] and tools from [32,

Date: May 14, 2018.

N.K. was partially supported through the EPSRC grant EP/N017412/1. T.P. was partially supported through the EPSRC grant EP/P000835/1.

Keywords: p -Bilaplacian; ∞ -Bilaplacian; Generalised solutions; Calculus of Variations in L^∞ ; Finite element method; Fully nonlinear equations; Young measures

11, 17], we are able to show that under additional regularity assumptions the approximation converges with specific rates that depend on p .

Making use of these convergence results and the uniqueness of solutions in one dimension from [21], extended to multi-spatial dimensions in [20] we are able to justify that approximations of the p -Bilaplacian for large p are “good” approximations to ∞ -Biharmonic functions. These functions are solutions of the ∞ -Bilaplacian which is the PDE

$$(1.2) \quad \Delta_\infty^2 u := (\Delta u)^3 |\mathbf{D}(\Delta u)|^2 = 0,$$

derived in [21] as the formal limit of the p -Bilaplacian (1.1) as $p \rightarrow \infty$. The ∞ -Bilaplacian is the prototypical example of a PDE from second order Calculus of Variations in L^∞ , arising as the analogue of the *Euler–Lagrange* equation associated with critical points of the supremal functional

$$(1.3) \quad \mathcal{J}[u; \infty] := \|\Delta u\|_{L^\infty(\Omega)}.$$

Variational problems in L^∞ are notoriously challenging. The first order case is reasonably well understood and was initiated in the sequence of works by Aronsson starting with [1]. In this case, the respective Euler–Lagrange equation associated with critical points of the functional

$$(1.4) \quad \mathcal{J}[u] = \|Du\|_{L^\infty(\Omega)},$$

is quasi-linear, second order and given by

$$(1.5) \quad \Delta_\infty u = (Du \otimes Du) : D^2 u = 0.$$

This equation is called the ∞ -Laplacian and can be derived through a p -approximation of the underlying $W^{1,p}$ energy functional, see [30, 25].

It can be shown that solutions to (1.2) can not, in general, be C^3 even when $d = 1$; in particular, the Dirichlet problem is not solvable in the class of classical solutions. For a more extensive discussion we refer to [21]. Hence, the development of a solution concept which can be interpreted in an appropriate weak sense is in order. In the case of the ∞ -Laplacian, the appropriate notion is that of the Crandall–Ishii–Lions notion of viscosity solutions [8]. For an introduction to this theory we refer to the monograph [23]. We note that in the framework of viscosity solutions we can obtain uniqueness of solution for the Dirichlet problem [19]. In the case of second order Calculus of Variations in L^∞ the viscosity solution concept for the resulting equations is no longer applicable since we do not have access to a maximum principle for third order PDEs like (1.2), from which the solution concept stems.

One possibility for a generalised solution concept to (1.2) is that of \mathcal{D} -solutions [24, 22, 21]. Roughly, this is a probabilistic approach where derivatives that do not exist classically are represented as limits of difference quotients into Young measures over a compactification of the space of derivatives. This solution concept has already borne substantial fruit in the first order vectorial case of Calculus of Variations in L^∞ , as well as for more general PDE systems. In the present second order setting it proves to be an appropriate notion as well, since absolute minimisers $u \in \mathcal{W}_g^{2,\infty}(\Omega)$ satisfying

$$(1.6) \quad \|\Delta u\|_{L^\infty(\Omega')} \leq \|\Delta v\|_{L^\infty(\Omega')} \quad \forall \Omega' \Subset \Omega \text{ and } v \in \mathcal{W}_g^{2,\infty}(\Omega'),$$

are indeed unique \mathcal{D} -solutions of (1.2). Note that the appropriate space to take minimisers is *not* $W_g^{2,\infty}(\Omega)$ but rather the larger space

$$(1.7) \quad \mathcal{W}_g^{2,\infty}(\Omega) := \left\{ u \in \bigcap_{p \in (1, \infty)} W_g^{2,p}(\Omega) : \Delta u \in L^\infty(\Omega) \right\}.$$

In [21] it has been shown that in one spatial dimension the problem does indeed have a unique absolutely minimising \mathcal{D} -solution and in [20] for higher spatial dimension.

The design of numerical schemes that are compatible with these solution concepts that are inherently incompatible with duality techniques is extremely difficult. Even for the well developed area of viscosity solutions most numerical schemes that exist which are compatible with the solution concept are based on the arguments of [4] which advocates approximations based on differences satisfying a discrete monotonicity property. The only other methodology in the design of numerical schemes for the ∞ -Laplacian is to make use of the variational principle from which the equation is derived. Galerkin approximations of the p -Laplacian

can then be shown to converge to the viscosity solution of the ∞ -Laplacian [30]. This method has also been used to characterise the nature of solutions to the variational ∞ -Laplace system [25]. This is also the approach we use here. We build a scheme convergent to the weak solution of the p -Bilaplacian and then justify its use as an approximation of ∞ -Biharmonic functions. This allows us significant insight as to the nature of non-classical solutions of the ∞ -Bilaplacian and to make various conjectures about their structure and behaviour.

The rest of the paper is set out as follows: In §2 we formalise notation and begin exploring some of the properties of the p -Bilaplacian. In particular, we reformulate the PDE as a saddle point type problem. We show inf-sup conditions for the underlying operators guarantee that the saddle point type problem is well posed, motivating the discretisation of this directly. In §3 we perform the discretisation for fixed p and show that discrete versions of the inf-sup conditions hold. A priori results for both primal and auxiliary variables are a consequence of this. Numerical experiments are given in §4 illustrating the behaviour of numerical approximations to this problem. In addition, we examine the solutions for large p and make various conjectures as to the structure of solutions in multiple spatial dimensions.

2. APPROXIMATION VIA THE p -BILAPLACIAN

In this section we describe how ∞ -Biharmonic functions can be approximated using p -Biharmonic functions. We give a brief introduction to the p -Bilaplacian problem, beginning by introducing the Sobolev spaces

$$(2.1) \quad L^p(\Omega) = \left\{ \phi \text{ measurable} : \int_{\Omega} |\phi|^p \, d\mathbf{x} < \infty \right\} \text{ for } p \in [1, \infty) \text{ and}$$

$$(2.2) \quad L^\infty(\Omega) = \{ \phi \text{ measurable} : \text{ess sup}_{\Omega} |\phi| < \infty \},$$

$$(2.3) \quad W^{l,p}(\Omega) = \{ \phi \in L^p(\Omega) : D^\alpha \phi \in L^p(\Omega), \text{ for } |\alpha| \leq l \} \text{ and } H^l(\Omega) := W^{l,2}(\Omega),$$

which are equipped with the following norms and semi-norms:

$$(2.4) \quad \|v\|_{L^p(\Omega)}^p := \int_{\Omega} |v|^p \, d\mathbf{x} \text{ for } p \in [1, \infty) \text{ and } \|v\|_{L^\infty(\Omega)} := \text{ess sup}_{\Omega} |v|$$

$$(2.5) \quad \|v\|_{W^{l,p}(\Omega)}^p := \sum_{|\alpha| \leq l} \|D^\alpha v\|_{L^p(\Omega)}^p$$

$$(2.6) \quad |v|_{W^{l,p}(\Omega)}^p := \sum_{|\alpha|=l} \|D^\alpha v\|_{L^p(\Omega)}^p$$

where $\alpha = \{\alpha_1, \dots, \alpha_d\}$ is a multi-index, $|\alpha| = \sum_{i=1}^d \alpha_i$ and derivatives D^α are understood in the weak sense. We pay particular attention to the case $l = 2$ and define

$$(2.7) \quad W_g^{2,p}(\Omega) := g + W_0^{2,p}(\Omega) = \{ \phi \in W^{2,p}(\Omega) : \phi|_{\partial\Omega} = g \text{ and } D\phi|_{\partial\Omega} = Dg \},$$

for a prescribed function $g \in W^{2,\infty}(\Omega)$, where the boundary condition is understood in the trace sense if $\partial\Omega \in C^{0,1}(\Omega)$. We note that if $p > d$, then the boundary condition is satisfied in the pointwise sense since $W_0^{2,p}(\Omega) \subseteq C^1(\bar{\Omega})$.

For the p -Bilaplacian, the action functional is given as

$$(2.8) \quad \mathcal{J}[u; p] = \int_{\Omega} |\Delta u|^p \, d\mathbf{x}.^1$$

We then look to find a minimiser over the space $W_g^{2,p}(\Omega)$, that is, to find $u \in W_g^{2,p}(\Omega)$ such that

$$(2.9) \quad \mathcal{J}[u; p] = \min_{v \in W_g^{2,p}(\Omega)} \mathcal{J}[v; p].$$

¹Typically $\mathcal{J}[u; p] = \frac{1}{p} \int |\Delta u|^p$. Note here the rescaling has no effect on the resultant Euler–Lagrange equations as the Lagrangian is independent of u .

If we assume temporarily that we have access to a smooth minimiser, i.e., $u \in C^4(\Omega)$, then, given that the Lagrangian is of second order, we have that the Euler–Lagrange equations are (in general) fourth order and read

$$(2.10) \quad \Delta(|\Delta u|^{p-2} \Delta u) = 0.$$

Note that, for $p = 2$, the PDE reduces to the Bilaplacian $\Delta^2 u = 0$. In general, the Dirichlet problem for the p -Bilaplacian is, given $g \in W^{2,\infty}(\Omega)$, to find u such that

$$(2.11) \quad \begin{cases} \Delta_p u := \Delta(|\Delta u|^{p-2} \Delta u) = 0, & \text{in } \Omega, \\ u = g, & \text{on } \partial\Omega, \\ Du = Dg, & \text{on } \partial\Omega. \end{cases}$$

2.1. Definition (weak solution). The problem (2.11) has a weak formulation. Consider the semilinear form

$$(2.12) \quad \mathcal{A}(u, v) := \int_{\Omega} (|\Delta u|^{p-2} \Delta u) \Delta v \, d\mathbf{x}.$$

Then, $u \in W_g^{2,p}(\Omega)$ is a *weak solution* of (2.11) if it satisfies

$$(2.13) \quad \mathcal{A}(u, v) = 0 \quad \forall v \in W_0^{2,p}(\Omega).$$

2.2. Proposition (coercivity of \mathcal{J}). Suppose that $u \in W_0^{2,p}(\Omega)$ and $f \in L^q(\Omega)$, where $\frac{1}{p} + \frac{1}{q} = 1$. We have that the action functional $\mathcal{J}[\cdot; p]$ is coercive over $W_0^{2,p}(\Omega)$, that is,

$$(2.14) \quad \mathcal{J}[u; p] \geq C |u|_{2,p}^p - \gamma,$$

for some $C > 0$ and $\gamma \geq 0$. Equivalently, we have that there exists a constant $C > 0$ such that

$$(2.15) \quad \mathcal{A}(v, v) \geq C |v|_{2,p}^p \quad \forall v \in W_0^{2,p}(\Omega).$$

2.3. Corollary (weak lower semicontinuity). The action functional \mathcal{J} is weakly lower semi-continuous over $W_g^{2,p}(\Omega)$. That is, given a sequence of functions $\{u_j\}_{j \in \mathbb{N}}$ which has a weak limit $u \in W_g^{2,p}(\Omega)$, we have

$$(2.16) \quad \mathcal{J}[u; p] \leq \liminf_{j \rightarrow \infty} \mathcal{J}[u_j; p].$$

Proof The proof of this fact is a straightforward extension of [10, Section 8.2 Thm 1] to second order Lagrangians, noting that \mathcal{J} is coercive (from Proposition 2.2) and convex. We omit the full details for brevity. \square

2.4. Corollary (existence and uniqueness). There exists a unique minimiser to the p -Dirichlet energy functional. Equivalently, there exists a unique (weak) solution $u \in W_g^{2,p}(\Omega)$ to the (weak form of the) Euler–Lagrange equations:

$$(2.17) \quad \mathcal{A}(u, v) = \int_{\Omega} |\Delta u|^{p-2} \Delta u \Delta v \, d\mathbf{x} = 0 \quad \forall v \in W_0^{2,p}(\Omega).$$

Proof Again, the result can be deduced by extending the arguments in [10, Section 8.2] or [7, Thm 5.3.1], again, noting the results of Propositions 2.2 and convexity. The full argument is omitted for brevity. \square

2.5. Theorem (the limit as $p \rightarrow \infty$). Let $(u_p)_1^\infty$ denote a sequence of weak solutions $u_p \in W_g^{2,p}(\Omega)$ to the p -Bilaplacian. Then, there exists a subsequence converging uniformly together with their derivatives to a (candidate ∞ -Biharmonic) function $u_\infty \in \mathcal{W}_g^{2,\infty}(\Omega)$. Namely,

$$(2.18) \quad u_{p_j} \rightarrow u_\infty \text{ in } C^1(\overline{\Omega}),$$

along a subsequence as $p \rightarrow \infty$.

Proof Let $u_p \in W_g^{2,p}(\Omega)$ denote the weak solution of (2.11). In view of Corollary 2.4, we know that u_p minimises the energy functional

$$(2.19) \quad \mathcal{J}[u_p] = \int_{\Omega} |\Delta u_p|^p \, d\mathbf{x}.$$

In particular,

$$(2.20) \quad \mathcal{J}[u_p] \leq \mathcal{J}[g],$$

where $g \in W^{2,\infty}(\Omega)$ is the associated boundary data to (2.11). Using this fact, we have

$$(2.21) \quad \|\Delta u_p\|_{L^p(\Omega)}^p = \mathcal{J}[u_p] \leq \mathcal{J}[g] = \|\Delta g\|_{L^p(\Omega)}^p,$$

and we may infer that

$$(2.22) \quad \|\Delta u_p\|_{L^p(\Omega)} \leq \|\Delta g\|_{L^p(\Omega)}.$$

Now fix a $k > d$ and take $p \geq k$. Then, by using Hölder's inequality with $r = \frac{p}{k}$ and $q = \frac{r}{r-1}$ such that $\frac{1}{r} + \frac{1}{q} = 1$, we obtain

$$(2.23) \quad \|\Delta u_p\|_{L^k(\Omega)}^k = \int_{\Omega} |\Delta u_p|^k \, d\mathbf{x} \leq \left(\int_{\Omega} 1^q \, d\mathbf{x} \right)^{1/q} \left(\int_{\Omega} |\Delta u_p|^p \, d\mathbf{x} \right)^{1/r}.$$

Hence

$$(2.24) \quad \|\Delta u_p\|_{L^k(\Omega)}^k \leq |\Omega|^{\frac{r-1}{r}} \|\Delta u_p\|_{L^p(\Omega)}^k = |\Omega|^{1-\frac{k}{p}} \|\Delta u_p\|_{L^p(\Omega)}^k$$

and we see

$$(2.25) \quad \|\Delta u_p\|_{L^k(\Omega)} \leq |\Omega|^{\frac{1}{k}-\frac{1}{p}} \|\Delta u_p\|_{L^p(\Omega)}.$$

By using the triangle inequality, a double application of the Poincaré inequality (since both $u = g$ and $Du = Dg$ on $\partial\Omega$) from Proposition 2.7 and the Calderon-Zygmund L^k estimates from Proposition 2.8, we have

$$(2.26) \quad \begin{aligned} \|u_p\|_{L^k(\Omega)} &\leq \|u_p - g\|_{L^k(\Omega)} + \|g\|_{L^k(\Omega)} \\ &\leq C'(k, \Omega) \|D^2 u_p - D^2 g\|_{L^k(\Omega)} + \|g\|_{L^k(\Omega)} \\ &\leq C(k, \Omega) \|\Delta u_p - \Delta g\|_{L^k(\Omega)} + \|g\|_{L^k(\Omega)}. \end{aligned}$$

By utilising the triangle inequality again, we have

$$(2.27) \quad \begin{aligned} \|u_p\|_{L^k(\Omega)} &\leq C \left(\|\Delta u_p\|_{L^k(\Omega)} + \|g\|_{W^{2,k}(\Omega)} \right) \\ &\leq C \left(|\Omega|^{\frac{1}{k}-\frac{1}{p}} \|\Delta u_p\|_{L^p(\Omega)} + \|g\|_{W^{2,k}(\Omega)} \right), \end{aligned}$$

by virtue of (2.25). Similarly, one may show that

$$(2.28) \quad \|Du_p\|_{L^k(\Omega)} \leq C \left(|\Omega|^{\frac{1}{k}-\frac{1}{p}} \|\Delta u_p\|_{L^p(\Omega)} + \|g\|_{W^{2,k}(\Omega)} \right).$$

Thus, in view of (2.22) we infer that

$$(2.29) \quad \|u_p\|_{W^{2,k}(\Omega)} \leq C \|g\|_{W^{2,k}(\Omega)}.$$

This means that for any $k > d$ we have the uniform bound

$$(2.30) \quad \sup_{p>k} \|u_p\|_{W^{2,k}(\Omega)} \leq C = C(k, \Omega).$$

By invoking standard weak compactness arguments, we may extract a sub-sequence $\{u_{p_j}\}_{j=1}^{\infty} \subset \{u_p\}_{p=1}^{\infty}$ and a function $u_{\infty} \in W^{2,k}(\Omega)$ such that, for any $k > n$,

$$(2.31) \quad u_{p_j} \rightharpoonup u_{\infty} \text{ weakly in } W^{2,k}(\Omega)$$

as $j \rightarrow \infty$ and

$$(2.32) \quad \begin{aligned} \|u_{\infty}\|_{W^{2,k}(\Omega)} &\leq \liminf_{j \rightarrow \infty} \|u_{p_j}\|_{W^{2,k}(\Omega)} \\ &\leq \liminf_{j \rightarrow \infty} C \|g\|_{W^{2,k}(\Omega)}. \end{aligned}$$

Since this is true for any fixed k , it is clear that $u_\infty \in \bigcap_{k \in (1, \infty)} W^{2,k}(\Omega)$. Further, by the weak lower semi-continuity of the L^k norm, from (2.25) we may infer $\Delta u_\infty \in L^\infty(\Omega)$ and hence $u_\infty \in \mathcal{W}_g^{2,\infty}(\Omega)$, therefore concluding the proof. \square

2.6. Remark (elementary properties). We will throughout this exposition use the notation p to denote the exponent appearing in the Lagrangian and q its conjugate exponent which satisfies

$$(2.33) \quad \frac{1}{p} + \frac{1}{q} = 1.$$

For a given $v \in L^p(\Omega)$ it then holds that

$$(2.34) \quad \left\| |v|^{p-1} \right\|_{L^q(\Omega)} = \|v\|_{L^p(\Omega)}^{p-1}.$$

2.7. Proposition (Poincaré inequality). *Let $\Omega \subset \mathbb{R}^d$ be a bounded domain. For any $p \in [1, \infty]$, there exists a constant $C = C(\Omega, p) > 0$ depending only on Ω and p such that*

$$(2.35) \quad \|u\|_{L^p(\Omega)} \leq C(\Omega, p) \|Du\|_{L^p(\Omega)},$$

for all $u \in W_0^{1,p}(\Omega)$.

2.8. Proposition (Calderon-Zygmund estimate [16, Cor 9.10]). *Let $\Omega \subset \mathbb{R}^d$ be a bounded domain. Then, for any $p \in (1, \infty)$, there is a constant $C = C(d, p) > 0$ depending only on d and p such that*

$$(2.36) \quad \|D^2 u\|_{L^p(\Omega)} \leq C(d, p) \|\Delta u\|_{L^p(\Omega)},$$

for all $u \in W_0^{2,p}(\Omega)$.

An immediate consequence of Propositions 2.7 and 2.8 above is that the norm $\|\cdot\|_{2,p}$ is equivalent to either of the seminorms $\|D^2(\cdot)\|_{L^p(\Omega)}$ and $\|\Delta(\cdot)\|_{L^p(\Omega)}$ over the space $W_0^{2,p}(\Omega)$.

2.9. Mixed formulation of the p -Bilaplacian. The mixed formulation we propose to analyse is based on the observation that if $\phi(t) = |t|^{p-2}t$, the inverse is well defined as $\phi^{-1}(t) = \text{sgn}(t)|t|^{1/(p-1)} = |t|^{q-2}t$. Using this we make the following choice of auxiliary variable

$$(2.37) \quad w = |\Delta u|^{p-2} \Delta u$$

from which we can infer that

$$(2.38) \quad |w|^{q-2} w = \Delta u.$$

This allows us to write the problem as the mixed system:

$$(2.39) \quad \begin{cases} -\Delta u = |w|^{q-2} w, \\ -\Delta w = 0. \end{cases}$$

The mixed formulation can be written in a strong form as: Find a pair $(u, w) \in W_g^{2,p}(\Omega) \times L^q(\Omega)$ such that

$$(2.40) \quad \begin{cases} a(w, \psi) + b(u, \psi) = 0, \\ b(\phi, w) = 0, \end{cases} \quad \forall (\psi, \phi) \in L^q(\Omega) \times W_0^{2,p}(\Omega),$$

where the semilinear form $a(w, \psi)$ and bilinear form $b(u, \psi)$ are given by

$$(2.41) \quad \begin{cases} a(w, \psi) := \int_{\Omega} |w|^{q-2} w \psi \, d\mathbf{x} \\ b(u, \psi) := \int_{\Omega} -\Delta u \psi \, d\mathbf{x} \end{cases}$$

Notice that the problem (1.1) has been reformulated in a mixed form. Although we already know that the problem has a unique solution as a consequence of Corollary 2.4, we will show that the equivalent mixed formulation also admits a unique solution since the methodology will be useful henceforth. We begin with the following result.

2.10. Proposition (Inf-sup stability of $b(\cdot, \cdot)$ over $W_0^{2,p}(\Omega)$). *For any $u_0 \in W_0^{2,p}(\Omega)$, the bilinear form $b(\cdot, \cdot)$ satisfies the following inf-sup property:*

$$(2.42) \quad \|\Delta u_0\|_{L^p(\Omega)} \leq C \sup_{0 \neq v \in L^q(\Omega)} \frac{b(u_0, v)}{\|v\|_{L^q(\Omega)}}.$$

Proof Fix $u_0 \in W_0^{2,p}(\Omega)$. Then, we certainly have that $|\Delta u_0|^{p-2} \Delta u_0 \in L^q(\Omega)$. Therefore, by choosing $v = |\Delta u_0|^{p-2} \Delta u_0$ we have

$$(2.43) \quad b(u_0, v) = \|\Delta u\|_{L^p(\Omega)}^p$$

and that

$$(2.44) \quad \|v\|_{L^q(\Omega)} = \left\| |\Delta u_0|^{p-1} \right\|_{L^q(\Omega)} = \|\Delta u_0\|_{L^p(\Omega)}^{p-1},$$

in view of the property given in Remark 2.6. Hence we have

$$(2.45) \quad b(u_0, v) = \|\Delta u_0\|_{L^p(\Omega)}^p = \|\Delta u_0\|_{L^p(\Omega)} \|v\|_{L^q(\Omega)},$$

which implies the desired result. \square

2.11. Theorem (The mixed formulation is well posed). *For every $g \in W^{2,\infty}(\Omega)$, there exists a unique pair (u, w) solving (2.40) that satisfies*

$$(2.46) \quad \|\Delta u\|_{L^p(\Omega)} + \|w\|_{L^q(\Omega)}^{q-1} \leq C \|\Delta g\|_{L^p(\Omega)}.$$

Proof The results of Proposition 2.10 show that, for $u_0 := u - g \in W_0^{2,p}(\Omega)$, we have

$$(2.47) \quad \begin{aligned} \|\Delta u_0\|_{L^p(\Omega)} &\leq \sup_{0 \neq v \in L^q(\Omega)} \frac{b(u_0, v)}{\|v\|_{L^q(\Omega)}} \\ &\leq \sup_{0 \neq v \in L^q(\Omega)} \frac{b(u, v)}{\|v\|_{L^q(\Omega)}} + \sup_{0 \neq v \in L^q(\Omega)} \frac{b(g, v)}{\|v\|_{L^q(\Omega)}} \\ &\leq \sup_{0 \neq v \in L^q(\Omega)} \frac{-a(w, v)}{\|v\|_{L^q(\Omega)}} + \sup_{0 \neq v \in L^q(\Omega)} \frac{b(g, v)}{\|v\|_{L^q(\Omega)}}. \end{aligned}$$

in view of (2.40). Now, by using Remark 2.6 we estimate

$$(2.48) \quad \begin{aligned} \|\Delta u_0\|_{L^p(\Omega)} &\leq \|w^{q-1}\|_{L^p(\Omega)} + \|\Delta g\|_{L^p(\Omega)} \\ &\leq \|w\|_{L^q(\Omega)}^{q-1} + \|\Delta g\|_{L^p(\Omega)}. \end{aligned}$$

Now take $\psi = w$ in (2.40). Then,

$$(2.49) \quad a(w, w) + b(u, w) = 0.$$

Set $\phi = u_0$ in (2.40). Then,

$$(2.50) \quad b(u_0, w) = 0$$

and in particular

$$(2.51) \quad a(w, w) + b(u, w) - b(u_0, w) = 0.$$

This in turn implies

$$(2.52) \quad a(w, w) + b(g, w) = 0,$$

or explicitly

$$(2.53) \quad \int_{\Omega} |w|^q - \Delta g w \, d\mathbf{x} = 0.$$

Hence

$$(2.54) \quad \begin{aligned} \|w\|_{L^q(\Omega)}^q &= \int_{\Omega} \Delta g w \, d\mathbf{x} \\ &\leq \|\Delta g\|_{L^p(\Omega)} \|w\|_{L^q(\Omega)}, \end{aligned}$$

and

$$(2.55) \quad \|w\|_{L^q(\Omega)}^{q-1} \leq \|\Delta g\|_{L^p(\Omega)},$$

which yields the desired result upon noting

$$(2.56) \quad \|\Delta u\|_{L^p(\Omega)} \leq \|\Delta u_0\|_{L^p(\Omega)} + \|\Delta g\|_{L^p(\Omega)}$$

and combining with (2.48). \square

2.12. Remark (Convergence to “weak” solutions to the ∞ -Bilaplacian). Theorem 2.5 guarantees convergence to a candidate ∞ -Harmonic function. The correct notion of weak solution to the limiting problem

$$(2.57) \quad \begin{cases} (\Delta u)^3 |\mathcal{D}(\Delta u)|^2 = 0, & \text{in } \Omega, \\ u = g, & \text{on } \partial\Omega, \\ \mathcal{D}u = \mathcal{D}g, & \text{on } \partial\Omega, \end{cases}$$

is that of \mathcal{D} -solutions [21, 20]. The solution is probabilistic in nature and interpreted in a weak sense. It is the only candidate ∞ -Biharmonic function which means Theorem 2.5 guarantees convergence of the sequence of p -Biharmonic functions to the unique ∞ -Biharmonic \mathcal{D} -solution.

3. DISCRETISATION OF THE p -BILAPLACIAN

In this section we describe a mixed finite element discretisation of the p -Bilaplacian. Let \mathcal{T} be a conforming triangulation of Ω , namely, \mathcal{T} is a finite family of sets such that

- (1) $K \in \mathcal{T}$ implies K is an open simplex (segment for $d = 1$, triangle for $d = 2$, tetrahedron for $d = 3$),
- (2) for any $K, J \in \mathcal{T}$ we have that $\overline{K} \cap \overline{J}$ is a full lower-dimensional simplex (i.e., it is either \emptyset , a vertex, an edge, a face, or the whole of \overline{K} and \overline{J}),
- (3) $\bigcup_{K \in \mathcal{T}} \overline{K} = \overline{\Omega}$.

The shape regularity constant of \mathcal{T} is defined as the number

$$(3.1) \quad \mu(\mathcal{T}) := \inf_{K \in \mathcal{T}} \frac{\rho_K}{h_K},$$

where ρ_K is the radius of the largest ball contained inside K and h_K is the diameter of K . An indexed family of triangulations $\{\mathcal{T}^n\}_n$ is called *shape regular* if

$$(3.2) \quad \mu := \inf_n \mu(\mathcal{T}^n) > 0.$$

We let \mathcal{E} be the skeleton (set of common interfaces) of the triangulation \mathcal{T} and say $e \in \mathcal{E}$ if e is on the interior of Ω and $e \in \partial\Omega$ if e lies on the boundary $\partial\Omega$.

We let $\mathbb{P}^k(\mathcal{T})$ denote the space of piecewise polynomials of degree $k \geq 2$ over the triangulation \mathcal{T} , that is,

$$(3.3) \quad \mathbb{P}^k(\mathcal{T}) = \{\phi \text{ such that } \phi|_K \in \mathbb{P}^k(K)\},$$

and introduce the *finite element space*

$$(3.4) \quad \mathbb{V} := \mathbb{P}^k(\mathcal{T}) \cap C^0(\Omega),$$

to be the usual space of continuous piecewise polynomial functions. We define jump operators for arbitrary scalar functions v and vectors \mathbf{v} over an edge e shared by elements K_1 and K_2 as $\llbracket v \rrbracket = v|_{K_1} \mathbf{n}_{K_1} + v|_{K_2} \mathbf{n}_{K_2}$, $\llbracket \mathbf{v} \rrbracket = \mathbf{v}|_{K_1} \cdot \mathbf{n}_{K_1} + \mathbf{v}|_{K_2} \cdot \mathbf{n}_{K_2}$ and when e is on $\partial\Omega$ we understand $\llbracket v \rrbracket = v|_K \mathbf{n}_{\partial\Omega}$ and $\llbracket \mathbf{v} \rrbracket = \mathbf{v}|_K \cdot \mathbf{n}_{\partial\Omega}$.

Further, we define $h : \Omega \rightarrow \mathbb{R}$ to be the piecewise constant *meshsize function* of \mathcal{T} given by

$$(3.5) \quad h(\mathbf{x}) := \max_{\overline{K} \ni \mathbf{x}} h_K.$$

A mesh is called quasi-uniform when there exists a positive constant C such that $\max_{x \in \Omega} h \leq C \min_{x \in \Omega} h$. In what follows we shall assume that all triangulations are shape-regular and quasi-uniform although the results may be extendable even in the non-quasi-uniform case using techniques developed in [9].

3.1. Definition (Ritz projection operators). The Ritz projection operator R is defined through requiring

$$(3.6) \quad \int_{\Omega} D(Rv) \cdot D\phi \, d\mathbf{x} = \int_{\Omega} Dv \cdot D\phi \, d\mathbf{x} \quad \forall \phi \in \mathbb{V} \cap H_0^1(\Omega),$$

and Rv coincides with an appropriate interpolant of v on the boundary. This operator satisfies the following approximation properties for quasi-uniform meshes [28]: for any $v \in W^{k+1,q}(\Omega)$, and $k \geq 2$

$$(3.7) \quad \|v - Rv\|_{L^q(\Omega)} + \|h(Dv - D(Rv))\|_{L^q(\Omega)} + \left(\sum_{K \in \mathcal{T}} \|h^2(\Delta v - \Delta(Rv))\|_{L^q(K)}^q \right)^{1/q} \leq Ch^{k+1} |v|_{k+1,q}.$$

The Neumann Ritz projection \bar{R} is defined through requiring orthogonality over a larger space

$$(3.8) \quad \int_{\Omega} D(\bar{R}w) \cdot D\psi \, d\mathbf{x} = \int_{\Omega} Dw \cdot D\psi \, d\mathbf{x} \quad \forall \psi \in \mathbb{V}$$

and requiring

$$(3.9) \quad \int_{\Omega} \bar{R}w \, d\mathbf{x} = \int_{\Omega} w \, d\mathbf{x}.$$

The results of [28] also imply that \bar{R} satisfies the same approximation properties as R .

3.2. Definition (Mesh-dependent norms). We introduce the mesh-dependent L^p - and $W^{2,p}$ -norms to be

$$(3.10) \quad \begin{aligned} \|w_h\|_{L_h^p(\Omega)}^p &:= \|w_h\|_{L^p(\Omega)}^p + \|h^{1/p} w_h\|_{L^p(\mathcal{E})}^p \\ \|w_h\|_{W_h^{2,p}(\Omega)}^p &:= \|\Delta_h w_h\|_{L^p(\Omega)}^p + \|h^{1/p-1} [Dw_h]\|_{L^p(\mathcal{E})}^p, \end{aligned}$$

where Δ_h denotes an elementwise Laplace operator.

3.3. Galerkin discretisation. Consider the space

$$(3.11) \quad \mathbb{V}_g := \{\phi \in \mathbb{V} : \phi|_{\partial\Omega} = Rg\}.$$

Then, we consider the Galerkin discretisation of (2.11), to find $(u_h, w_h) \in \mathbb{V}_g \times \mathbb{V}$ such that

$$(3.12) \quad \begin{aligned} a(w_h, \psi) + b_h(u_h, \psi) &= 0 \\ b_h(\phi, w_h) &= 0, \quad \forall (\psi, \phi) \in \mathbb{V} \times \mathbb{V}_0, \end{aligned}$$

where the bilinear form $a(\cdot, \cdot)$ is given in (2.41), $b_h(\cdot, \cdot)$ is a consistent discretisation of $b(\cdot, \cdot)$ given by

$$(3.13) \quad b_h(u_h, \psi) = - \sum_{K \in \mathcal{T}} \int_K \Delta u_h \psi \, d\mathbf{x} + \int_{\mathcal{E}} [Du_h] \psi \, ds.$$

Notice that the method is equivalent to finding $(u_h, w_h) \in \mathbb{V}_g \times \mathbb{V}$ such that

$$(3.14) \quad \begin{aligned} \int_{\Omega} |w_h|^{q-2} w_h \psi + Du_h \cdot D\psi \, d\mathbf{x} &= \int_{\partial\Omega} Dg \cdot \mathbf{n} \psi \, ds \\ \int_{\Omega} Dw_h \cdot D\phi \, d\mathbf{x} &= 0, \quad \forall (\psi, \phi) \in \mathbb{V} \times \mathbb{V}_0. \end{aligned}$$

Hence the Ritz projection operator from Definition 3.1 is the b_h -orthogonal projection onto \mathbb{V}_g , that is, for $v \in H_g^1(\Omega)$

$$(3.15) \quad b_h(Rv - v, \phi) = 0 \quad \forall \phi \in \mathbb{V}_0.$$

3.4. Remark. The reason for defining the mesh-dependent norms as we do is to ensure the boundedness property

$$(3.16) \quad |b_h(u_h, v_h)| \leq \|u_h\|_{W_h^{2,p}(\Omega)} \|v_h\|_{L_h^q(\Omega)}.$$

The scaling in the edge terms is chosen so that for arbitrary $v_h \in \mathbb{V}$ each mesh-dependent norm is equivalent to the continuous counterpart, that is $\|v_h\|_{L_h^p(\Omega)} \sim \|v_h\|_{L^p(\Omega)}$ for example.

3.5. Lemma. Assume the mesh is quasi-uniform, then the bilinear form $b_h(\cdot, \cdot)$ satisfies the following inf-sup property: for any $\Phi \in \mathbb{V}_0$,

$$(3.17) \quad \|\Phi\|_{W_h^{2,p}(\Omega)} \leq C \sup_{0 \neq v_h \in \mathbb{V}_0} \frac{b_h(\Phi, v_h)}{\|v_h\|_{L_h^q(\Omega)}}.$$

Proof The proof of this fact takes inspiration from [29] (see also [14] and [12] for related ideas). We begin by showing that there exists a function v that is discrete but not an element of \mathbb{V}_0 such that

$$(3.18) \quad b_h(\Phi, Rv) \geq C \|\Phi\|_{W_h^{2,p}(\Omega)}^p$$

and then showing the discrete stability estimate that $\|Rv\|_{L_h^q(\Omega)} \leq C \|\Phi\|_{W_h^{2,p}(\Omega)}^{p-1}$.

To begin we denote b_K as the cubic a posteriori bubble function. This is a function that is \mathbb{P}^3 , positive over K , extended by zero outside of K and satisfies that $\|b_K\|_{L^\infty(K)} = 1$. Now take v_1 such that $v_1|_K = -b_K |\Delta\Phi|^{p-2} \Delta\Phi$. Notice that $v_1 \in W_0^{1,q}(\Omega)$ and that $v_1|_e = 0$ for all $e \in \mathcal{E} \cup \partial\Omega$. Then through the equivalence of norms over finite dimensional linear spaces.

$$(3.19) \quad \frac{1}{C} \sum_{K \in \mathcal{T}} \|\Delta\Phi\|_{L^p(K)}^p \leq \sum_{K \in \mathcal{T}} \int_K b_K |\Delta\Phi|^p d\mathbf{x} = b_h(\Phi, v_1) = b_h(\Phi, Rv_1).$$

Now let b_e be the edge bubble function that vanishes over all vertices of \mathcal{T} . Again this is a polynomial that is positive over K , extended by zero outside of the two elements sharing $e \in \mathcal{E}$ and satisfies $\|b_e\|_{L^\infty(e)} = 1$. Define $v_e : e \rightarrow \mathbb{R}$ such that $v_e = h^{1-p} \|\llbracket D\Phi \rrbracket\|^{p-2} \llbracket D\Phi \rrbracket$ on the face e and extended by a constant on the direction normal to e . Set $v_2 := \sum_{e \in \mathcal{E}} b_e v_e$ then we have $v_2 \in W_0^{1,q}(\Omega)$ and

$$(3.20) \quad \begin{aligned} b_h(\Phi, v_2) &= \sum_{K \in \mathcal{T}} \int_K -\Delta\Phi v_2 d\mathbf{x} + \int_{\mathcal{E}} \llbracket D\Phi \rrbracket v_2 ds \\ &= \sum_{K \in \mathcal{T}} \int_K -\Delta\Phi v_2 d\mathbf{x} + \int_{\mathcal{E}} b_e h^{1-p} \|\llbracket D\Phi \rrbracket\|^p ds. \end{aligned}$$

Now equivalence of norms shows there exists a constant $C > 0$ independent of Φ and h such that

$$(3.21) \quad \begin{aligned} \frac{1}{C} \left\| h^{1/p-1} \llbracket D\Phi \rrbracket \right\|_{L^p(\mathcal{E})}^p &\leq \int_{\mathcal{E}} b_e h^{1-p} \|\llbracket D\Phi \rrbracket\|^p ds \\ &= b_h(\Phi, v_2) + \sum_{K \in \mathcal{T}} \int_K \Delta\Phi v_2 d\mathbf{x} \\ &\leq b_h(\Phi, Rv_2) + \left(\sum_{K \in \mathcal{T}} \int_K |\Delta\Phi|^p d\mathbf{x} \right)^{1/p} \|v_2\|_{L^q(\Omega)}. \end{aligned}$$

Young's inequality with ϵ shows that

$$(3.22) \quad \begin{aligned} \left(\sum_{K \in \mathcal{T}} \int_K |\Delta\Phi|^p d\mathbf{x} \right)^{1/p} \|v_2\|_{L^q(\Omega)} &\leq C(\epsilon) \left(\sum_{K \in \mathcal{T}} \int_K |\Delta\Phi|^p d\mathbf{x} \right) + \epsilon \|v_2\|_{L^q(\Omega)}^q \\ &\leq C(\epsilon) \left(\sum_{K \in \mathcal{T}} \int_K |\Delta\Phi|^p d\mathbf{x} \right) + C\epsilon \left\| h^{1/p-1} \llbracket D\Phi \rrbracket \right\|_{L^p(\mathcal{E})}^p \end{aligned}$$

in view of the definition of v_2 . Now substituting (3.22) into (3.21) and choosing ϵ appropriately small we see that

$$(3.23) \quad \begin{aligned} \left\| h^{1/p-1} \llbracket D\Phi \rrbracket \right\|_{L^p(\mathcal{E})}^p &\leq C \left(b_h(\Phi, Rv_2) + \left(\sum_{K \in \mathcal{T}} \int_K |\Delta\Phi|^p d\mathbf{x} \right) \right) \\ &\leq C(b_h(\Phi, Rv_2) + b_h(\Phi, Rv_1)), \end{aligned}$$

by (3.19). Hence with $v = v_1 + v_2$ we have shown (3.18).

We must now show the stability bound. To begin we show a stability result for the Ritz projection. With $z \in W^{2,p}(\Omega) \cap H_0^1(\Omega)$ solving the problem

$$(3.24) \quad -\Delta z = |Rv|^{q-2} Rv,$$

we see

$$(3.25) \quad \|Rv\|_{L^q(\Omega)}^q = b_h(z, Rv) = b_h(z, Rv - v) + b_h(z, v) = b_h(z - z_h, Rv - v) + b_h(z, v),$$

with z_h chosen as the Clément interpolant of z . Now using the definition of z and approximation properties of z_h

$$(3.26) \quad \begin{aligned} \|Rv\|_{L^q(\Omega)}^q &\leq C \|h(D(Rv) - Dv)\|_{L^q(\Omega)} \|z\|_{W^{2,p}(\Omega)} + \int_{\Omega} |Rv|^{q-2} Rv v \, dx \\ &\leq C \|Rv\|_{L^q(\Omega)}^{q-1} \left(\|h(D(Rv) - Dv)\|_{L^q(\Omega)} + \|v\|_{L^q(\Omega)} \right). \end{aligned}$$

Using the $W^{1,q}$ stability of R from [28] we have

$$(3.27) \quad \|Rv\|_{L^q(\Omega)} \leq C \left(\|hDv\|_{L^q(\Omega)} + \|v\|_{L^q(\Omega)} \right).$$

Notice we have not used the super-approximation ideas from [29, 12] and are working on quasi-uniform meshes only. Now for $v = v_1 + v_2$ defined above we are able to use inverse inequalities to see that

$$(3.28) \quad \|Rv\|_{L^q(\Omega)} \leq C(p) \|v\|_{L^q(\Omega)},$$

and through the definition of v_1 and v_2 we have

$$(3.29) \quad \|v\|_{L^q(\Omega)} \leq C \|\Phi\|_{W_h^{2,p}(\Omega)}^{p-1}.$$

Hence

$$(3.30) \quad \|Rv\|_{L_h^q(\Omega)} \|\Phi\|_{W_h^{2,p}(\Omega)} \leq C \|Rv\|_{L^q(\Omega)} \|\Phi\|_{W_h^{2,p}(\Omega)} \leq C \|v\|_{L^q(\Omega)} \|\Phi\|_{W_h^{2,p}(\Omega)} \leq C \|\Phi\|_{W_h^{2,p}(\Omega)}^p \leq C b_h(\Phi, Rv)$$

and certainly

$$(3.31) \quad \|\Phi\|_{W_h^{2,p}(\Omega)} \leq C \frac{b_h(\Phi, Rv)}{\|Rv\|_{L_h^q(\Omega)}} \leq \sup_{v_h \in \mathbb{V}_0} C \frac{b_h(\Phi, v_h)}{\|v_h\|_{L_h^q(\Omega)}},$$

concluding the proof. \square

3.6. Theorem (existence and uniqueness of solution to (3.12)). *There exists a unique pair $(u_h, w_h) \in \mathbb{V}_g \times \mathbb{V}$ solving (3.12). They satisfy the stability bound*

$$(3.32) \quad \|u_h\|_{W_h^{2,p}(\Omega)} + \|w_h\|_{L^q(\Omega)}^{q-1} \leq C \|\Delta g\|_{L^p(\Omega)}.$$

Note that since $g \in W^{2,\infty}(\Omega)$, the right hand side of (3.32) is finite.

Proof The proof of this mirrors that of Theorem 2.11. We begin by noting that for $\psi = w_h$ we have

$$(3.33) \quad a(w_h, w_h) + b_h(u_h, w_h) = 0.$$

Now for $\phi = u_{h,0} := u_h - Rg$ we see that

$$(3.34) \quad b_h(u_h - Rg, w_h) = 0,$$

hence

$$(3.35) \quad 0 = a(w_h, w_h) + b_h(Rg, w_h).$$

Now, by definition, we obtain

$$(3.36) \quad \begin{aligned} \|w_h\|_{L^q(\Omega)}^q &\leq \|Rg\|_{W_h^{2,p}(\Omega)} \|w_h\|_{L_h^q(\Omega)} \\ &\leq C \|\Delta g\|_{L^p(\Omega)} \|w_h\|_{L^q(\Omega)}, \end{aligned}$$

by Remark 3.4 and Lemma 3.5 and hence

$$(3.37) \quad \|w_h\|_{L^q(\Omega)}^{q-1} \leq \|\Delta g\|_{L^p(\Omega)}.$$

The result follows because

$$\begin{aligned}
\|u_{h,0}\|_{W_h^{2,p}(\Omega)} &\leq C \sup_{0 \neq v_h \in \mathbb{V}_0} \frac{b_h(u_{h,0}, v_h)}{\|v_h\|_{L_h^q(\Omega)}} \\
&\leq C \left(\sup_{0 \neq v_h \in \mathbb{V}_0} \frac{b_h(u_h, v_h)}{\|v_h\|_{L_h^q(\Omega)}} + \sup_{0 \neq v_h \in \mathbb{V}_0} \frac{b_h(Rg, v_h)}{\|v_h\|_{L_h^q(\Omega)}} \right) \\
(3.38) \quad &\leq C \left(\sup_{0 \neq v_h \in \mathbb{V}_0} \frac{-a(w_h, v_h)}{\|v_h\|_{L_h^q(\Omega)}} + \sup_{0 \neq v_h \in \mathbb{V}_0} \frac{b_h(g, v_h)}{\|v_h\|_{L_h^q(\Omega)}} \right) \\
&\leq C \left(\| |w_h|^{q-1} \|_{L^p(\Omega)} + \|\Delta g\|_{L^p(\Omega)} \right) \\
&\leq C \left(\|w_h\|_{L^q(\Omega)}^{q-1} + \|\Delta g\|_{L^p(\Omega)} \right)
\end{aligned}$$

by the discrete inf-sup condition in Lemma 3.5 and the same argument as in the proof of Theorem 2.11. Since

$$\begin{aligned}
\|u_h\|_{W_h^{2,p}(\Omega)} &\leq \|u_{h,0}\|_{W_h^{2,p}(\Omega)} + \|Rg\|_{W_h^{2,p}(\Omega)} \\
(3.39) \quad &\leq C \left(\|u_{h,0}\|_{W_h^{2,p}(\Omega)} + \|\Delta g\|_{L^p(\Omega)} \right)
\end{aligned}$$

combining (3.36), (3.38) and (3.39) concludes the proof. \square

Next we state some technical properties that will be used in the theorem that follows.

3.7. Lemma (Properties of $a(\cdot, \cdot)$, cf. [32, Prop 3.1]). *With $w \in L^q(\Omega)$ and $w_h, v_h \in \mathbb{V}$, for any $p \geq 2$, there exist constants*

(1) $C_1 > 0$ such that

$$(3.40) \quad C_1 \frac{\|w - w_h\|_{L^q(\Omega)}^2}{\|w\|_{L^q(\Omega)}^{2-q} + \|w_h\|_{L^q(\Omega)}^{2-q}} \leq a(w, w - w_h) - a(w_h, w - w_h).$$

(2) $C_2 > 0$ such that

$$(3.41) \quad C_2 \int_{\Omega} \left| |w|^{q-2} w - |w_h|^{q-2} w_h \right| |w - w_h| \, d\mathbf{x} \leq a(w, w - w_h) - a(w_h, w - w_h).$$

(3) $C_3 > 0$ such that

$$(3.42) \quad a(w, w - v_h) - a(w_h, w - v_h) \leq C_3 \left(\int_{\Omega} \left| |w|^{q-2} w - |w_h|^{q-2} w_h \right| |w - w_h| \, d\mathbf{x} \right)^{1/p} \|w - v_h\|_{L^q(\Omega)}.$$

3.8. Theorem (Approximability of the numerical schemes). *Let $(u, w) \in W_g^{k+1,p}(\Omega) \times W^{k+1,q}(\Omega)$ be the unique solution of (2.40) and $(u_h, w_h) \in \mathbb{V}_g \times \mathbb{V}$ be the finite element approximation satisfying (3.12). Then, the following error estimate holds*

$$(3.43) \quad \|w - w_h\|_{L^q(\Omega)} + \|u - u_h\|_{W_h^{2,p}(\Omega)}^{p-1} \leq C \left(h^{\frac{q}{2}(k+1)} |w|_{W^{k+1,q}(\Omega)}^{q/2} + h^{k+1} |w|_{W^{k+1,q}(\Omega)} + h^{k-1} |u|_{W^{k+1,p}(\Omega)} \right).$$

Proof We begin by noting the Galerkin orthogonality results

$$\begin{aligned}
(3.44) \quad &b_h(\phi, w - w_h) = 0 \quad \forall \phi \in \mathbb{V}_0, \\
&a(w, \psi) - a(w_h, \psi) + b_h(u - u_h, \psi) = 0 \quad \forall \psi \in \mathbb{V},
\end{aligned}$$

in view of (2.40) and (3.12).

Now using Lemma 3.7 we have

$$\begin{aligned}
(3.45) \quad &\frac{C_1 \|w - w_h\|_{L^q(\Omega)}^2}{2(\|w\|_{L^q(\Omega)}^{2-q} + \|w_h\|_{L^q(\Omega)}^{2-q})} + \frac{C_2}{2} \int_{\Omega} \left| |w|^{q-2} w - |w_h|^{q-2} w_h \right| |w - w_h| \, d\mathbf{x} \leq a(w, w - w_h) - a(w_h, w - w_h)
\end{aligned}$$

Now using the semilinearity of $a(\cdot, \cdot)$ we have, for $\chi \in \mathbb{V}$ denoting some approximation of w to be chosen, that

$$(3.46) \quad \begin{aligned} a(w, w - w_h) - a(w_h, w - w_h) &= a(w, w - \chi) - a(w_h, w - \chi) + a(w, \chi - w_h) - a(w_h, \chi - w_h) \\ &= \underbrace{a(w, w - \chi) - a(w_h, w - \chi)}_{=:I} + \underbrace{a(w, \chi - w_h) - a(w_h, \chi - w_h)}_{=:II}, \end{aligned}$$

in view of (3.44). We proceed to bound these terms separately, starting with I.

Making use of Lemma 3.7

$$(3.47) \quad a(w, w - \chi) - a(w_h, w - \chi) \leq C_3 \left(\int_{\Omega} \left| |w|^{q-2} w - |w_h|^{q-2} w_h \right| |w - w_h| \, d\mathbf{x} \right)^{1/p} \|w - \chi\|_{L^q(\Omega)}.$$

Young's inequality with ϵ states for $a, b, \epsilon > 0$

$$(3.48) \quad ab \leq \frac{1}{p}(\epsilon a)^p + \frac{1}{q} \left(\frac{b}{\epsilon} \right)^q,$$

which, upon applying to (3.47), shows

$$(3.49) \quad a(w, w - \chi) - a(w_h, w - \chi) \leq \frac{\epsilon^p}{p} \int_{\Omega} \left| |w|^{q-2} w - |w_h|^{q-2} w_h \right| |w - w_h| \, d\mathbf{x} + \frac{C_3^q}{q\epsilon^q} \|w - \chi\|_{L^q(\Omega)}^q.$$

Now choosing $\epsilon = \left(\frac{C_3^q}{2} \right)^{1/p}$ and we have

$$(3.50) \quad a(w, w - \chi) - a(w_h, w - \chi) \leq \frac{C_2}{2} \int_{\Omega} \left| |w|^{q-2} w - |w_h|^{q-2} w_h \right| |w - w_h| \, d\mathbf{x} + C(q) \|w - \chi\|_{L^q(\Omega)}^q.$$

Notice we have picked ϵ such that the first term on the right hand side of (3.50) will cancel with the second term on the left hand side of (3.45).

To control II we pick χ such that

$$(3.51) \quad b_h(\phi, \chi) = 0 \quad \forall \phi \in \mathbb{V}_0.$$

An example of such an operator is the Neumann Ritz projection operator, $\bar{R}w$, given in Definition 3.1. With this choice of χ , noting the definition of w_h from (3.12), it is clear that

$$(3.52) \quad b_h(\phi, w_h - \chi) = 0 \quad \forall \phi \in \mathbb{V}_0,$$

and hence

$$(3.53) \quad b_h(u - u_h, w_h - \chi) = b_h(u - u_h - R(u - u_h), w_h - \chi) = b_h(u - Ru, w_h - \chi).$$

Now making use of the boundedness of $b_h(\cdot, \cdot)$ we have

$$(3.54) \quad \begin{aligned} b_h(u - u_h, w_h - \chi) &\leq \|u - Ru\|_{W_h^{2,p}(\Omega)} \|w_h - \chi\|_{L_h^q(\Omega)} \\ &\leq C \|u - Ru\|_{W_h^{2,p}(\Omega)} \|w_h - \chi\|_{L^q(\Omega)} \\ &\leq \frac{C}{4\epsilon} \|u - Ru\|_{W_h^{2,p}(\Omega)}^2 + \epsilon \|w_h - \chi\|_{L^q(\Omega)}^2 \\ &\leq \frac{C}{4\epsilon} \|u - Ru\|_{W_h^{2,p}(\Omega)}^2 + 2\epsilon \left(\|w - w_h\|_{L^q(\Omega)}^2 + \|w - \chi\|_{L^q(\Omega)}^2 \right). \end{aligned}$$

Substituting (3.50) and (3.54) into (3.45) and choosing ϵ small enough we see

$$(3.55) \quad \|w - w_h\|_{L^q(\Omega)}^2 \leq C \left(\|w - \chi\|_{L^q(\Omega)}^q + \|u - Ru\|_{W_h^{2,p}(\Omega)}^2 + \|w - \chi\|_{L^q(\Omega)}^2 \right),$$

allowing us to use the approximability of R and \bar{R} concluding the proof of the auxiliary variable.

To show a bound for the primal variable we make use of the inf-sup condition from Lemma 3.5, noting that in view of Galerkin orthogonality and the definition of R we have

$$(3.56) \quad \begin{aligned} 0 &= a(w, \phi) - a(w_h, \phi) + b_h(u - u_h, \phi) \\ &= a(w, \phi) - a(w_h, \phi) + b_h(Ru - u_h, \phi) \quad \forall \phi \in \mathbb{V}_0. \end{aligned}$$

It is then clear that

$$\begin{aligned}
\|Ru - u_h\|_{W_h^{2,p}(\Omega)} &\leq \sup_{0 \neq \phi \in \mathbb{V}_0} \frac{b_h(Ru - u_h, \phi)}{\|\phi\|_{L_h^q(\Omega)}} \\
&= \sup_{0 \neq \phi \in \mathbb{V}_0} \frac{a(w_h, \phi) - a(w, \phi)}{\|\phi\|_{L_h^q(\Omega)}} \\
(3.57) \quad &\leq C_3 \sup_{0 \neq \phi \in \mathbb{V}_0} \frac{\left(\int_{\Omega} |w|^{p-2} w - |w_h|^{p-2} w_h |w - w_h| \, d\mathbf{x} \right)^{1/p}}{\|\phi\|_{L_h^q(\Omega)}} \\
&\leq C_3 C \left(\int_{\Omega} |w|^{p-2} w - |w_h|^{p-2} w_h |w - w_h| \, d\mathbf{x} \right)^{1/p},
\end{aligned}$$

through the equivalence of the L^q -norm and its discrete counterpart. Now by Lemma 3.7 and Young's inequality with ϵ we have

$$\begin{aligned}
(3.58) \quad C_2 \int_{\Omega} |w|^{p-2} w - |w_h|^{p-2} w_h |w - w_h| \, d\mathbf{x} &\leq a(w, w - w_h) - a(w_h, w - w_h) \\
&\leq C_3 \left(\int_{\Omega} |w|^{p-2} w - |w_h|^{p-2} w_h |w - w_h| \, d\mathbf{x} \right)^{1/p} \|w - w_h\|_{L^q(\Omega)} \\
&\leq \frac{\epsilon^p}{p} \int_{\Omega} |w|^{p-2} w - |w_h|^{p-2} w_h |w - w_h| \, d\mathbf{x} + \frac{C_3^q}{q\epsilon^q} \|w - w_h\|_{L^q(\Omega)}^q.
\end{aligned}$$

The particular choice $\epsilon = \left(\frac{pC_2}{2}\right)^{1/p}$ then shows that

$$(3.59) \quad \int_{\Omega} |w|^{p-2} w - |w_h|^{p-2} w_h |w - w_h| \, d\mathbf{x} \leq C \|w - w_h\|_{L^q(\Omega)}^q.$$

Substituting (3.59) into (3.57) results in

$$(3.60) \quad \|Ru - u_h\|_{W_h^{2,p}(\Omega)} \leq C \|w - w_h\|_{L^q(\Omega)}^{q/p}.$$

The result follows from the fact

$$(3.61) \quad \|u - u_h\|_{W_h^{2,p}(\Omega)} \leq \|Ru - u_h\|_{W_h^{2,p}(\Omega)} + \|Ru - u\|_{W_h^{2,p}(\Omega)}$$

and using the approximation properties of the Ritz projection, concluding the proof. \square

3.9. Remark (Optimality of the bounds). Notice that the rates trail off as p gets large. A similar phenomena was noticed when constructing methods for the p -Laplacian [7, Thm 5.3.5] where for a conforming piecewise linear approximation, u_h , the error behaved like

$$(3.62) \quad \|u - u_h\|_{W^{1,p}(\Omega)} \leq Ch^{1/(p-1)}.$$

An analysis based on quasi-norms [5] was then introduced to rectify this. It may be possible to use these techniques to show optimal error bounds for the p -Bilaplacian based on the quasi-norm

$$(3.63) \quad \|u\|_{v,p}^p := \int_{\Omega} |\Delta u|^2 (|\Delta u| + |\Delta v|)^{p-2} \, d\mathbf{x}.$$

We shall not push this point further in this work however. Instead, in order to try to characterise the limiting problem, we shall focus on convergence under minimal regularity.

We begin by defining the semilinear form

$$(3.64) \quad c((u, w), (\phi, \psi)) := a(w, \psi) + b_h(u, \psi) + b_h(\phi, w),$$

then the discrete mixed form of the Bilaplacian can be written, equivalently to (3.12), as seeking $(u_h, w_h) \in \mathbb{V}_g \times \mathbb{V}$ such that

$$(3.65) \quad c((u_h, w_h), (\phi, \psi)) = 0 \quad \forall (\phi, \psi) \in \mathbb{V}_0 \times \mathbb{V}.$$

3.10. Theorem (Convergence under minimal regularity). *Let (u_h, w_h) be a sequence of finite element solutions of (3.12) indexed by the mesh parameter h and let also $u \in W_g^{2,p}(\Omega)$ be the solution of the p -Bilaplacian. Then we have*

- $u_h \rightarrow u$ strongly in L^p as $h \rightarrow 0$,
- $w_h \rightharpoonup w$ weakly in L^q as $h \rightarrow 0$.

Proof The stability result given in Theorem 3.6 allows us to infer that the sequence (u_h, w_h) is bounded uniformly in h . This means, up to a subsequence, that there exists a $(u^*, w^*) \in W_g^{2,p}(\Omega) \times L^q(\Omega)$ such that $u_h \rightarrow u^*$ strongly in $L^p(\Omega)$ and $w_h \rightharpoonup w^*$ weakly in $L^q(\Omega)$.

Now suppose $v_1 \in C^\infty(\Omega)$. Take $(\phi, \psi) = (0, \bar{R}v_1)$ in (3.65). Then,

$$(3.66) \quad 0 = c((u_h, w_h), (0, \bar{R}v_1)) = a(w_h, \bar{R}v_1) + b_h(u_h, \bar{R}v_1).$$

Since $u_h \rightarrow u^*$ and by the properties of the projection \bar{R} given in Definition 3.1 we have that

$$(3.67) \quad b_h(u_h, \bar{R}v_1) \rightarrow b(u^*, v_1).$$

Also, since $w_h \rightharpoonup w^*$ and $\bar{R}v_1 \rightarrow v_1$ strongly we have

$$(3.68) \quad a(w_h, \bar{R}v_1) \rightarrow a(w^*, v_1).$$

Hence

$$(3.69) \quad a(w_h, \bar{R}v_1) + b_h(u_h, \bar{R}v_1) \rightarrow a(w^*, v_1) + b(u^*, v_1).$$

Now suppose $v_2 \in C_0^\infty(\Omega)$ and take $(\phi, \psi) = (Rv_2, 0)$ in (3.65), then

$$(3.70) \quad b_h(w_h, Rv_2) = 0.$$

By the same arguments we have

$$(3.71) \quad b_h(Rv_2, w_h) \rightarrow b(v_2, w^*).$$

Using density of $C_0^\infty(\Omega) \times C^\infty(\Omega)$ functions in $W_0^{2,p}(\Omega) \times L^q(\Omega)$ shows that (u^*, w^*) must solve the Bilaplacian and since the solution was unique, the whole sequence $(u_h, w_h) \rightarrow (u, w)$. \square

3.11. Corollary. *Let $u_{h,p} \in \mathbb{V}_g$ be the Galerkin solution of (3.12) and let u_∞ denote a candidate ∞ -Biharmonic function. Then, along a subsequence we have*

$$(3.72) \quad u_{h,p_j} \rightarrow u_\infty \in C^0(\bar{\Omega}) \text{ as } p \rightarrow \infty \text{ and } h \rightarrow 0.$$

3.12. Remark. Since there exists a unique subsequential p -Biharmonic limit u_∞ to the ∞ -Bilaplacian on Ω the whole sequence must converge to this function, that is

$$(3.73) \quad u_{h,p} \rightarrow u_\infty \in C^0(\bar{\Omega}) \text{ as } p \rightarrow \infty \text{ and } h \rightarrow 0.$$

4. NUMERICAL EXPERIMENTS

In this section we summarise numerical experiments validating the analysis done in previous sections.

4.1. Test 1: Benchmarking a 2-dimensional problem. We begin by benchmarking the scheme against a known solution of the p -Biharmonic problem. To do this we introduce a source term into the problem

$$(4.1) \quad \begin{cases} \Delta(|\Delta u|^{p-2} \Delta u) = f, & \text{in } \Omega, \\ u = g, & \text{on } \partial\Omega, \\ Du = Dg, & \text{on } \partial\Omega. \end{cases}$$

This allows us to pick a function g and construct the appropriate source term such that g solves (4.1). For these tests we choose

$$(4.2) \quad u(x, y) = \frac{1}{\pi^2} \sin(\pi x) \sin(\pi y).$$

We take $\Omega = [-1, 1]^2$ and discretise the domain with a sequence of concurrently refined criss-cross type meshes.

The nonlinear system of equations generated are solved using a damped Newton method initialised by solving the 2-Bilaplacian with corresponding boundary data and forcing. The damping parameter is chosen as $\frac{1}{p-2}$. The results are presented in Figure 1.

4.2. Test 2: Characterising ∞ -Harmonic functions in 1-dimension. In this experiment we illustrate some of the properties of ∞ -Biharmonic functions. The results illustrate that for practical purposes, as one would expect, the approximation of p -Biharmonic functions for large p gives good resolution of candidate ∞ -Biharmonic functions.

We consider the Dirichlet problem for the p -Bilaplacian for $d = 1$ with the boundary data given by the values of the cubic function

$$(4.3) \quad g(x) = \frac{1}{120}(4x-3)(2x-1)(4x-1)$$

on $[0, 1]$. We simulate the p -Bilaplacian for increasing values of p and present the results in Figure 2 indicating that in the limit the ∞ -Biharmonic function should be piecewise quadratic.

4.3. Test 3: Characterising ∞ -Harmonic functions in 2-dimensions. Now we illustrate some of the complicated behaviour of the p -Bilaplacian for $d = 2$:

$$(4.4) \quad \begin{cases} \Delta(|\Delta u|^{p-2}\Delta u) = 0, & \text{in } \Omega = [-1, 1]^2, \\ u = g, & \text{on } \partial\Omega, \\ Du = Dg, & \text{on } \partial\Omega, \end{cases}$$

where g is prescribed as

$$(4.5) \quad g(x, y) = \frac{1}{m20} \cos(m\pi x) \cos(m\pi y),$$

for various values of m . We simulate the p -Bilaplacian for increasing values of p and present the results in Figures 3, 4 and 5 indicating that in the limit the ∞ -Biharmonic function should be piecewise quadratic however the behaviour is quite unexpected and complicated interface patterns emerge even with this relatively simple boundary data.

5. CONCLUSION

In this work we constructed a numerical method for the approximation of solutions of the p -Bilaplacian equation. We were able to analytically show convergence of the numerical approximation and, in particular, to the solution of the limiting problem of the ∞ -Bilaplacian. This is particularly challenging as it is a third order fully nonlinear PDE that is not in divergence form.

We have shown numerically that, for fixed p , our method converges with rates that are better than the analysis predicted. This is well documented in the case of similar lower order problems and can be improved by using appropriate quasi-norms. We have utilised the numerical method to make various interesting observations on the structure of ∞ -Biharmonic functions in that they are piecewise quadratic over the domain with particularly complicated structures for the interfaces.

REFERENCES

- [1] G. Aronsson. “Minimization problems for the functional $\sup_x F(x, f(x), f'(x))$ ”. In: *Ark. Mat.* 6 (1965), 33–53 (1965).
- [2] I. Babuška and J Osborn. “Analysis of finite element methods for second order boundary value problems using mesh dependent norms”. In: *Numerische Mathematik* 34.1 (1980), pp. 41–62.
- [3] I. Babuška, J Osborn, and J. Pitkäranta. “Analysis of mixed methods using mesh dependent norms”. In: *Mathematics of Computation* 35.152 (1980), pp. 1039–1062.
- [4] G. Barles and P. E. Souganidis. “Convergence of approximation schemes for fully nonlinear second order equations”. In: *Asymptotic Anal.* 4.3 (1991), pp. 271–283.


- [5] J. W. Barrett and W. Liu. “Quasi-norm error bounds for the finite element approximation of a non-Newtonian flow”. In: *Numerische Mathematik* 68.4 (1994), pp. 437–456.
- [6] A. Cangiani, E. H. Georgoulis, T. Pryer, and O. J. Sutton. “A posteriori error estimates for the virtual element method”. In: *Numerische mathematik* 137.4 (2017), pp. 857–893.
- [7] P. G. Ciarlet. *The finite element method for elliptic problems*. Studies in Mathematics and its Applications, Vol. 4. Amsterdam: North-Holland Publishing Co., 1978, pp. xix+530.
- [8] M. G. Crandall, H. Ishii, and P.-L. Lions. “User’s guide to viscosity solutions of second order partial differential equations”. In: *Bull. Amer. Math. Soc. (N.S.)* 27.1 (1992), pp. 1–67.
- [9] L. Diening and C. Kreuzer. “Linear convergence of an adaptive finite element method for the p -Laplacian equation”. In: *SIAM J. Numer. Anal.* 46.2 (2008), pp. 614–638.
- [10] L. C. Evans. *Partial differential equations*. Vol. 19. Graduate Studies in Mathematics. Providence, RI: American Mathematical Society, 1998, pp. xviii+662.
- [11] M. Farhloul. “A mixed finite element method for a nonlinear Dirichlet problem”. In: *IMA Journal of Numerical Analysis* 18.1 (1998), pp. 121–132.
- [12] E. Georgoulis, C. Makridakis, and T. Pryer. “Babuška-Osborn techniques in discontinuous Galerkin methods: L^2 -norm error estimates for unstructured meshes”. In: *To appear in SIAM Journal of Numerical Analysis* (2018).
- [13] E. H. Georgoulis and P. Houston. “Discontinuous Galerkin methods for the biharmonic problem”. In: *IMA J. Numer. Anal.* 29.3 (2009), pp. 573–594.
- [14] E. H. Georgoulis and T. Pryer. “Analysis of discontinuous Galerkin methods using mesh-dependent norms and applications to problems with rough data”. In: *Calcolo* 54.4 (2017), pp. 1533–1551.
- [15] E. H. Georgoulis and T. Pryer. “Recovered finite element methods”. In: *Computer Methods in Applied Mechanics and Engineering* (2018).
- [16] D. Gilbarg and N. S. Trudinger. *Elliptic Partial Differential Equations of Second Order*. Second. Berlin: Springer-Verlag, 1983, pp. xiii+513.
- [17] V. Girault and P.-A. Raviart. *Finite element methods for Navier-Stokes equations: theory and algorithms*. Vol. 5. Springer Science & Business Media, 2012.
- [18] T. Gyulov and G. Moroşanu. “On a class of boundary value problems involving the p -biharmonic operator”. In: *Journal of Mathematical Analysis and Applications* 367.1 (2010), pp. 43–57.
- [19] R. Jensen. “Uniqueness of Lipschitz extensions: minimizing the sup norm of the gradient”. In: *Arch. Rational Mech. Anal.* 123.1 (1993), pp. 51–74.
- [20] N. Katzourakis and R. Moser. “Existence, Uniqueness and Structure of Second Order absolute minimisers”. In: *ArXiv* <https://arxiv.org/abs/1701.03348> (2018).
- [21] N. Katzourakis and T. Pryer. “Second order L^∞ variational problems and the ∞ -Polylaplacian”. In: *Advances in Calculus of Variations* (2018).
- [22] N. Katzourakis. “Absolutely minimising generalised solutions to the equations of vectorial calculus of variations in L^∞ ”. In: *Calculus of Variations and Partial Differential Equations* 56.1 (2017), p. 15.
- [23] N. Katzourakis. *An introduction to viscosity solutions for fully nonlinear PDE with applications to calculus of variations in L^∞* . Springer Briefs in Mathematics. Springer, Cham, 2015, pp. xii+123.
- [24] N. Katzourakis. “Generalised solutions for fully nonlinear PDE systems and existence–uniqueness theorems”. In: *Journal of Differential Equations* 263.1 (2017), pp. 641–686.
- [25] N. Katzourakis and T. Pryer. “On the numerical approximation of ∞ -harmonic mappings”. In: *Non-linear differential equations and applications* (2016).
- [26] A. Lasis and E. Süli. “Poinaré-type inequalities for broken Sobolev spaces”. In: *Tech Report*. (2003).
- [27] A. Lazer and P. McKenna. “Large-amplitude periodic oscillations in suspension bridges: some new connections with nonlinear analysis”. In: *Siam Review* 32.4 (1990), pp. 537–578.
- [28] H. Li. “The W_p^1 stability of the Ritz projection on graded meshes”. In: *Mathematics of Computation* 86.303 (2017), pp. 49–74.
- [29] C. G. Makridakis. “On the Babuška–Osborn approach to finite element analysis: L^2 estimates for unstructured meshes”. In: *Numerische Mathematik* (2018), pp. 1–14.
- [30] T. Pryer. “An a posteriori analysis of some inconsistent, nonconforming Galerkin methods approximating elliptic problems”. In: *arXiv preprint arXiv:1505.04318* (2015).

- [31] T. Pryer. “Discontinuous Galerkin methods for the p-biharmonic equation from a discrete variational perspective”. In: *Electronic Transactions on Numerical Analysis* 41 (2014), pp. 328–349.
- [32] D. Sandri. “Sur l’approximation numérique des écoulements quasi-newtoniens dont la viscosité suit la loi puissance ou la loi de Carreau”. In: *RAIRO Modél. Math. Anal. Numér.* 27.2 (1993), pp. 131–155.
- [33] L. Beirão da Veiga, F. Brezzi, A. Cangiani, G. Manzini, L. D. Marini, et al. “Basic principles of virtual element methods”. In: *Math. Models Methods Appl. Sci.* 23.1 (2013), pp. 199–214.


NIKOS KATZOURAKIS DEPARTMENT OF MATHEMATICS AND STATISTICS, WHITEKNIGHTS, UNIVERSITY OF READING, READING RG6 6AX, UK N.Katzourakis@reading.ac.uk.

TRISTAN PRYER DEPARTMENT OF MATHEMATICS AND STATISTICS, WHITEKNIGHTS, UNIVERSITY OF READING, READING RG6 6AX, UK T.Pryer@reading.ac.uk.


FIGURE 1. **Test 1:** Benchmarking results for the mixed finite element approximation to (4.1). We test the cases $p = 2, \dots, 7$ for polynomials of degree $k = 2$. The results show that the convergence rates as predicted in the analysis are achieved for the primal variable. Note that in the case $p > 2$ convergence rates are both higher than predicted for both primal *and* auxiliary variable. Notice also that as p increases the auxiliary variable converges at a faster rate.




(a) The 2-Bilaplacian.




(b) The 3-Bilaplacian.




(c) The 4-Bilaplacian.



(d) The 5-Bilaplacian.




(e) The 6-Bilaplacian.




(f) The 7-Bilaplacian.


FIGURE 2. **Test 2:** A mixed finite element approximations to an ∞ -Biharmonic function using p -Biharmonic functions for various p for the problem given by (4.3). Notice that as p increases, u'' tends to a piecewise constant up to Gibbs oscillations. This is an indication the solution is indeed piecewise quadratic. Also there is only one breaking point in the solution, the location and size of this discontinuity was fully characterised in [21].




(a) The approximation to u , the solution of the 4-Bilaplacian.




(b) The approximation to u , the solution of the 202-Bilaplacian.




(c) The approximation to u'' , the Laplacian of the solution of the 4-Bilaplacian.



(d) The approximation to u'' , the Laplacian of the solution of the 12-Bilaplacian.




(e) The approximation to u'' , the Laplacian of the solution of the 42-Bilaplacian.




(f) The approximation to u'' , the Laplacian of the solution of the 202-Bilaplacian.


FIGURE 3. **Test 3a:** A mixed finite element approximations to an ∞ -Biharmonic function using p -Biharmonic functions for various p for the problem given by (4.4) and (4.5) with $m = 1$. Notice that as p increases, Δu tends to be piecewise constant. This is an indication the solution satisfies the Poisson equation with piecewise constant right hand side albeit with an extremely complicated solution pattern that clearly warrants further investigation.




(a) The approximation to u , the solution of the 4-Bilaplacian.




(b) The approximation to u , the solution of the 142-Bilaplacian.




(c) The approximation to Δu , the Laplacian of the solution of the 4-Bilaplacian.



(d) The approximation to Δu , the Laplacian of the solution of the 42-Bilaplacian.




(e) The approximation to Δu , the Laplacian of the solution of the 68-Bilaplacian.




(f) The approximation to Δu , the Laplacian of the solution of the 142-Bilaplacian.


FIGURE 4. **Test 3b:** A mixed finite element approximations to an ∞ -Biharmonic function using p -Biharmonic functions for various p for the problem given by (4.4) and (4.5) with $m = 2$. Notice that as p increases, Δu tends to be piecewise constant. This is an indication the solution satisfies the Poisson equation with piecewise constant right hand side albeit with an extremely complicated solution pattern that clearly warrants further investigation.




(a) The approximation to u , the solution of the 4-Bilaplacian.




(b) The approximation to u , the solution of the 142-Bilaplacian.




(c) The approximation to Δu , the Laplacian of the solution of the 4-Bilaplacian.



(d) The approximation to Δu , the Laplacian of the solution of the 42-Bilaplacian.




(e) The approximation to Δu , the Laplacian of the solution of the 68-Bilaplacian.




(f) The approximation to Δu , the Laplacian of the solution of the 142-Bilaplacian.

FIGURE 5. **Test 3c:** A mixed finite element approximations to an ∞ -Biharmonic function using p -Biharmonic functions for various p for the problem given by (4.4) and (4.5) with $m = 3$. Notice that as p increases, Δu tends to be piecewise constant. This is an indication the solution satisfies the Poisson equation with piecewise constant right hand side albeit with an extremely complicated solution pattern that clearly warrants further investigation.




(a) The approximation to u , the solution of the 4-Bilaplacian.




(b) The approximation to u , the solution of the 142-Bilaplacian.




(c) The approximation to Δu , the Laplacian of the solution of the 4-Bilaplacian.



(d) The approximation to Δu , the Laplacian of the solution of the 42-Bilaplacian.



(e) The approximation to Δu , the Laplacian of the solution of the 68-Bilaplacian.



(f) The approximation to Δu , the Laplacian of the solution of the 142-Bilaplacian.