

A critical assessment of the long term changes in the wintertime surface Arctic Oscillation and Northern Hemisphere storminess in the ERA20C reanalysis

Article

Published Version

Creative Commons: Attribution 3.0 (CC-BY)

Open Access

Bloomfield, H. C. ORCID: <https://orcid.org/0000-0002-5616-1503>, Shaffrey, L. C. ORCID: <https://orcid.org/0000-0003-2696-752X>, Hodges, K. I. ORCID: <https://orcid.org/0000-0003-0894-229X> and Vidale, P. L. ORCID: <https://orcid.org/0000-0002-1800-8460> (2018) A critical assessment of the long term changes in the wintertime surface Arctic Oscillation and Northern Hemisphere storminess in the ERA20C reanalysis. *Environmental Research Letters*, 13 (9). 094004. ISSN 1748-9326 doi: 10.1088/1748-9326/aad5c5 Available at <https://centaur.reading.ac.uk/78311/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

To link to this article DOI: <http://dx.doi.org/10.1088/1748-9326/aad5c5>

Publisher: Institute of Physics

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online

LETTER • **OPEN ACCESS**

A critical assessment of the long-term changes in the wintertime surface Arctic Oscillation and Northern Hemisphere storminess in the ERA20C reanalysis

To cite this article: H C Bloomfield *et al* 2018 *Environ. Res. Lett.* **13** 094004

View the [article online](#) for updates and enhancements.

Environmental Research Letters



LETTER

OPEN ACCESS

RECEIVED
23 April 2018

REVISED
24 July 2018

ACCEPTED FOR PUBLICATION
25 July 2018

PUBLISHED
24 August 2018

Original content from this work may be used under the terms of the [Creative Commons Attribution 3.0 licence](#).

Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.



A critical assessment of the long-term changes in the wintertime surface Arctic Oscillation and Northern Hemisphere storminess in the ERA20C reanalysis

H C Bloomfield^{1,2} , L C Shaffrey², K I Hodges² and P L Vidale^{1,2}

¹ Department of Meteorology, University of Reading, Reading, RG6 6BB, United Kingdom

² National Centre for Atmospheric Science, University of Reading, United Kingdom

E-mail: h.c.bloomfield@reading.ac.uk

Keywords: ERA20C, reanalysis trends, Arctic Oscillation, extratropical cyclones

Abstract

This study investigates the robustness of the long-term changes in the wintertime surface Arctic Oscillation (AO) in the ERA20C reanalysis. A statistically significant trend in the AO is found in ERA20C over the period 1900–2010. These long-term changes in the AO are not found in two other observational datasets. The long-term change in the AO in ERA20C is associated with statistically significant negative trend (approximately -6 hPa per century) in mean-sea level pressure (MSLP) over the Northern Hemisphere polar regions. This is not seen in the HADSLP2 observational dataset, suggesting that the trends in the ERA20C AO index may be spurious. The spurious long-term changes in MSLP and the AO index in ERA20C result in a strengthening of the meridional MSLP gradient in ERA20C. The strengthening of the meridional MSLP gradient is consistent with increases in wintertime storminess in Northern Europe and the NH high latitudes.

1. Introduction

Global atmospheric reanalyses are a common tool for the evaluation of climate models and for studies of long-term climate variability. Spurious trends are known to occur in reanalysis products, even for the modern satellite period, due to changes in observation quantity and quality (Bengtsson *et al* 2004). Similar questions have been raised with regards to century-long reanalyses (e.g. Wang *et al* 2013), which are primarily based on surface observational networks. Two century-long reanalyses are currently used: the ECMWF 20th century reanalysis (ERA20C; Poli *et al* 2016) and the second version of the National Oceanic and Atmospheric Administration Twentieth Century reanalysis (NOAA-20CR; Compo *et al* 2011). Century-long reanalyses have been used to study a number of phenomena including wind power generation (Bett *et al* 2015), European wave climate (Paris *et al* 2014, Ulazia *et al* 2017) East Asian surface air temperatures (Liu *et al* 2017) and North Russian snow cover (Wegmann *et al* 2017).

A number of studies have used ERA20C to investigate long-term changes in the Northern Hemisphere (NH) atmospheric circulation. Poli *et al* (2016) have compared the time evolution of teleconnection indices from NOAA-20CR and ERA20C, including the North Atlantic Oscillation (NAO). The NAO is an index which describes fluctuations in sea-level pressure in the North Atlantic Ocean, commonly calculated by taking the pressure difference between Iceland and the Azores (Hurrell *et al* 2003). Poli *et al* (2016) found the NAO to be consistent between NOAA-20CR and ERA20C from 1950 to the present. There were large discrepancies between the two reanalyses in the first half of the century, with ERA20C tending to produce lower values of the NAO index at the start of the century compared to NOAA-20CR. However, Poli *et al* (2016) did not compare the reanalyses to observations and so did not consider which of the two reanalyses was the most realistic.

Belleflamme *et al* (2015) also found discrepancies between ERA20C and NOAA-20CR, with ERA20C overestimating high pressure systems over the

Beaufort Sea and underestimating low pressure systems over the Arctic Ocean in the early half of the reanalysis compared to NOAA-20CR. Similarly, Dell'Aquila *et al* (2016) showed that 2–7 day bandpass 500 hPa geopotential height variance in ERA20C was substantially lower in 1900–1930 than in 1980–2010. The differences between 1900–1930 and 1980–2010 were much smaller in the NOAA-20CR reanalyses. Although Belleflamme *et al* (2015) and Dell'Aquila *et al* (2016) compared the ERA20C and NOAA-20CR reanalyses, they did not compare with observational datasets.

Befort *et al* (2016) and Varino *et al* (2018) found that there was a significant positive trend in ERA20C in the number of extratropical cyclones in the NH polar and European regions. Befort *et al* (2016) did not find significant positive trends in extratropical cyclone number in the NOAA-20CR reanalyses. Varino *et al* (2018) investigated the relationship between decadal modes of climate variability and extratropical storminess. Given these dynamical links were considered realistic, Varino *et al* (2018) concluded that confidence could be placed in the trend in extratropical cyclone number in ERA20C. Other observational studies have concluded that trends over the 20th Century in storminess and extratropical cyclone number are statistically insignificant over Europe (e.g. Alexanderson *et al* 2000, Feser *et al* 2015).

A key aspect of the NH atmospheric circulation that has received less attention in century-long reanalyses is the Arctic Oscillation (AO). The wintertime surface AO is defined as the dominant mode of variability of the NH atmospheric circulation and its variability is associated with changing weather patterns over Europe and North America (Thompson and Wallace 2000).

Previous studies have shown that extremes in the NAO index (the dominant mode of variability of the North Atlantic and Europe; Hurrell *et al* 2003) are associated with extreme temperature and precipitation anomalies over Europe (Yiou and Nogaj 2004, Cattiaux *et al* 2010). The NAO also shows significant correlation with and significant wave height anomalies. This is most noticeable in the Eastern North Atlantic (Carretero *et al* 1998, Woolf *et al* 2002, Dodet *et al* 2010, Bertin *et al* 2013). A strong positive correlation is seen in Northern latitudes and significant negative correlation is seen in southern latitudes (Dodet *et al* 2010). This suggests a link between the large scale modes of atmospheric circulation and regional weather extremes which can impact on major population centres. It is therefore important to understand the behaviour of the large scale modes of atmospheric variability.

The long-term changes in AO from observational datasets have been previously calculated. The first to do this was Thompson and Wallace (2000), using sea level pressure data from 1900 to 1997. The AO has also been calculated in Allan and Ansell (2006) for the

period 1850–2003 using the HADSLP2. Both studies find the AO experiences multiple positive and negative extended phases. A predominantly negative AO state is present from 1960 to 1985, with a positive AO in the 1990s, and returning to a more negative phase in the 2000s. Neither study found a significant long-term trend in the AO.

Feldstein (2002) examined the AO index from 1899 to 2000 with an extended version of the dataset from Trenberth and Paolino (1980), finding a linear trend in the AO index between 1967 and 1997, in excess of the level expected by internal variability. Feldstein (2002) show all of the interannual variability of the annular mode in the first 60 years can be explained by atmospheric internal variability, whereas this is not possible in the later period.

The AO has previously been calculated in the ERA20C reanalysis by Liu *et al* (2017), who examined the relationship between the AO and East Asian surface air temperatures. A positive trend in the normalised AO index was found in Liu *et al* (2017) (≈ 0.06 standard deviations decade⁻¹). However Liu *et al* (2017) did not compare the AO to other century-long reanalyses or observational datasets.

It is clear from previous studies that long-term changes in the atmospheric circulation have been seen in ERA20C. Other studies have hypothesised that long-term changes may be due to the sparse observational network in the early years of the reanalysis compared to the later period (Belleflamme *et al* 2015, Befort *et al* 2016, Dell'Aquila *et al* 2016, Poli *et al* 2016). Similar hypothesis have been made following studies in the Southern Hemisphere using the NCEP-NCAR reanalysis (Kalnay *et al* 1996). Spurious trends have been observed in surface pressure field Hines *et al* (2000) and the number of cyclone tracks (Simmonds and Keay 2000). The majority of these spurious trends coincide with areas where surface pressure observations are most sparse (50N–70N; Hines *et al* 2000, Simmonds and Keay 2000).

Although, these studies have suggested that these trends are not real, they have not made direct comparisons between ERA20C and observational datasets. To a certain extent, observational uncertainty can be partially assessed by using an ensemble reanalysis (e.g. the NOAA-20CR reanalyses has 56 members). ERA20C, however, has only a single member making such assessment difficult. A new coupled version of ERA20C (available at: <https://ecmwf.int/en/forecasts/datasets/reanalysis-datasets/cera-20c>), has ten ensemble members which allows for the robustness of long-term changes to be assessed.

In this study, the robustness of long-term changes in the wintertime surface AO and the NH storminess in ERA20C and the recently available Coupled ERA20C (CERA20C) are investigated. Results from CERA20C are included to provide a comparison to ERA20C and to assess ensemble uncertainty. The paper is structured as follows. Section 2 summarises

the datasets analysed in this study (section 2.1), the methods for calculating the AO (section 2.2), and wintertime storminess (section 2.3). Section 3 presents the results for the representation of the AO in ERA20C and two other ECMWF reanalysis products (section 3.1), with details of how this relates to the mean-sea level pressure (MSLP) patterns in each reanalysis (section 3.2) and wintertime storminess (section 3.3). Section 4 concludes and discusses the implications of the key findings.

2. Methods

2.1. Datasets

The main dataset used in this study is the ERA20C reanalysis which spans the period 1900–2010. ERA20C is available every 3 hours at 1.25° horizontal resolution. ERA-20C was created as part of the ERA-CLIM project (see <http://era-clim.eu>), and is based on ECMWF's Integrated Forecast System. ERA20C uses a four-dimensional variational data assimilation scheme with 24 h window and a spectral resolution of T159 (Poli *et al* 2016). Only surface pressure data (archived in the International Surface Pressure Databank, ISPD; Cram *et al* 2015) and surface marine winds (available from the Internal Comprehensive Ocean-Atmosphere Data set, ICOADS; Woodruff *et al* 2011) are assimilated into ERA20C. The HADISST2.1 sea surface temperature and sea ice concentration datasets are used as boundary conditions (Titchner and Rayner 2014). For further details on ERA20C see Poli *et al* (2016).

Comparisons between ERA20C and other ECMWF twentieth century reanalyses for the period of 1900–2000 are discussed later in this paper and include:

- (i) *CERA20C*: The Coupled European reanalysis System, which assimilates both atmosphere and ocean observations, allowing for interaction between the atmosphere and ocean. CERA20C has ten ensemble members with 3 hourly data available, covering the period 1900–2010 at the same model resolution as ERA20C. The ten member ensemble provides an assessment of both model and observation uncertainty throughout the 20th century. CERA20C was initialised from the uncoupled reanalyses ERA-20C and the 20th Century Ocean reanalysis. See Laloyaux *et al* (2016) for details of the CERA assimilation system.
- (ii) *ERA20CM*: A ten member ensemble of free-running model integrations using the same model as in ERA20C, covering the period 1900–2010 at the same horizontal resolution as ERA20C. The ensemble does not assimilate any observations, however: sea-surface temperature and the sea-ice cover are prescribed by an ensemble of realisations from HadISST2. ERA20CM provides an


estimate of uncertainty in the model climate (see Hersbach *et al* 2015 for further details).

HADSLP2 is used for comparison with ERA20C. HADSLP2 is a 5° gridded dataset spanning the period 1850–2004. It is created based on digitised terrestrial observations of surface pressure as well as sea-level pressure observations from ICOADS. An empirical orthogonal function (EOF)-based smoother is used to interpolate over data-sparse regions (Allan and Ansell 2006). HADSLP2 has previously been used for twentieth century reanalysis validation by Compo *et al* (2011). All observations undergo extensive quality control before inclusion in the dataset (as discussed in Allan and Ansell 2006). However, there are still data-sparse regions in HADSLP2, particularly in the early period of the dataset and at high latitudes. Over the ocean the surface pressure data that ERA20C has the option to assimilate should be similar to that used in HADSLP2 (as both are using ICOADS as an input). However, there may be larger differences in land observations, as Allan and Ansell (2006) use a large set of terrestrial observations, whereas ERA20C use all that are available from the ISPD.

In Allan and Ansell (2006) it is noted that HADSLP2 includes a series of quality control checks. First the stations were checked for internal consistency, and stations from multiple sources are converted into a single time-series. Nearest-neighbour and climatology checks were also performed. Large issues in the sources of MSLP data were highlighted from the quality control procedure. Allan and Ansell (2006) state that the measurement and sampling errors are particularly large in the high southern latitudes, where the number of observations is very low.

The monthly-mean AO index created using the method from Thompson and Wallace (2000) is also used in this study (available at: http://atmos.colostate.edu/davet/ao/Data/ao_index.html). This index covers the period 1899–2002 and is generated by projecting the AO pattern onto monthly-mean MSLP anomalies. Sea level pressure data created for 1899–1997 using an extended version of the dataset, and the method of Trenberth and Paolino (1980). The NCEP-NCAR reanalysis (Kalnay *et al* 1996) is used from 1997 to 2002. The dataset of Trenberth and Paolino (1980) spans the period 1899–1977, and includes MSLP from eight different sources which can lead to discontinuities when changing from one data source to another. A discontinuity may be present in 1997 when the data source is changed to the NCEP-NCAR reanalysis.

In Trenberth and Paolino (1980) a time series of sea level pressure data is deemed *well behaved* if there were no long-term trends or discontinuities, and if there were no data points greater than three standard deviations from the long-term mean. If a time series contained issues and was close to another available data source a correction was established, otherwise the



problems were documented and data set to missing values. Trenberth and Paolino (1980) note a number of problems in the dataset, the first of which is in the high latitudes, arising due to sparse data coverage. The sea level pressures are shown to be biased high until the end of 1923 at latitudes greater than 70°N.

2.2. Calculating the wintertime surface AO

The wintertime surface AO is calculated as in Thompson and Wallace (2000) by taking the first EOF of the normalised NH monthly-mean MSLP data from 20 to 90°N. To ensure equal area weighting within the EOF analysis, the normalised MSLPs are weighted by the square root of the cosine of latitude (as in Thompson and Wallace 2000). The October–March AO value is then calculated as the mean of the six monthly values. This period is chosen for analysis as it is the winter storm season used in Befort *et al* (2016). All datasets are interpolated onto a 5° grid before calculation of the AO (as this is the resolution of the coarsest product, HADSLP2). The spatial patterns of the AO are shown in subsequent figures as the correlation coefficient between the AO index and the MSLP at each grid box.

2.3. Wintertime cyclone tracking

Extratropical cyclones are identified in the reanalysis datasets by an automated cyclone tracking algorithm based on the method described by Hodges (1995, 1999), used to create the European Wind storm catalogue (Roberts *et al* 2014). This method provides the 3 hourly tracks of the locations of maximum 850 hPa relative vorticity (at T42 resolution). This method means that cyclone numbers are not impacted by the background MSLP. Only cyclones North of 30°N are included, to exclude tropical cyclones. For a feature to be tracked it must reach a relative vorticity maximum greater than $1 \times 10^{-5} \text{ s}^{-1}$ and last for a minimum of 2 days. The tracking algorithm has been applied to ERA20C, and individual members of CERA20C and ERA20CM for the period 1900–2000, from October to March. It is important to note that the number of storms identified in this study is dependent on the tracking algorithm used (see Ulbrich *et al* 2013 and Rudeva *et al* 2014). It has been shown in Hodges *et al* (2017) that the chosen method for this study has been used to successfully identify tropical cyclones in six reanalysis products, which match well to tropical cyclone track archives. As Hodges *et al* (2017)




Figure 2. Power spectrum density for the October–March AO indices from Thompson and Wallace (2000) (black), HADSLP2 (red) and ERA20C (blue) (a) for periods 0–50 years (b) for period 0–10 years. The 95% confidence spectrum for white noise is given as the grey dashed line.

state that nearly every cyclone is found in all six reanalysis over the period 1979–2012 we are confident the method can accurately represent storm counts.

3. Results

3.1. The representation of the wintertime surface AO in ERA20C, CERA20C and ERA20CM

Figure 1(a) shows the AO time-series from Thompson and Wallace (2000), ERA20C, and HADSLP2 over the common time period of 1900–2000. The ERA20C, HADSLP2 and Thompson and Wallace (2000) AO indices compare well in the second half of the century. The agreement of the HADSLP2 AO index with the AO index from Thompson and Wallace (2000) gives confidence that if a bias is present in HADSLP2, then it is the same as that in Thompson and Wallace (2000). Figures 1(c) and (d) show the spatial pattern of the AO for ERA20C and HADSLP2 respectively. The general agreement between the two subplots shows that ERA20C is able to capture the spatial structure of the AO well over the 20th century. The main difference is the magnitude of the pattern (which is larger in HADSLP2).

Figure 1(b) shows the differences between ERA20C and the two other AO indices. In ERA20C, the AO tends to be too negative on average from 1900 to 1950 and then moderately too positive on average from 1950 to 2000. A significant positive trend (defined using a Mann-Kendal test with $p \leq 0.05$) is present in the time-series of the AO in ERA20C. No significant trends are present in HADSLP2 or Thompson and Wallace (2000) over the same time period.

The power spectrum density from the ERA20C, HADSLP2 and the Thompson and Wallace (2000) AO indices are shown in figure 2 along with the 95% confidence spectrum for white noise. Figure 2(b) shows that at short periods (less than 10 years) the observational datasets have a similar power spectrum density to ERA20C. Statistically significant peaks occur at approximately 2.5 years in all three datasets. However, at larger periods (greater than 40 years) ERA20C is the only dataset with statistically significant power. This is primarily due to the positive trend in the AO index through the century in ERA20C.

ERA20C was part of a suite of experiments including ERA20CM and CERA20C for which the analysis in this paper has been repeated (see section 2.1 for details of the experiments). CERA20C, exhibits very similar trends in the AO to ERA20C (see figure A1). In the




Table 1. Pearsons correlation coefficient between the ERA20C AO index and the regional-mean wintertime MSLP for regions defined in figure 2(a). All p -values are less than 0.01.

Region	Correlation coefficient between ERA20C AO index and regional MSLP
Extra-tropics	−0.47
Northern Europe	0.40
Polar Region	−0.95
North Atlantic	0.46
North Pacific	0.29


period from 1950 to 2000 the AO index is very similar to ERA20C and there is very little spread in the ten member ensemble from CERA20C, which suggests the AO index value is well constrained. However, there is much more spread in the ensemble from 1900 to 1950. There are also more discrepancies between the CERA20C ensemble members and ERA20C where both the atmosphere and ocean are less strongly constrained. The spatial map of the AO for ERA20CM looks similar to that for ERA20C and CERA20C (see figure A1). However, ERA20CM does not show a positive trend in AO in nine out of ten ensemble members (and the trend is only weakly positive in the one member where it is seen; not shown). This suggests that the positive trend seen in the reanalyses may be related to the use

of observations in the data assimilation, rather than the boundary conditions used in ERA20C or ERA20CM.

3.2. Long-term changes in wintertime MSLP in ERA20C, CERA20C and ERA20CM

Figures 3(a)–(c) shows October–March MSLP data for ERA20C for the early period (1900–1930), late period (1970–2000) and the difference between the two periods. The figures shows a statistically significant decrease in MSLP of approximately 6 hPa in the polar regions in the late period compared to the early period (results significant to 2 sample t test $p \leq 0.05$). A statistically significant approximately 2 hPa increase in pressure is seen over the North Pacific and over central and southern Europe. Figures 3(d)–(f) shows the October–March MSLP data averaged over the same time periods from HADSLP2. In HADSLP2 only small significant changes in mean sea-level pressure are seen over the poles (approximately 1 hPa), and no significant decreases are seen in central and southern Europe or the North Pacific. Hence the pattern of the difference in MSLP from early period to late period is very different in HADSLP2 compared to ERA20C.

The differences in October–March MSLP shown in figure 2(c) project strongly on to the spatial pattern of the wintertime surface AO in figure 1(c). The




correlation coefficient between the ERA20C AO index and the winter-mean MSLP in each region are given in table 1. The extremely strong negative correlation between the AO index and MSLP in the polar region ($r = -0.95$, p value < 0.01) suggests that the trend in the AO index seen in ERA20C (figures 1(a), (b)) is consistent with the higher MSLP in the early period compared to the late period in the high latitudes.

Figure 4 shows the mean October–March MSLP averaged over the NH extra-tropics, North Pacific, North Atlantic, Polar region and Northern Europe (defined in figure 3(a)) for HADSLP2 and ERA20C. Figure 4 shows that HADSLP2 does not have a statistically significant trend in MSLP in any of the chosen regions. However, over the whole extra-tropics there is a statistically significant negative trend in MSLP in ERA20C (figure 4(a)). This is dominated by a large negative trend in MSLP over the polar region (figure 4(d)). Significant trends are not present in the North Pacific (figure 4(b)) or North Atlantic (figure 4(c)) in ERA20C.

The October–March MSLP differences between the early and late period for the ensemble-mean of CERA20C show a similar pattern to ERA20C (compare figure A2 with figure 3). However, the magnitude of the anomalies in the Polar region are smaller. No significant differences are seen between the early and late period in the October–March MSLP in the ensemble-mean of the free-running ERA20CM (see figure A2). This suggests that the long-term change in the AO index in ERA20C between the beginning and end of the century is likely due to changes in the atmospheric observing system rather than a trend produced by the boundary conditions.

3.3. Wintertime storminess in ERA20C, CERA20C and ERA20CM

The AO/NAO have been related to NH storm tracks by Nie *et al* (2008), Luo *et al* (2007) and Luo *et al* (2011) who showed strong positive winter AO anomalies are associated with increased synoptic scale wave activity in the North Atlantic and therefore an intensification



of the North Atlantic Storm track. When the AO is in its positive phase there is an increased meridional pressure gradient in the North Atlantic and North Pacific associated with stronger winds circulating the North Pole (Thompson and Wallace 2000). The stronger winds result in colder air being confined to polar regions, which is associated with regions of increased meridional temperature gradient, providing the baroclinicity needed for storm development and growth (Thompson and Wallace 1998).

The differences in the October–March MSLP over the NH, shown in figure 3, result in changes in the location and strength of the surface meridional pressure gradients over the course of the century. The surface meridional pressure gradients for ERA20C are shown in figures 5(a)–(c), for 1900–1930, 1970–2000 and the difference between the two periods. The reduction in MSLP at polar latitudes in ERA20C from 1970 to 2000 (see figure 3(c)) results in a statistically significant increase in surface meridional pressure gradient over North Atlantic and European sector. In the North Pacific there is a statistically significant increase in meridional MSLP gradient at $\approx 60^\circ\text{N}$ (figure 5(c)) and a smaller reduction in MSLP gradient at $\approx 30^\circ\text{N}$.

These statistically significant changes are not seen in HADSLP2 (not shown).

Figures 5(d)–(f) shows the number of storms from October–March for the 1900–1930, 1970–2000 and the difference between the two periods for ERA20C. The increases in ERA20C's surface meridional MSLP gradient shown in figure 5(c) are consistent with increases in wintertime storminess over the North Atlantic and European sector and the high-latitude North Pacific. However, increasing storminess over North America and Eurasia seen in ERA20C is less consistent with changes in meridional MSLP gradients.

Figure 6 shows the long-term changes in the number of storms for ERA20C in the regions used in figure 4. Statistically significant increases in the number of storms are seen in the extra-tropics which are mostly due to increased numbers of storms over Northern Europe and the Polar region. The analysis here suggests that changes in North Atlantic storminess and high-latitude North Pacific storminess seen in previous studies (Befort *et al* 2016, Varino *et al* 2018) are consistent with the spurious trends in MSLP seen in ERA20C.




Figure 6 also shows the changes in the number of storms for the ensemble mean of CERA20C and the free-running ERA20CM. The significant positive trends in the number of storms that are seen in ERA20C are also present in CERA20C in Northern Europe and the Polar region. However, they are not seen in these regions for ERA20CM. It is interesting to note that in Northern Europe and the Polar region the number of storms in the free-running ERA20CM is much higher than in ERA20C for the early period. This suggests that without any observations the model is potentially able to represent the average number of storms more accurately than when the model has only sparse observations, highlighting the difficulties when assimilating sparse observations.

4. Conclusions

The aim of this study is to evaluate the long-term changes in the wintertime surface AO in ERA20C, and

associated changes in NH wintertime MSLP and storminess. The key results are as follows:

- (i) A statistically significant long-term change is seen in the wintertime surface AO in ERA20C for the period 1900–2000 (as measured by a Mann-Kendal test $p < 0.05$). A statistically significant long-term change is not present in the AO index calculated from two observational datasets (Thompson and Wallace 2000, Allan and Ansell 2006).
- (ii) The long-term change in the AO in ERA20C is due to a spurious decrease in MSLP in the polar region (latitudes $>60^\circ\text{N}$). Again this long-term change in MSLP is not seen in the two observational datasets which have been quality controlled.
- (iii) The long-term changes in MSLP correspond to changes in the meridional pressure gradients over Northern Europe and the Polar region, which are consistent with increases in wintertime storminess

seen in ERA20C over the North Atlantic and North Pacific.

The long-term changes in MSLP in polar regions are consistent with Belleflamme *et al* (2015) who showed that at the start of the century ERA20C has less low pressure systems over the Arctic Ocean and more high pressure systems located over the Beaufort Sea (when compared to NOAA-20CR). This study has extended the work of Belleflamme *et al* (2015) by comparing to gridded observations in order to show that the polar region is the region of largest spurious change in ERA20C.

This study provides evidence for the increasing trend in wintertime surface AO index seen in Liu *et al* (2017), by demonstrating significant differences in the AO time series generated using ERA20C before and after 1950 compared to two observational datasets. In this study the observational datasets are considered as truth. However, there are notable discrepancies between them, particularly in the early period, when observations are sparse (see figure 1). Trenberth and Paolino (1980) comment that there are anomalously high sea level pressures at high latitudes in their dataset before 1923, which may explain some of these differences. The key point for this study is that although both observational AO indices are different, they both do not contain a long-term change.

The increasing long-term change in NH storminess is consistent with that found in Bafort *et al* (2016) and Varino *et al* (2018) for ERA20C. The consistency between the number of storms in ERA20C and ERA-interim from 1980 to 2000 suggests that ERA20C has a good representation of storms for this period.

The long-term changes in ERA20C MSLP are most pronounced at high latitudes. This is an area where Poli *et al* (2016) have documented there are

particularly large increases in the number of observations that are assimilated into the reanalysis over time, with sparse observations in the early period. Simmonds and Keay (2000) and Hines *et al* (2000) find similar problems in the southern hemisphere in the early period of the NCEP-NCAR re-analysis where observations are more sparse than in the later period.


The fact that long-term changes in the AO, MSLP and wintertime storminess are not present in free-running ERA20CM (which does not assimilate surface pressure or wind speeds; see figures A1, A2 and 6) supports previous studies that have suggested long-term changes may be due to the assimilation of increasing numbers of observations (Belleflamme *et al* 2015, Bafort *et al* 2016, Dell'Aquila *et al* 2016, Poli *et al* 2016).


This study has shown that a significant long-term positive trend is seen in the ERA20C AO index, which is not seen in other century-long observational datasets. The ERA20C reanalysis should therefore be used with caution when examining the effects of long-term climate change at latitudes greater than 60°N, as there may be spurious trends in this region.

Acknowledgments

Bloomfield was funded as part of the Wind Storms Information Services project, a Copernicus Climate Change Services programme project funded by the European Union. Vidale and Hodges were funded as part of National Centre for Atmospheric Science. Shaffrey was funded by the WINDSURFER project, part of the ERA4CS, an ERA-NET initiated by JPI Climate with co-funding by the European Union (Grant 690462).

Appendix





ORCID iDs

H C Bloomfield  <https://orcid.org/0000-0002-5616-1503>

References

- Alexanderson H, Tuomenvirta H, Schmith T and Iden K 2000 Trends of storms in NW Europe derived from an updated pressure data set *Clim. Res.* **14** 71–3
- Allan R and Ansell T 2006 A new globally complete monthly historical gridded mean sea level pressure dataset (HadSLP2): 18502004 *J. Clim.* **19** 5816–42
- Befort D, Wild S, Kruschke T, Ulbrich U and Leckebusch G 2016 Different long-term trends of extra-tropical cyclones and windstorms in ERA-20C and NOAA-20CR reanalyses *Atmos. Sci. Lett.* **17** 586–95
- Belleflamme A, Fettweis X and Erpicum M 2015 Recent summer Arctic atmospheric circulation anomalies in a historical perspective *Cryosphere* **9** 53–64
- Bengtsson L, Hagemann S and Hodges K 2004 Can climate trends be calculated from reanalysis data? *J. Geophys. Res.: Atmos.* **109** 1–8
- Bertin X, Prouteau E and Letetrel C 2013 A significant increase in wave height in the North Atlantic Ocean over the 20th century *Glob. Planet. Change* **106** 77–83
- Bett P, Thornton H and Clark R 2015 Using the twentieth century reanalysis to assess climate variability for the European wind industry *Theor. Appl. Climatol.* **127** 61–80
- Carretero J *et al* 1998 Changing waves and storms in the Northeast Atlantic? *Bull. Am. Meteorol. Soc.* **79** 741–60
- Cattiaux J, Vautard R, Cassou C, Yiou P, Masson-Delmotte V and Codron F 2010 Winter 2010 in Europe: a cold extreme in a warming climate *Geophys. Res. Lett.* **37** 1–4
- Compo G *et al* 2011 The twentieth century reanalysis project *Q. J. R. Meteorol. Soc.* **137** 1–28
- Cram T A *et al* 2015 The international surface pressure databank version 2 *Geosci. Data J.* **2** 31–46
- Dell'Aquila A, Corti S, Weisheimer A, Hersbach H, Peubey C, Poli P, Berrisford P, Dee D and Simmons A 2016 Benchmarking Northern Hemisphere midlatitude atmospheric synoptic variability in centennial reanalysis and numerical simulations *Geophys. Res. Lett.* **43** 5442–9
- Dodet G, Bertin X and Taborda R 2010 Wave climate variability in the North-East Atlantic Ocean over the last six decades *Ocean Modelling* **31** 120–31
- Feldstein S 2002 The recent trend and variance increase of the annular mode *J. Clim.* **15** 88–94
- Feser F, Barcikowska M, Krueger O, Schenk F, Weisse R and Xia L 2015 Storminess over the North Atlantic and Northwestern Europe—a review *Q. J. R. Meteorol. Soc.* **141** 350–82
- Hersbach H, Peubey C, Simmons A, Berrisford P, Poli P and Dee D 2015 ERA-20CM: a twentieth-century atmospheric model ensemble *Q. J. R. Meteorol. Soc.* **141** 2350–75
- Hines K, Bromwich D and Marshall G 2000 Artificial surface pressure trends in the NCEP-NCAR reanalysis over the Southern Ocean and Antarctica *J. Clim.* **13** 3940–52
- Hodges K, Cobb A and Vidale P 2017 How well are tropical cyclones represented in reanalysis datasets? *J. Clim.* **30** 5243–64
- Hodges K I 1995 Feature tracking on the unit-sphere *Mon. Weather Rev.* **123** 3458–65
- Hodges K I 1999 Adaptive constraints for feature tracking *Mon. Weather Rev.* **126** 3458–65

- Hurrell J, Kushnir Y, Otterson G and Visbeck M 2003 An overview of the North Atlantic Oscillation *J. Clim.* **134** 263–73
- Kalnay E *et al* 1996 The NCEP/NCAR 40-year reanalysis project *Bull. Am. Meteorol. Soc.* **77** 437–71
- Laloyaux P, Balmaseda M, Dee D, Mogensen K and Janssen P 2016 A coupled data assimilation system for climate reanalysis *Q. J. R. Meteorol. Soc.* **142** 65–78
- Liu Y, Shengping H, Li F, Wang H and Zhu Y 2017 Interdecadal change between the Arctic Oscillation and East Asian Climate during 1900–2015 winters *Int. J. Climatol.* **37** 4791–802
- Luo D, Diao Y and Feldstein S 2011 The variability of the Atlantic storm track activity and North Atlantic Oscillations: a link between intraseasonal and interannual variability *J. Atmos. Sci.* **68** 577–601
- Luo D, Lupo A and Wan H 2007 Dynamics of eddy-driven low-frequency dipole modes: I. A simple model of North Atlantic Oscillations *J. Atmos. Sci.* **64** 3–28
- Nie J, Wang P, Yang W and Tan B 2008 Northern hemisphere storm tracks in strong AO anomaly winters *Atmos. Sci. Lett.* **9** 153–9
- Paris F, Lecacheux S, Idier D and Charles E 2014 Assessing wave climate trends in the Bay of Biscay through an intercomparison of wave hindcasts and reanalyses *Ocean Dyn.* **64** 1247–67
- Poli P *et al* 2016 ERA20C: an atmospheric reanalysis of the twentieth century *J. Clim.* **29** 4083–97
- Roberts J F, Champion A, Dawkins L, Hodges K, Shaffrey L, Stephenson D, Stringer M, Thornton H and Youngman B 2014 The XWS open access catalogue of extreme European windstorms from 1979 to 2012 *Nat. Hazards Earth Syst. Sci.* **14** 2487–501
- Rudeva I, Gulev S, Simmonds I and Tilinina N 2014 The sensitivity of characteristics of cyclone activity to identification procedures in tracking algorithms *Tellus A* **66** 1–20
- Simmonds I and Keay K 2000 Variability of Southern Hemisphere Extratropical Cyclone Behavior, 1958–97 *J. Clim.* **13** 550–61
- Thompson D and Wallace J 1998 The arctic oscillation signature in the wintertime geopotential height and temperature fields *Geophys. Res. Lett.* **25** 1297–300
- Thompson D and Wallace J 2000 Annual modes in the extratropical circulation: I. Month-to-month variability *J. Clim.* **13** 1000–15
- Titchner H and Rayner N 2014 The Met Office Hadley Centre sea ice and sea surface temperature data set, version 2: I. Sea ice concentrations *J. Geophys. Res. Atmos.* **119** 2864–89
- Trenberth K and Paolino D 1980 The Northern Hemisphere Sea-level pressure data set: trends, errors and discontinuities *Mon. Weather Rev.* **108** 855–72
- Ulazia A, Penalba M, Ibarra-Berastegui G, Ringwood J and Saenz J 2017 Wave energy trends over the Bay of Biscay and the consequences for wave energy converters *Energy* **141** 624–34
- Ulbrich U *et al* 2013 Are Greenhouse Gas Signals of Northern Hemisphere winter extra-tropical cyclone activity dependent on the identification and tracking algorithm? *Meteorol. Z.* **22** 61–8
- Varino F, Arbogast P, Joly B, Riviere G, Fandeur M, Bovy H and Granier J 2018 Northern Hemisphere extratropical winter cyclones variability over the 20th Century derived from ERA20C reanalysis *Clim. Dyn.* (<https://doi.org/10.1007/s00382-018-4176-5>)
- Wang X, Feng Y, Compo G, Swail V, Zwiers F, Allan R and Sardeshmukh P 2013 Trends and low frequency variability of extra-tropical cyclone activity in the ensemble of twentieth century reanalysis *Clim. Dyn.* **40** 2775–800
- Wegmann M, Orsolini Y, Dutra E, Bulygina O, Sterin A and Bronnimann S 2017 Eurasian snow depth in long-term climate reanalyses *Cryosphere* **11** 923–35
- Woodruff S D *et al* 2011 ICOADS Release 2.5: extensions and enhancements to the surface marine meteorological archive *Int. J. Climatol.* **31** 951–67
- Woolf D, Challenor P and Cotton P 2002 Variability and predictability of the North Atlantic wave climate *J. Geophys. Res.: Oceans* **107** 1–12
- Yiou P and Nogaj M 2004 Extreme climatic events and weather regimes over the North Atlantic: when and where? *Geophys. Res. Lett.* **31** 1–4