University of Reading

Higher vegetable protein consumption, assessed by an isoenergetic macronutrient exchange model, is associated with a lower presence of overweight and obesity in the web-based Food4me European study

Article

Accepted Version

Navas-Carretero, S., San-Cristobal, R., Livingstone, K. M., Celis-Morales, C., Marsaux, C. F., Macready, A. L. ORCID: https://orcid.org/0000-0003-0368-9336, Fallaize, R. ORCID: https://orcid.org/0000-0003-3734-6489, O'Donovan, C. B., Forster, H., Woolhead, C., Moschonis, G., Lambrinou, C. P., Jarosz, M., Manios, Y., Daniel, H., Gibney, E. R., Brennan, L., Walsh, M. C., Drevon, C. A., Gibney, M., Saris, W. H. M., Lovegrove, J. A. ORCID: https://orcid.org/0000-0001-7633-9455, Mathers, J. C. and Martinez, J. A. (2019) Higher vegetable protein consumption, assessed by an isoenergetic macronutrient exchange model, is associated with a lower presence of overweight and obesity in the web-based Food4me European study. International Journal of Food Sciences and Nutrition, 70 (2). pp. 240-253. ISSN 1465-3478 doi: 10.1080/09637486.2018.1492524 Available at https://centaur.reading.ac.uk/78383/

University of Reading

It is advisable to refer to the publisher's version if you intend to cite from the work. See Guidance on citing.

To link to this article DOI: http://dx.doi.org/10.1080/09637486.2018.1492524

Publisher: Informa Healthcare

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the End User Agreement.

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online

HIGHER VEGETABLE PROTEIN CONSUMPTION, ASSESSED BY AN ISOENERGETIC MACRONUTRIENT EXCHANGE MODEL, IS ASSOCIATED WITH A LOWER PRESENCE OF OVERWEIGHT AND OBESITY IN THE WEB-BASED FOOD4ME EUROPEAN STUDY

Santiago Navas-Carretero[1,2*†], Rodrigo San-Cristobal[1*], Katherine M Livingstone[3], Carlos Celis-Morales[3], Cyril F Marsaux[4], Anna L Macready[5], Rosalind Fallaize[5], Clare B O'Donovan[6], Hannah Forster[6], Clara Woolhead[6], George Moschonis[7], Christina P. Lambrinou[7], Miroslaw Jarosz[8], Yannis Manios[7], Hannelore Daniel[9], Eileen R Gibney[6], Lorraine Brennan[6], Marianne C Walsh[6], Christian A Drevon[10], Mike Gibney[6], Wim HM Saris[4], Julie A Lovegrove[5], John C Mathers[3], J. Alfredo Martinez[1,2,11,12] on behalf of the Food4Me Study.

AUTHOR AFFILIATIONS

[1]University of Navarra, Centre for Nutrition Research, Department of Nutrition, Food Science and Physiology, Pamplona 31008, Spain.

[2]CIBER Fisiopatología Obesidad y Nutrición (CIBERobn), Instituto de Salud Carlos III, Madrid 28023, Spain.

[3]Human Nutrition Research Centre, Institute of Cellular Medicine, Newcastle University, Newcastle Upon Tyne NE1 7RU, United Kingdom.

[4]Department of Human Biology, NUTRIM School for Nutrition and Translational Research in Metabolism. Maastricht University Medical Centre, Maastricht 6200MD, The Netherlands

[5]Hugh Sinclair Unit of Human Nutrition and Institute for Cardiovascular and Metabolic Research, University of Reading, Reading RG6 6AA, United Kingdom.

[6]UCD Institute of Food and Health, UCD School of Agriculture and Food Science, University College Dublin, Belfield, Dublin 4, Republic of Ireland

[7]Department of Nutrition and Dietetics, Harokopio University of Athens, Athens 17671, Greece.

[8] Institute of Food and Nutrition (IZZ), Warszaw 02-903, Poland.

[9]ZIEL Research Center of Nutrition and Food Sciences, Biochemistry Unit, Technische Universität München, Munich 85354, Germany.

[10]Department of Nutrition, Institute of Basic Medical Sciences, Faculty of Medicine, University of Oslo, Oslo 0317, Norway.

[11]Instituto de Investigaciones Sanitarias de Navarra (IDisNa), Pamplona 31008, Spain.

[12]Institute IMDEA Food, Madrid 28049, Spain.

* Both authors contributed equally to this work.

† Corresponding author: Dr. Santiago Navas-Carretero. University of Navarra. C/Irunlarrea, 1. 31008 Pamplona. E-mail: snavas@unav.es. Tel: +34 948 42 56 00 ext. 806623; Fax: +34 948 42 57 49

SHORT TITLE: Vegetable protein intake and risk of obesity

Abstract

The objective was to evaluate differences in macronutrient intake and to investigate the possible association between consumption of vegetable protein and the risk of

overweight/obesity, within the Food4Me randomised, online intervention. Differences in macronutrient consumption among the participating countries grouped by EU Regions (Western Europe, British Isles, Eastern Europe and Southern Europe) were assessed. Relation of protein intake, within isoenergetic exchange patterns, from vegetable or animal sources with risk of overweight/obesity was assessed through the multivariate nutrient density model and a multivariate-adjusted logistic regression.

A total of 2413 subjects who completed the Food4Me screening were included, with self-reported data on age, weight, height, physical activity and dietary intake.

As success rates on reducing overweight/obesity are very low, form a public health perspective, the elaboration of policies for increasing intakes of vegetable protein and reducing animal protein and sugars, may be a method of combating overweight/obesity at a population level.

Trial registration: The Food4Me trial was registered as a RCT (NCT01530139) at clinicaltrials.gov (http://clinicaltrials.gov/show/NCT01530139).

Keywords

Food4Me Study; Macronutrient intake; Overweight and Obesity; Protein quality; Vegetable protein.

Introduction

The current worldwide increase in overweight and obesity prevalence represents a serious threat to health and quality of life in most societies (Ng et al,. 2014). The growing concern for being healthy is taking a lead role in the understanding of social culture and population welfare (Kumanyika et al,. 2008). This issue is supported by research evidence, in which according to the recent report of GBD Risk Factors, dietary factors are one of the leading risk factors of non-communicable diseases, and importantly, are high preventable (Phillips et al,. 2014, Cecchini et al,. 2010, G. B. D. Risk Factors Collaborators 2016).

Lifestyle choices, mainly dietary and physical activity patterns, are key factors for preventing the onset of obesity features, type 2 diabetes, cardiovascular diseases, accompanying comorbidities and cancer (Mathers 2015, Raguso et al,. 2006, Milte and McNaughton 2015, Grosso et al,. 2017). Recent studies on the relationship between

protein consumption and prevention/management of obesity, as well as interactions between macronutrient intake and age, sex and physical activity, have revealed controversial or inconsistent results (Riddle et al,. 2016, Levine et al,. 2014, Hernandez-Alonso et al,. 2016, Larsen et al,. 2010, Navas-Carretero et al,. 2016).

In this context, the development of new epidemiological concepts and tools may contribute to envisage public policies and to help health professionals modify risk factors of disease onset to implement adequate therapeutic approaches during all life stages (Srikanthan and Karlamangla 2014, Stenholm, Rantanen, et al,. 2007, Stenholm, Sainio, et al,. 2007, Woods et al,. 2016). Precision nutrition data, based on genetics, clinical history, likes/dislikes, lifestyles, perinatal nutrition, social/cultural constraints, etc., tries to harmonize accurate prediction equations and algorithms with sensitive procedures to give more individualised dietary advice for health management (Forster et al,. 2016, Bauer et al,. 2014). Furthermore, the increasing use of internet, as well as the reduced timetables of working families to attend to the health professional practice, creates a necessity to develop procedures that help professionals to acquire and provide valid information from users, in order to prescribe personalised and valuable nutritional advice (San-Cristobal et al,. 2013, Gibney and Walsh 2013, Fallaize et al,. 2015). Moreover, in research and primary care there is a growing concern about the relative lack of success of public health policies in developed as well as developing countries, and the necessity to combine personalised with public health nutrition advices (Hafekost et al,. 2013, Kumanyika et al,. 2008), where the role of energy-yielding macronutrients on the obesity epidemic needs to be ascertained, although the intake of micronutrients and other food derived compounds will receive attention in the future, when more precise information about micronutrients, food processing effects, etc., is available.

To address such issues, the current analysis used data from the Food4Me study (www.food4me.org), which is an European, randomised controlled trial, designed to investigate the effect of different types of personalised nutrition advice using a web-based intervention platform (Celis-Morales, Livingstone, Marsaux, et al,. 2015).

In this context, the authors hypothesized that an increased consumption of protein from vegetable origin instead of protein from animal origin, using models of isocaloric nutrient exchange, would be associated with a reduction in overweight/obesity. Therefore, the aim of the present study was to evaluate the differences in macronutrient

intake among participating centres, with populations presenting different features, in the Food4Me study, and to assess the role of the substitution of animal and vegetable protein sources by other macronutrients, on the prevalence of overweight or obesity.

Subjects and methods

Study design

The current investigation is an ancillary cross-sectional analysis from the Food4Me randomised controlled trial (Food4Me 2011), including data from 2413 participants. Food4Me was a Pan-European project designed for evaluating the opportunities and challenges of personalised nutrition (Food4Me 2011). The Food4Me study included four arms, conducted in 7 European Countries. The intervention was designed to emulate a real-life web-based personalised nutrition service. The extended protocol of the present intervention has been described elsewhere (Celis-Morales, Livingstone, Marsaux, et al,. 2015).

Ethical approval and participant recruitment

Ethical approval for the study was obtained from all the local research ethics committees at each participating centre. All participants interested in the study were required to sign online consent forms at two stages in the screening process: before sending the first data and before completing the screening food frequency questionnaire (FFQ).

Recruitment for the study was conducted between August 2012 and August 2013, in 7 European centres, as an internet-based personalised nutrition intervention. The recruitment centres for the Food4Me study were as follows: University College Dublin (Ireland); Maastricht University (The Netherlands); University of Navarra (Spain); Harokopio University (Greece); University of Reading (United Kingdom, UK); National Food and Nutrition Institute (Poland); and Technical University of Munich (Germany). To facilitate recruitment, each centre used one or more of the following strategies: Local and National advertising of the study, via newspapers, radio, the internet, posters, flyers or social media; press releases through the internet (online news media), radio, newspaper; and word of mouth (Livingstone et al,. 2016). Participants had to voluntarily register their details on the Food4Me website, participant section (Food4Me 2011), which was designed for the purposes of the study.

Inclusion and exclusion criteria

The Food4Me inclusion criteria were set as wide as possible to include the most representative populations excluding known diseases or prescribed diets. Inclusion criteria were: age > 18 years; to have internet access at home or easily accessible; to be healthy (self-perceived) and without any known food allergy/intolerance.

The exclusion criteria were: pregnancy or planning to be pregnant the next 6 months; following a prescribed diet, including for weight loss, in the last 3 months; to have very limited or no access to the internet; suffering from diabetes, coeliac disease, Crohn's disease, or any other metabolic conditions implying an alteration of nutritional requirements such as known non-controlled thyroid disorders or food allergies and intolerances.

Additionally, for the specific analysis in the current study, all those participants who reported energy intakes lower than 1.1-fold of the calculated basal metabolic rate (1.1*BMR) , or higher than 5000 kcal/day as well as those who provided unrealistic data on height or weight were excluded.

Data collection

Once the interested individuals had registered on the Food4me website, and signed the informed consent forms, they were then asked to fill in online-based questionnaires.

Participants were requested to provide age and sex, ease of internet access, pregnancy, food intolerances and/or allergies, in order to automatically proceed to the inclusion/exclusion of the participants. In addition, self-reported socio-demographic information, health-related issues, anthropometrics data and dietary intakes were collected.

Dietary intake evaluation

An online food frequency questionnaire (FFQ) was completed, where participants provided details of the frequency and portion size of 157 commonly-used food items they had consumed within the previous month, as described elsewhere (Forster et al,. 2014).

This questionnaire provided the list of food items, categorised in food groups, and for each item, the participants had to mark the frequency they had consumed it, expressed as times per month/week/day. In relation to the estimated portion size, for each item, appropriate photographs were shown, in order to help the participant to fill in the most realistic weight of the portion, according to these pictures.

Intakes of foods and nutrients were then computed in real time, using a common food composition database, with a small number of local food-item differences (McCance et al,. 2002). Both the online FFQ and the food database were developed and validated specifically for the Food4Me trial (Forster et al,. 2014, Fallaize et al,. 2014, Marshall et al,. 2016). In the current analyses, the focus was specifically set on macronutrient and energy intakes of the participants, and the percentual exchange of one macronutrient by another.

Anthropometric and physical activity assessment

Body weight and height were self-measured and self-reported via internet following the online instructions available at the website. These self-reported data evidenced good agreement with standard methods, as it was demonstrated in a face-to-face validation of a sample of the Food4Me participants (Celis-Morales, Livingstone, Woolhead, et al,. 2015). Physical activity was assessed with the self-reported occupational and non-occupational activities using the Baecke questionnaire (Marsaux et al,. 2016). Participants were asked to categorise their occupational activity as light (e.g. administrative and managerial), moderate (e.g. sales worker) or heavy (e.g. equipment operator) and their non-occupational activity as sedentary (no activity or little walking/cycling/exercise), moderately active (intense exercise 20-45 minutes, at least twice per week) or very active (intense exercise during at least one hour daily) as published (Marsaux et al,. 2016).

Statistical analyses

Results from descriptive analyses are presented as means and Standard Deviation (SD) for continuous variables or as percentages for categorical variables. Data were analysed using Stata (version 12; Statacorp LP, College Station, TX, USA).

To facilitate the analytical process, participating centres were grouped according to the United Nations Composition of geographical regions, geographical sub-regions, and

196 selected economic and other groupings (Nations). Thus, Ireland and the United
197 Kingdom were analysed together as "British Isles", Germany and The Netherlands were
198 grouped as "Western Europe", Poland was representative of "Eastern Europe", and
199 finally Spain and Greece were grouped as "Southern Europe".

200 To assess the influence of specific variables and to minimize biases, subjects were
201 dichotomised. Thus, participants were categorised by the median age in those below 40
202 years old and those who were 40 years or older. To study the influence of BMI and sex,
203 dichotomised categorical variables were used (under or above 25 kg/m^2, and
204 men/women, respectively).

205 To detect differences between regions, as well as between categorized variables, one
206 factor ANOVA and multivariate linear regression analyses were performed. Results
207 were considered statistically significant for p values <0.05.

208 The multivariate nutrient density model developed by Hu et al (Hu et al,. 1999) was
209 performed to evaluate the influence of protein intake and the isoenergetic substitution of
210 protein sources on the risk of being overweight/obese. In this study, presence or absence
211 of overweight/obesity was considered the event outcome and dependent variable. A
212 multivariate-adjusted logistic regression was performed to calculate the Prevalence Risk
213 Ratio (PRR) of being overweight, using as independent variables the macronutrient
214 proportions (% of total energy intake), excluding protein for which the isoenergetic
215 exchange was analysed, and adjusting for total energy intake, geographical region, age,
216 sex and physical activity level (Moslehi et al,. 2015). This approach allowed assessment
217 of the effect (risk) of the exchange of 1% (total energy intake) of protein by 1% (total
218 energy intake) of other macronutrients on being overweight or obese.

219 Sensitivity analysis for over and under estimation of food intake was performed, using
220 the cut-offs proposed by Goldberg and Black (Black 2000) based on the ratio of
221 reported energy intake/estimated energy requirements, to prevent biases in the risk
222 estimation model.

223 **Results**

224 A total of 5562 subjects across Europe were interested in receiving personalized
225 nutrition advice and registered on the Food4Me website (Figure 1), while 2402 did not
226 complete the whole screening process. Subsequently, these 3160 individuals signed the

two informed consents, filled in the two screening questionnaires and the food frequency questionnaire. Finally, 2413 participants were considered eligible as they provided valid data and fulfilled inclusion criteria.

Participant characteristics

Women represented the 64.69% of the total sample, with the biggest difference in Eastern Europe, where 75% of registered participants were women (Table 1). Therefore, data analyses were adjusted for sex. Anthropometric measures showed slight, but significant differences between regions ($p<0.001$). Notably, height was significantly greater in Western Europe compared with the other regions by 4cm on average. Differences in BMI were also statistically significant, with participants in Western and Eastern Europe presenting the lowest BMI, whereas overweight/obesity was more prevalent in the British Isles and Southern Europe.

In relation to food intake, no differences in energy consumption (kcal/day) were observed, with mean energy intake ranging from 2558 to 2608 kcal/day across regions. However, macronutrient distribution was significantly different ($p<0.001$) across European regions (Table 1). Southern European participants consumed the highest fat content in their diet ($p<0.001$), mainly attributable to olive oil, as MUFA intake was higher (15.2±3.8 %E, $p<0.001$) and SFA intake was lower (13.5±3.1 %E) in southern European participants compared with other EU Regions. Protein intake was significantly higher in Southern Europe (18.6±4.1 %E) compared with Western Europe (16.0±2.9 %E), Eastern Europe (16.8±3.3 %E) and the British Isles (16.3±3.1 %E). Southern Europeans consumed the largest proportion of animal protein (12.2%E) and the lowest percentage of vegetable protein (4.7%), compared with other regions. Western Europe and British Isles participants consumed more dietary fibre (33 g/day), while Eastern Europeans showed the highest intake of carbohydrates (49.2±7.1 %E) and simple sugars (22.0±6.6 %E) compared to the other regions.

When participants were grouped according to age (Table 2), the youngest participants were lighter and had lower BMI (70.6 kg and 24.1 kg/m^2 *vs* 76.9 kg and 26.4 kg/m^2; $p<0.001$ in both measures), as well as higher physical activity level (1.53 *vs* 1.50 arbitrary units, $p<0.001$). Dietary intake was similar between age groups, and there were only very highly significant differences ($p<0.001$) for intake of PUFA and dietary fibre, which was higher in the group > 40 years-old (Table 2). Also, a significant difference

(p=0.02) in protein intake was observed, as younger participants consumed more than older subjects.

As expected, when men were compared with women, significant differences (p<0.001) were found in all the analysed outcomes, concerning anthropometry, physical activity level and dietary intake (table 2), except for protein and carbohydrates intake.

Finally, regression analysis of BMI as dependent variable and protein intake (%E) as the independent variable, adjusted for age, sex, country of origin, total energy intake and physical activity level, showed a weak but significant positive association ($\beta = 0.12$; $p < 0.001$). Comparing participants with BMI below 25 kg/m^2 (n=1322) with those presenting overweight or obesity (BMI $\geq$ 25 kg/m^2; n=1091), this second group were significantly older and practised significantly less physical activity (table 2). In relation to nutrient intake, those presenting overweight reported higher energy intakes (p<0.001). In addition, the overweight group showed a slight but significantly higher proportion of total fat (p=0.046) and lower carbohydrates (p<0.001) in their diets. Also, those subjects with overweight reported significantly higher intakes of total protein (p<0.001).

Contribution of food groups to reported protein intakes

When the sources of protein were evaluated separately (Figure 2a), animal products were the major contributors to dietary protein intake in all regions, although differences between regions were highly significant (p<0.001) for the main protein sources: meat, fish and eggs (33.5% Western Europe, 37.4% Eastern Europe; 39.5% British Isles; and 46.7% Southern Europe); dairy products (19.4% Western Europe; 19.9% Eastern Europe; 17.1% British Isles; 16.6% Southern Europe); bread, grains, pasta and rice (27.8% Western Europe; 25.7% Eastern Europe; 22.8% British Isles; 18.8% Southern Europe) or vegetables (7.6% Western Europe; 5.9% Eastern Europe; 9.1% British Isles; 7.8% Southern Europe).

Moreover, protein sources in overweight/obese subjects, differed significantly from those reported by participants with BMI < 25 kg/m^2 (Figure 2b). Although meat, fish and eggs were the major contributors for protein in both groups, subjects with lower BMI consumed less animal products than overweight/obese subjects (37.9% *vs* 42.9%, p<0.001, respectively), while the percentages of cereals, grains, pasta and rice (24.0%),

and fruits and vegetables (8.3%) were significantly higher among normal weight subjects than in overweight participants (22% of protein from cereals and 7.1% from fruits and vegetables, $p<0.001$).

Influence of isoenergetic protein exchange on the risk of overweight/obesity

No relevant changes in relative risks were found within the sensitivity analyses performed according to available cut-offs.

When the isocaloric nutrient exchange analysis was performed, according to the method described in the statistics section, the focus was set on evaluating the effect of increasing vegetable protein consumption in isolation from other macronutrients. Therefore, the Prevalence Risk Ratio of increasing 1% of total energy from vegetable protein, and the reduction in 1% of total energy of other nutrients is given (Figure 3). The risk of overweight/obesity was reduced if vegetable protein replaced animal protein ($p=0.012$), also observed in relation to sugar consumption ($p=0.006$). The calculated Prevalence Risk Ratio of replacing 1%E of sugar by 1%E of vegetable protein was of 0.940 (95%CI: 0.900-0.982), i.e. a reduction of about 6% in the risk of being overweight/obese. With regards to total fat exchange, although not significant, the substitution of total fat by vegetable protein showed a marginal trend to clearly protect against obesity ($p=0.079$).

The comparison of subjects categorised by tertiles for animal and vegetable protein consumption showed significant differences in BMI (Figure 4), with the highest animal protein consumption (above 11.4% of total energy consumption), associated to the highest BMI, and those participants consuming less vegetable protein had BMI > 25 kg/m^2.

Discussion

A major finding was that a higher vegetable protein intake is associated with reduced risk of developing overweight or obesity, whereas a positive association was found for animal protein consumption and the risk of overweight/obesity. Moreover, the current results revealed that self-reported data, collected via the Internet may be useful for acquiring nutritional and socio-demographic information in large cohorts. Although it cannot be completely confirmed whether this is an accurate reflection of the real food intake, this knowledge will enable the future elaboration of a more accurate and directed

personalised advice. It must be drawn to one's attention that the FFQ used within the present study had already been validated (Fallaize et al,. 2014, Forster et al,. 2014, Marshall et al,. 2016), with respect to the accuracy in estimating changes in dietary habits, being this feasible and simple tool to estimate nutrient intake for screening, which may be considered more a strength than a limitation (Forster et al,. 2016).

In agreement with previous studies on dietary habits, Food4Me online participants from Western Europe, Eastern Europe and the British Isles, reported a larger consumption of saturated fats, whereas Greece and Spain presented the largest intakes of MUFA, possibly because their main fat staple source is traditionally olive oil (Osler and Schroll 1997, Lasheras et al,. 2000, Trichopoulou et al,. 2003). Furthermore, differences in meat and fish consumption were reflected in protein intake (larger in southern countries), and in agreement with other studies, reduced intake of grain and potatoes was associated with lower carbohydrates intake in Southern European countries (Naska et al,. 2006, Trichopoulou et al,. 2007). However, it should be noted that dietary patterns are changing in Mediterranean and other countries towards more westernized patterns (San-Cristobal et al,. 2015, da Silva et al,. 2009, Vardavas et al,. 2010).

In relation to differences found between men and women in nutrients intake, it is interesting to highlight that women in the Food4Me study seemed to present "a priori" a less favourable macronutrient profile, according to their reported total fat, saturated fat and sugars consumption, with a lower intake of dietary fibre. These differences seemed inconsistent given that women had a lower BMI and a significantly higher percentage of men were overweight or obese. In this context, the misreporting of overweight/obese subjects may have played a role (Jessri et al,. 2016). When normal weight subjects were compared to overweight/obese participants, differences in sugar consumption were observed, being higher in the normal weight group, which may be the result of an inverse causality phenomena, indicating that the expected cause-effect response has not been observed due to a bias in reporting, as previously noted in other studies (Santiago et al,. 2015, Santiago et al,. 2013). Also, it must be pointed out that a previous analysis of dietary patterns in the Spanish Food4Me participants identified four dietary patterns, which were consistently associated with subjects' weight-status (San-Cristobal et al,. 2015). In the Spanish cohort, the "compensatory" pattern, characterised by an overconsumption of both beneficial and detrimental food items, was related to the

highest BMI, and a similar association might occur in this larger sample (San-Cristobal et al,. 2015).

The association observed between protein consumption and BMI has also been reported in several life-stages in previous studies (Alkerwi et al,. 2015, Hernandez-Alonso et al,. 2016, Lin et al,. 2015). The PREDIMED trial (Hernandez-Alonso et al,. 2016) as well as the ORISCAV-LUX study (Alkerwi et al,. 2015) showed an association between higher risk of obesity and death with higher total and animal protein consumption, but apparently did not report data on protein intake from vegetable sources. With a similar approach, the HELENA study reported an association between total and animal protein consumption and higher risk of obesity in European adolescents (Lin et al,. 2015). Interestingly, the HELENA study evidenced a protective effect of vegetable protein consumption in the development of obesity among young Europeans, which is in agreement with our data.

In other studies, the dietary intake of protein is positively associated with percent body fat in middle-aged and older adults (Vinknes et al,. 2011), and that cysteine intake may be the causal factor (Elshorbagy et al,. 2012). Thus, it would be of interest to examine the amino acid pattern of the protein sources of our European populations to achieve new insights. Nevertheless, and taking into account the origin of protein, vegetable protein is accompanied by many other micronutrients and compounds which may play a role in metabolism, and these effects must not be disregarded, given the possibility of interaction of phytochemical activity with energy metabolism [50].

In another trial carried out with patients presenting features of the metabolic syndrome, within the RESMENA study (Zulet et al,. 2011, de la Iglesia et al,. 2014), it was observed that protein quality may have an important impact in overweight/obesity, but also in related diseases (Lopez-Legarrea et al,. 2014). In this context, consuming vegetable protein sources under energy restriction was specifically associated to a reduction of inflammatory markers, which allowed to hypothesize that obesity could also be tackled through this anti-inflammatory process. The positive effect of diets differing in macronutrient composition on weight loss has also been previously shown by this research group (Abete et al,. 2009). In a study with hypoenergetic diets, additional benefits in consuming high- legumes or high-protein (30%E), were observed (7% and 8% weight loss, respectively, compared to 5% weight loss with a control diet)

and mitochondrial oxidation, which led the authors to conclude that an increase in energy expenditure led to a higher basal metabolic rate in the volunteers (Abete et al,. 2009). These researchers also observed how inflammatory and lipid markers, and blood pressure improved after the nutritional intervention enriched in protein (Hermsdorff et al,. 2011).

Focusing on weight maintenance, some nutritional interventions revealed better or similar responses to weight control with diets containing higher protein proportions (Larsen et al,. 2010, Navas-Carretero et al,. 2016, Keogh et al,. 2007, Brinkworth et al,. 2004, Clifton et al,. 2014). In this context, nutritional interventions such as the DIOGENES study (Larsen et al,. 2010) have shown that diets with higher protein content (30%E) and lower glycemic index may have a marginal effect on maintaining the weight loss at 6 months (Larsen et al,. 2010) and at 12 months (Aller et al,. 2014). Indeed, a more thorough analysis of these results led to the conclusion that gender may also need to be considered as another factor to integrate in the complex process leading to precision nutrition, in order to prescribe the best possible dietary patterns for each subject (Navas-Carretero et al,. 2016).

Other studies have reported similar positive results, when analysing weight loss and maintenance on higher protein diets compared with control diets as well as on cardiovascular risk markers (Keogh et al,. 2007, Brinkworth et al,. 2004). In any case, it must be considered that in most of these trials and nutritional interventions, the effectiveness of good adherence to the prescribed diet is essential to achieve successful and sustainable weight loss and maintenance results (Clifton et al,. 2014).

Different sources of protein have been investigated through intervention studies and epidemiological cohorts showing distinct health responses depending on the animal and vegetable protein intake, where animal protein sources are associated with increased risk of developing obesity-related diseases (Lin et al,. 2015). However, white meats or fish products have not been often related to these outcomes (Battaglia Richi et al,. 2015), or as in RESMENA study, fish-protein has been correlated, as well as vegetable protein, with positive effects on inflammation (Lopez-Legarrea et al,. 2014). The results of the present analysis, suggest that the risk of overweight or obesity was lower when higher amounts of protein from vegetable origin are consumed, whereas animal protein (in general) has been associated with an increased risk for overweight or obesity.

Interestingly, substitution of 1% E from sugars by vegetable protein demonstrated a lower prevalence of obesity, in agreement with studies reporting that refined carbohydrates may be implicated in the obesity epidemics (Stanhope 2016). The same trend, although with a marginal statistical evidence, was found with the exchange of 1%E from fat by vegetable protein confirming the benefit of vegetable protein increase in the diet (Lopez-Legarrea et al,. 2014, Feskens et al,. 2014).

Current data point to the differential effect and potential interactions of isolated nutrients. Moreover, it shows the importance of the nutrient sources and provides an opportunity for further investigation of dietary patterns. This strategy would take into account the combination of nutrients in a food matrix, behavioural influences, and interaction between different genes (Bauer et al,. 2014, Fallaize et al,. 2013, Kelly et al,. 2016, San-Cristobal et al,. 2015), leading personalised nutrition to the next step of precision nutrition, by considering lifestyle, social environment, and clinical features among others (Ferguson et al,. 2016).

Data collection may be considered a weak point in the present study, given that all measurements were self-reported. However, as mentioned previously, the FFQ used to obtain the analysed data have been validated to ensure the accuracy of measurements (Fallaize et al,. 2014, Forster et al,. 2014, Marshall et al,. 2016). In addition, checking of anthropometrical and genetic markers as proxy for identity was carried out at each intervention centre in a random subsample of the volunteers enrolled in the intervention study (Celis-Morales, Livingstone, Woolhead, et al,. 2015). These results showed a strong validity and agreement between the self-reported data and data collected by trained researchers (Celis-Morales, Livingstone, Woolhead, et al,. 2015). It is also worth mentioning as a strength of the study, the nutrient substitution model used in the analyses, which has been widely used since it was developed (Hu et al,. 1999) and allows to study the association of the substitution of isolated nutrients, as well as some interactions, while energy intake is kept constant, as previously reported (Moslehi et al,. 2015, Skilton et al,. 2008, Vergnaud et al,. 2013).

Nevertheless, some limitations that the current analysis might present must also be mentioned. In this sense, the consideration of "crude" isolated macronutrients on overweight or obesity might disregard some synergistic effects of micronutrients contained in foods that are source of vegetable or animal protein, such as

phytochemicals, vitamins or fibre, and minerals, respectively. It must also be noted that the use of BMI as a marker of adiposity, may underestimate cases of high adipose tissue within a normal weight. However, and acknowledging the limitations, the measure of BMI is still widely used as a proxy screening tool in population studies [10, 47, 48].

The need of developing valid, feasible, effective and economic personalised medicine strategies will also have an impact on precision nutrition, with individual and public health perspectives. In this context, the Food4Me study hypothesized that web-based contact with subjects interested in improving their nutritional status is feasible, and internet-based personalised nutritional advice (Celis-Morales, Livingstone, Marsaux, et al,. 2015) may be a future tool for preventing and managing non-communicable diseases, although the impact of these type of interventions in subjects suffering from specific diseases needs to be assessed. Taking into account differences in macronutrient intakes among countries, a "One size fits all" strategy may be inappropriate, while more specific messages, such as increasing vegetable protein consumption, may be easier to deliver. Indeed, personalising nutritional advice based on the phenotype of individuals, as well as their previous dietary habits may advance our understanding of precision nutrition, because dietary habits differ substantially in European regions (Livingstone et al,. 2016). Although from a public health nutrition point of view, general recommendations are advisable, the need for combining general messages with nutrient-specific targets depending on the region is becoming urgent to reduce the epidemic of obesity and accompanying diseases, such as diabetes, hypertension and dislipemia (Pavlovic et al,. 2007, Jankovic et al,. 2015, Kirwan et al,. 2016), and messages stressing the role of protein and the possible effects depending on the protein quality and sources may be beneficial in this public health actions.

Conclusion

In conclusion, the present results shed light on the differential role of protein quality in the occurrence of overweight/obesity, stressing the importance of increasing vegetable protein sources in our diet, in substitution of animal protein and simple sugars. Differences found in macronutrient intakes depending on region of origin, sex, age and physical activity also point to the importance of personalised nutrition in targeting successful messages for a healthier lifestyles.

Acknowledgements

Not applicable

Funding

This work was supported by the European Commission under the Food, Agriculture, Fisheries and Biotechnology Theme of the 7th Framework Programme for Research and Technological Development [grant agreement: 265494].

The research leading to these results has received funding from "la Caixa" Banking Foundation (RS-C was granted for the PhD work).

The European Commission had no role in the design, analysis or writing of this article.

Competing interests

The authors declare that they have no conflict of interest concerning this research

Authorship

Author responsibilities were as follows: SNC and RSC drafted the paper and performed the statistical analysis for the manuscript. SNC and RSC are joint first authors. JAM was the responsible of Spanish centre of intervention. MG, JCM, JAM, CCM, MCW, ERG, LB, WHMS, HD, CAD, JAL, YM and IT, contributed to the research design. SNC, RSC, KML, CFM, CO'D, HF, CW, ALM, RF, GM, CPL, MJ, and AS conducted the intervention. All authors contributed to a critical review of the manuscript during the writing process. All authors approved the final version to be published.

Ethical Standards Disclosure

All procedures performed in the study were in accordance with the ethical standards of the corresponding research committees of each of the seven participating centres, and with the 1964 Helsinki declaration and its later amendments or comparable ethical standards. Written informed consent was obtained from all individual participants included in the study.

The Research Ethics Committees evaluating the study protocol were those with the appropriate authority in each study site: University College Dublin, Ireland; University of Maastricht, Netherlands; Universidad de Navarra, Spain; Harokopio University, Greece; The University of Reading, United Kingdom; National Food and Nutrition Institute, Poland; Technische Universitaet Muenchen, German.

Being the study coordinator Ireland, the relevant Health Authority in Food4Me study was the Research Ethics Committee of Ireland.

The Food4Me trial was registered as a RCT (NCT01530139) at clinicaltrials.gov (http://clinicaltrials.gov/show/NCT01530139).

References

Abete, I., Parra, D. & Martinez, J. A. (2009). Legume-, fish-, or high-protein-based hypocaloric diets: effects on weight loss and mitochondrial oxidation in obese men. *J Med Food,* **12,** 100-8.

Alkerwi, A., Sauvageot, N., Buckley, J. D., Donneau, A. F., Albert, A., Guillaume, M. & Crichton, G. E. (2015). The potential impact of animal protein intake on global and abdominal obesity: evidence from the Observation of Cardiovascular Risk Factors in Luxembourg (ORISCAV-LUX) study. *Public Health Nutr,* **18,** 1831-8.

Aller, E. E., Larsen, T. M., Claus, H., Lindroos, A. K., Kafatos, A., Pfeiffer, A., Martinez, J. A., Handjieva-Darlenska, T., Kunesova, M., Stender, S., Saris, W. H., Astrup, A. & van Baak, M. A. (2014). Weight loss maintenance in overweight subjects on ad libitum diets with high or low protein content and glycemic index: the DIOGENES trial 12-month results. *Int J Obes (Lond),* **38,** 1511-7.

Battaglia Richi, E., Baumer, B., Conrad, B., Darioli, R., Schmid, A. & Keller, U. (2015). Health Risks Associated with Meat Consumption: A Review of Epidemiological Studies. *Int J Vitam Nutr Res,* **85,** 70-8.

Bauer, D. C., Gaff, C., Dinger, M. E., Caramins, M., Buske, F. A., Fenech, M., Hansen, D. & Cobiac, L. (2014). Genomics and personalised whole-of-life healthcare. *Trends Mol Med,* **20,** 479-86.

Black, A. E. (2000). Critical evaluation of energy intake using the Goldberg cut-off for energy intake:basal metabolic rate. A practical guide to its calculation, use and limitations. *Int J Obes Relat Metab Disord,* **24,** 1119-30.

Brinkworth, G. D., Noakes, M., Keogh, J. B., Luscombe, N. D., Wittert, G. A. & Clifton, P. M. (2004). Long-term effects of a high-protein, low-carbohydrate diet on weight control and cardiovascular risk markers in obese hyperinsulinemic subjects. *Int J Obes Relat Metab Disord,* **28,** 661-70.

Cecchini, M., Sassi, F., Lauer, J. A., Lee, Y. Y., Guajardo-Barron, V. & Chisholm, D. (2010). Tackling of unhealthy diets, physical inactivity, and obesity: health effects and cost-effectiveness. *Lancet,* **376,** 1775-84.

Celis-Morales, C., Livingstone, K. M., Marsaux, C. F., Forster, H., O'Donovan, C. B., Woolhead, C., Macready, A. L., Fallaize, R., Navas-Carretero, S., San-Cristobal, R., Kolossa, S., Hartwig, K., Tsirigoti, L., Lambrinou, C. P., Moschonis, G., Godlewska, M., Surwillo, A., Grimaldi, K., Bouwman, J., Daly, E. J., Akujobi, V., O'Riordan, R., Hoonhout, J., Claassen, A., Hoeller, U., Gundersen, T. E., Kaland, S. E., Matthews, J. N., Manios, Y., Traczyk, I., Drevon, C. A., Gibney, E. R., Brennan, L., Walsh, M. C., Lovegrove, J. A., Alfredo Martinez, J., Saris, W. H., Daniel, H., Gibney, M. & Mathers, J. C. (2015). Design and baseline characteristics of the Food4Me study: a web-based randomised controlled trial of personalised nutrition in seven European countries. *Genes Nutr,* **10,** 450.

Celis-Morales, C., Livingstone, K. M., Woolhead, C., Forster, H., O'Donovan, C. B., Macready, A. L., Fallaize, R., Marsaux, C. F., Tsirigoti, L., Efstathopoulou, E., Moschonis, G., Navas-Carretero, S., San-Cristobal, R., Kolossa, S., Klein, U. L., Hallmann, J., Godlewska, M., Surwillo, A., Drevon, C. A., Bouwman, J., Grimaldi, K., Parnell, L. D., Manios, Y., Traczyk, I., Gibney, E. R., Brennan, L., Walsh, M. C., Lovegrove, J. A., Martinez, J. A., Daniel, H., Saris, W. H., Gibney, M. & Mathers, J. C. (2015). How reliable is internet-based self-reported identity, socio-demographic and obesity measures in European adults? *Genes Nutr,* **10,** 476.

Clifton, P. M., Condo, D. & Keogh, J. B. (2014). Long term weight maintenance after advice to consume low carbohydrate, higher protein diets--a systematic review and meta analysis. *Nutr Metab Cardiovasc Dis,* **24,** 224-35.

Collaborators, G. B. D. R. F. (2016). Global, regional, and national comparative risk assessment of 79 behavioural, environmental and occupational, and metabolic risks or clusters of risks, 1990-2015: a systematic analysis for the Global Burden of Disease Study 2015. *Lancet,* **388,** 1659-1724.

da Silva, R., Bach-Faig, A., Raido Quintana, B., Buckland, G., Vaz de Almeida, M. D. & Serra-Majem, L. (2009). Worldwide variation of adherence to the Mediterranean diet, in 1961-1965 and 2000-2003. *Public Health Nutr,* **12,** 1676-84.

de la Iglesia, R., Lopez-Legarrea, P., Abete, I., Bondia-Pons, I., Navas-Carretero, S., Forga, L., Martinez, J. A. & Zulet, M. A. (2014). A new dietary strategy for long-term treatment of the metabolic syndrome is compared with the American Heart Association (AHA) guidelines: the MEtabolic Syndrome REduction in NAvarra (RESMENA) project. *Br J Nutr,* **111,** 643-52.

Elshorbagy, A. K., Kozich, V., Smith, A. D. & Refsum, H. (2012). Cysteine and obesity: consistency of the evidence across epidemiologic, animal and cellular studies. *Curr Opin Clin Nutr Metab Care,* **15,** 49-57.

Fallaize, R., Forster, H., Macready, A. L., Walsh, M. C., Mathers, J. C., Brennan, L., Gibney, E. R., Gibney, M. J. & Lovegrove, J. A. (2014). Online dietary intake estimation: reproducibility and validity of the Food4Me food frequency questionnaire against a 4-day weighed food record. *J Med Internet Res,* **16,** e190.

Fallaize, R., Macready, A. L., Butler, L. T., Ellis, J. A., Berezowska, A., Fischer, A. R., Walsh, M. C., Gallagher, C., Stewart-Knox, B. J., Kuznesof, S., Frewer, L. J., Gibney, M. J. & Lovegrove, J. A. (2015). The perceived impact of the National Health Service on personalised nutrition service delivery among the UK public. *Br J Nutr,* **113,** 1271-9.

Fallaize, R., Macready, A. L., Butler, L. T., Ellis, J. A. & Lovegrove, J. A. (2013). An insight into the public acceptance of nutrigenomic-based personalised nutrition. *Nutr Res Rev,* **26,** 39-48.

Ferguson, L. R., De Caterina, R., Gorman, U., Allayee, H., Kohlmeier, M., Prasad, C., Choi, M. S., Curi, R., de Luis, D. A., Gil, A., Kang, J. X., Martin, R. L., Milagro, F. I., Nicoletti, C. F., Nonino, C. B., Ordovas, J. M., Parslow, V. R., Portillo, M. P., Santos, J. L., Serhan, C. N., Simopoulos, A. P., Velazquez-Arellano, A., Zulet, M. A. & Martinez, J. A. (2016). Guide and Position of the International Society of Nutrigenetics/Nutrigenomics on Personalised Nutrition: Part 1 - Fields of Precision Nutrition. *J Nutrigenet Nutrigenomics,* **9,** 12-27.

Feskens, E. J., Sluik, D. & Du, H. (2014). The Association Between Diet and Obesity in Specific European Cohorts: DiOGenes and EPIC-PANACEA. *Curr Obes Rep,* **3,** 67-78.

Food4Me [internet]. 2011. [cited 15/02/2016]. Available from: http://www.food4me.org/

Forster, H., Fallaize, R., Gallagher, C., O'Donovan, C. B., Woolhead, C., Walsh, M. C., Macready, A. L., Lovegrove, J. A., Mathers, J. C., Gibney, M. J., Brennan, L. & Gibney, E. R. (2014). Online dietary intake estimation: the Food4Me food frequency questionnaire. *J Med Internet Res,* **16,** e150.

Forster, H., Walsh, M. C., Gibney, M. J., Brennan, L. & Gibney, E. R. (2016). Personalised nutrition: the role of new dietary assessment methods. *Proc Nutr Soc,* **75,** 96-105.

Gibney, M. J. & Walsh, M. C. (2013). The future direction of personalised nutrition: my diet, my phenotype, my genes. *Proc Nutr Soc,* **72,** 219-25.

Grosso, G., Bella, F., Godos, J., Sciacca, S., Del Rio, D., Ray, S., Galvano, F. & Giovannucci, E. L. (2017). Possible role of diet in cancer: systematic review and multiple meta-analyses of dietary patterns, lifestyle factors, and cancer risk. *Nutr Rev,* **75,** 405-419.

Hafekost, K., Lawrence, D., Mitrou, F., O'Sullivan, T. A. & Zubrick, S. R. (2013). Tackling overweight and obesity: does the public health message match the science? *BMC Med,* **11,** 41.

Hermsdorff, H. H., Zulet, M. A., Abete, I. & Martinez, J. A. (2011). A legume-based hypocaloric diet reduces proinflammatory status and improves metabolic features in overweight/obese subjects. *Eur J Nutr,* **50,** 61-9.

Hernandez-Alonso, P., Salas-Salvado, J., Ruiz-Canela, M., Corella, D., Estruch, R., Fito, M., Aros, F., Gomez-Gracia, E., Fiol, M., Lapetra, J., Basora, J., Serra-Majem, L., Munoz, M. A., Buil-Cosiales, P., Saiz, C. & Bullo, M. (2016). High dietary protein intake is associated with an increased body weight and total death risk. *Clin Nutr,* **35,** 496-506.

Hu, F. B., Stampfer, M. J., Rimm, E., Ascherio, A., Rosner, B. A., Spiegelman, D. & Willett, W. C. (1999). Dietary fat and coronary heart disease: a comparison of approaches for adjusting for total energy intake and modeling repeated dietary measurements. *Am J Epidemiol,* **149,** 531-40.

Jankovic, N., Geelen, A., Streppel, M. T., de Groot, L. C., Kiefte-de Jong, J. C., Orfanos, P., Bamia, C., Trichopoulou, A., Boffetta, P., Bobak, M., Pikhart, H., Kee, F., O'Doherty, M. G., Buckland, G., Woodside, J., Franco, O. H., Ikram, M. A., Struijk, E. A., Pajak, A., Malyutina, S., Kubinova, R., Wennberg, M., Park, Y., Bueno-de-Mesquita, H. B., Kampman, E. & Feskens, E. J. (2015). WHO guidelines for a healthy diet and mortality

631 from cardiovascular disease in European and American elderly: the CHANCES project.
632 *Am J Clin Nutr,* **102,** 745-56.
633 Jessri, M., Lou, W. Y. & L'Abbe, M. R. (2016). Evaluation of different methods to handle
634 misreporting in obesity research: evidence from the Canadian national nutrition
635 survey. *Br J Nutr,* **115,** 147-59.
636 Kelly, S., Martin, S., Kuhn, I., Cowan, A., Brayne, C. & Lafortune, L. (2016). Barriers and
637 Facilitators to the Uptake and Maintenance of Healthy Behaviours by People at Mid-
638 Life: A Rapid Systematic Review. *PLoS One,* **11,** e0145074.
639 Keogh, J. B., Luscombe-Marsh, N. D., Noakes, M., Wittert, G. A. & Clifton, P. M. (2007). Long-
640 term weight maintenance and cardiovascular risk factors are not different following
641 weight loss on carbohydrate-restricted diets high in either monounsaturated fat or
642 protein in obese hyperinsulinaemic men and women. *Br J Nutr,* **97,** 405-10.
643 Kirwan, L., Walsh, M. C., Brennan, L., Gibney, E. R., Drevon, C. A., Daniel, H., Lovegrove, J. A.,
644 Manios, Y., Martinez, J. A., Saris, W. H., Traczyk, I., Mathers, J. C. & Gibney, M. (2016).
645 Comparison of the portion size and frequency of consumption of 156 foods across
646 seven European countries: insights from the Food4ME study. *Eur J Clin Nutr,* **70,** 642-4.
647 Kumanyika, S. K., Obarzanek, E., Stettler, N., Bell, R., Field, A. E., Fortmann, S. P., Franklin, B. A.,
648 Gillman, M. W., Lewis, C. E., Poston, W. C., 2nd, Stevens, J. & Hong, Y. (2008).
649 Population-based prevention of obesity: the need for comprehensive promotion of
650 healthful eating, physical activity, and energy balance: a scientific statement from
651 American Heart Association Council on Epidemiology and Prevention, Interdisciplinary
652 Committee for Prevention (formerly the expert panel on population and prevention
653 science). *Circulation,* **118,** 428-64.
654 Larsen, T. M., Dalskov, S. M., van Baak, M., Jebb, S. A., Papadaki, A., Pfeiffer, A. F., Martinez, J.
655 A., Handjieva-Darlenska, T., Kunesova, M., Pihlsgard, M., Stender, S., Holst, C., Saris, W.
656 H. & Astrup, A. (2010). Diets with high or low protein content and glycemic index for
657 weight-loss maintenance. *N Engl J Med,* **363,** 2102-13.
658 Lasheras, C., Fernandez, S. & Patterson, A. M. (2000). Mediterranean diet and age with respect
659 to overall survival in institutionalized, nonsmoking elderly people. *Am J Clin Nutr,* **71,**
660 987-92.
661 Levine, M. E., Suarez, J. A., Brandhorst, S., Balasubramanian, P., Cheng, C. W., Madia, F.,
662 Fontana, L., Mirisola, M. G., Guevara-Aguirre, J., Wan, J., Passarino, G., Kennedy, B. K.,
663 Wei, M., Cohen, P., Crimmins, E. M. & Longo, V. D. (2014). Low protein intake is
664 associated with a major reduction in IGF-1, cancer, and overall mortality in the 65 and
665 younger but not older population. *Cell Metab,* **19,** 407-17.
666 Lin, Y., Mouratidou, T., Vereecken, C., Kersting, M., Bolca, S., de Moraes, A. C., Cuenca-Garcia,
667 M., Moreno, L. A., Gonzalez-Gross, M., Valtuena, J., Labayen, I., Grammatikaki, E.,
668 Hallstrom, L., Leclercq, C., Ferrari, M., Gottrand, F., Beghin, L., Manios, Y., Ottevaere,
669 C., Van Oyen, H., Molnar, D., Kafatos, A., Widhalm, K., Gomez-Martinez, S., Prieto, L. E.,
670 De Henauw, S. & Huybrechts, I. (2015). Dietary animal and plant protein intakes and
671 their associations with obesity and cardio-metabolic indicators in European
672 adolescents: the HELENA cross-sectional study. *Nutr J,* **14,** 10.
673 Livingstone, K. M., Celis-Morales, C., Navas-Carretero, S., San-Cristobal, R., O'Donovan, C. B.,
674 Forster, H., Woolhead, C., Marsaux, C. F., Macready, A. L., Fallaize, R., Kolossa, S.,
675 Tsirigoti, L., Lambrinou, C. P., Moschonis, G., Godlewska, M., Surwillo, A., Drevon, C. A.,
676 Manios, Y., Traczyk, I., Gibney, E. R., Brennan, L., Walsh, M. C., Lovegrove, J. A., Alfredo
677 Martinez, J., Saris, W. H., Daniel, H., Gibney, M. & Mathers, J. C. (2016). Profile of
678 European adults interested in internet-based personalised nutrition: the Food4Me
679 study. *Eur J Nutr,* **55,** 759-69.
680 Lopez-Legarrea, P., de la Iglesia, R., Abete, I., Navas-Carretero, S., Martinez, J. A. & Zulet, M. A.
681 (2014). The protein type within a hypocaloric diet affects obesity-related
682 inflammation: the RESMENA project. *Nutrition,* **30,** 424-9.

683 Marsaux, C. F., Celis-Morales, C., Hoonhout, J., Claassen, A., Goris, A., Forster, H., Fallaize, R.,
684 Macready, A. L., Navas-Carretero, S., Kolossa, S., Walsh, M. C., Lambrinou, C. P.,
685 Manios, Y., Godlewska, M., Traczyk, I., Lovegrove, J. A., Martinez, J. A., Daniel, H.,
686 Gibney, M., Mathers, J. C. & Saris, W. H. (2016). Objectively Measured Physical Activity
687 in European Adults: Cross-Sectional Findings from the Food4Me Study. *PLoS One,* **11,**
688 e0150902.
689 Marshall, S. J., Livingstone, K. M., Celis-Morales, C., Forster, H., Fallaize, R., O'Donovan, C. B.,
690 Woolhead, C., Marsaux, C. F., Macready, A. L., Navas-Carretero, S., San-Cristobal, R.,
691 Kolossa, S., Tsirigoti, L., Lambrinou, C. P., Moschonis, G., Godlewska, M., Surwillo, A.,
692 Drevon, C. A., Manios, Y., Traczyk, I., Martinez, J. A., Saris, W. H., Daniel, H., Gibney, E.
693 R., Brennan, L., Walsh, M. C., Lovegrove, J. A., Gibney, M. & Mathers, J. C. (2016).
694 Reproducibility of the Online Food4Me Food-Frequency Questionnaire for Estimating
695 Dietary Intakes across Europe. *J Nutr,* **146,** 1068-75.
696 Mathers, J. C. (2015). Impact of nutrition on the ageing process. *Br J Nutr,* **113 Suppl,** S18-22.
697 McCance, R. A., Widdowson, E. M., Royal Society of Chemistry (Great Britain), Food Standards
698 Agency (Great Britain) & AFRC Institute of Food Research (2002). The composition of
699 foods. Pp. XV, 537 p. Cambridge (UK) London, Royal Society of Chemistry ;.
700 Milte, C. M. & McNaughton, S. A. (2015). Dietary patterns and successful ageing: a systematic
701 review. *Eur J Nutr*.
702 Moslehi, N., Ehsani, B., Mirmiran, P., Hojjat, P. & Azizi, F. (2015). Association of Dietary
703 Proportions of Macronutrients with Visceral Adiposity Index: Non-Substitution and Iso-
704 Energetic Substitution Models in a Prospective Study. *Nutrients,* **7,** 8859-70.
705 Naska, A., Fouskakis, D., Oikonomou, E., Almeida, M. D., Berg, M. A., Gedrich, K., Moreiras, O.,
706 Nelson, M., Trygg, K., Turrini, A., Remaut, A. M., Volatier, J. L. & Trichopoulou, A.
707 (2006). Dietary patterns and their socio-demographic determinants in 10 European
708 countries: data from the DAFNE databank. *Eur J Clin Nutr,* **60,** 181-90.
709 Nations, U. U. [internet]. [cited 05/11/2015]. Available from:
710 http://millenniumindicators.un.org/unsd/methods/m49/m49regin.htm
711 Navas-Carretero, S., Holst, C., Saris, W. H., van Baak, M. A., Jebb, S. A., Kafatos, A., Papadaki,
712 A., Pfeiffer, A. F., Handjieva-Darlenska, T., Hlavaty, P., Stender, S., Larsen, T. M., Astrup,
713 A. & Martinez, J. A. (2016). The Impact of Gender and Protein Intake on the Success of
714 Weight Maintenance and Associated Cardiovascular Risk Benefits, Independent of the
715 Mode of Food Provision: The DiOGenes Randomized Trial. *J Am Coll Nutr,* **35,** 20-30.
716 Ng, M., Fleming, T., Robinson, M., Thomson, B., Graetz, N., Margono, C., Mullany, E. C.,
717 Biryukov, S., Abbafati, C., Abera, S. F., Abraham, J. P., Abu-Rmeileh, N. M., Achoki, T.,
718 AlBuhairan, F. S., Alemu, Z. A., Alfonso, R., Ali, M. K., Ali, R., Guzman, N. A., Ammar, W.,
719 Anwari, P., Banerjee, A., Barquera, S., Basu, S., Bennett, D. A., Bhutta, Z., Blore, J.,
720 Cabral, N., Nonato, I. C., Chang, J. C., Chowdhury, R., Courville, K. J., Criqui, M. H.,
721 Cundiff, D. K., Dabhadkar, K. C., Dandona, L., Davis, A., Dayama, A., Dharmaratne, S. D.,
722 Ding, E. L., Durrani, A. M., Esteghamati, A., Farzadfar, F., Fay, D. F., Feigin, V. L.,
723 Flaxman, A., Forouzanfar, M. H., Goto, A., Green, M. A., Gupta, R., Hafezi-Nejad, N.,
724 Hankey, G. J., Harewood, H. C., Havmoeller, R., Hay, S., Hernandez, L., Husseini, A.,
725 Idrisov, B. T., Ikeda, N., Islami, F., Jahangir, E., Jassal, S. K., Jee, S. H., Jeffreys, M., Jonas,
726 J. B., Kabagambe, E. K., Khalifa, S. E., Kengne, A. P., Khader, Y. S., Khang, Y. H., Kim, D.,
727 Kimokoti, R. W., Kinge, J. M., Kokubo, Y., Kosen, S., Kwan, G., Lai, T., Leinsalu, M., Li, Y.,
728 Liang, X., Liu, S., Logroscino, G., Lotufo, P. A., Lu, Y., Ma, J., Mainoo, N. K., Mensah, G.
729 A., Merriman, T. R., Mokdad, A. H., Moschandreas, J., Naghavi, M., Naheed, A., Nand,
730 D., Narayan, K. M., Nelson, E. L., Neuhouser, M. L., Nisar, M. I., Ohkubo, T., Oti, S. O.,
731 Pedroza, A., et al. (2014). Global, regional, and national prevalence of overweight and
732 obesity in children and adults during 1980-2013: a systematic analysis for the Global
733 Burden of Disease Study 2013. *Lancet,* **384,** 766-81.

734 Osler, M. & Schroll, M. (1997). Diet and mortality in a cohort of elderly people in a north
735 European community. *Int J Epidemiol,* **26,** 155-9.
736 Pavlovic, M., Prentice, A., Thorsdottir, I., Wolfram, G. & Branca, F. (2007). Challenges in
737 harmonizing energy and nutrient recommendations in Europe. *Ann Nutr Metab,* **51,**
738 108-14.
739 Phillips, K., Wood, F. & Kinnersley, P. (2014). Tackling obesity: the challenge of obesity
740 management for practice nurses in primary care. *Fam Pract,* **31,** 51-9.
741 Raguso, C. A., Kyle, U., Kossovsky, M. P., Roynette, C., Paoloni-Giacobino, A., Hans, D., Genton,
742 L. & Pichard, C. (2006). A 3-year longitudinal study on body composition changes in the
743 elderly: role of physical exercise. *Clin Nutr,* **25,** 573-80.
744 Riddle, E. S., Stipanuk, M. H. & Thalacker-Mercer, A. E. (2016). Amino acids in healthy aging
745 skeletal muscle. *Front Biosci (Elite Ed),* **8,** 326-50.
746 San-Cristobal, R., Milagro, F. I. & Martinez, J. A. (2013). Future challenges and present ethical
747 considerations in the use of personalized nutrition based on genetic advice. *J Acad*
748 *Nutr Diet,* **113,** 1447-54.
749 San-Cristobal, R., Navas-Carretero, S., Celis-Morales, C., Brennan, L., Walsh, M., Lovegrove, J.
750 A., Daniel, H., Saris, W. H., Traczyk, I., Manios, Y., Gibney, E. R., Gibney, M. J., Mathers,
751 J. C. & Martinez, J. A. (2015). Analysis of Dietary Pattern Impact on Weight Status for
752 Personalised Nutrition through On-Line Advice: The Food4Me Spanish Cohort.
753 *Nutrients,* **7,** 9523-37.
754 Santiago, S., Sayon-Orea, C., Babio, N., Ruiz-Canela, M., Marti, A., Corella, D., Estruch, R., Fito,
755 M., Aros, F., Ros, E., Gomez-Garcia, E., Fiol, M., Lapetra, J., Serra-Majem, L., Becerra-
756 Tomas, N., Salas-Salvado, J., Pinto, X., Schroder, H. & Martinez, J. A. (2015). Yogurt
757 consumption and abdominal obesity reversion in the PREDIMED study. *Nutr Metab*
758 *Cardiovasc Dis*.
759 Santiago, S., Zazpe, I., Marti, A., Cuervo, M. & Martinez, J. A. (2013). Gender differences in
760 lifestyle determinants of overweight prevalence in a sample of Southern European
761 children. *Obes Res Clin Pract,* **7,** e391-400.
762 Skilton, M. R., Laville, M., Cust, A. E., Moulin, P. & Bonnet, F. (2008). The association between
763 dietary macronutrient intake and the prevalence of the metabolic syndrome. *Br J Nutr,*
764 **100,** 400-7.
765 Srikanthan, P. & Karlamangla, A. S. (2014). Muscle mass index as a predictor of longevity in
766 older adults. *Am J Med,* **127,** 547-53.
767 Stanhope, K. L. (2016). Sugar consumption, metabolic disease and obesity: The state of the
768 controversy. *Crit Rev Clin Lab Sci,* **53,** 52-67.
769 Stenholm, S., Rantanen, T., Alanen, E., Reunanen, A., Sainio, P. & Koskinen, S. (2007). Obesity
770 history as a predictor of walking limitation at old age. *Obesity (Silver Spring),* **15,** 929-
771 38.
772 Stenholm, S., Sainio, P., Rantanen, T., Koskinen, S., Jula, A., Heliovaara, M. & Aromaa, A.
773 (2007). High body mass index and physical impairments as predictors of walking
774 limitation 22 years later in adult Finns. *J Gerontol A Biol Sci Med Sci,* **62,** 859-65.
775 Trichopoulou, A., Bamia, C., Norat, T., Overvad, K., Schmidt, E. B., Tjonneland, A., Halkjaer, J.,
776 Clavel-Chapelon, F., Vercambre, M. N., Boutron-Ruault, M. C., Linseisen, J., Rohrmann,
777 S., Boeing, H., Weikert, C., Benetou, V., Psaltopoulou, T., Orfanos, P., Boffetta, P.,
778 Masala, G., Pala, V., Panico, S., Tumino, R., Sacerdote, C., Bueno-de-Mesquita, H. B.,
779 Ocke, M. C., Peeters, P. H., Van der Schouw, Y. T., Gonzalez, C., Sanchez, M. J.,
780 Chirlaque, M. D., Moreno, C., Larranaga, N., Van Guelpen, B., Jansson, J. H., Bingham,
781 S., Khaw, K. T., Spencer, E. A., Key, T., Riboli, E. & Trichopoulos, D. (2007). Modified
782 Mediterranean diet and survival after myocardial infarction: the EPIC-Elderly study. *Eur*
783 *J Epidemiol,* **22,** 871-81.
784 Trichopoulou, A., Costacou, T., Bamia, C. & Trichopoulos, D. (2003). Adherence to a
785 Mediterranean diet and survival in a Greek population. *N Engl J Med,* **348,** 2599-608.

786 Vardavas, C. I., Linardakis, M. K., Hatzis, C. M., Saris, W. H. & Kafatos, A. G. (2010).
787 Cardiovascular disease risk factors and dietary habits of farmers from Crete 45 years
788 after the first description of the Mediterranean diet. *Eur J Cardiovasc Prev Rehabil,* **17,**
789 440-6.
790 Vergnaud, A. C., Norat, T., Mouw, T., Romaguera, D., May, A. M., Bueno-de-Mesquita, H. B.,
791 van der, A. D., Agudo, A., Wareham, N., Khaw, K. T., Romieu, I., Freisling, H., Slimani,
792 N., Perquier, F., Boutron-Ruault, M. C., Clavel-Chapelon, F., Palli, D., Berrino, F.,
793 Mattiello, A., Tumino, R., Ricceri, F., Rodriguez, L., Molina-Montes, E., Amiano, P.,
794 Barricarte, A., Chirlaque, M. D., Crowe, F. L., Orfanos, P., Naska, A., Trichopoulou, A.,
795 Teucher, B., Kaaks, R., Boeing, H., Buijsse, B., Johansson, I., Hallmans, G., Drake, I.,
796 Sonestedt, E., Jakobsen, M. U., Overvad, K., Tjonneland, A., Halkjaer, J., Skeie, G.,
797 Braaten, T., Lund, E., Riboli, E. & Peeters, P. H. (2013). Macronutrient composition of
798 the diet and prospective weight change in participants of the EPIC-PANACEA study.
799 *PLoS One,* **8,** e57300.
800 Vinknes, K. J., de Vogel, S., Elshorbagy, A. K., Nurk, E., Drevon, C. A., Gjesdal, C. G., Tell, G. S.,
801 Vollset, S. E. & Refsum, H. (2011). Dietary intake of protein is positively associated with
802 percent body fat in middle-aged and older adults. *J Nutr,* **141,** 440-6.
803 Woods, N. F., Rillamas-Sun, E., Cochrane, B. B., La Croix, A. Z., Seeman, T. E., Tindle, H. A.,
804 Zaslavsky, O., Bird, C. E., Johnson, K. C., Manson, J. E., Ockene, J. K., Seguin, R. A. &
805 Wallace, R. B. (2016). Aging Well: Observations From the Women's Health Initiative
806 Study. *J Gerontol A Biol Sci Med Sci,* **71 Suppl 1,** S3-S12.
807 Zulet, M. A., Bondia-Pons, I., Abete, I., de la Iglesia, R., Lopez-Legarrea, P., Forga, L., Navas-
808 Carretero, S. & Martinez, J. A. (2011). The reduction of the metabolyc syndrome in
809 Navarra-Spain (RESMENA-S) study: a multidisciplinary strategy based on
810 chrononutrition and nutritional education, together with dietetic and psychological
811 control. *Nutr Hosp,* **26,** 16-26.

812

Table 1. Demographic anthropometric and dietary characteristics of subjects in the Food4Me studyby European Regions at screening.

	Total		Western Europe		British Isles		Southern Europe		Eastern Europe		p[1]
	Mean	SD	Mean	SD	Mean	SD	Mean	SD	Mean	SD	
n (n females)	2413 (1561)		742 (462)		518 (341)		900 (568)		253 (190)		
Age (years)	40.2	13.0	46.0	14.1^{c}	37.8	13.1^{a,b}	38.1	10.3^{b}	35.7	12.7^{a}	<0.001
Height (m)	1.71	0.09	1.74	0.09^{b}	1.70	0.09^{a}	1.69	0.09^{a}	1.69	0.08^{a}	<0.001
Weight (kg)	73.6	15.5	74.8	14.5^{a}	72.9	15.7^{a,b}	74.0	16.1^{a}	70.4	15.7^{b}	<0.001
BMI (kg/m^2)	25.2	4.7	24.7	4.0^{a}	25.2	4.7^{a}	25.9	5.0^{b}	24.6	4.9^{a}	<0.001
BMI status											
Underweight (< 18.5 kg/m^2)	2.3%		2.4%		2.1%		1.8%		4.3%		
Normal weight (18.5 – 24.99 kg/m^2)	52.5%		56.6%		55.0%		46.9%		54.9%		<0.001[2]
Overweight (25 – 29.9 kg/m^2)	31.1%		30.5%		28.6%		34.2%		26.9%		
Obesity (> 30 kg/m^2)	14.1%		10.5%		14.3%		17.1%		13.8%		
Physical activity factor (AU)	1.51	0.10	1.52	0.10^{b}	1.53	0.10^{b}	1.50	0.10^{a}	1.50	0.11^{a}	<0.001
Energy (kcal/d)	2602	797	2609	744	2632	823	2592	823	2558	802	0.641
Total fat (% E)	35.7	6.4	35.4	6.5^{a}	35.6	6.2^{a,b}	36.4	6.6^{b}	34.5	5.7^{a}	<0.001
SFA (% E)	14.0	3.4	14.2	3.6^{b}	14.2	3.4^{b}	13.5	3.1^{a}	14.5	3.6^{b}	<0.001
MUFA (% E)	13.8	3.4	13.0	2.9^{b}	13.2	3.0^{b}	15.2	3.8^{c}	11.9	2.2^{a}	<0.001
PUFA (% E)	5.6	1.4	5.9	1.4^{b}	5.8	1.5^{b}	5.2	1.3^{a}	5.8	1.6^{b}	<0.001
Protein (% E)	17.1	3.6	16.0	2.9^{a}	16.3	3.1^{a,b}	18.6	4.1^{c}	16.8	3.3^{c}	<0.001
Animal Protein (%E)	10.3	4.3	8.7	3.6^{a}	9.5	3.6^{b}	12.2	4.7^{c}	9.9	3.8^{b}	<0.001
Vegetable Protein (%E)	5.1	1.7	5.5	1.7^{c}	5.0	1.5^{b}	4.7	1.7^{a}	5.1	1.7^{b}	<0.001
Total carbohydrates (% E)	46.4	8.2	46.9	8.0^{b}	47.1	8.1^{b}	44.7	8.5^{a}	49.2	7.1^{c}	<0.001
Simple sugars (% E)	21.3	6.4	20.8	5.9^{a}	22.0	6.2^{b}	21.1	6.7^{a,b}	22.0	6.6^{b}	0.002
Dietary fibre (g)	30.8	13.3	33.4	14.2^{b}	33.1	13.6^{b}	27.3	11.8^{a}	30.8	12.6^{b}	<0.001

Values are expressed as mean and SD (Standard Deviation) or percentages. Western = Netherlands and Germany; British isles = United Kingdom and Ireland; BMI: Body Mass Index; AU = Arbitrary Units; SFA = Saturated fatty acids; MUFA = Monounsaturated fatty acids; PUFA = Polyunsaturated fatty acids; %E = % of total energy intake. [1]Differences between European Regions analysed by one way ANOVA. [2] Chi-square p-value for distribution. Different superscript letters mean significant differences among regions ($p<0.05$) in Tukey post-hoc analysis.

Table 2. Differences among Food4Me screenees according to age, sex and BMI.

	<40 years		≥40 years		p[1]	Male		Female		p[1]	BMI<25 kg/m^2		BMI>25kg/m^2		p[1]
	Mean	SD	Mean	SD		Mean	SD	Mean	SD		Mean	SD	Mean	SD	
n (n females)	1236 (826)		1177 (735)			852		1561			1322 (958)		1091 (603)		
Age (years)	29.4	5.7	51.5	8.0	<0.001	42.0	13.6	39.2	12.6	<0.001	37.4	13.0	43.5	12.3	<0.001
Height (m)	1.71	0.09	1.70	0.09	0.602	1.79	0.07	1.66	0.07	<0.001	1.70	0.09	1.71	0.09	0.118
Weight (kg)	70.6	14.9	76.8	15.6	<0.001	83.8	13.4	68.1	13.7	<0.001	64.0	9.2	85.3	13.5	<0.001
BMI (kg/m^2)	24.1	4.3	26.4	4.7	<0.001	26.3	4.0	24.7	4.9	<0.001	22.0	1.89	29.2	3.9	<0.001
BMI status*															
Under-weight	3.2%		1.4%			0.5%		3.3%							
Normal Weight	62.8%		41.6%		<0.001[2]	42.3%		58.0%		<0.001[2]	NA				
Overweight	25.2%		37.2%			41.5%		25.4%							
Obesity	8.7%		19.8%			15.7%		13.3%							
Physical activity factor (AU)*	1.53	0.11	1.50	0.09	<0.001	1.54	0.12	1.50	0.08	<0.001	1.53	0.10	1.50	0.10	<0.001
Energy (kcal/d)	2599	808	2606	785	0.617	2888	758	2446	774	<0.001	2481	774	2750	800	<0.001
Total fat (% E)	35.6	6.1	35.7	6.7	0.702	34.8	6.5	36.2	6.3	<0.001	35.4	6.3	36.1	6.5	0.046
SFA (% E)	13.9	3.2	14.0	3.5	0.8019	13.5	3.4	14.2	3.4	<0.001	13.8	3.3	14.1	3.4	0.0168
MUFA (% E)	13.8	3.3	13.7	3.5	0.4697	13.4	3.3	14.0	3.5	<0.001	13.5	3.4	14.0	3.4	0.001
PUFA (% E)	5.5	1.3	5.8	1.5	<0.001	5.5	1.4	5.7	1.4	0.001	5.6	1.4	5.6	1.4	0.5703
Protein (% E)	17.3	3.7	16.9	3.5	0.0205	17.2	3.6	17.1	3.7	0.572	16.9	3.5	17.4	3.7	<0.001
Animal Protein (%E)	10.5	4.3	10.1	4.3	0.010	10.4	4.3	10.3	4.3	0.553	9.9	4.2	10.8	4.4	<0.001
Vegetable Protein (%E)	5.0	1.6	5.1	1.7	0.616	5.1	1.8	5.0	1.6	0.098	5.2	1.7	4.9	1.7	<0.001
Total carbohydrates (% E)	46.7	7.9	46.1	8.6	0.0531	46.1	8.4	46.5	8.1	0.359	47.1	8.1	45.5	8.3	<0.001
Simple sugars (% E)	21.5	6.1	21.1	6.6	0.1457	20.2	5.9	21.9	6.5	<0.001	21.8	6.2	20.7	6.5	<0.001
Dietary fibre (g/day)	29.7	13.1	32.0	13.5	<0.001	32.5	13.9	29.9	12.9	<0.001	31.1	13.8	30.5	12.8	0.300

Values are expressed as mean and SD (Standard Deviation).

BMI = Body Mass Index; AU = Arbitrary Units; SFA = Saturated fatty acids; MUFA = Monounsaturated fatty acids; PUFA = Polyunsaturated fatty acids; %E = % of total energy intake; NA=Not applicable. [1]Differences between groups analysed by one way ANOVA. [2] Chi-square p-value for distribution.

* BMI Categories: Underweight (< 18.5 kg/m2); Normal weight (18.5 – 24.99 kg/m2); Overweight (25 – 29.9 kg/m2); Obesity (> 30 kg/m2).

FIGURE TITLES AND FOOTNOTES

Figure 1. Flow-chart for the participants in the online Food4Me screening included in the present study.

Figure 2. Contribution of different Food Groups to protein intake in each EU Region (a), and divided by BMI (<25 kg/m^2 *vs* >25 kg/m^2) (b).

[Footnote] Protein intake differed between regions and BMI in meat, fish and eggs ($p<0.001$); cereal, grain, pasta and rice ($p<0.001$), as well as in dairy products and fruits and vegetables ($p<0.001$ in both comparisons).

Figure 3. Prevalence Risk Ratio (PRR) of overweight or obesity in Food4Me screenees.

[Footnote] Calculation of PRR was performed according to the isoenergetic substitution (1 %E) of macronutrients by vegetable protein, following the nutrient-density model (n=2413).

Figure 4. Differences in BMI by tertiles of animal protein and vegetable protein intake, in the Food4Me Screenees

[Footnote] Different superscript letters represent significant mean differences between tertiles ($p<0.05$) in Tukey post-hoc analysis.