

# *Up and away: ontogenic transference as a pathway for aerial dispersal of microplastics*

Article

Accepted Version

Al-Jaibachi, Rana, Cuthbert, Ross N. and Callaghan, Amanda (2018) Up and away: ontogenic transference as a pathway for aerial dispersal of microplastics. *Biology Letters*, 14 (9). 20180479. ISSN 1744-957X doi: <https://doi.org/10.1098/rsbl.2018.0479> Available at <https://centaur.reading.ac.uk/78783/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

To link to this article DOI: <http://dx.doi.org/10.1098/rsbl.2018.0479>

Publisher: The Royal Society

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

[www.reading.ac.uk/centaur](http://www.reading.ac.uk/centaur)

**CentAUR**

Central Archive at the University of Reading

Reading's research outputs online

**BIOLOGY  
LETTERS**

**Up and away: ontogenic transference as a pathway for  
aerial dispersal of microplastics**

Journal:	<i>Biology Letters</i>
Manuscript ID	RSBL-2018-0479.R1
Article Type:	Research
Date Submitted by the Author:	21-Aug-2018
Complete List of Authors:	Aljaibachi, Rana; University of Reading, School of Biological Sciences Cuthbert, Ross; Queen's University Belfast, Biological Sciences Callaghan, Amanda; University of Reading, School of Biological Sciences
Subject:	Ecology < BIOLOGY, Environmental Science < BIOLOGY
Categories:	Community Ecology
Keywords:	Food chain, Culex pipiens, ontogeny, Malpighian tubules, microplastics

SCHOLARONE™  
Manuscripts

1 **Up and away: ontogenic transference as a pathway for aerial dispersal of**  
2 **microplastics**

3 Rana Al-Jaibachi<sup>1</sup>, Ross N. Cuthbert<sup>1,2</sup>, Amanda Callaghan<sup>1</sup>

4

5 <sup>1</sup>*Ecology and Evolutionary Biology, School of Biological Sciences, University of Reading,*  
6 *Harborne Building, Reading RG6 6AS, England*

7 <sup>2</sup>*Institute for Global Food Security, School of Biological Sciences, Medical Biology Centre,*  
8 *Queen's University Belfast, Belfast BT9 7BL, Northern Ireland*

9

10 \*Corresponding author: e-mail, [a.callaghan@reading.ac.uk](mailto:a.callaghan@reading.ac.uk)

11

12

13

14

15

16

17

18

19

20

21

22

23

24

25

26

27

28 **Abstract**

29 Microplastics (MPs) are ubiquitous pollutants found in marine, freshwater and terrestrial  
30 ecosystems. With so many MPs in aquatic systems it is inevitable that they will be ingested  
31 by aquatic organisms, and be transferred up through the food chain. However, to date, no  
32 study has considered whether MPs can be transmitted by means of ontogenic transference i.e.  
33 between life stages that utilise different habitats. Here, we determine whether fluorescent  
34 polystyrene beads could transfer between *Culex* mosquito life stages and, particularly, could  
35 move into the flying adult stage. We show for the first time that MPs can be transferred  
36 ontogenically from a feeding (larva) into a non-feeding (pupa) life stage and subsequently  
37 into the adult terrestrial life stage. However, transference is dependent on particle size, with  
38 smaller 2µm MPs transferring readily into pupae and adult stages, whilst 15µm MPs  
39 transferred at a significantly reduced rate. Microplastics appear to accumulate in the  
40 Malpighian tubule renal excretion system. The transfer of MPs to the adults represents a  
41 potential aerial pathway to contamination of new environments. Thus, any organism that  
42 feeds on terrestrial life phases of freshwater insects could be impacted by MPs found in  
43 aquatic ecosystems.

44

45 **Keywords**

46 Food chain: ontology; life stage; Malpighian tubules, microplastics; *Culex pipiens*

47

48

49

50

51

52

53

54 **Introduction**

55 Microplastics (MPs) are ubiquitous pollutants found in marine, freshwater and terrestrial  
56 ecosystems [1–3]. There is little doubt that plastic and MP pollution is a major  
57 environmental concern globally. Despite this, there is relatively little research into the impact  
58 of MPs on freshwater ecosystems, with most research concentrating on marine systems and  
59 organisms [2]. MPs have been defined as plastic particles smaller than 5mm in size [4,5].  
60 However, this simple description covers a wide range of types, including, among others,  
61 polypropylene, polyethylene and polystyrene MPs entering the environment in different  
62 shapes and sizes, including fibres, pellets and cosmetic beads [6,7]. MPs are categorised  
63 based on their origin as primary or secondary types, depending on whether they were  
64 released into the environment as MPs (primary) or have degraded to that size in the  
65 environment (secondary) [8,9]. Microplastics pass through terrestrial environments in  
66 household wastewater [2,10]. Rivers can subsequently deliver MPs into the sea and lakes,  
67 where they can be found in high concentrations [11–13].

68

69 Microplastics are ingested by aquatic organisms, and can be transferred through the food  
70 chain in both freshwater and marine environments [14–18]. However, to date no study has  
71 considered whether MPs can be transmitted by means of ontogenic transference i.e. between  
72 life stages that utilise different habitats. Freshwater environments are inhabited by insects that  
73 spend their juvenile stages in water but their adult stages in the terrestrial environment. Such  
74 insects include mayflies, dragonflies, midges and mosquitoes, most of which are eaten by  
75 terrestrial vertebrates. This raises the potential for MPs to enter terrestrial ecosystems from  
76 freshwater habitats aerially *via* transference to adult invertebrate life stages. Here, we thus  
77 determine whether 2 and 15µm fluorescent polystyrene beads could transfer between insect

78 life stages and, particularly, could move into the flying adult stage. Fluorescent beads were  
79 selected to enable MPs to be easily detected in the non-feeding stages and also to allow an  
80 investigation of location within the body during metamorphosis. The *Culex pipiens* mosquito  
81 complex was selected as a model for this study given their worldwide distribution and broad  
82 habitat preference [19]. Mosquitoes develop through four feeding larval instars and a non-  
83 feeding pupal stage, and finally emerge into a flying adult.

84

### 85 **Materials and methods**

86 For additional details of all methods and analyses, see the electronic supplementary material.

87 Two types of MPs were used: a 2µm fluorescent yellow-green carboxylate-modified  
88 polystyrene (density 1.050g/cm<sup>3</sup>, excitation 470nm; emission 505nm, Sigma-Aldrich, UK)  
89 and a 15.45±1.1µm fluorescent dragon green polystyrene (density 1.06 g/cm<sup>3</sup> (5x10<sup>6</sup>  
90 particles/ml, excitation 480nm; emission 520nm, Bangs Laboratories Inc., USA). Four  
91 treatments were used; a control with no microplastics, a treatment of 8x10<sup>5</sup> 2µm particles/ml,  
92 a treatment of 8x10<sup>2</sup> 15µm particles/ml, and a 1:1 mixture of both treatments. Each replicate  
93 (five per treatment) contained ten 3<sup>rd</sup> instar *C. pipiens* larvae in a 50ml glass beaker filled  
94 with 50ml of tap water. The control and all treatments contained 100mg of pelleted guinea  
95 pig food. Treatments were assigned randomly to a position on the laboratory bench to reduce  
96 experimental error.

97 One random individual was removed from each beaker when every mosquito had moulted  
98 into the 4<sup>th</sup> instar, and again when they pupated or emerged as adults. All samples were then  
99 placed in separate 1.5ml Eppendorf tubes and stored at -20 °C prior to examination.

100 Microplastics were extracted from mosquitoes by homogenization and filtration. The filter  
101 membrane was examined using an epi-fluorescent microscope (Zeiss Axioskop) under a 20x  
102 lens to count the number of fluorescent MPs. Adults were further dissected under a binocular

103 stereo microscope (0.7X-4.5X) to extract the gut and quantify the numbers of MPs under the  
104 epi-fluorescent microscope [20].

105 All data were analyzed using the statistical software R v3.4.2 [21]. Microplastic counts were  
106 analysed using generalized linear models (GLMs) assuming a quasi-Poisson distribution.

107 Uptake of microplastics was examined with respect to ‘particle size’, ‘treatment’ and ‘life  
108 stage’. We performed model simplification via stepwise removal of non-significant effects.

109 Tukey tests were used post hoc for multiple comparisons.

110

## 111 **Results**

112 No MPs were found in control groups of any mosquito life stage. Densities of MPs were  
113 significantly different between life stages ( $F_{2, 56}=160.42$ ,  $P<0.001$ ), with MP numbers  
114 significantly falling as mosquitoes moved between successive ontogenic levels (all  $P<0.001$ )  
115 (Figure 1, Table S1, S2). Microplastic transference to adults was confirmed by fluorescent  
116 microscopy where the beads were detected in the adult abdomen, specifically inside the  
117 Malpighian tubules (Figure 2).

118 Significantly more 2 $\mu$ m particles were found in mosquito life stages than 15 $\mu$ m particles  
119 overall ( $F_{1, 58}=303.98$ ,  $P<0.001$ ). Microplastics uptake was also significantly greater overall  
120 in mixed exposure treatments ( $F_{1, 55}=6.00$ ,  $P=0.02$ ). Although 2 $\mu$ m particles were transferred  
121 to adults in all instances, we found no transference of 15 $\mu$ m particles following single  
122 treatment exposures. However, in the mixed MPs treatment, transference to adults of both  
123 2 $\mu$ m and 15 $\mu$ m particles was evidenced (Figure 1).

124

## 125 **Discussion**

126 Here, we show for the first time that MPs can be transferred ontogenically from a feeding  
127 (larval) into a non-feeding (pupal) life stage and subsequently into the flying (adult) life



128 stage. Transference through to adults was found in both MP sizes, although the larger 15µm  
129 MPs were not ingested as readily as the 2µm MPs. Dissection of mosquito adults showed that  
130 2µm MPs accumulated in the renal excretion system of Malpighian tubules which, unlike the  
131 gut, pass from larvae to adult stages without visible reorganization [22]. This has been  
132 demonstrated previously to provide a physical transport system between stages during  
133 metamorphosis for *Pseudomonas* bacteria and seems to be important for ontogenic  
134 transmission from larvae to adults [23].

135 Few 15µm MPs were transferred into adults suggesting that MP size is an important factor in  
136 ontogenic transfer which could be related to the transfer and accumulation of MPs in the  
137 Malpighian tubes. Although the translocation mechanism of MPs to the Malpighian tubules  
138 is unclear in mosquitoes, analysis of fish, fiddler crab and marine mussels has demonstrated  
139 that MPs can be translocated from gastrointestinal tracts into other tissues in a wide range of  
140 phyla [24, 25,26]. Malpighian tubules have an entry point to the gut between the mid- and  
141 hindgut of mosquitoes, but the flow of fluid is from the Malpighian tubules to the hindgut  
142 [27]. Diptera are known to produce structures called concretions in the Malpighian tubules  
143 which have been shown to sequester heavy metals [28]. However, it is unlikely that this  
144 pathway would operate with a solid MP.

145 Our results have important implications since any aquatic life stage that is able to consume  
146 MPs and transfer them to their terrestrial life stage is a potential vector of MPs onto novel  
147 aerial and terrestrial habitats. Ingestion of MP-contaminated organisms by terrestrial  
148 organisms is not new [29]. Indeed, the widespread distribution of MPs in marine  
149 environments has meant that animals such as fish and shellfish sold for human consumption  
150 are contaminated with a range of plastics with a consequent transference of MPs between  
151 trophic levels [24]. Unlike MP fibres, which are common in the air and atmosphere, there  
152 has been no evidence for MPs being transported into the air [24]. We have demonstrated here

153 that species with aquatic and terrestrial life stages can harbour MPs through their life history.  
154 Adults are predated on emergence by many animals including dipteran flies Empididae and  
155 Dolichopodidae, whilst resting predominantly by spiders and in flight they are the prey of  
156 dragonflies, damselflies, birds (such as swallows and swifts) and bats (31). Where many  
157 insects are emerging from a highly contaminated site, the possibility of contamination of  
158 these predators could be high. Whilst mosquitoes were used here as a model organism, any  
159 freshwater insect that can ingest MPs will likely equally transmit plastics into a terrestrial  
160 adult stage. This has implications for organisms that feed on adult mosquitoes with aerial and  
161 terrestrial animals accordingly open to MP exposure and transference would appear to occur  
162 at a higher rate for smaller MPs.

### 163 **References**

- 164 1. Sighicelli M, Pietrelli L, Lecce F, Iannilli V, Falconieri M, Coscia L, Di Vito S, Nuglio  
165 S, Zampetti G. 2018 Microplastic pollution in the surface waters of Italian Subalpine  
166 Lakes. *Environ. Pollut.* **236**, 645–651. (doi:10.1016/j.envpol.2018.02.008)
- 167 2. Wagner M, Lambert S. 2018 *Freshwater Microplastics*. Cham: Springer International  
168 Publishing. (doi:10.1007/978-3-319-61615-5)
- 169 3. Mason SA, Welch V, Neratko J. 2018 Synthetic polymer contamination in bottled  
170 water. *Fredonia State Univ. New York* , 1–17.
- 171 4. Eriksen M, Lebreton LCM, Carson HS, Thiel M, Moore CJ, Borerro JC, Galgani F,  
172 Ryan PG, Reisser J. 2014 Plastic Pollution in the World's Oceans: More than 5  
173 Trillion Plastic Pieces Weighing over 250,000 Tons Afloat at Sea. *PLoS One* **9**,  
174 e111913. (doi:10.1371/journal.pone.0111913)
- 175 5. Imhof HK, Ivleva NP, Schmid J, Niessner R, Laforsch C. 2013 Contamination of  
176 beach sediments of a subalpine lake with microplastic particles. *Curr. Biol.* **23**, 1–15.  
177 (doi:10.1016/j.cub.2013.09.001)
- 178 6. Andrady AL, Neal MA. 2009 Applications and societal benefits of plastics. *Philos.*

- 179 *Trans. R. Soc. Lond. B. Biol. Sci.* **364**, 1977–1984. (doi:10.1098/rstb.2008.0304)
- 180 7. Rocha-Santos T, Duarte AC. 2014 A critical overview of the analytical approaches to  
181 the occurrence, the fate and the behavior of microplastics in the environment. *Trends*  
182 *Anal. Chem.* **65**, 47–53. (doi:10.1016/j.trac.2014.10.011)
- 183 8. Moore CJ. 2008 Synthetic polymers in the marine environment: A rapidly increasing,  
184 long-term threat. *Environ. Res.* **108**, 131–139. (doi:10.1016/j.envres.2008.07.025)
- 185 9. Barnes DKA, Galgani F, Thompson RC, Barlaz M. 2009 Accumulation and  
186 fragmentation of plastic debris in global environments. *Philos. Trans. R. Soc. Lond. B.*  
187 *Biol. Sci.* **364**, 1985–98. (doi:10.1098/rstb.2008.0205)
- 188 10. Mason SA, Garneau D, Sutton R, Chu Y, Ehmann K, Barnes J, Fink P, Papazissimos  
189 D, Rogers DL. 2016 Microplastic pollution is widely detected in US municipal  
190 wastewater treatment plant effluent. *Environ. Pollut.* **218**, 1045–1054.  
191 (doi:10.1016/j.envpol.2016.08.056)
- 192 11. Eriksen M, Mason S, Wilson S, Box C, Zellers A, Edwards W, Farley H, Amato S.  
193 2013 Microplastic pollution in the surface waters of the Laurentian Great Lakes. *Mar.*  
194 *Pollut. Bull.* **77**, 177–182. (doi:10.1016/j.marpolbul.2013.10.007)
- 195 12. Fischer EK, Paglialonga L, Czech E, Tamminga M. 2016 Microplastic pollution in  
196 lakes and lake shoreline sediments - A case study on Lake Bolsena and Lake Chiusi  
197 (central Italy). *Environ. Pollut.* **213**, 648–657. (doi:10.1016/j.envpol.2016.03.012)
- 198 13. Su L, Xue Y, Li L, Yang D, Kolandhasamy P, Li D, Shi H. 2016 Microplastics in  
199 Taihu Lake, China. *Environ. Pollut.* **216**, 711–719. (doi:10.1016/j.envpol.2016.06.036)
- 200 14. Aljaibachi R, Callaghan A. 2018 Impact of polystyrene microplastics on *Daphnia*  
201 *magna* mortality and reproduction in relation to food availability. *PeerJ* **6**, e4601.  
202 (doi:10.7717/peerj.4601)
- 203 15. Cole M, Lindeque P, Fileman E, Halsband C, Goodhead R, Moger J, Galloway TS.

- 204 2013 Microplastic ingestion by zooplankton. *Environ. Sci. Technol.* **47**, 6646–6655.  
205 (doi:10.1021/es400663f)
- 206 16. Scherer C, Brennholt N, Reifferscheid G, Wagner M. 2017 Feeding type and  
207 development drive the ingestion of microplastics by freshwater invertebrates. *Sci. Rep.*  
208 **7**, 17006. (doi:10.1038/s41598-017-17191-7)
- 209 17. Sussarellu R *et al.* 2016 Oyster reproduction is affected by exposure to polystyrene  
210 microplastics. *Proc. Natl. Acad. Sci.* **113**, 2430–2435. (doi:10.1073/pnas.1519019113)
- 211 18. Messinetti S, Mercurio S, Parolini M, Sugni M, Pennati R. 2018 Effects of polystyrene  
212 microplastics on early stages of two marine invertebrates with different feeding  
213 strategies. *Environ. Pollut.* **237**, 1080–1087. (doi:10.1016/j.envpol.2017.11.030)
- 214 19. Dow JA, Maddrell SH, Görtz A, Skaer NJ, Brogan S, Kaiser K. 1994 The malpighian  
215 tubules of *Drosophila melanogaster*: a novel phenotype for studies of fluid secretion  
216 and its control. *J. Exp. Biol.* **197**, 421–8.
- 217 20. Coleman J, Juhn J, James AAA. 2007 Dissection of Midgut and Salivary Glands from  
218 *Ae. aegypti* Mosquitoes. *J. Vis. Exp.* , 2007. (doi:10.3791/228)
- 219 21. R Development Core Team. 2017 R: A Language and Environment for Statistical  
220 Computing. *R Found. Stat. Comput. Vienna Austria* (doi:10.1038/sj.hdy.6800737)
- 221 22. Clements AN. 1992 *The biology of mosquitoes. Volume 1: Development, nutrition and*  
222 *reproduction*. London: Chapman and Hall.
- 223 23. Chavshin AR, Oshaghi MA, Vatandoost H, Yakhchali B, Zarenejad F, Terenius O.  
224 2015 Malpighian tubules are important determinants of *Pseudomonas* transstadial  
225 transmission and longtime persistence in *Anopheles stephensi*. *Parasites and Vectors*  
226 **8**, 1–7. (doi:10.1186/s13071-015-0635-6)
- 227 24. Von Moos N, Burkhardt-Holm P, Köhler A. 2012 Uptake and Effects of Microplastics  
228 on Cells and Tissue of the Blue Mussel *Mytilus edulis* L. after an Experimental

- 229 Exposure. *Environ. Sci. Technol.* **46**, 11327–11335.
- 230 25. Brennecke D, Ferreira EC, Costa TMM, Appel D, da Gama BAP, Lenz M. 2015  
231 Ingested microplastics (>100µm) are translocated to organs of the tropical fiddler crab  
232 *Uca rapax*. *Mar. Pollut. Bull.* **96**, 491–495. (doi:10.1016/j.marpolbul.2015.05.001)
- 233 26. Avio CG, Gorbi S, Regoli F. 2015 Experimental development of a new protocol for  
234 extraction and characterization of microplastics in fish tissues: First observations in  
235 commercial species from Adriatic Sea. *Mar. Environ. Res.* **111**, 18–26.  
236 (doi:10.1016/j.marenvres.2015.06.014)
- 237 27. Piermarini PM. 2016 *Renal Excretory Processes in Mosquitoes*. 1st edn. Elsevier Ltd.  
238 (doi:10.1016/bs.aiip.2016.04.003)
- 239 28. Leonard EM, Pierce LM, Gillis PL, Wood CM, O'Donnell MJ. 2009 Cadmium  
240 transport by the gut and Malpighian tubules of *Chironomus riparius*. *Aquat. Toxicol.*  
241 **92**, 179–186. (doi:10.1016/j.aquatox.2009.01.011)
- 242 29. Huerta Lwanga E, Gertsen H, Gooren H, Peters P, Salánki T, Van Der Ploeg M,  
243 Besseling E, Koelmans AA, Geissen V. 2016 Microplastics in the Terrestrial  
244 Ecosystem: Implications for *Lumbricus terrestris* (Oligochaeta, Lumbricidae).  
245 *Environ. Sci. Technol.* (doi:10.1021/acs.est.5b05478)
- 246 30. Dris R, Gasperi J, Mirande C, Mandin C, Guerrouache M, Langlois V, Tassin B. 2017  
247 A first overview of textile fibers, including microplastics, in indoor and outdoor  
248 environments. *Environ. Pollut.* (doi:10.1016/j.envpol.2016.12.013)
- 249 31. Medlock JM, Snow, KR. 2008. Natural predators and parasites of British mosquitoes:  
250 a review. *European Mosquito Bulletin*, 25 (2008), 1-11.

251

252 **Figure legends**

253 **Figure 1.** Uptake counts of microplastics (MP) across larval (a, b), pupal (c, d) and adult (e,  
254 f) *Culex* mosquito stages following single (a, c, e) and mixed (b, d, f) exposures to 2µm and  
255 15µm beads. Means are ±SE ( $n=5$  per experimental group).

256

257 **Figure 2.** Epi-fluorescent microscope images showing fluorescent microplastic particles  
258 within (A) the abdomen of an adult mosquito before dissection, and (B) the abdominal  
259 Malpighian tubules following dissection.

260

## 261 **Ethics**

262 Ethics committee approval was not required.

## 263 **Data accessibility**

264 Data files are available in online supplementary material.

## 265 **Author contribution**

266 All authors provided substantial contributions to conception and design, or acquisition of  
267 data, or analysis and interpretation of data; were involved in drafting the article or revising it  
268 critically for important intellectual content; approved the final version to be published; and  
269 agree to be accountable for all aspects of the work in ensuring that questions related to the  
270 accuracy or integrity of any part of the work are appropriately investigated and resolved.

271

## 272 **Competing interests**

273 We declare we have no competing interests.


## 274 **Funding**

275 A.C. is funded by the University of Reading. R.A. is self-funded and R.N.C. is funded  
276 through the Department for the Economy, Northern Ireland.

## 277 **Acknowledgements**

278 We thank Natali Ortiz-Perea for assisting with mosquito colony rearing.

279



Uptake counts of microplastics (MP) across larval (a, b), pupal (c, d) and adult (e, f) *Culex* mosquito stages following single (a, c, e) and mixed (b, d, f) exposures to 2µm and 15µm beads. Means are  $\pm$ SE (n=5 per experimental group).

115x144mm (300 x 300 DPI)

