

Self-assembled micellar structures of Lipopeptides with variable number of attached lipid chains revealed by atomistic molecular dynamics simulations

Article

Supplemental Material

Zhao, L., Tu, Y., Fang, H., Hamley, I. W. and Wang, Z. (2018) Self-assembled micellar structures of Lipopeptides with variable number of attached lipid chains revealed by atomistic molecular dynamics simulations. *Journal of Physical Chemistry B*, 122 (41). pp. 9605-9615. ISSN 1520-6106 doi: <https://doi.org/10.1021/acs.jpcc.8b07877> Available at <https://centaur.reading.ac.uk/79751/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

To link to this article DOI: <http://dx.doi.org/10.1021/acs.jpcc.8b07877>

Publisher: American Chemical Society

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online

Supporting Information

Self-Assembled Micellar Structures of Lipopeptides with Variable Number of Attached Lipid Chains Revealed by Atomistic Molecular Dynamics Simulations

Liang Zhao[†], Yusong Tu^{*†}, Haiping Fang[‡], Ian Hamley[§] and Zuowei Wang^{*¶}

[†]College of Physical Science and Technology, Yangzhou University, Jiangsu, 225009, China

[‡]Division of Interfacial Water and Key Laboratory of Interfacial Physics and Technology, Shanghai Institute of Applied Physics,
Chinese Academy of Sciences, Shanghai 201800, China

[§]Department of Chemistry, University of Reading, Whiteknights, Reading, RG6 6AD, United Kingdom

[¶]School of Mathematical, Physical and Computational Sciences, University of Reading, Whiteknights, Reading RG6 6AX,
United Kingdom

*Corresponding Author.

E-mail: ystu@yzu.edu.cn, zuowei.wang@reading.ac.uk

Table of Contents

PS 1. Definitions of Moment of Inertia Tensor, Relative Anisotropy Shape and Body Reference Frame

PS 2. Comparison of Atomic Density Functions of Different Lipid Chains in the Same Lipopeptide Molecule

PS 3. Probability Distributions of the Length of Lipid Chains in Lipopeptide Molecules

PS 4. Probability Distribution of Angles (Including Pam₃CSK4 Molecules at the Curved Side Edges of Flattened Wormlike Micelle)

PS1. Definitions of Moment of Inertia Tensor, Relative Anisotropy Shape and Body Reference Frame

The moment of inertia tensor of a self-assembled structure, consisting of n atoms with masses m_i , is defined by a matrix I

$$I = \sum_{i=1}^n \begin{bmatrix} m_i(y_i^2 + z_i^2) & -m_i x_i y_i & -m_i x_i z_i \\ -m_i x_i y_i & m_i(x_i^2 + z_i^2) & -m_i y_i z_i \\ -m_i x_i z_i & -m_i y_i z_i & m_i(x_i^2 + y_i^2) \end{bmatrix}, \quad (1)$$

where x_i , y_i and z_i are the atomic coordinates with respect to the center of mass (COM) of a self-assembled structure. The eigenvalues of I are denoted by I_{xx} , I_{yy} and I_{zz} .

The relative shape anisotropy K^2 is of the form

$$K^2 = \frac{3}{2} \frac{\lambda_1^2 + \lambda_2^2 + \lambda_3^2}{(\lambda_1 + \lambda_2 + \lambda_3)^2} - \frac{1}{2}, \quad (2)$$

where λ_1 , λ_2 and λ_3 are the recombination of eigenvalues of I , given by the following formulae

$$\lambda_1 = \frac{1}{2}(I_{yy} + I_{zz} - I_{xx})$$

$$\lambda_2 = \frac{1}{2}(I_{xx} + I_{zz} - I_{yy})$$

$$\lambda_3 = \frac{1}{2}(I_{xx} + I_{yy} - I_{zz}).$$

The body reference frame is identified by the principal axes obtained by diagonalizing the moment of inertia tensor I .

PS2. Comparison of Atomic Density Functions of Different Lipid Chains in the Same Lipopeptide Molecule

To further compare the ADFs of different lipid chains in the same Pam₂CSK4 and Pam₃CSK4 molecules, we plot the ADFs of selected components of Pam^(1, 2) for Pam₂CSK4 and Pam⁽¹⁻³⁾ for Pam₃CSK4 in Figure S1. Figure S1a shows that the ADFs of “CT”, “CM” and “CH” in Pam^(1, 2) of Pam₂CSK4 almost overlap with each other, an indication of statistical equivalency of the two lipid chains in the same molecule. Similar distributions of the three lipid chains Pam⁽¹⁻³⁾ in Pam₃CSK4 can also be seen in Figure S1b.




Figure S1. Atomic density functions (ADFs) of selected components of (a) the lipid chains Pam^(1, 2) in Pam₂CSK4 and (b) Pam⁽¹⁻³⁾ in Pam₃CSK4. “CT”, “CM” and “CH” denote the head and middle methylene groups and terminal methyl group on a lipid chain, respectively.

PS 3. Probability Distributions of the Length of Lipid Chains in Lipopeptide Molecules




Figure S2. Probability distributions of the length of lipid chains of (a) PamCSK4, (b) Pam₂CSK4 and (c) Pam₃CSK4.

The narrow and sharp distributions of lipid chains on lipopeptides indicates that the lipid chains are straight and their lengths are of about 1.5 nm.

PS 4. Probability Distribution of Angles (Including Pam₃CSK4 Molecules at the Curved Side Edges of Flattened Wormlike Micelle)




Figure S3. Probability distribution of the angle (a) between the vector of lipid chains and the normal of flattened wormlike micelle of Pam₃CSK4, and (b) between any two lipid chains on the same molecule.

When the Pam₃CSK4 molecules at the edge of the flattened wormlike micelle are considered, peaks around 90° appear in the Figure S3a and two peaks around 20° and 160° still exist. This indicates that molecules at the edge are almost distributed vertical to the normal of a flattened wormlike micelle, and the inclusion of these molecules has little impact on the recognition of bilayer stacking structures inner the flattened wormlike micelle.