

They say coconut oil can aid weight loss, but can it really?

Article

Accepted Version

Clegg, M. E. (2017) They say coconut oil can aid weight loss, but can it really? European Journal of Clinical Nutrition, 71 (10). pp. 1139-1143. ISSN 0954-3007 doi: 10.1038/ejcn.2017.86 Available at <https://centaur.reading.ac.uk/80389/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

To link to this article DOI: <http://dx.doi.org/10.1038/ejcn.2017.86>

Publisher: Nature Publishing Group

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online

1 They say coconut oil can aid weight loss, but can it really?

2
3 **Miriam E Clegg**

4
5 Functional Food Centre, Department of Sport and Health Sciences, Faculty of Health and
6 Life Sciences, Oxford Brookes University, Gypsy Lane, Oxford OX3 0BP, UK

7
8 *Corresponding author: Miriam Clegg, Functional Food Centre, Department of Sport and
9 Health Sciences, Faculty of Health and Life Sciences, Oxford Brookes University, Gypsy
10 Lane, Oxford OX3 0BP, UK

11 Email: mclegg@brookes.ac.uk; Ph: +44 1865 484365

12
13
14 **Running title:** Coconut oil for weight loss

15
16 **Keywords:** Coconut oil, MCT oil, satiety, energy expenditure, body weight

17
18 **Conflict of interest.**

19 The author declares no conflict of interest.

Abstract

There has in recent years, been much media speculation and consumer interest in the beneficial satiating properties of consuming coconut oil and it's potential to aid weight loss. However the media has primarily cited studies using MCT oil. The current perspective looks at the research that is available on coconut oil. It examines if and how MCT related research can be applied to coconut oil and if there is potential for coconut oil to aid weight loss. The current report indicates a lack of consistent evidence on the topic of coconut oil, satiety and weight loss. Given both the publicity and the increased consumption of coconut oil further research, particularly long term clinical trials, in this area are warranted.

Coconut oil has gained considerable popularity in recent years with coconut oil in food and beverages accounting for 26% of food and drink new product launches in 2012, this is an increase from 15% in 2008 (1). Coconut oil exports across Asia have also grown 3.3% annually over the past five years, according to the Asian Pacific Coconut Community (2). This is not surprising given the numerous ways that coconut oil has been identified by the media, to potentially improve our lives. Articles are wide ranging promoting adding it to stir-fries, baking with it and even adding it to coffee (3, 4). Many media articles promote the consumption of coconut oil for weight loss, advocating similar health benefits to that of medium chain triglycerides (MCT). This has contributed to an increase in intake of coconut oil in recent years (5). Coconut oil is said to aid weight loss through a combination of increased energy expenditure and satiety induced by MCT. MCT are dietary triglycerides with fatty acids chains that are 6 to 12 carbon atoms in length (6). MCT have a smaller molecular weight than long chain triglycerides. This allows them to be more rapidly and completely hydrolysed, and absorbed when there are decreased intraluminal concentrations of pancreatic enzymes and bile salts (7). During digestion, MCT are converted to medium-chain fatty acids (MCFA) and transported directly in the portal venous system to the liver, as opposed to being transported as chylomicrons in the lymphatic system like LCT (8). MCT therefore bypass peripheral tissues, including adipose tissue, which makes them less likely to be deposited into the adipose tissue via the actions of hormone-sensitive lipase (6). Finally, MCFA can cross the mitochondrial membrane of the liver and muscle independently of the acylcarnitine transfer system, this makes them a more readily available energy source that is likely to be utilised more rapidly (9).

It has been proposed that MCT can affect satiety via a number of mechanisms; however a lot is still unknown. Potential mechanisms include the production of ketones due to the increased acetyl-CoA influx which is necessary to oxidize fatty acids (10). Furthermore, Van Wymelbeke et al (11) and Rolls et al (12) indicate that the increase in satiety maybe due to the rapid rate of absorption of MCT. Where LCT result in two peaks during absorption; the

initial peak at the point of ingestion and a second delayed peak at the beginning of the next meal, MCT are fully absorbed at the point of ingestion (13). Hence, MCT may contribute to satiation due to complete absorption mechanism.

The evidence for both increases in diet induced thermogenesis (14-24) and reduced food intake (11, 12, 25-29) following the consumption of MCT has been well documented, with interventions using MCT oil indicating that there is potential for it to help aid weight loss (10, 30-33) (Table 1). However it needs to be emphasised that MCT oil and coconut oil are not the same thing. Lauric acid (carbon chain length 12) is found in much larger quantities in coconut oil, making up 47.7 % of the total fat, where no lauric acid is found in MCT oil. Other MCFA in coconut oil are capric acid (C10– 5.5%), caprylic acid (C8-7.6%) and caproic acid (C6 – 0.52%) (34). There is some debate as to whether lauric acid is a MCT or not and this is demonstrated in how it is utilized in the body. Unlike with pure MCT oil containing fatty acids of shorter carbon length (C6-C10), only twenty to thirty percent of lauric acid is taken directly to the liver to be used as energy via the portal vein (35). This means that in total only ~23.16% of the coconut oil contains MCTs that is absorbed and metabolised in the same way as pure MCT oil.

Studies on satiety and MCT have shown that 3g is not sufficient to have an effect on satiety (26, 27). However this is in contrast to a study by Rolls et al (12) that showed a dose as low as 2.9g (100kcal containing 24% MCT) reduced food intake in dieters but had no effect in non-dieters. This obviously has practical implications as it is dieters that are most likely to want to see the satiating effects. The other studies that have shown an effect have used much higher doses of ~25g (11, 25, 29). Studies looking at energy expenditure have shown doses of 5g have the ability to increase postprandial thermogenesis (18). However, similar to satiety the majority of other studies tended to use much larger doses of greater than 20g (17, 21, 36). For weight loss, similarly doses of 5g of MCT for 12 week resulted in significant decreases in body fat, subcutaneous and visceral fat (31).

113

114 This shows that doses as low as 5g and perhaps 3g may have an effect on satiety and body
115 weight; however the majority of the research has used MCT amounts much higher than this.
116 Nonetheless, if a dose as low of 5g MCT per day is sufficient to have beneficial effects on
117 weight management then 21.6g of coconut oil would be required to obtain sufficient amounts
118 of MCFA. Current UK guidelines limit the intake of saturated to a maximum of just 21g in
119 females and 31g in males (19-64 years) (37). Hence at the lowest dose of coconut oil known
120 to have an effect on body weight, people will have reached or almost reached their total
121 saturated fat intake for the day.

122

123 Two studies examining the effects of coconut oil compared to LCTs reported no increase in
124 satiety and no effect on food intake (38, 39) (Table 2). Poppit et al (39) found no difference in
125 visual analogue scale ratings of satiety or differences in ad libitum food intake at lunch
126 following the consumption of either coconut oil (containing 10g MCT), high short chain
127 triglyceride (3g SCT, 7g MCT) (from soft fraction milk fat) or long chain triglycerides (from
128 tallow). Rizzo et al (38) found that in a dinner meal following ice-cream with varying
129 quantities of coconut oil there was trend towards reduced consumption with the coconut oil,
130 however this was compensated for later when there was a significant increase in snack
131 consumption resulting in no overall difference between the ice-creams. The amounts of
132 coconut oil used here are 7.5g coconut oil (high dose) consisting of only ~4.8 g MCT
133 (carbons 6-12). This study should be commended on giving the coconut in a realistic form
134 (ice-cream) and amount. However compared to the amounts seen in the many of the MCT
135 studies the dose is very low. The authors conclude that the differences and trends observed
136 may reflect the differences in the absorption and metabolism of the two ice-creams. A third
137 study published across three papers providing 14 days of coconut oil, found no effect on
138 total energy expenditure or thermic effect of feeding. However they did find an increase in
139 basal metabolic rate after 7 days but not 14 days and an increase in endogenous long chain

saturated fatty acid oxidation after 14 days (19, 20, 24). In this study, caprylic and capric acid made up 7.9% of the energy intake of the diet and lauric acid made up 17.7%.

Currently there appears to be a lack of research in this area and to the best of the author's knowledge there are no studies looking at the effect of coconut oil on weight loss and none comparing it to MCT oil. Further work is needed in this area to confirm these preliminary calculations however indications would suggest that the use of coconut oil as a practical means of increasing satiety are not credible. The research available on the use of coconut oil on satiety and energy expenditure is limited and particularly there have been no long term clinical trials looking at the effects on weight loss. Given both the publicity and the increased consumption of coconut oil further research in this area is warranted.

References

1. Mintel. Tropical storm in the food and drink market: launches of coconut water quintuple over the past five years 2013 [Available from: <http://www.mintel.com/press-centre/food-and-drink/launches-of-coconut-water-quintuple-over-the-past-five-years>.
2. Mahr K. Why the Coconut Craze Isn't Helping Farmers 2012 [Available from: <http://business.time.com/2012/12/05/why-the-coconut-craze-isnt-helping-farmers/>.
3. Waters J. Bulletproof coffee: is adding butter to your brew a step too far? 2014 [Available from: <https://www.theguardian.com/lifeandstyle/wordofmouth/2014/nov/25/bulletproof-coffee-is-adding-butter-to-your-morning-coffee-a-step-too-far>.
4. Sacks K. 6 Ways to Cook With Coconut Oil: Epicurious; 2015 [Available from: <http://www.epicurious.com/ingredients/how-to-cook-with-coconut-oil-article>.
5. Lockyer S, Stanner S. Coconut oil - a nutty idea? Nutrition Bulletin. 2016;41:42-54.
6. Bach AC, Babayan VK. Medium-chain triglycerides: an update. The American journal of clinical nutrition. 1982;36(5):950-62.

- 167 7. Fernandes J, van de KJ, Weijers HA. Differences in absorption of the various fatty
168 acids studied in children with steatorrhea. The Journal of clinical investigation. 1962;41:488-
169 94.
- 170 8. Bloom B, Chaikoff IL, Reinhardt. Intestinal lymph as pathway for transport of
171 absorbed fatty acids of different chain lengths. Am J Physiol. 1951;166(2):451-5.
- 172 9. Williamson JR, Browning ET, Scholz R, Kreisberg RA, Fritz IB. Inhibition of fatty acid
173 stimulation of gluconeogenesis by (+)-decanoylcarnitine in perfused rat liver. Diabetes.
174 1968;17(4):194-208.
- 175 10. Tsuji H, Kasai M, Takeuchi H, Nakamura M, Okazaki M, Kondo K. Dietary medium-
176 chain triacylglycerols suppress accumulation of body fat in a double-blind, controlled trial in
177 healthy men and women. J Nutr. 2001;131(11):2853-9.
- 178 11. Van Wymelbeke V, Louis-Sylvestre J, Fantino M. Substrate oxidation and control of
179 food intake in men after a fat-substitute meal compared with meals supplemented with an
180 isoenergetic load of carbohydrate, long-chain triacylglycerols, or medium-chain
181 triacylglycerols. The American journal of clinical nutrition. 2001;74(5):620-30.
- 182 12. Rolls BJ, Gnizak N, Summerfelt A, Laster LJ. Food intake in dieters and nondieters
183 after a liquid meal containing medium-chain triglycerides. The American journal of clinical
184 nutrition. 1988;48(1):66-71.
- 185 13. Fielding BA, Callow J, Owen RM, Samra JS, Matthews DR, Frayn KN. Postprandial
186 lipemia: the origin of an early peak studied by specific dietary fatty acid intake during
187 sequential meals. The American journal of clinical nutrition. 1996;63(1):36-41.
- 188 14. Alexandrou E, Herzberg GR, White MD. High-level medium-chain triglyceride feeding
189 and energy expenditure in normal-weight women. Can J Physiol Pharmacol. 2007;85(5):507-
190 13.
- 191 15. Binnert C, Pachiaudi C, Beylot M, Hans D, Vandermander J, Chantre P, et al.
192 Influence of human obesity on the metabolic fate of dietary long- and medium-chain
193 triacylglycerols. The American journal of clinical nutrition. 1998;67(4):595-601.

194 16. Dulloo AG, Fathi M, Mensi N, Girardier L. Twenty-four-hour energy expenditure and
195 urinary catecholamines of humans consuming low-to-moderate amounts of medium-chain
196 triglycerides: a dose-response study in a human respiratory chamber. *European journal of*
197 *clinical nutrition*. 1996;50(3):152-8.

198 17. Flatt JP, Ravussin E, Acheson KJ, Jequier E. Effects of dietary fat on postprandial
199 substrate oxidation and on carbohydrate and fat balances. *The Journal of clinical*
200 *investigation*. 1985;76(3):1019-24.

201 18. Kasai M, Nosaka N, Maki H, Suzuki Y, Takeuchi H, Aoyama T, et al. Comparison of
202 diet-induced thermogenesis of foods containing medium- versus long-chain triacylglycerols.
203 *J Nutr Sci Vitaminol (Tokyo)*. 2002;48(6):536-40.

204 19. Papamandjaris AA, White MD, Raeini-Sarjaz M, Jones PJ. Endogenous fat oxidation
205 during medium chain versus long chain triglyceride feeding in healthy women. *Int J Obes*
206 *Relat Metab Disord*. 2000;24(9):1158-66.

207 20. Papamandjaris AA, White MD, Jones PJ. Components of total energy expenditure in
208 healthy young women are not affected after 14 days of feeding with medium-versus long-
209 chain triglycerides. *Obesity research*. 1999;7(3):273-80.

210 21. Seaton TB, Welle SL, Warenko MK, Campbell RG. Thermic effect of medium-chain
211 and long-chain triglycerides in man. *The American journal of clinical nutrition*.
212 1986;44(5):630-4.

213 22. St-Onge MP, Ross R, Parsons WD, Jones PJ. Medium-chain triglycerides increase
214 energy expenditure and decrease adiposity in overweight men. *Obesity research*.
215 2003;11(3):395-402.

216 23. St-Onge MP, Bourque C, Jones PJ, Ross R, Parsons WE. Medium- versus long-
217 chain triglycerides for 27 days increases fat oxidation and energy expenditure without
218 resulting in changes in body composition in overweight women. *Int J Obes Relat Metab*
219 *Disord*. 2003;27(1):95-102.

220 24. White MD, Papamandjaris AA, Jones PJ. Enhanced postprandial energy expenditure
 221 with medium-chain fatty acid feeding is attenuated after 14 d in premenopausal women. The
 222 American journal of clinical nutrition. 1999;69(5):883-9.

223 25. Coleman H, Quinn P, Clegg ME. Medium-chain triglycerides and conjugated linoleic
 224 acids in beverage form increase satiety and reduce food intake in humans. Nutrition
 225 research (New York, NY. 2016;36(6):526-33.

226 26. Kovacs EM, Westerterp-Plantenga MS, de Vries M, Brouns F, Saris WH. Effects of 2-
 227 week ingestion of (-)-hydroxycitrate and (-)-hydroxycitrate combined with medium-chain
 228 triglycerides on satiety and food intake. Physiol Behav. 2001;74(4-5):543-9.

229 27. Kovacs EM, Westerterp-Plantenga MS, Saris WH. The effects of 2-week ingestion of
 230 (--)hydroxycitrate and (--)hydroxycitrate combined with medium-chain triglycerides on
 231 satiety, fat oxidation, energy expenditure and body weight. Int J Obes Relat Metab Disord.
 232 2001;25(7):1087-94.

233 28. Stubbs RJ, Harbron CG. Covert manipulation of the ratio of medium- to long-chain
 234 triglycerides in isoenergetically dense diets: effect on food intake in ad libitum feeding men.
 235 Int J Obes Relat Metab Disord. 1996;20(5):435-44.

236 29. Van Wymelbeke V, Himaya A, Louis-Sylvestre J, Fantino M. Influence of medium-
 237 chain and long-chain triacylglycerols on the control of food intake in men. The American
 238 journal of clinical nutrition. 1998;68(2):226-34.

239 30. Han JR, Deng B, Sun J, Chen CG, Corkey BE, Kirkland JL, et al. Effects of dietary
 240 medium-chain triglyceride on weight loss and insulin sensitivity in a group of moderately
 241 overweight free-living type 2 diabetic Chinese subjects. Metabolism: clinical and
 242 experimental. 2007;56(7):985-91.

243 31. Nosaka N, Maki H, Suzuki Y, Haruna H, Ohara A, Kasai M, et al. Effects of
 244 margarine containing medium-chain triacylglycerols on body fat reduction in humans. J
 245 Atheroscler Thromb. 2003;10(5):290-8.

32. St-Onge MP, Bosarge A. Weight-loss diet that includes consumption of medium-chain triacylglycerol oil leads to a greater rate of weight and fat mass loss than does olive oil. *The American journal of clinical nutrition*. 2008;87(3):621-6.
33. St-Onge MP, Jones PJ. Greater rise in fat oxidation with medium-chain triglyceride consumption relative to long-chain triglyceride is associated with lower initial body weight and greater loss of subcutaneous adipose tissue. *Int J Obes Relat Metab Disord*. 2003;27(12):1565-71.
34. Orsavova J, Misurcova L, Ambrozova JV, Vicha R, Mlcek J. Fatty Acids Composition of Vegetable Oils and Its Contribution to Dietary Energy Intake and Dependence of Cardiovascular Mortality on Dietary Intake of Fatty Acids. *Int J Mol Sci*. 2015;16(6):12871-90.
35. Denke MA, Grundy SM. Comparison of effects of lauric acid and palmitic acid on plasma lipids and lipoproteins. *The American journal of clinical nutrition*. 1992;56(5):895-8.
36. Clegg ME, Golsorkhi M, Henry CJ. Combined medium-chain triglyceride and chilli feeding increases diet-induced thermogenesis in normal-weight humans. *European journal of nutrition*. 2013;52(6):1579-85.
37. Public Health England. Government Dietary Recommendations London2016
[Available from:
https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/547050/government_dietary_recommendations.pdf.
38. Rizzo G, Masic U, Harrold JA, Norton JE, Halford JC. Coconut and sunflower oil ratios in ice cream influence subsequent food selection and intake. *Physiol Behav*. 2016;164(Pt A):40-6.
39. Poppitt SD, Strik CM, MacGibbon AK, McArdle BH, Budgett SC, McGill AT. Fatty acid chain length, postprandial satiety and food intake in lean men. *Physiol Behav*. 2010;101(1):161-7.